



# Complete Genome Sequences of 17 Clinical *Campylobacter jejuni* Strains from Chile

V. Bravo,<sup>a,b</sup> L. Porte,<sup>c</sup>  T. Weitzel,<sup>c</sup> C. Varela,<sup>c</sup>  C. J. Blondel,<sup>b,e</sup>  N. Gonzalez-Escalona<sup>d</sup>

<sup>a</sup>Unidad de Microbiología, Escuela de Medicina, Facultad de Ciencias Médicas, Universidad de Santiago de Chile, Santiago, Chile

<sup>b</sup>Instituto de Ciencias Biomédicas, Facultad de Medicina y Facultad de Ciencias de la Vida, Universidad Andrés Bello, Santiago, Chile

<sup>c</sup>Laboratorio Clínico, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana, Universidad del Desarrollo, Santiago, Chile

<sup>d</sup>Center for Food Safety and Applied Nutrition, Office of Regulatory Science, Division of Microbiology, U.S. Food and Drug Administration, College Park, Maryland, USA

<sup>e</sup>Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad Autónoma de Chile, Santiago, Chile

**ABSTRACT** *Campylobacter jejuni* is the leading cause of bacterial foodborne disease worldwide. Here, we report the complete annotated genomes and plasmid sequences of 17 *Campylobacter jejuni* strains isolated from patients with gastroenteritis in Santiago, Chile.

*Campylobacter* spp. are now considered the most common bacterial cause of human gastroenteritis worldwide. Of the 25 *Campylobacter* species, *Campylobacter jejuni* is the most common species associated with human disease (1–3). In Chile, *Campylobacter jejuni* infections are emerging as an important cause of foodborne illnesses (4–7). In this study, we obtained the complete genome sequences of 17 *C. jejuni* strains using a combination of long (Oxford Nanopore) and short (Illumina) reads.

These 17 strains, all belonging to different sequence types, were selected for genome closing out of 81 total strains. These strains were isolated in the laboratory from stool samples from patients with gastroenteritis as previously described (6, 8). These strains were exempt from institutional review board (IRB) review. Strains were grown overnight in Mueller-Hinton 5% sheep blood agar plates at 42°C under microaerobic conditions, and genomic DNA was extracted using the DNeasy blood and tissue kit (Qiagen). DNA quality and quantity were assessed using a NanoDrop spectrophotometer and a Qubit fluorometer (Thermo Scientific, Waltham, MA, USA), respectively, following the manufacturer's instructions. The long reads for each strain were generated in a MinION sequencer (Nanopore Technologies, Oxford, UK). The sequencing library was prepared using the rapid barcoding sequencing kit (SQK-RBK004). Every 10 libraries were pooled and run in a FLO-MIN106 (R9.4.1) flow cell, according to the manufacturer's instructions, for 48 h. The run was base called live using default settings (MinKNOW v19.06.8, Guppy v3.0.7). Default parameters were used for all software unless otherwise specified. In total, we conducted 2 Nanopore runs. The sequencing outputs for each of the 2 runs were 2.50 Gb (quality score, 12.41;  $N_{50}$ , 3,957 bp; total reads, 300,000) and 7.2 Gb (quality score, 11.46;  $N_{50}$ , 6,000 bp; total reads, 569,349), for an estimated average genome coverage of 60 to 90×. Reads of <5,000 bp and with a quality score of <7 were discarded for downstream analysis.

The short-read sequencing libraries for each strain were prepared using 100 ng DNA per strain according to the manufacturer's instructions using the Nextera DNA flex kit (Illumina, San Diego, CA, USA) for the MiSeq sequencing and 1 ng DNA for the Nextera XT kit for the NextSeq sequencing. Strains were sequenced using a MiSeq sequencer and a NextSeq sequencer (Illumina). For the MiSeq sequencing, we used a MiSeq v3 kit using 2 × 250-bp paired-end chemistry, according to the manufacturer's instructions, with >100× average coverage. For the NextSeq sequencing, we used a NextSeq

**Citation** Bravo V, Porte L, Weitzel T, Varela C, Blondel CJ, Gonzalez-Escalona N. 2020. Complete genome sequences of 17 clinical *Campylobacter jejuni* strains from Chile. *Microbiol Resour Announc* 9:e00535-20. <https://doi.org/10.1128/MRA.00535-20>.

**Editor** David Rasko, University of Maryland School of Medicine

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to N. Gonzalez-Escalona, [narjol.gonzalez-escalona@fda.hhs.gov](mailto:narjol.gonzalez-escalona@fda.hhs.gov).

**Received** 9 May 2020

**Accepted** 25 June 2020

**Published** 23 July 2020

**TABLE 1** Metadata for the 17 *C. jejuni* strains reported in this study<sup>a</sup>

CFSAN no. <sup>b</sup>	GenBank accession no. for:		GC content (%)	Hybrid assembly genome coverage (x)	Nanopore sequencing depth (x)	MiSeq sequencing depth (x)	Illumina SRA no. (no. of reads)	Nanopore SRA no. (no. of reads)	Yr of isolation	ST <sup>c</sup>
	Chromosome	Plasmid								
CFSAN093260	CP040618 (1,627,779)		30.56	70	72	145	SRR10860968 (1,022,308)	SRR11620707 (4,572)	2018	52
CFSAN093259	CP040617 (1,688,111)		30.51	64	31	135	SRR10860969 (981,408)	SRR11620706 (5,234)	2018	50
CFSAN093257	CP040616 (1,720,666)		30.47	76	20	168	SRR10860971 (1,250,338)	SRR11620698 (2,472)	2018	1359
CFSAN093256	CP040615 (1,653,603)		30.53	84	95	184	SRR10860973 (1,334,060)	SRR11620697 (9,032)	2018	10196
CFSAN093246	CP040613 (1,680,823)	CP040614 (71,914)	30.54	73	20	150	SRR10860981 (1,224,214)	SRR11620696 (2,637)	2018	8938
CFSAN093241	CP040612 (1,662,100)		30.51	66	90	140	SRR10860987 (995,550)	SRR11620695 (9,485)	2018	475
CFSAN093238	CP040611 (1,745,502)		30.35	76	54	160	SRR10860990 (1,187,310)	SRR11620694 (5,587)	2018	353
CFSAN093227	CP040610 (1,824,459)		30.16	66	46	133	SRR10860972 (1,072,404)	SRR11620693 (4,855)	2017	607
CFSAN093226	CP040608 (1,681,003)	CP040609 (39,913)	30.52	71	60	149	SRR10860983 (1,086,354)	SRR11620692 (7,883)	2018	21
CFSAN093224	CP040607 (1,625,593)		30.56	88	78	187	SRR10860995 (1,288,270)	SRR11620691 (7,887)	2018	6091
CFSAN096296 <sup>d</sup>	CP047484 (1,635,710)		30.54	296	20	675	SRR10859473 (7,707,678)	SRR11620705 (2,243)	2019	50
CFSAN096297 <sup>d</sup>	CP047482 (1,664,471)	CP047483 (48,209)	30.42	210	79	424	SRR10859482 (5,183,902)	SRR11620704 (7,953)	2018	257
CFSAN096301 <sup>d</sup>	CP047481 (1,661,307)		30.48	271	200	553	SRR10859574 (6,233,036)	SRR11620703 (48,095)	2019	52
CFSAN096302 <sup>d</sup>	CP047480 (1,721,182)		30.47	249	97	510	SRR10859489 (5,972,014)	SRR11620702 (11,058)	2018	1359
CFSAN096305 <sup>d</sup>	CP047479 (1,721,265)		30.44	260	58	520	SRR10859603 (6,157,576)	SRR11620701 (6,781)	2019	222
CFSAN096306 <sup>d</sup>	CP047478 (1,650,107)		30.51	139	267	285	SRR10859609 (3,157,756)	SRR11620700 (47,192)	2019	50
CFSAN096307 <sup>d</sup>	CP047477 (1,720,216)		30.42	81	30	165	SRR10859583 (1,902,526)	SRR11620699 (3,198)	2018	1359

<sup>a</sup>All strains were isolated in Santiago de Chile, Chile, from stools of patients with gastroenteritis.

<sup>b</sup>CFSAN, Center for Food Safety and Applied Nutrition.

<sup>c</sup>ST, sequence type (<https://pubmlst.org/campylobacter/>).

<sup>d</sup>Sequenced using NextSeq.

500/550 high-output kit v2.5 (300 cycles) using  $2 \times 150$ -bp paired-end chemistry, according to the manufacturer's instructions, with  $>300\times$  average coverage. The reads were trimmed with Trimmomatic v0.36 (9).

The final complete genome sequence (comprising the chromosome and plasmid, when present) for each strain was obtained using a previously described pipeline (10), except that Flye v2.6 (11) was used instead of Canu v1.7 (12) for long-read *de novo* assembling. The genomes were confirmed as circular closed by finding the contig end overlap, which was then manually trimmed. Each closed genome was rotated to start at the *dnaA* gene.

**Data availability.** The complete genome sequences reported here have been deposited in NCBI GenBank under the accession numbers listed in Table 1.

## ACKNOWLEDGMENTS

This study was supported by funding from the MCMi Challenge Grants Program proposal number 2018-646 (N.G.-E.), the FDA Foods Program Intramural Funds (N.G.-E.), and REDI170269 from CONICYT (C.J.B.). C.J.B. is a Howard Hughes Medical Institute (HHMI)-Gulbenkian international research scholar (grant number 55008749).

## REFERENCES

1. Kaakoush NO, Castano-Rodriguez N, Mitchell HM, Man SM. 2015. Global epidemiology of *Campylobacter* infection. Clin Microbiol Rev 28: 687–720. <https://doi.org/10.1128/CMR.00006-15>.
2. Igwaran A, Okoh AI. 2019. Human campylobacteriosis: a public health concern of global importance. Heliyon 5:e02814. <https://doi.org/10.1016/j.heliyon.2019.e02814>.
3. Burnham PM, Hendrixson DR. 2018. *Campylobacter jejuni*: collective components promoting a successful enteric lifestyle. Nat Rev Microbiol 16:551–565. <https://doi.org/10.1038/s41579-018-0037-9>.
4. Collado L, Gutierrez M, Gonzalez M, Fernandez H. 2013. Assessment of the prevalence and diversity of emergent campylobacteria in human stool samples using a combination of traditional and molecular methods. Diagn Microbiol Infect Dis 75:434–436. <https://doi.org/10.1016/j.diagmicrobio.2012.12.006>.
5. Collado L, Munoz N, Porte L, Ochoa S, Varela C, Munoz I. 2018. Genetic diversity and clonal characteristics of ciprofloxacin-resistant *Campylobacter jejuni* isolated from Chilean patients with gastroenteritis. Infect Genet Evol 58:290–293. <https://doi.org/10.1016/j.meegid.2017.12.026>.
6. Porte L, Varela C, Haecker T, Morales S, Weitzel T. 2016. Impact of changing from staining to culture techniques on detection rates of *Campylobacter* spp. in routine stool samples in Chile. BMC Infect Dis 16:196. <https://doi.org/10.1186/s12879-016-1546-7>.
7. Levican A, Ramos-Tapia I, Briceno I, Guerra F, Mena B, Varela C, Porte L. 2019. Genomic analysis of Chilean strains of *Campylobacter jejuni* from human faeces. Biomed Res Int 2019:1902732. <https://doi.org/10.1155/2019/1902732>.
8. Bravo V, Varela C, Porte L, Weitzel T, Kastanis GJ, Balkey M, Blondel CJ, Gonzalez-Escalona N. 2020. Draft whole-genome sequences of 51 *Campylobacter jejuni* and 12 *Campylobacter coli* clinical isolates from Chile. Microbiol Resour Announc 9:e00072-20. <https://doi.org/10.1128/MRA.00072-20>.
9. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
10. Gonzalez-Escalona N, Kastanis GJ, Timme R, Roberson D, Balkey M, Tallent SM. 2020. Closed genome sequences of 28 foodborne pathogens from the CFSAN verification set, determined by a combination of long and short reads. Microbiol Resour Announc 9:e00152-20. <https://doi.org/10.1128/MRA.00152-20>.
11. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol 37:540–546. <https://doi.org/10.1038/s41587-019-0072-8>.
12. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27:722–736. <https://doi.org/10.1101/gr.215087.116>.