

POSTER PRESENTATION

Open Access

Carbapenemase-producing *Acinetobacter baumannii* in a surgical intensive care unit – take time for tracking

T Götting^{1*}, W Ebner¹, D Jonas¹, A Serr¹, G Häcker¹, M Dettenkofer²

From 3rd International Conference on Prevention and Infection Control (ICPIC 2015)
Geneva, Switzerland. 16-19 June 2015

Introduction

Although the prevalence of carbapenemase-producing *Acinetobacter baumannii* (CP-AB) in Germany is still low an increasing number of outbreaks has been reported in recent years. In August 2014 we observed a cluster of 4 patients with CP-AB (OXA-23) on our surgical intensive care unit (SICU).

Objectives

After excluding an environmental source we performed epidemiological analyses to gain insight into the spread of CP-AB in our university hospital.

Methods

To characterize the CP-AB isolates molecular typing using amplified fragment length polymorphism was conducted. A genotypic comparison with strains collected in 2013 and the first half of 2014 was carried out. All new patient isolates of CP-AB were genotyped prospectively.

Results

Molecular typing revealed that 3 out of the 4 SICU-isolates were genotypically identical. Although temporally and locally coinciding, the 4th isolate of the patient cluster did not match. This 4th strain however was identical to a strain previously identified in a patient from the neurological department in July 2014. A physical link between these two patients could not be identified. Even more surprisingly, another isolate of CB-AB from a neurosurgical patient isolated in September 2014 was found to be identical with the three isolates of the SICU cluster, despite the absence of detectable temporal or local link to the SICU.

Conclusion

Isolates of CP-AB from six patients fell into two clusters of two and four strains, respectively.

In this case patient transfer within the hospital cannot explain these unexpected genotypical relationships between isolates from three departments. Transmission of CP-AB may have occurred in the context of consultants' visits or when patients were temporarily moved to a diagnostic unit such as endoscopy or radiology.

In the context of clusters of CP-AB isolated from several patients, genotyping provides the opportunity to follow the movement of these highly resistant bacteria within a hospital. Surprising genotyping results should engender efforts to analyse the pathways of transmission.

Disclosure of interest

None declared.

Authors' details

¹University Medical Center Freiburg, Freiburg, Germany.

²Gesundheitsverbund Landkreis Konstanz, Singen, Germany.

Published: 16 June 2015

doi:10.1186/2047-2994-4-S1-P137

Cite this article as: Götting et al.: Carbapenemase-producing *Acinetobacter baumannii* in a surgical intensive care unit – take time for tracking. *Antimicrobial Resistance and Infection Control* 2015 **4**(Suppl 1):P137.

¹University Medical Center Freiburg, Freiburg, Germany

Full list of author information is available at the end of the article