

Draft Genome Sequence of a Potential Nitrate-Dependent Fe(II)-Oxidizing Bacterium, *Aquabacterium parvum* B6

Xiaoxin Zhang,^{a,b} Fang Ma,^a Ulrich Szewzyk^b

State Key Laboratory of Urban Water Resource and Environment, School of Municipal and Environmental Engineering, Harbin Institute of Technology, Harbin, China^a; Department of Environmental Microbiology, Technical University of Berlin, Berlin, Germany^b

***Aquabacterium parvum* B6 is a potential nitrate-dependent Fe(II)-oxidizing bacterium. The genes related to its denitrifying mechanism and iron metabolisms were unknown. We present the draft genome of *Aquabacterium parvum* B6, which could provide further insight into the nitrate-dependent Fe(II)-oxidizing mechanism of strain B6.**

Received 2 December 2015 Accepted 7 December 2015 Published 28 January 2016

Citation Zhang X, Ma F, Szewzyk U. 2016. Draft genome sequence of a potential nitrate-dependent Fe(II)-oxidizing bacterium, *Aquabacterium parvum* B6. *Genome Announc* 4(1):e01651-15. doi:10.1128/genomeA.01651-15.

Copyright © 2016 Zhang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Fang Ma, mafang@hit.edu.cn, or Ulrich Szewzyk, ulrich.szewzyk@tu-berlin.de.

Nitrate-dependent Fe(II)-oxidizing (NDFO) bacteria, conducting nitrate-dependent Fe(II)-oxidation, include both autotrophic and heterotrophic organisms (1, 2). In previous studies, several *Aquabacterium* spp. have been mentioned as capable of NDFO. A ferrous iron-oxidizing, nitrate-reducing bacterium, *Aquabacterium* strain BrG2, was obtained from freshwater sediments (3). The strain is facultative anaerobic, and the Fe(II) oxidation could be accelerated significantly with the presence of an organic co-substrate (4). A short rod-shaped *Aquabacterium fontiphilum* sp. nov., isolated from springwater in Taiwan, was also reported to accomplish nitrate reduction via utilizing nitrate serves as an alternative electron acceptor (5). Moreover, isolates from the sediments in the littoral of Lake Constance closely related to the *Aquabacterium* cluster were also able to conduct NDFO (6). To provide clues for future investigations of the mechanism and metabolism of *Aquabacteria*, as well as to facilitate further studies with the genus and species, we report here a draft genome assembly of *Aquabacterium parvum*.

Aquabacterium parvum strain B6, characterized approximately two decades ago, is a Gram-negative, rod-shaped bacterium, isolated from a Berlin drinking water distribution system (7). The strain is available from the German Collection of Microorganisms and Cell Cultures (Deutsche Sammlung von Mikroorganismen und Zellkulturen, DSMZ 11968). Draft genome data for strain B6 were determined using the Illumina HiSeq2000 system. The reads were assembled into 619 large contigs using Velvet version 1/2/10 (8). The contig N_{50} is approximately 10.7 kb, and the largest contig assembled is approximately 50.0 kb. The draft sequence consists of 4,592,999 bases with a mean G+C percentage of 65.3%. Annotation of open reading frames was performed by using the RAST annotation server (9) and the NCBI Prokaryotic Genome Annotation Pipeline (10). A total of 4,019 coding sequences and 63 structural RNAs were predicted. There are 429 subsystems in the genome, as determined by the RAST server. Comparison with the genome sequences available at the RAST server suggested that the closest neighbor of strain B6 is *Leptothrix cholodnii* SP6 (score 533), followed by *Methylibium petroleiphilum* PM1 (score 528),

Alicyclophilus denitrificans BC (score 389), and *Acidovorax* sp. JS42 (score 383).

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [LFRI00000000](https://www.ncbi.nlm.nih.gov/nuclink/LFRI00000000). The version described in this paper is the first version, [LFRI01000000](https://www.ncbi.nlm.nih.gov/nuclink/LFRI01000000).

ACKNOWLEDGMENTS

We acknowledge the contributions of the Department of Environmental Microbiology, TU Berlin, for supplying the strain used in this study.

This work was supported by the National Natural Science Foundation of China (no. 51178139) and a doctoral scholarship from the Chinese Scholarship Council (CSC).

REFERENCES

- Chakraborty A, Roden EE, Schieber J, Picardal F. 2011. Enhanced growth of *Acidovorax* sp. strain 2an during nitrate-dependent Fe(II) oxidation in batch and continuous-flow systems. *Appl Environ Microbiol* 77:8548–8556. <http://dx.doi.org/10.1128/AEM.06214-11>.
- Weber KA, Hedrick DB, Peacock AD, Thrash JC, White DC, Achenbach LA, Coates JD. 2009. Physiological and taxonomic description of the novel autotrophic, metal-oxidizing bacterium, *Pseudogulbenkiania* sp. strain 2002. *Appl Microbiol Biotechnol* 83:555–565. <http://dx.doi.org/10.1007/s00253-009-1934-7>.
- Buchholz-Cleven BEE, Rattunde B, Straub KL. 1997. Screening for genetic diversity of isolates of anaerobic Fe(II)-oxidizing bacteria using DGGE and whole-cell hybridization. *Syst Appl Microbiol* 20:301–309. [http://dx.doi.org/10.1016/S0723-2020\(97\)80077-X](http://dx.doi.org/10.1016/S0723-2020(97)80077-X).
- Straub KL, Schönhuber WA, Buchholz-Cleven BEE, Schink B. 2004. Diversity of ferrous iron-oxidizing, nitrate-reducing bacteria and their involvement in oxygen-independent iron cycling. *Geomicrobiol J* 21: 371–378. <http://dx.doi.org/10.1080/01490450490485854>.
- Lin MC, Jiang SR, Chou JHA, Arun AB, Young C-C, Chen W-M. 2009. *Aquabacterium fontiphilum* sp. nov., isolated from spring water. *Int J Syst Evol Microbiol* 59:681–685. <http://dx.doi.org/10.1099/ijs.0.000745-0>.
- Benz M, Brune A, Schink B. 1998. Anaerobic and aerobic oxidation of ferrous iron at neutral pH by chemoheterotrophic nitrate-reducing bacteria. *Arch Microbiol* 169:159–165. <http://dx.doi.org/10.1007/s002030050555>.
- Kalmbach S, Manz W, Wecke J, Szewzyk U. 1999. *Aquabacterium* gen. nov., with description of *Aquabacterium citratiphilum* sp. nov., *Aquabacterium parvum* sp. nov. and *Aquabacterium commune* sp. nov., three *in situ* dominant

- bacterial species from the Berlin drinking water system. *Int J Syst Bacteriol* 49:769–777. <http://dx.doi.org/10.1099/00207713-49-2-769>.
8. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 9. Aziz RK, Bartels D, Best AA, Dejongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 10. Pruitt KD, Tatusova T, Klimke W, Maglott DR. 2009. NCBI reference sequences: current status, policy and new initiatives. *Nucleic Acids Res* 37(suppl 1):D32–D36. <http://dx.doi.org/10.1093/nar/gkn721>.