

POSTER PRESENTATION

Open Access

Workload associated with mrsa control in surgery: a prospective study alongside a controlled clinical trial

S Bahrami^{1*}, AS Lee², S Harbarth³, S Malhotra-Kumar⁴, C Brun-Buisson⁵, I Durand-Zaleski¹, MOSAR WP9 and WP4 working groups

From 3rd International Conference on Prevention and Infection Control (ICPIC 2015) Geneva, Switzerland. 16-19 June 2015

Introduction

Controversies regarding control of endemic methicillin-resistant *Staphylococcus aureus* (MRSA) stem in part from the paucity of data available on the actual costs of implementing MRSA control strategies.

Objectives

To estimate the workloads associated with screening and hand hygiene (HH) promotion interventions for MRSA control.

Methods

Prospective evaluation of costs associated with MRSA control interventions, alongside the MOSAR–Surgery multicenter intervention trial [1], emphasizing the workloads associated with interventions. Interventions under study were (1) universal MRSA screening with contact precautions and decolonization (4 hospitals) and (2) enhanced HH promotion (4 hospitals); 2 hospitals implemented a combined strategy using targeted MRSA screening.

Workloads were estimated from a hospital perspective, using a top-down approach, distinguishing infection control (IC) and ward staff duties, set-up and routine activities, and excluding research-driven tasks.

Results

In the screening arm (13 wards; mean, 27.8 ± 10.4 beds), set-up required 2.6 ± 0.58 weeks of work from the IC team, and the mean annual workload was 20.5 weeks or 8.62 ± 4.39 weeks for a 10-bed ward. In the HH

promotion arm (13 wards, 87.8 ± 79.5 beds), set-up required 2.77 ± 1.13 weeks, and mean annual workload was 12.1 weeks (3.29 ± 3.72 per 10 beds). In the combined arm (7 wards, 44.9 ± 24.6 beds), set-up required 7.0 ± 1.93 weeks, and mean annual workload was 23.7 weeks (5.94 ± 1.84 weeks per 10 beds). The burden on ward staff was relatively limited in most wards.

Conclusion

Workload associated with the MRSA screening strategy is relatively homogenous and predictable. Investment in the HH promotion strategy showed large variations between centers and was maximal for the successful combined intervention. Further research is needed on the optimal implementation of HH promotion interventions.

Disclosure of interest

S. Bahrami: None declared, A. Lee: None declared, S. Harbarth Grant/Research support from: Geneva University Hospitals, B. Braun, Pfizer, Speaker's bureau of: BioMerieux, Pfizer, Conflict with: Scientific Advisory Board, Destiny Pharma, DaVolterra, BioMérieux, S. Malhotra-Kumar: None declared, C. Brun-Buisson: None declared, I. Durand-Zaleski: None declared.

Authors' details

¹AP-HP, Paris, France. ²Royal Prince Alfred Hospital, Sydney, Australia. ³University of Geneva Hospitals, Switzerland. ⁴University of Antwerp, Belgium. ⁵Inserm U657, Paris, France.

Published: 16 June 2015

¹AP-HP, Paris, France

Full list of author information is available at the end of the article

Reference

1. Lee AS, Cooper BS, Malhotra-Kumar S, *et al*: *BMJ Open* 2013, **3**:e003126.

doi:10.1186/2047-2994-4-S1-P188

Cite this article as: Bahrami *et al.*: Workload associated with mrsa control in surgery: a prospective study alongside a controlled clinical trial. *Antimicrobial Resistance and Infection Control* 2015 **4**(Suppl 1):P188.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

