



Evolution of metazoan morphological disparity

Bradley Deline^{a,1}, Jennifer M. Greenwood^b, James W. Clark^b, Mark N. Puttick^{b,c}, Kevin J. Peterson^d, and Philip C. J. Donoghue^{b,1}

^aDepartment of Geoscience, University of West Georgia, Carrollton, GA 30118; ^bSchool of Earth Sciences, University of Bristol, BS8 1TQ Bristol, United Kingdom; ^cDepartment of Biology and Biochemistry, University of Bath, BA2 7AY Bath, United Kingdom; and ^dDepartment of Biological Science, Dartmouth College, Hanover, NH 03755

Edited by Neil H. Shubin, The University of Chicago, Chicago, IL, and approved August 7, 2018 (received for review June 26, 2018)

The animal kingdom exhibits a great diversity of organismal form (i.e., disparity). Whether the extremes of disparity were achieved early in animal evolutionary history or clades continually explore the limits of possible morphospace is subject to continuing debate. Here we show, through analysis of the disparity of the animal kingdom, that, even though many clades exhibit maximal initial disparity, arthropods, chordates, annelids, echinoderms, and mollusks have continued to explore and expand the limits of morphospace throughout the Phanerozoic, expanding dramatically the envelope of disparity occupied in the Cambrian. The “clumpiness” of morphospace occupation by living clades is a consequence of the extinction of phylogenetic intermediates, indicating that the original distribution of morphologies was more homogeneous. The morphological distances between phyla mirror differences in complexity, body size, and species-level diversity across the animal kingdom. Causal hypotheses of morphologic expansion include time since origination, increases in genome size, protein repertoire, gene family expansion, and gene regulation. We find a strong correlation between increasing morphological disparity, genome size, and microRNA repertoire, but no correlation to protein domain diversity. Our results are compatible with the view that the evolution of gene regulation has been influential in shaping metazoan disparity whereas the invasion of terrestrial ecospace appears to represent an additional gestalt, underpinning the post-Cambrian expansion of metazoan disparity.

Metazoa | disparity | evolution | morphology | Cambrian explosion

The diversity of animal organismal form (i.e., disparity) is decidedly nonrandom; members of one phylum share “bodyplan” characteristics distinct from those of other phyla, suggesting that only a very small subset of the universe of possible bodyplans has been realized. Paleontological analyses have suggested that the limits on organismal disparity were realized early in animal evolutionary history (1–3), inspiring the view that fundamental innovation has been precluded subsequently by gene regulatory developmental constraints (4–6), and that the evolutionary processes underlying the emergence of animals are nonuniformitarian (7, 8). However, this perspective is based largely on the timing of appearance of Linnean rank taxa in the fossil record (8, 9), assuming they provide an effective proxy for organismal disparity. Attempts to capture disparity by using morphometry have borne out the hypothesis of maximal initial disparity (3, 10), but this approach is limited practically to analysis at low taxonomic levels and it is not clear that the results can be generalized to higher taxa, including the phylum and kingdom levels. Here we attempt to map metazoan disparity within an empirical morphospace based on a large sampling of discrete characters from across the breadth of extant metazoan diversity. We use this to explore the impact of extinction on morphospace occupation and the relationship between organismal disparity and other phenomena such as complexity, body size, diversity, and Linnean rank. We then undertake quantitative tests of hypotheses of causality, including random variation, genome size, protein diversity, and gene regulatory complexity.

Mapping Metazoan Morphospace

The construction of a morphospace is dependent on methodology and the selection of relevant features. However, with a significantly large data source, the distances among taxa within morphospace will begin to approximate the evolutionary scale of the differences. Spatial landmark analysis is precluded at high taxonomic rank such as phylum because, by definition, phyla share few morphological homologies. Discrete characters provide a suitable alternative given that there are no practical limits to their scalability (2, 3), and comparative analyses have shown that continuous and discrete character datasets can capture the same phenomenon (11–14). The use of discrete characters produces results that have nonmetric properties (15–17), but this approach can and has been used to elucidate broad patterns of similarities and clustering within multidimensional space (18), particularly in formulating the hypotheses we seek to test. To test between competing hypotheses for the evolution of disparity—whether the limits of disparity were established early or have continued to expand throughout evolutionary history—we compiled a cladistic character matrix derived from Peter Ax’s *Phylogenetic System* (19–21). This constitutes a single, densely sampled synthetic overview of character distribution among metazoans, including all phyla, by an individual who was not a taxonomic specialist in any of the groups that could perhaps be considered overrepresented in the dataset. Ax’s taxonomic sampling is not uniform across metazoans, but the number of characters and taxa within a phylum is representative of its in-traphylum diversity (ref. 22; Spearman’s correlation, $\rho = 0.821$, $P < 0.001$). Therefore, we do not consider that any clades are

Significance

We attempt to quantify animal “bodyplans” and their variation within Metazoa. Our results challenge the view that maximum variation was achieved early in animal evolutionary history by nonuniformitarian mechanisms. Rather, they are compatible with the view that the capacity for fundamental innovation is not limited to the early evolutionary history of clades. We perform quantitative tests of the principal hypotheses of the molecular mechanisms underpinning the establishment of animal bodyplans and corroborate the hypothesis that animal evolution has been permitted or driven by gene regulatory evolution.

Author contributions: B.D., K.J.P., and P.C.J.D. designed research; B.D., J.M.G., J.W.C., M.N.P., K.J.P., and P.C.J.D. performed research; M.N.P. contributed new reagents/analytic tools; B.D., J.M.G., and J.W.C. analyzed data; and B.D., J.M.G., J.W.C., M.N.P., K.J.P., and P.C.J.D. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence may be addressed. Email: bdeline@westga.edu or phil.donoghue@bristol.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1810575115/-DCSupplemental.

Published online September 4, 2018.

significantly underrepresented or overrepresented in the dataset. We coded 1,767 characters for 212 extant, terminal taxa, including 34 animal phyla, most commonly down to the Linnean rank of order (*SI Appendix, Table S1*). The characters encompass all aspects of morphology (cellular, developmental, sexual, and skeletal and soft-tissue anatomy), including those minimally defining each clade, comprising 915 characters that are shared among the operational taxa (homologies and homoplasies) and 852 unique (i.e., autapomorphic) characters (*SI Appendix, Table S2* and *Dataset S1*).

We mapped the relationships between features to identify characters that are nonapplicable rather than absent, which can be differentiated analytically by using Gower's similarity metric (23–25). We subjected these data to a nonmetric multidimensional scaling (NMDS) analysis, a noneigenvector-based multivariate method that attempts to optimize the fit between the data and a preselected number of axes (Fig. 1). The use of a nonmetric ordination technique has several advantages (e.g., the ability to handle large amounts of absent data), but the distances between organisms may not be directly Euclidean, which may alter measurements of disparity. To control for the choice of ordination technique, we repeated all analyses by using principal coordinate analysis. The choice of ordination had no impact on the morphospace or any of the presented results (*SI Appendix, Fig. S1*). The distances between taxa using linear (i.e., principal coordinate analysis) and nonlinear (i.e., NMDS) methods are strongly and linearly correlated (Mantel test, $R^2 = 0.9764$, $P = 0.001$), indicating that, even though the NMDS ordination is built by using a nonlinear method, it has linear properties. Absolute distances within the space can still be subject to nonmetric artifacts such that the distances between taxa should be taken as a qualitative metric of the overall distribution of metazoan morphologic diversity. An analysis of the stress (representing the goodness of fit) indicates that the majority of variance in the data is captured by the first two axes (*SI Appendix, Fig. S2*).

Concerned that different treatments of nonapplicable data (25) could significantly alter the structure of the ordination, we structured our data to reflect different coding strategies (Fig. 2). Treating nonapplicable characters as absent (Fig. 2*A*) or missing (Fig. 2*B*) produced statistically similar ordinations to the main analysis (Fig. 1) in which inapplicable characters are assigned a distinct state conferring distance (Mantel test, $R = 0.907$, $P = 0.001$). However, these differ in the (relative) displacement of the nonmetazoan eukaryotic outgroup, Porifera and Placozoa, into the central area of morphospace (Fig. 2*A*). Treating nonapplicable characters as absent or missing also increased intraphylum disparity at the cost of interphylum disparity (Fig. 2*A* and *B*). However, the similarity in results from the different treatments of the data indicate the strength of the underlying structure of the data. These results indicate that the structure of disparity is robust to ordination and coding strategies.

The position of taxa based on the first two axes is presented in Fig. 1*A*. Most of this variation is based on shared characters, as analysis of a dataset excluding autapomorphies has no significant impact on the structure of the morphospace ($R^2 > 0.92$, $P = 0.001$; *SI Appendix, Fig. S3*). Phyla differ dramatically in the position and areal extent of their envelope of disparity. Although most taxa are clustered along both axes, the nonmetazoan eukaryote outgroup and Porifera plot separately from eumetazoans, principally for their lack of shared eumetazoan characters, even though neither of these groups occupy a large area of morphospace. Chordata (Fig. 1*B*), Arthropoda (Fig. 1*C*), and, to a lesser extent, Annelida, Echinodermata, and Mollusca, are much more disparate, each occupying larger ranges of morphospace than all other phyla combined, and defining the extremities of morphospace on the two principal axes. A Q-mode analysis of the distribution of characters (Fig. 2*C*) shows that the characters


that describe intraphylum features load at the extremities of both axes. Superphylum- and phylum-level characters occupy approximately one fourth of the area of the lower-level characters and vary primarily along the first axis.

From Modern to Historical Disparity

Central to the thesis of maximal initial disparity in animal evolution was the discovery of distinct bodyplans among Cambrian Konservat-Lagerstätten (1), which were assigned historically to extinct phyla, classes, or orders. Thus, by comparing only living taxa, it could be argued that we have captured only net historical disparity. Therefore, we coded a phylogenetically diverse and representative sample of Cambrian taxa, principally the earliest representatives of ordinal level clades (26). This entailed coding 70 fossil taxa for the existing character set and adding 111 mostly autapomorphic characters. Coding these fossil taxa was potentially problematic in that most of the characters (54.1%) are not preserved, and therefore unknown. On average, only 8.6% of the characters were coded as applicable, resulting in the fossil taxa appearing more constrained and skewed toward lower values on the second axis, making the Cambrian taxa appear less complex (Fig. 2*D*); we interpret this result as an artifact of the great volume of data missing for the fossils. There are two possible solutions to accommodating fossil species. One approach is to subsample our dataset for fossilizable characters based on known examples of fossilized features or the anatomical nature of the character (1,000 characters). NMDS analysis of this subsampled dataset results in a plot of morphospace occupation with the same broad structure (Mantel test, $R = 0.974$, $P = 0.001$; Fig. 2*E*) as that recovered from analysis of the entire dataset (Fig. 1*A*). However, the resulting morphospace accentuates the relative disparity of vertebrates and arthropods while diminishing the relative disparity of all other phyla (nonbilaterians especially), individually and in combination, exaggerating the significance of skeletal and gross anatomical characters that are fossilized in instances of routine and exceptional preservation. Few of these characters are representative of bodyplans more generally, which are defined on the basis of soft-tissue, cellular, tissue, organ, and developmental characters that are not usually fossilized. Thus, restricting the analysis to only fossilizable characters cannot be considered to capture organismal disparity within metazoans in any meaningful way. These results are of concern because they suggest that the majority of disparity analyses, which have been applied principally to fossil groups, may have limited inferential power. This is because they are constrained to characterization of fossilizable characters that may be otherwise unrepresentative of phenotypic evolution.

An alternative approach to including fossil species exploits their known phylogenetic position among living and fossil relatives to infer character states that are lost during fossilization. There are obviously assumptions inherent in inferring missing data, including missing secondary reversals in soft tissues, the potential of differential evolutionary rates between preservable and nonpreservable characters, or limiting the coded fossil autapomorphies to preservable characteristics. However, given the rarity of reversals of superphylum-level nonfossilizable characters in extant taxa and the observation that autapomorphies contribute little to the construction of the morphospace (*SI Appendix, Fig. S3*), these assumptions are likely to have a minor impact on the projection of fossil taxa into the morphospace defined by the living species. The approach of inferring missing data likely strengthens the phylogenetic signal in the morphospace. However, a comparison with the taphonomically culled dataset (Fig. 2*E*) indicates a similar and robust placement of the fossil taxa within morphospace.

To implement this approach, we derived a consensus, time-scaled phylogenetic tree for the operational taxa, which differed from Ax's original phylogenetic hypothesis (*SI Appendix, Figs.*



EVOLUTION

EARTH, ATMOSPHERIC
AND PLANETARY SCIENCES

Fig. 1. (A) Morphospace encompassing Ax's 212 operational taxa representing 34 phyla. The character matrix was analyzed by using NMDS. (B) Morphospace of the Chordata, which includes 26 vertebrata taxa, 2 urochordate taxa, and a single cephalochordate taxon grouped by class. (C) Morphospace of the 94 arthropod taxa included in the study grouped by subphylum.

tempo of clade disparity: the majority of phylum-level clades exhibit maximal disparity achieved by the Cambrian (Fig. 3, gray), compatible with previous studies at low taxonomic rank (18, 31, 32), whereas others exhibit a progressive exploration of morphospace: principally arthropods and chordates (corroborating refs. 33–35), but also annelids, echinoderms, and mollusks. Thus, the envelope of disparity explored by Kingdom Metazoa has increased through geological time.

Relationship Between Disparity and Complexity, Body Size, and Diversity. Having codified metazoan disparity, we next attempted to understand the relationship between morphology and other primary biologic metrics. To achieve this, taxonomic rank had to be normalized to the phylum level. The morphologic position of phyla was determined by using two methods: (i) including the crown ancestor of each phylum in the preexisting morphospace and (ii) independently analyzing the modeled characters for the crown ancestor of each phylum (*SI Appendix, Fig. S6*). The first method is influenced by differences in diversity or disparity between phyla, whereas the second disregards those differences between phyla in an effort to control for potential sampling biases. As expected, the two morphospaces differ in the uniqueness of arthropods and chordates, which alter the strength of correlation between morphology and some other datasets. However, these two methods maintain the structure of the morphospace (Mantel test, $R = 0.946$ $P = 0.001$; *SI Appendix, Fig. S6*). Thus, differences in sampling are not the controlling factor in correlations between morphology and other datasets. In all of the following tests of correlation, we undertook parallel analyses by using both methods, the results of which were similar; we present only those results based on the independent ordination of the character sets inferred for the phylum crown-ancestors. The distances between the morphological position of the phyla are then considered as a qualitative measure of the overall similarity of the phyla and can be used as a guide to compare morphology to other primary biological metrics.

The concept of disparity has been linked to, and sometimes even conflated with, the concept of organismal complexity (36). To explore their relationship, we compiled a new dataset of metazoan cell type diversity, the only widely accepted proxy for morphological complexity (e.g., ref. 11). We questioned whether the two phenomena were correlated by using a nonlinear (Spearman) Mantel test to compare pairwise distances derived from the morphology and complexity datasets (Table 1). The possibility that the two datasets are uncorrelated can be rejected at a high level of significance (Table 1). This relationship can be rationalized because only simple body plans are possible with few cell types, compatible with the view that expansion in cell type diversity has underpinned the expansion of metazoan disparity (37).

Morphologic distances between phyla were also compared with compilations of minimum, maximum, and range in body size within each phylum (38). Body size is correlated with many ecological and evolutionary traits and has increased by 16 orders of magnitude during the history of life (39). Morphologic distance correlates significantly with maximum body size and range of body size, but not with minimum body size (Table 1). This correlation reflects the greater physical demands and adaptive solutions to body form required by larger body size. It also suggests that there is a threshold in body size below which broad phenotypic disparity may not be possible, perhaps linked to the greater diversity of cell types that characterize organisms that achieve large body size (40, 41).

The relationship between diversity and disparity has been an area of intense study in deciphering the meaning of the two metrics (42) as well as the use of higher-level diversity as a proxy for disparity (9). Many metrics have been used to calculate disparity from constructed morphospaces, but the average squared distance between taxa within morphospace shows the greatest

Table 1. Statistical comparison of morphology against other biologic features

Feature	Mantel r	P value	No. of phyla
No. of cell types	0.3387	0.01	29
Minimum body size	0.09537	0.248	28
Maximum body size	0.4875	0.004	28
Range in body size	0.4862	0.007	28
Species level diversity	0.3678	0.007	32
Genome length	0.278	0.028	22
Protein (superfamily)	0.2512	0.141	12
Protein (family)	0.1701	0.218	12
Protein (architecture)	−0.288	0.886	12
microRNA	0.39	0.005	24

The positions of phyla were calculated in two manners. First, the modeled character suites for the ancestral nodes of the individual phyla were projected onto the morphospace constructed with Ax's 212 operational taxa. This method includes the disparity contained within phyla to some degree and therefore could be potentially biased by differential sampling. Second, the modeled character suites for the ancestral nodes were independently ordinated by using NMDS. This method treats each phylum equally and disregards the synapomorphies contained within each phylum such that the structure of the data are not controlled by differential sampling. The results were the same, so only the later is presented. Correlations between matrices were analyzed by using Mantel tests (Spearman), which compare the rank-order distance between phyla within the two matrices. The number of phyla included in the test varies with the availability of data (lists of phyla included in the different comparisons are presented in *SI Appendix, Table S3*). Multivariate datasets (miRNA and proteins) were analyzed in a similar manner to the morphological dataset (NMDS). Body size data are from McClain and Boyer (38). Genome length data were accessed from the animal genome size database

stability with smaller sample sizes (43). Differences in modern species-level diversity within phyla (22) correlate to morphologic distances between phyla (Table 2), and the number of species within a phylum correlates strongly to the disparity contained within (Fig. 4A and Table 2). This indicates that, at a higher taxonomic level, these metrics are closely related. However, a comparison of phylum-level diversity and disparity through time (based on origination data from ref. 29) indicates that there is no correlation between these two aspects of variance. Indeed, the relationship is static through time (Fig. 4B), indicating that the number of phyla provides a poor measure of metazoan disparity. However, our results cannot reject equivalence between our measure of disparity and diversity measured by counts of Linnean ranks below the phylum level. In sum, Linnean rank taxonomic measures of disparity have overestimated the scale of early metazoan disparity and therefore the phenomenon to be explained by intrinsic and extrinsic causal factors.

Testing Hypotheses of Causality. Distilling the phenomenon of animal disparity is one thing; establishing its causality is another. Explanations encompass intrinsic causes, including expansions in genome size (44), the diversification of protein domains and domain architectures (45), the origin of a “developmental toolkit” of transcription factors and cell signaling molecules (46), and the evolutionary assembly of gene regulatory networks (GRNs) (5), although it has also been argued that the exploration of metazoan morphospace is largely a time-dependent random walk (36). To test among these hypotheses, we compiled datasets of phylum origination dates (29), protein domains and their architectures (47), average genome sizes (48), and microRNAs (49) to serve in proxy for the diversity of GRNs. The multivariate protein and microRNA data were also analyzed by using NMDS. Our tests are limited to correlation with the use of a Mantel test. The position of a phylum within morphospace was taken as the position of its crown ancestor (*SI Appendix, Fig. S6*).

Table 2. Statistical comparison of morphology against diversity and origination

Diversity/origination	Spearman's ρ	P value	No. of phyla
Species level diversity	0.698	<0.001	32
Species level diversity (excl. 0s)	0.865	<0.001	16
Origination	0.041	0.905	11
Origination (excl. 0s)	0.047	0.903	9

The positions of phyla were calculated as described in Table 1. The number of phyla included in the test varies with the availability of data (lists of phyla included in the different comparisons are presented in [SI Appendix, Table S3](#)). Morphological disparity was calculated as the average squared distance between taxon within a phylum. Diversity was compiled by Chapman (22), and origination data are from Erwin et al. (29)

To explore whether the observed patterns in phylum-level morphologic differences could be the result of accumulated random evolutionary processes over time (36), we compared origination time to within-phylum disparity (Fig. 4C). In the absence of morphological constraint or stabilizing selection, clades that originated earlier would have more time to accumulate random changes and therefore to accumulate a higher level of disparity. No significant trend is seen between disparity and origination time (Table 2), indicating that, even though random processes may be contributing to disparity, there must be other, dominant factors that cause the vast difference in morphologic expansion between phyla regardless of origination time.

To test among the intrinsic causes of disparity ([SI Appendix, Tables S3 and S4](#)), we first compared differences in genome size (c-value) to the distances between phyla within morphospace. Genome size has been linked to several biological features such as organismal complexity, metabolic rates, and developmental rate (50), and we can reject the hypothesis that genome size and morphological complexity are uncorrelated at a high level of significance (Table 1). Genome size is not correlated to the number of genes (50), suggesting that the link between genome size and morphology might instead be effected largely through gene regulation, rather than the number of genes per se. Although the expansions of some protein-domain superfamilies have been correlated with the evolution of complexity (51), our data show no overall correlation between disparity and the repertoire of protein domains or architectures (Table 1 and [SI Appendix, Fig. S7](#)). In contrast, the number of miRNA families correlates significantly with morphological disparity (Table 1 and [SI Appendix, Fig. S8](#)), lending correlative support to the hypothesis that expansions in gene regulatory complexity underlie the evolution of metazoan morphological complexity (5, 7, 52).

Discussion

Our mapping of metazoan morphospace has shown that, even though some animal phyla demonstrate maximal initial disparity, others, notably chordates, arthropods, annelids, and mollusks, have progressively expanded on the limits of phylum and kingdom level morphospace post-Cambrian. This contradicts the generally held view of maximal initial disparity for the animal kingdom, which has been based largely on two assumptions: (i) that Linnean ranks serve as an effective proxy for organismal disparity and (ii) that patterns of morphospace occupation captured by continuous or discrete character observations at low taxonomic rank reflect the same phenomenon at phylum and kingdom levels. These are assumptions we refute. Not all studies have returned a pattern of maximal initial disparity. Indeed, disparity analyses based on continuous (53) and discrete characters (18) have identified instances in which maximal disparity is achieved in the middle period or late in the evolutionary history of lineages. Nevertheless, recent compilations have shown that, more often than not, maximal disparity is achieved early in the

evolutionary history clades (10, 18). However, it does not follow logically that, at the comparatively low taxonomic level at which these studies have been conducted, a pattern of maximal initial disparity will scale to a self-similar pattern at the highest taxonomic levels. Nevertheless, disparity at low and high taxonomic levels is linked hierarchically such that, regardless of whether its zenith is achieved early or late, increasing disparity at low taxonomic




Fig. 4. Comparisons between taxonomic diversity and morphologic disparity. Disparity is calculated as the average squared distance between taxa within the morphospace. Intra-phylum disparity is calculated by using the distance between Ax 's operational taxa within the morphospace for each phylum, and inter-phylum disparity is calculated by using the distance between phyla ([SI Appendix, Fig. S5](#)). (A) Intra-phyla disparity compared with estimates of modern diversity. If there was a single operational taxon within a phylum, the disparity is given a value of 0. (B) Comparison between metazoan diversity and inter-phylum disparity for 4 time intervals for the 11 phyla with well-resolved origination dates ([SI Appendix, Table S5](#)). For each time bin, the number of phyla that were present during the beginning of the interval were tallied (i.e., metazoan diversity) so, for example, 3 of the 11 phyla originated before the beginning of the Ediacaran, whereas an additional 3 phylum-level crown groups had appeared by the beginning of the Cambrian. Inter-phylum disparity was then calculated for the phyla present during that interval. (C) Intra-phyla disparity compared with the age of the phylum. Origination data are from Erwin et al. (29) and diversity data are from Chapman (22). Error bars are calculated as the SE of 1,000 bootstrap replicates.




Fig. 5. Distribution of taxa within morphospace based on fossilizable and nonfossilizable characters. (A) Morphospace based on 912 fossilizable characters. (B) Morphospace based on 878 characters unlikely to be observable in fossil organisms. (C) Intrametazoan and intraphylum disparity based on ordinations of only fossilizable characters and only nonfossilizable characters. Phyla above red line have higher disparity within nonpreserved soft-tissue characters, whereas those below have higher disparity within preserved gross anatomical or skeletal characters.

levels must scale to gross disparity increasing progressively at the highest taxonomic levels.

The clumpiness of metazoan morphospace occupation has been alternately argued to be a consequence of (i) gene regulatory and/or other constraints such as integration or modularity, (ii) extinction, and (iii) incomplete exploration of morphospace. Our analyses demonstrate that, even though many (but not all) modern clades occupy discrete regions of morphospace, the inclusion of Cambrian taxa indicates that phylogenetic intermediates of living clades occupy concomitantly intermediate regions of morphospace. Combined with our inferences of the course of metazoan phylogeny through morphospace, these results indicate that the morphological discreteness of modern clades is largely a consequence of the extinction of phylogenetic intermediates. Nevertheless, the results of our analyses including

fossils and phylogenetic ancestors indicate that the majority of the empirical morphospace circumscribed by our dataset has not been explored in metazoan evolutionary history. This may be because insufficient time has elapsed for all of these theoretical phenotypes to have been realized during metazoan phylogeny. However, there is a high level of phylogenetic linkage among many of the phenotypic characters in the dataset (Dataset S1): approximately half of the characters in the dataset are contingent directly on just 20 characters, and almost all of the characters are contingent ultimately on just a few characters (epithelia; ontogenesis; somatic differentiation). This reflects a high level of phylogenetically rooted developmental constraint underpinning the distribution of characters and their possible combinations. Thus, it is likely that the majority of the theoretical morphologies represented by unoccupied volumes of morphospace are unrealizable because the implied character combinations are not possible (e.g., ref. 54).

Evidently, the heterogeneity in the phylogenetic linkage of characters reflects the fact that some have contributed more than others to the realization of metazoan bodyplans. These characters can be identified based simply on their contingent linkage, and also through ordination of the phenotypic characters based on their distribution among species rather than through ordination of the species based on their distribution among characters (Fig. 2C). This reveals that the large distance of morphospace that separates nonmetazoans from metazoans is influenced strongly by those characters with a heavy contingent burden that evolved early within the metazoan stem-lineage. Crown metazoan characters, such as a defined head, stomochord, protoceol, ecdysis, segmentation, and features of nephridial cells, all load strongly along NMDS axis one, indicating that these characters are important in the establishment of metazoan body plans.

Undoubtedly, it would be useful to explore the impact of including a greater number of fossil taxa in future analyses of




Fig. 6. The role of transitioning to terrestrial environments in the post-Cambrian exploration of morphospace. The phylogenetic pathway within morphospace as well as the area occupied during the Cambrian is shown as in Fig. 3. Organisms that are exclusively aquatic (marine, brackish, freshwater, or parasitic) are colored in blue, whereas taxonomic groups that have some terrestrial representation are colored in red. Terrestrial groups are included within arthropods, chordates, onychophorans, tardigrades, gastropods, rotifers, annelids, nematodes, nemerteans, and Platyhelminthes.

