



Increased Arterial Stiffness in Adolescents With Obesity

Global Pediatric Health
Volume 6: 1–8
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2333794X19831297
journals.sagepub.com/home/gph


Yvette E. Lentferink, MD¹, Lisa A. J. Kromwijk, MD¹,
Marloes P. van der Aa, MD, PhD¹, Catherijne A. J. Knibbe, PharmD, PhD¹,
and Marja M. J. van der Vorst, MD, PhD¹ 

Abstract

Increased arterial stiffness (AS) is an early sign of cardiovascular disease. Influence of weight, puberty, and insulin resistance (IR) on AS in adolescents is unclear. Therefore, this study compared AS, assessed with pulse wave velocity (PWV) and augmentation index (Alx), of adolescents with and without obesity and evaluated the influence of puberty and IR on AS. Sixty-two lean and 61 adolescents with obesity were included. Significantly higher PWV was observed in adolescents with obesity (4.1 ms^{-1} [2.4 to 5.6 ms^{-1}] vs 3.6 ms^{-1} [0.4 to 6.1 ms^{-1}]; $P = .01$), while Alx was not significantly different. However, significantly higher Alx was observed in adolescents with obesity and IR (3.0 [-17.5% to 28.5%] vs -3.0 [-19.0% to 13.0%]; $P = .01$). For Tanner stages, no differences were observed. The higher PWV in adolescents with obesity and higher Alx in adolescents with obesity and IR both indicate an increased AS. Consequently, measurement of AS should be considered in adolescents with obesity and IR as part of cardiovascular risk assessment.

Keywords

obesity, adolescents, arterial stiffness, puberty, insulin resistance

Received May 2, 2018. Received revised December 8, 2018. Accepted for publication December 14, 2018.

Introduction

Metabolic and cardiovascular complications are more frequently seen in pediatric populations due to the increasing incidence of obesity and shift of obesity toward a younger age.¹⁻³ Since cardiovascular complications are the leading cause of death worldwide, awareness for cardiovascular risk assessment during childhood is important.⁴ However, measurements for early cardiovascular changes in children with obesity are rarely performed during routine clinical care, potentially causing underestimation and undertreatment of obesity-related cardiovascular complications.⁵ However, this is essential as it is known that early detection, evaluation, and treatment of cardiovascular complications can result in reversibility of cardiac abnormalities.⁵

Arterial stiffness (AS) is an early sign of cardiovascular disease.⁶⁻⁸ Stiffening of the arteries is a physiological process as a result of aging.⁹ It is the result of decreasing amount of elastin fibers and an overproduction of abnormal collagen in the arterial wall induced by

an inflammatory milieu.⁸ In addition, other factors such as hypertension and dyslipidemia may accelerate this process.¹⁰ It has also been suggested that AS is influenced by gender and puberty, caused by differences in sex steroids.¹¹ Estrogens are assumed to decrease AS while androgens are proposed to promote it, causing higher AS in boys compared with girls.¹¹ Ethnicity might also influence AS, since it has been shown that South-Asians and Africans have higher AS in comparison with Caucasians.¹² Moreover, weight and insulin resistance (IR) are suggested to have an influence on AS, although literature shows conflicting results.^{10,13}

AS can be measured using pulse wave velocity (PWV) and augmentation index (AIx), which are

¹St. Antonius Hospital, Nieuwegein, Netherlands

Corresponding Author:

Marja M. J. van der Vorst, St Antonius Hospital, Koekoekslaan 1, Nieuwegein, 3435 CM, Netherlands.
Email: m.van.der.vorst@antoniusziekenhuis.nl



increased in case of atherosclerosis.^{14,15} Applanation tonometry is considered to be the gold standard for measuring PWV and AIx.¹⁰ PWV can be determined at multiple sites of the arterial system. However, central measurements from carotid to femoral artery are preferred, since this has the strongest correlation with cardiovascular morbidity and mortality.^{16,17}

As there is limited and conflicting evidence on the influence of weight, puberty, and IR on AS in adolescents, the aim of this study is to compare AS of adolescents with obesity with lean adolescents and to determine the influence of puberty and IR.

Methods

Population

For this cross-sectional study, data were obtained through 2 separate study protocols, and both were approved by the Medical Ethical Committee of the St. Antonius Hospital, Nieuwegein/Utrecht, the Netherlands (lean participants: NL39370.100.12; participants with obesity: NL34811.100.11). All study procedures were in accordance with the Declaration of Helsinki and the Medical Research Involving Human Subjects Act (WMO) of the Netherlands. Written informed consent was obtained from all participants and their parents/caregivers. Lean participants (hereafter referred to as controls) were recruited between June 2012 and February 2016 during preoperative assessment at the outpatient clinic of the Anesthesia Department of the St. Antonius Hospital. Controls were eligible for inclusion if they were 10 to 20 years of age, were lean (body mass index standard deviation score [BMI-sds] >-1.1 and ≤ 1.1),^{18,19} of Caucasian descent, and scheduled for minor surgery. Exclusion criteria were syndromal disorders and chronic somatic diseases.

Participants with obesity (hereafter referred to as patients) were participating in a randomized controlled trial on metformin versus placebo,²⁰ in which PWV and AIx measurements were performed using applanation tonometry between August 2011 and January 2017. Only the first PWV and AIx measurements, before start of any study medication, were used in the current study. They were eligible for inclusion if they were 10 to 16 years of age, suffered from obesity (BMI-sds >2.3),^{18,19} and were of Caucasian descent. Patients with syndromal disorders and/or chronic somatic diseases (especially diabetes mellitus type 2) were excluded.

Measurements

During a physical examination by the attending physician, the following measurements were performed:

assessment of pubertal stage using Tanner stage classification and measurement of weight, height, waist circumference, and blood pressure. Pubertal stage was categorized as prepubertal (T1), pubertal (T2-T4), or postpubertal (T5). Weight and height were measured using calibrated measuring equipment with an accuracy of 0.05 kg and 0.1 cm, respectively. BMI was calculated as weight divided by height-squared (kg/m^2), BMI-sds and height-sds were calculated using the TNO growth calculator for professionals.²¹ Blood pressure was measured in supine position from the right arm.

Bio-impedance measurements were obtained with the bio-impedance scale (Tanita Body Composition Analyzer BC-420MA, Tanita Corporation, Tokyo, Japan). This noninvasive measurement is determining in a single frequency mode the total body water (kg), fat mass, and fat free mass.

Blood samples were taken for fasting plasma glucose, fasting plasma insulin, total cholesterol, high-density lipoprotein, low-density lipoprotein, and triglyceride. IR was calculated using Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) ($[\text{fasting plasma glucose (mmol/L)} \times \text{fasting plasma insulin (mU/L)/22.5}]$)²² and defined IR as HOMA-IR ≥ 3.4 .²³

AS was measured noninvasively using the SphygmoCor (Model SCOR-Px, Software version 7.01; AtCor Medical Pvt Ltd, Sydney, Australia), measuring both the carotid-femoral PWV and the AIx with applanation tonometry. For this measurement, pressure is applied to appanate (flatten) the artery using a special probe creating a signal that approximates arterial pressure.^{10,24} For the PWV, waveforms of the right carotid artery and the right femoral artery were obtained. The waveforms of the carotid and femoral arteries were recorded sequentially gated by the R-wave on a simultaneously recorded electrocardiogram. In addition, the distance from the carotid artery to femoral artery was measured to the nearest 0.1 cm using a tapeline. PWV was subsequently calculated by the software as the distance from the carotid-to-distal path length divided by the lapsed time between the “feet” of the 2 waveforms (carotid and femoral arteries) reported in ms^{-1} .^{10,25,26} A higher PWV indicates higher AS, since this method is based on the theory that a pulse wave generated by contraction of the left ventricle travels at a speed determined by the size, shape, and properties of the artery, which is altered by stiffening of the arteries.^{10,25} AIx was measured at the right radial artery. It represents the pulse wave reflection, which is derived from the difference between the main outgoing wave and the reflected wave of the central arterial waveform (ie, augmentation pressure). The AIx is calculated by dividing the augmentation pressure by the pulse pressure (systolic blood

Table 1. Population Characteristics^a.

	Patients (N = 61)	Controls (N = 62)	P ^b
Age (years)	13.4 (10.2 to 16.8)	15.0 (10.1 to 19.5)	.01
Male, n (%)	17 (27.9)	30 (48.4)	.02
Height (cm)	162.6 (145.7 to 179.2)	166.5 (143.0 to 192.0)	.02
Height-sds	0.2 (-1.9 to 2.9)	0.24 (-1.36 to 1.71)	.71
Weight (kg)	82.7 (58.6 to 117.8)	54.0 (31.5 to 86.2)	<.01
BMI (kg/m ²)	30.8 (25.3 to 44.1)	18.6 (14.8 to 24.0)	<.01
BMI-sds	3.3 (2.4 to 4.4)	0.1 (-1.0 to 1.0)	<.01
Fat (%)	41.2 (27.1 to 54.7)	17.6 (5.5 to 33.6)	<.01
Tanner stage, n (%)			.30
Prepubertal	9 (14.8)	12 (19.4)	
Pubertal	40 (65.8)	42 (67.7)	
Postpubertal	11 (18.0)	6 (9.7)	
SBP (mm Hg)	120 (92 to 135)	114 (92 to 153)	<.01
DBP (mm Hg)	67 (46 to 84)	66 (40 to 84)	.41
FPG (mmol/L)	4.8 (4.0 to 5.5)	4.7 (3.1 to 6.5)	.62
FPI (mmol/L)	19.0 (1.0 to 48.0)	6.8 (0.8 to 26.1)	<.01
HOMA-IR	4.1 (0.2 to 11.5)	1.4 (0.2 to 6.6)	<.01
IR, n (%)	39 (63.9)	4 (6.5)	<.01
Cholesterol (mmol/L)	4.7 (2.8 to 7.8)	3.9 (2.3 to 5.5)	<.01
LDL (mmol/L)	2.8 (1.6 to 5.8)	2.1 (0.8 to 3.9)	<.01
HDL (mmol/L)	1.20 (0.70 to 1.80)	1.45 (0.89 to 2.18)	<.01
Triglyceride (mmol/L)	1.4 (0.5 to 3.4)	1.0 (0.3 to 3.0)	<.01

Abbreviations: sds, standard deviation score; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; FPG, fasting plasma glucose; FPI, fasting plasma insulin; HOMA-IR, Homeostatic Model Assessment for Insulin Resistance; LDL, low-density lipoprotein; HDL, high-density lipoprotein.

^aData presented as median with range or frequency with percentage.

^bP values represents the differences between healthy controls and study participants (Student's *t* test). Bold values represent significant *P* values.

pressure – diastolic blood pressure) and expressed as percentage.^{10,25} A higher AIX indicates stiffer arteries. Since the AIX is influenced by heart rate, all values were automatically adjusted to a heart rate of 75 beats per minute.

Statistical Analysis

Statistical analysis was performed using IBM SPSS Statistics, version 24 (IBM SPSS Statistics, Chicago, IL). Continuous parameters were reported as median with range. Categorical data were expressed as frequencies with percentage. The Student's *t* test or Mann-Whitney *U* test and χ^2 test were used to compare controls and patients. As it can be assumed that most participants with IR are part of the patient subgroup, the analysis with regard to the influence of IR on AS was limited to the patient subgroup, so that BMI can be excluded as a potential confounder. The Student's *t* test or Mann-Whitney *U* test and χ^2 test were used to compare the no-IR and IR subgroup. The Kruskal-Wallis test and χ^2 test were used to compare the different Tanner stages. Correlations were determined using the Spearman *R*

test. An α level of 5% was considered significant for all statistical tests.

Results

Table 1 shows the population characteristics of the study population. Patients and controls were comparable regarding pubertal stages. Patients were significantly younger, smaller, and less frequently of the male sex ($P = .01$; $P = .02$; $P = .02$, respectively). In addition, patients had by definition a significant higher weight, BMI, and BMI-sds (all $P < .01$). Moreover, all metabolic parameters were significantly higher in the patients compared with controls (all $P < .01$; Table 1).

As shown in Figure 1, PWV was significantly higher in patients compared with controls (4.1 ms^{-1} [2.4 to 5.6 ms^{-1}] vs 3.6 ms^{-1} [0.4 to 6.1 ms^{-1}]; $P = .01$). For AIX, no significant difference was observed (1.0% [-19.0% to 28.5%] vs -1.0% [-33.5% to 29.0%]; $P = .38$).

When stratifying the total population according to Tanner stage, significant differences were observed for sex, age, height, weight, and BMI, while no differences were observed for BMI-sds and metabolic parameters,




Figure 1. Boxplot with median and interquartile range of (A) PWV (pulse wave velocity) and (B) AIx (augmentation index) of controls (light gray) compared with patients (dark gray).




Figure 2. Boxplot with median and interquartile range of (A) PWV (pulse wave velocity) and (B) AIx (augmentation index) of the total population stratified by pubertal stage; prepubertal (light gray, $n = 21$), pubertal (medium gray, $n = 83$), and postpubertal (dark gray, $n = 17$) participants.

which is shown in Supplementary Table 1 (available online). In the prepubertal subgroup, participants were significantly more frequently male ($P < .01$). And at the successive puberty stages, participants had a significant higher age, height, weight, and BMI (all $P < .01$; Supplementary Table 1).

With regard to the PWV and AIx, no significant differences between the Tanner stages were observed in the total population: (PWV: prepubertal = 3.8 ms^{-1} [0.4 to 5.2 ms^{-1}] vs pubertal = 4.2 ms^{-1} [0.6 to 6.1 ms^{-1}] vs postpubertal = 3.9 ms^{-1} [1.4 to 4.9 ms^{-1}]; $P = .17$); (AIx: prepubertal = -1.0% [-18.25% to 23.5%] vs pubertal = 1.0% [-33.5% to 29.0%] vs postpubertal = -2.5% [-27.5% to 4.5%]; $P = .12$), which is shown in Figure 2. When

stratifying the controls and patients separately according to Tanner stage, all results with regard to population characteristics and PWV and AIx measurements were comparable to the results observed in the total population and therefore only shown in Supplementary Table 1.

Patients with obesity, with and without IR, were comparable regarding age, sex, pubertal stage, weight, and all of the laboratory values except triglyceride and by definition HOMA-IR (Table 2). Patients with IR had a significantly higher AIx (3.0% [-17.5% to 28.5%] vs -3.0% [-19.0% to 13.0%]; $P = .01$), while no difference was observed for PWV (4.1 ms^{-1} [2.4 to 5.6 ms^{-1}] vs 4.2 ms^{-1} [2.5 to 5.5 ms^{-1}]; $P = .64$) between patients with and without IR as shown in Figure 3.

Table 2. Characteristics of Patients With Obesity Stratified by Presence of IR^a.

	No IR (N = 22)	IR (N = 39)	<i>P</i> ^b
Age (years)	14.2 (10.2 to 16.6)	12.7 (10.3 to 16.8)	.16
Male (%)	8 (36.4)	9 (23.1)	.27
Tanner stage, n (%)			.32
Prepubertal	3 (14.3)	6 (15.4)	
Pubertal	12 (57.1)	28 (71.8)	
Postpubertal	6 (28.6)	5 (12.8)	
Height (cm)	163.0 (145.7 to 177.6)	161.5 (147.4 to 179.2)	.56
Height-sds	-0.08 (-1.86 to 2.21)	0.23 (-1.40 to 2.94)	.19
Weight (kg)	82.6 (53.2 to 117.8)	82.4 (58.6 to 114.4)	.96
BMI (kg/m ²)	31.46 (23.46 to 44.09)	31.11 (25.39 to 42.22)	.29
BMI-sds	3.38 (2.42 to 4.15)	3.25 (2.39 to 4.42)	.72
Fat (%)	39.0 (28.1 to 51.0)	41.5 (33.6 to 54.7)	.05
SBP (mm Hg)	118 (109 to 135)	120 (92 to 135)	.83
DBP (mm Hg)	67 (46.83)	65 (46-85)	.74
FPG (mmol/L)	4.6 (4.0 to 5.0)	4.8 (4.4 to 5.5)	<.01
FPI (mmol/L)	11.0 (1.0 to 16.0)	25.0 (16.0 to 48.0)	<.01
HOMA-IR	2.3 (0.2 to 3.3)	5.3 (3.4 to 11.5)	<.01
Cholesterol (mmol/L)	4.5 (2.8 to 7.2)	4.5 (3.2 to 7.8)	.60
LDL (mmol/L)	2.7 (1.6 to 5.3)	2.7 (1.6 to 5.8)	.92
HDL (mmol/L)	1.13 (0.70 to 1.77)	1.14 (0.80 to 1.80)	.82
Triglyceride (mmol/l)	1.1 (0.5 to 3.4)	1.5 (0.6 to 2.3)	.04

Abbreviations: IR, insulin resistance; sds = standard deviation score; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; FPG, fasting plasma glucose; FPI, fasting plasma insulin; HOMA-IR, Homeostatic Model Assessment for Insulin Resistance; LDL, low-density lipoprotein; HDL, high-density lipoprotein.

^aData presented as median with range or frequency with percentage.

^b*P* values represents the differences between the no-IR and IR subgroups (Student's *t* test). Bold values represent significant values.




Figure 3. Boxplot with median and interquartile range of (A) PWV (pulse wave velocity) and (B) Alx (augmentation index) of patients with obesity with IR (light gray, n = 22) compared with obesity without IR (dark gray, n = 39).

For the total populations, significant correlations were observed between PWV and BMI, BMI-sds, and fat percentage ($r = 0.22$, $P = .04$; $r = 0.31$, $P < .01$; $r = 0.24$, $P = .03$, respectively). No correlation was observed between PWV and HOMA-IR ($r = 0.16$,

$P = .17$); age ($r = -0.13$, $P = .26$); Tanner stage ($r = -0.10$, $P = .39$), nor for blood pressure or lipid spectrum. Significant correlations were observed between Alx and Tanner stage ($r = -0.26$, $P = .01$) and HOMA-IR ($r = 0.23$, $P = .02$). No correlation was observed between

AIx and BMI ($r = 0.09$, $P = .37$); BMI-sds ($r = 0.10$, $P = .34$); age ($r = 0.17$, $P = .08$); fat percentage ($r = 0.16$, $P = .12$), nor for blood pressure or lipid spectrum.

Discussion

Arterial stiffness is considered an early sign of cardiovascular disease,⁶⁻⁸ which can be measured noninvasively using PWV and AIx. It has been suggested that AS in pediatric patients is influenced by sex steroids, weight, and IR.^{10,11,13} However, in pediatric populations, literature on these topics is scarce and conflicting.^{10,11,13} Therefore, AS of lean and obese adolescents of Caucasian descent was compared in this study and the influence of puberty and IR on AS evaluated.

In the current study, a significantly higher PWV was observed in adolescents with obesity. In contrast, no significant difference in AIx was observed between patients and controls. Our findings regarding the effect of obesity on PWV are in line with some previous studies in adolescents (aged 10-24 years) and adults.^{10,27-31} However, other studies reported a lower PWV in children/adolescents with obesity compared with lean controls.^{10,32-34} It is known that the arterial diameter and compliance increases physiologically with growth and development, resulting in a decreased peripheral resistance.⁵ It has been suggested that these adaptations are greater in children with obesity, resulting in a lower PWV in comparison with those with normal weight.^{5,10} However, obesity is also associated with a higher arterial pressure caused by an increased preload and afterload due to increased metabolic demand.⁵ Consequently, it can be anticipated that PWV is higher in populations with obesity if the arterial pressure surpasses the (greater) physiological adaptations.^{5,10} The higher PWV (and thus AS) observed in our patients with obesity implicates that the physiological adaptations have been exceeded, indicating that they may already be exposed to an increased risk to develop cardiovascular diseases. This observation is strengthened by the fact that the patients subgroup consisted of significantly more girls, and was significantly younger, both factors that normally will be "protective," as it is known that PWV is higher in boys and increases with age.^{11,17,35} The significant higher systolic blood pressure and disturbed lipid spectrum in our population of obese adolescents (Table 1) may also explain the current findings and support the hypothesis that our patients are already at higher risk to develop cardiovascular disease. The difference with studies reporting a lower PWV in children/adolescents with obesity might be explained by a lower age and blood pressure in comparison with our patients, so that the physiological adaptations were

still sufficient to mask the negative consequences of obesity.³²⁻³⁴

With regard to AIx and obesity, literature is conflicting, as some did show an association whereas others did not.^{10,13,25,34} As we did not observe a difference for AIx nor a correlation between BMI and AIx, it might be suggested that AIx is not a discriminating measurement to determine AS in children with obesity.

As it has been described that estrogens might decrease AS and that androgens might promote AS, differences between Tanner stages were expected as sex steroids are changing during puberty.¹¹ However, in the current study no significant differences for PWV nor AIx were observed between Tanner stages in the entire population nor for the lean and obese populations separately (Figure 2; Supplementary Table 1). It is known that girls with obesity have relatively higher androgen levels and are consequently prone for an increased AS, which might have influenced the results.³⁶ However, we could only speculate on the theory, since no analysis stratified by sex was performed, since the number of participants assigned to a subgroup stratified by sex and Tanner stage was too small to analyze. In addition, the observed difference in BMI may have limited the influence of Tanner stage on AS. Since BMI increases physiologically with age in children, this factor could also not be excluded in analysis performed solely in controls or patients. To the best of authors knowledge, only one study has been performed with specific interest into the effect of Tanners stage on AS.¹¹ This study only included prepubescent and postpubescent subjects and focused on differences in sex; therefore, comparison with the results of the current study was not possible.¹¹ In the current study, no differences were observed for Tanner stage, which might suggest that AS is not influenced by puberty.

IR is associated with obesity and usually the first sign of a disturbed glucose metabolism.³⁷ Apart from being a risk factor for development of type 2 diabetes mellitus, IR has also been suggested as an independent risk factor for development of cardiovascular diseases by potentiating the onset of dyslipidemia.^{37,38} Therefore, an additional analysis into the influence of IR in patients with obesity was performed in the current study. A significant higher AIx but not PWV was observed in patients with IR, and only AIx was correlated with IR. Several studies investigated this association between IR and AS.^{10,13,25,39-42} Some observed an association between PWV and IR,^{10,39,40,42} while others did not.^{10,13,25,41} One of the studies incorporated AIx in their analysis and showed an association between IR and AIx.¹³ However, the association between IR and AIx disappeared after adjustment for other cardiovascular risk factors such as age, gender, BMI, and blood pressure.¹³ From the above, it can be concluded that IR has a role in increasing AS. However, the

association might be influenced by the duration and severity of IR besides other factors such as obesity. Nevertheless, based on the results of the current study, measurement of AS in pediatric populations with obesity and IR should be considered, in order to evaluate early signs of cardiovascular diseases. This is important as cardiovascular diseases are the leading cause of death worldwide,⁴ and especially since it is known that early detection, evaluation, and treatment of cardiovascular complications can result in reversibility of cardiac abnormalities.⁵ However, prospective trials have to be performed to evaluate the predictive value of measurement of AS with PWV and AIx, the ability to use this tool in routine clinical care, as well as the cost-effectiveness.

To the best of authors' knowledge, this is the first study concerning the influence of weight, Tanner stage, and IR on AS, measured with PWV and AIx using applanation tonometry, in a population of lean and obese adolescents of Caucasian descent. Consequently, there is no influence of ethnic background as confounder in the current study. However, certain limitations should be mentioned. First, applanation tonometry used to measure PWV and AIx requires a skilled operator and can be difficult to measure in subjects with obesity. To overcome this limitation, measurements of all participants were performed by a trained staff. Second, the influence of Tanner stage on AS was not observed, which might have been caused by the limited number of participants and/or the fact that no analysis stratified by sex could be performed.

Conclusion

Higher PWV in adolescents with obesity and higher AIx in adolescents with obesity and IR were observed, both indicating an increased AS. Consequently, measurement of AS should be considered in adolescents with obesity and IR as part of cardiovascular risk assessment.

Author Contributions

YL: has collected study data, analyzed the data and wrote the manuscript. LK: collected study data and revised the manuscript. MA: collected study data and revised the manuscript. CK: was involved in the conception and design of the study and revised the manuscript. MVDV: was involved with conception and design of the study, supervised data collection, revised the manuscript, and is the guarantor of this work and, as such, had full access to all data in the study and takes responsibility for the integrity of the data and accuracy of the data analysis. All authors read and approved the final manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Marja M. J. van der Vorst  <https://orcid.org/0000-0001-9581-3044>

Supplemental Material

Supplemental material for this article is available online.

References

1. Han JC, Lawlor DA, Kimm SY. Childhood obesity. *Lancet*. 2010;375:1737-1748.
2. Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents: identification, associated health risks, and treatment approaches: a scientific statement from the American Heart Association. *Circulation*. 2013;128:1689-1712.
3. Lobstein T, Jackson-Leach R. Planning for the worst: estimates of obesity and comorbidities in school-age children in 2025. *Pediatr Obes*. 2016;11:321-325.
4. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents; National Heart, Lung, and Blood Institute. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: summary report. *Pediatrics*. 2011;128(suppl 5):S213-S256.
5. Cote AT, Harris KC, Panagiotopoulos C, Sandor GG, Devlin AM. Childhood obesity and cardiovascular dysfunction. *J Am Coll Cardiol*. 2013;62:1309-1319.
6. Mozos I, Malainer C, Horbańczuk J, et al. Inflammatory markers for arterial stiffness in cardiovascular diseases. *Front Immunol*. 2017;8:1058.
7. Safar ME, Blacher J, Jankowski P. Arterial stiffness, pulse pressure, and cardiovascular disease—is it possible to break the vicious circle? *Atherosclerosis*. 2011;218:263-271.
8. Zieman SJ, Melenovsky V, Kass DA. Mechanisms, pathophysiology, and therapy of arterial stiffness. *Arterioscler Thromb Vasc Biol*. 2005;25:932-943.
9. Mackenzie IS, Wilkinson IB, Cockcroft JR. Assessment of arterial stiffness in clinical practice. *QJM*. 2002;95:67-74.
10. Panchangam C, Merrill ED, Raghuvver G. Utility of arterial stiffness assessment in children. *Cardiol Young*. 2018;28:362-376.
11. Ahimastos AA, Formosa M, Dart AM, Kingwell BA. Gender differences in large artery stiffness pre- and post puberty. *J Clin Endocrinol Metab*. 2003;88:5375-5380.
12. Snijder MB, Stronks K, Agyemang C, Busschers WB, Peters RJ, van den Born BJ. Ethnic differences in arterial stiffness the Helius study. *Int J Cardiol*. 2015;191:28-33.
13. Urbina EM, Gao Z, Khoury PR, Martin LJ, Dolan LM. Insulin resistance and arterial stiffness in healthy adolescents and young adults. *Diabetologia*. 2012;55:625-631.
14. Cheung YF. Arterial stiffness in the young: assessment, determinants, and implications. *Korean Circ J*. 2010;40:153-162.

15. Tomiyama H, Yamashina A. Non-invasive vascular function tests: their pathophysiological background and clinical application. *Circ J*. 2010;74:24-33.
16. Tsuchikura S, Shoji T, Kimoto E, et al. Central versus peripheral arterial stiffness in association with coronary, cerebral and peripheral arterial disease. *Atherosclerosis*. 2010;211:480-485.
17. Reusz GS, Csepregal O, Temmar M, et al. Reference values of pulse wave velocity in healthy children and teenagers. *Hypertension*. 2010;56:217-224.
18. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes*. 2012;7:284-294.
19. Hirasing RA, Fredriks AM, van Buuren S, Verloove-Vanhorick SP, Wit JM. Increased prevalence of overweight and obesity in Dutch children, and the detection of overweight and obesity using international criteria and new reference diagrams [in Dutch]. *Ned Tijdschr Geneeskd*. 2001;145:1303-1308.
20. van der Aa MP, Elst MA, van de Garde EM, van Mil EG, Knibbe CA, van der Vorst MM. Long-term treatment with metformin in obese, insulin-resistant adolescents: results of a randomized double-blinded placebo-controlled trial. *Nutr Diabetes*. 2016;6:e228.
21. TNO. De TNO groeicalculator voor professionals—op basis van de vijfde landelijke groeistudie. <https://www.tno.nl/nl/aandachtsgebieden/gezond-leven/roadmaps/youth/groeicalculator-voor-professionals/>. Accessed February 6, 2019.
22. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28:412-419.
23. van der Aa MP, Fazeli Farsani S, Kromwijk LA, et al. How to screen obese children at risk for type 2 diabetes mellitus? *Clin Pediatr (Phila)*. 2014;53:337-342.
24. Stoner L, Young JM, Fryer S. Assessments of arterial stiffness and endothelial function using pulse wave analysis. *Int J Vasc Med*. 2012;2012:903107.
25. Shah AS, Gao Z, Urbina EM, Kimball TR, Dolan LM. Prediabetes: the effects on arterial thickness and stiffness in obese youth. *J Clin Endocrinol Metab*. 2014;99:1037-1043.
26. Laurent S, Cockcroft J, Van Bortel L, et al; European Network for Non-invasive Investigation of Large Arteries. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J*. 2006;27:2588-2605.
27. Recio-Rodriguez JI, Gomez-Marcos MA, Patino-Alonso MC, et al. Abdominal obesity vs general obesity for identifying arterial stiffness, subclinical atherosclerosis and wave reflection in healthy, diabetics and hypertensive. *BMC Cardiovasc Disord*. 2012;12:3.
28. Scuteri A, Orru M, Morrell CH, et al. Associations of large artery structure and function with adiposity: effects of age, gender, and hypertension. The SardiNIA Study. *Atherosclerosis*. 2012;221:189-197.
29. Dangardt F, Chen Y, Berggren K, Osika W, Friberg P. Increased rate of arterial stiffening with obesity in adolescents: a five-year follow-up study. *PLoS One*. 2013;8:e57454.
30. Sakuragi S, Abhayaratna K, Gravenmaker KJ, et al. Influence of adiposity and physical activity on arterial stiffness in healthy children: the lifestyle of our kids study. *Hypertension*. 2009;53:611-616.
31. Urbina EM, Kimball TR, Khoury PR, Daniels SR, Dolan LM. Increased arterial stiffness is found in adolescents with obesity or obesity-related type 2 diabetes mellitus. *J Hypertens*. 2010;28:1692-1698.
32. Dangardt F, Osika W, Volkmann R, Gan LM, Friberg P. Obese children show increased intimal wall thickness and decreased pulse wave velocity. *Clin Physiol Funct Imaging*. 2008;28:287-293.
33. Charakida M, Jones A, Falaschetti E, et al. Childhood obesity and vascular phenotypes: a population study. *J Am Coll Cardiol*. 2012;60:2643-2650.
34. Lurbe E, Torro I, Garcia-Vicent C, Alvarez J, Fernández-Fornoso JA, Redon J. Blood pressure and obesity exert independent influences on pulse wave velocity in youth. *Hypertension*. 2012;60:550-555.
35. Hidvégi EV, Illyés M, Benczúr B, et al. Reference values of aortic pulse wave velocity in a large healthy population aged between 3 and 18 years. *J Hypertens*. 2012;30:2314-2321.
36. Solorzano CMB, Helm KD, Patrie JT, et al. Increased adrenal androgens in overweight peripubertal girls. *J Endocr Soc*. 2017;1:538-552.
37. Castro AV, Kolka CM, Kim SP, Bergman RN. Obesity, insulin resistance and comorbidities? Mechanisms of association. *Arq Bras Endocrinol Metabol*. 2014;58:600-609.
38. Steinberger J, Daniels SR; American Heart Association Atherosclerosis, Hypertension, and Obesity in the Young Committee (Council on Cardiovascular Disease in the Young); American Heart Association Diabetes Committee (Council on Nutrition, Physical Activity, and Metabolism). Obesity, insulin resistance, diabetes, and cardiovascular risk in children: an American Heart Association scientific statement from the Atherosclerosis, Hypertension, and Obesity in the Young Committee (Council on Cardiovascular Disease in the Young) and the Diabetes Committee (Council on Nutrition, Physical Activity, and Metabolism). *Circulation*. 2003;107:1448-1453.
39. Czernichow S, Bertrais S, Blacher J, et al. Metabolic syndrome in relation to structure and function of large arteries: a predominant effect of blood pressure. A report from the SU.VI.MAX. Vascular Study. *Am J Hypertens*. 2005;18(9 pt 1):1154-1160.
40. Kasayama S, Saito H, Mukai M, Koga M. Insulin sensitivity independently influences brachial-ankle pulse-wave velocity in non-diabetic subjects. *Diabet Med*. 2005;22:1701-1706.
41. Schack-Nielsen L, Mølgaard C, Larsen D, et al. Arterial stiffness in 10-year-old children: current and early determinants. *Br J Nutr*. 2005;94:1004-1011.
42. McCloskey K, Sun C, Pezic A, et al. The effect of known cardiovascular risk factors on carotid-femoral pulse wave velocity in school-aged children: a population based twin study. *J Dev Orig Health Dis*. 2014;5:307-313.