

# SCIENTIFIC REPORTS



OPEN

## Identification of Liver Epithelial Cell-derived Ig Expression in $\mu$ chain-deficient mice

Wenwei Shao<sup>1</sup>, Chi Zhang<sup>1</sup>, Enyang Liu<sup>2</sup>, Long Zhang<sup>2</sup>, Junfan Ma<sup>1</sup>, Zhu Zhu<sup>1</sup>, Xiaoting Gong<sup>1</sup>, Zhihai Qin<sup>3</sup> & Xiaoyan Qiu<sup>1,2</sup>

Received: 20 October 2015

Accepted: 07 March 2016

Published: 29 March 2016

Growing evidence indicates that B cells are not the only source of immunoglobulin (Ig). To investigate this discovery further, we used  $\mu$ MT mice, which have a disruption of the first transmembrane exon of the  $\mu$  heavy chain and do not express the membrane form of IgM. These mice lack mature B cells and thus serve as a good model to explore Ig expression by liver epithelial cells. We found that Ig heavy chains ( $\mu$ ,  $\delta$ ,  $\gamma$  and  $\alpha$ ) and light chains ( $\kappa$  and  $\lambda$ ) were expressed in sorted liver epithelial cells of  $\mu$ MT mice. Surprisingly, each heavy chain class showed its respective variable region sequence characteristics in their variable region, instead of sharing the same VDJ usage, which suggests that class switching does not occur in liver epithelial cells. Moreover, the  $\gamma$  and  $\alpha$  chains, but not the  $\mu$  and  $\delta$  chains, showed mutations in the variable region, thus indicating that different classes of Ig have different activities. Our findings support the concept that non-B cells, liver epithelial cells here, can produce different classes of Ig.

Immunoglobulin (Ig) is one of the classic immune molecules and plays an important role in the body's immune response. Currently, B lineage lymphocytes are considered to be the only source of Ig. However, since 20 years ago, a series of studies by our group proved that many non-B cancer cells, especially epithelial cancer cells, can also express Ig, including IgG, IgA and IgM<sup>1–9</sup>. Moreover, epithelial cancer cell-expressed IgG showed growth factor-like activity, which can promote cancer progression<sup>1,2,10</sup>. Subsequently, these unusual findings were confirmed by other researchers<sup>11–19</sup>. Recently, several types of Ig, including IgG, IgA and IgM, have been found in normal non-B cells, including epithelial cells, germ cells, neurons, endothelial cells, and even monocytes<sup>4,5,13,20,21</sup>. Moreover, normal epithelial cell-derived IgG, IgA and IgM showed characteristic antibody activity<sup>9,22</sup>. All of these studies have challenged the classical concept that B cells are the only source of Ig.

The functional membrane form of the IgM heavy chain ( $\mu$ ) is thought to be essential for B cell differentiation. B cells that lack  $\mu$  chains are eliminated by the body<sup>23</sup>. The B cell-deficient  $\mu$ MT mice contain a disruption of the first transmembrane exon of the  $\mu$  heavy chain and thus do not express the membrane form of IgM. These mice lack mature B cells due to a developmental block at the pro-B cell stage, after which B cells undergo apoptosis<sup>24</sup>.  $\mu$ MT mice are considered to be a suitable model to explore infection or tumor immunity in the absence of B cells and Ig production due to the lack of B cells<sup>25–30</sup>. However, growing studies have found that  $\mu$ MT mice contain both Ig, including IgA, IgG and IgE, and Ig-producing cells<sup>31–34</sup>. With regard to our previous finding that Ig can be found in many non-B cells in human and mice, we hypothesized that the Ig in  $\mu$ MT mice is mainly produced by non-B cells instead of the residual a small population of B cells as described above.

In this study, B cell-deficient  $\mu$ MT mice were used as a model to verify our hypothesis. We first identified IgM and IgA expression in several non-immune tissues, including liver, lung and kidney. The levels of IgM and IgA in these tissues were similar to those found in wild type (WT) mice, whereas the levels of serum Ig in the  $\mu$ MT mice were much lower than those in WT mice. Subsequently, we analyzed IgM, IgG, IgA, and IgD heavy and light chain transcripts and protein in sorted liver epithelial cells and found that liver epithelial cells could express different classes of Ig. Moreover, the liver epithelial cell-derived Ig transcripts displayed distinct characteristics compared with B cell-derived Ig transcripts.

<sup>1</sup>Department of Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing, 100191, China. <sup>2</sup>Key Laboratory of Medical Immunology, Ministry of Health, Beijing, 100191, China. <sup>3</sup>Department of Immunology, Institute of Biophysics Chinese Academy of Sciences, Beijing, 100101, China. Correspondence and requests for materials should be addressed to Z.Q. (email: zhihai@ibp.ac.cn) or X.Q. (email: qiuxy@bjmu.edu.cn)

## Results

**No mature B cells and low levels of serum Ig were detected in  $\mu$ MT mice.** We first detected whether there was residual mature B cells in  $\mu$ MT mice. In peripheral blood, there were no cells that stained with the B220 B cell marker (Fig. 1A). We then analyzed B cell development in the bone marrow (BM) of  $\mu$ MT mice. Unlike their WT counterparts (BALB/c mice),  $\mu$ MT mice contained neither pre-B cells (CD43<sup>-</sup> B220<sup>+</sup>) nor mature B cells (B220<sup>+</sup> IgM<sup>+</sup>) in the BM (Fig. 1B,C). B cell development from pro-B to pre-B is prevented in  $\mu$ MT mice. Therefore, we used  $\mu$ MT mice as a B cell-deficient model.

Next, we measured the concentrations of IgM and IgG in the serum of  $\mu$ MT mice by ELISA, using the serum of WT mice as a control. The serum IgM and IgG levels in the  $\mu$ MT mice were reduced nearly 100-fold compared with WT mice (Fig. 1D).

**Detection of IgM and Ig $\kappa$  in non-B cells in multiple tissues of  $\mu$ MT mice by immunostaining.** To determine whether Ig was widely present in non-B cells of the  $\mu$ MT mice, we performed immunohistochemistry using anti-mouse IgM and anti-mouse Ig $\kappa$  chain antibodies. Although there was hardly any positive staining in the spleen (Fig. 2A), non-B cells (especially epithelial cells) in multiple tissues, including liver, lung, kidney, and small intestine, displayed a positive staining pattern that was equivalent to WT mice (Fig. 2B,C). In addition, the fact that Ig was widely present in non-B cells was confirmed by Western blotting (Fig. 2D).

**Different classes of Ig were found in livers of  $\mu$ MT mice.** To further explore the expression of Ig, we analyzed different classes of Ig in livers of  $\mu$ MT mice using a Rapid Mouse Ig Isotyping Array. First, blood was drawn from the eye canthus to obtain serum, and then the mice were perfused to remove blood cells and serum Ig. Protein was extracted from the liver, and the Ig in the liver lysate and the serum were quantitated by fluorescence with Rapid Mouse Ig Isotyping Array. As expected, there were low levels of Ig in the serum of  $\mu$ MT mice compared with WT mice. However, the levels of IgM, IgG1, IgG2b and Ig $\kappa$  in the  $\mu$ MT liver lysate were the same as those in the WT liver lysate. Moreover, the levels of IgA, IgD, Ig $\lambda$  and IgG2a were much higher in the  $\mu$ MT liver lysate than in the WT liver lysate (Fig. 3A). We also confirmed this characteristic of IgM and IgG in the liver lysate by ELISA (Fig. 3B).


Subsequently, to confirm that liver cells of  $\mu$ MT mice could produce Ig, we analyzed the  $\mu$  and  $\kappa$  chains in the  $\mu$ MT liver lysate by mass spectrometry (MS). An SDS-PAGE band of liver lysate of approximately 75 kDa, which was stained with a goat anti-mouse IgM antibody by Western blotting, contained 11 peptides that correspond to the mouse Ig  $\mu$  heavy chain (Fig. 3C). Similarly, a band of approximately 26 kDa, which was detected by an anti-mouse Ig  $\kappa$  antibody by Western blotting, yielded 1 peptide that corresponded to a  $\kappa$  chain variable region and 1 peptide that corresponded to a  $\kappa$  chain constant region (Fig. 3D). In addition, 1 peptide corresponding to the  $\lambda$  chain constant region was found in the 26 kDa band (Fig. 3E).

**IgM and IgG were secreted by liver epithelial cells in  $\mu$ MT mice.** To verify that the Ig found in the liver was produced by epithelial cells, we performed flow cytometric sorting of liver epithelial cells using the epithelial marker CK18 (Fig. 4A). We cultured the sorted cells and used ELISPOT to determine whether IgM or IgG was being secreted. In fact, sorted liver epithelial cells from both  $\mu$ MT and WT mice could secrete IgM and IgG (Fig. 4B). Moreover, the levels of IgM and IgG were comparable between  $\mu$ MT and WT mice (Fig. 4B).

**Ig heavy and light chain transcripts in sorted liver epithelial cells of  $\mu$ MT mice.** To confirm the presence of  $\mu$  and  $\kappa$  chain transcripts in the liver cells of  $\mu$ MT mice, we performed Northern blot analysis using the DNA fragments of Ig $\mu$  or Ig $\kappa$  constant region as probes, the Ig $\mu$  and  $\kappa$  chain transcripts were significantly found in liver tissues of  $\mu$ MT mice (Fig. 5A). Subsequently, the Ig $\mu$  and  $\kappa$  chain transcripts were also determined in the liver epithelial cells of  $\mu$ MT mice with digoxigenin (DIG)-labeled RNA fragments of Ig $\mu$  CH3-CH4, or J $\kappa$ -C $\kappa$  as probes by *in situ* hybridization (Fig. 5B).


We used RT-PCR to analyze the expression of Ig in the sorted liver epithelial cells. The variable and constant regions of the different classes of heavy chains and light chains were amplified using a series of primers (Table S1). We detected rearranged transcripts of heavy chains ( $\mu$ ,  $\delta$ ,  $\gamma$ , and  $\alpha$ ) and light chains ( $\kappa$  and  $\lambda$ ) in cDNA prepared from the sorted liver epithelial cells of  $\mu$ MT mice (Fig. 5C). However, we did not detect transcripts of the Ig $\epsilon$  chain. We also did not detect transcripts of CD20, a B cell marker, which confirmed that there was no B cell contamination in the cDNA libraries. The identities of all of the PCR products were confirmed by DNA sequencing, and these sequences have been submitted to the GenBank database (GenBank Accession Numbers: KT726173–KT726219).

**Ig variable region in sorted liver epithelial cells of  $\mu$ MT mice displayed distinct sequence characteristics.** We cloned 86 VDJ rearrangements (from the heavy chain variable region) and 35 V $\kappa$ J $\kappa$  rearrangements (from the  $\kappa$  chain variable region) that were expressed by the sorted liver epithelial cells from 5  $\mu$ MT mice. Sequence analysis showed that 82 of the 86 VDJ rearrangements and all 35 V $\kappa$ J $\kappa$  rearrangements were products of functional Ig rearrangements (Table 1). The analysis of Ig repertoires showed that similar to our previous findings<sup>6</sup>, in the same sorted epithelial cell population, the V, D, and J usage and VDJ sequences were dissimilar among the  $\mu$ ,  $\gamma$ ,  $\alpha$  and  $\delta$  heavy chains. In fact, each class of heavy chain showed its own unique VDJ pattern (Table 1). The results further suggested that the liver epithelial cells did not possess a class switching mechanism. Interestingly, the  $\mu$ ,  $\gamma$  and  $\kappa$  transcripts showed a restricted clonal VDJ pattern, whereas the  $\alpha$  and  $\delta$  transcripts displayed VDJ diversity patterns similar to those exhibited by B cell-derived Ig and distinguished from Ig $\mu$  and Ig $\gamma$  in liver epithelial cells (Table 1). To exclude an influence of the primers, we also amplified variable regions of the  $\mu$  and  $\kappa$  chains from WT mice spleen cells and found the rearrangement patterns to be diverse (Table 2). In addition, we analyzed mutations in the representative VDJ and V $\kappa$ J $\kappa$  rearrangements (Fig. 6A). The primers for the  $\lambda$  light




**Figure 1. Identification of B cells and Igs in  $\mu$ MT mice.** (A) B lymphocyte and T lymphocyte in peripheral blood of WT mice and  $\mu$ MT mice were detected with PE anti-mouse B220 and FITC anti-mouse CD3 by FACS. (B) Pro-B cell (CD43<sup>+</sup> B220<sup>low</sup>) and pre-B cell (CD43<sup>-</sup> B220<sup>+</sup>) in BM of WT mice and  $\mu$ MT mice were detected with PE anti-mouse CD43 and PE/Cy7 anti-mouse B220 by FACS. The cells in the upper left circle were pre-B cells, and the cells in the bottom right circle were pro-B cells. (C) Mature B cell (B220<sup>+</sup> IgM<sup>+</sup>) in BM of WT mice and  $\mu$ MT mice were detected with FITC anti-mouse IgM and PE/Cy7 anti-mouse B220 by FACS. The circle showed mature B cells. (D) The concentrations of IgM and IgG in serum from WT mice and  $\mu$ MT mice were detected by ELISA. \*\*\*P < 0.001, \*\*P < 0.01, ns P > 0.05.

chain were designed specific to V $\lambda$ 1/J $\lambda$ 1, and the products were confirmed to be V $\lambda$ 1/J $\lambda$ 1 by DNA sequencing (data not shown).




**Figure 2. IgM and Ig $\kappa$  immunostaining in non B cells in multiple tissues of  $\mu$ MT mice by immunohistochemistry.** (A) IgM and Ig $\kappa$  were detected in spleen of  $\mu$ MT mice and WT mice with anti-mouse IgM or Ig $\kappa$  antibodies by immunohistochemistry. (B) IgM was detected in liver (a,e), lung (b,f), small intestine (c,g) and kidney (d,h) of  $\mu$ MT mice and WT mice with goat anti-mouse IgM antibodies by immunohistochemistry. (C) Ig $\kappa$  was detected in liver (a,d), lung (b,e) and small intestine (c,f) of  $\mu$ MT mice and WT mice with goat anti-mouse Ig $\kappa$  antibodies by immunohistochemistry. (D) IgM and Ig $\kappa$  were detected in tissues of  $\mu$ MT mice and WT mice with goat anti-mouse IgM or Ig $\kappa$  antibodies by western blot.

N/P addition in the V-D, D-J and V-J junctions is a classical feature of B cell-derived Ig, so we analyzed N/P addition in Ig transcripts derived from the liver epithelial cells. Markedly, 93% of the VDJ rearrangements had classical N/P addition, whereas only 5% of the V $\kappa$ J $\kappa$  rearrangements had N/P addition (Fig. 6B). Interestingly, we found that 86% of the V $\kappa$ J $\kappa$  rearrangements used J $\kappa$ 2 and that VDJ rearrangements frequently used JH4 and JH2 in Ig expressed by the liver epithelial cells of  $\mu$ MT mice (Fig. 6C).



**Figure 3.** IgM, IgG, IgD, IgA, Ig  $\kappa$  and Ig  $\lambda$  detection in serum and liver cells. (A) Ig expression level in serum and liver lysate of WT mice and  $\mu$ MT mice were measured by Mouse Ig Isotyping Array. WT serum was diluted 100 times, while serum from  $\mu$ MT mice was diluted 50 times. The statistical analysis was performed under the diluted condition of serum and the dilution difference was not taken into account. Igs were detected in 30  $\mu$ g liver protein from WT mice or  $\mu$ MT mice. (B) The concentrations of IgM in serum and liver lysate from WT mice and  $\mu$ MT mice were detected by ELISA. \*\*\* $P < 0.001$ , \*\* $P < 0.01$ , ns  $P > 0.05$ . (C) Liver lysate from  $\mu$ MT mice were separated by Superdex 200 and SDS-PAGE. Ig $\kappa$  was detected by goat anti-mouse Ig $\kappa$  antibody. Mass spectrometry analysis of a ~75 kDa band, which can be detected by Western blot, revealed several peptide that match mice Ig $\mu$  heavy chain protein. (D) Liver lysate from  $\mu$ MT mice were separated by Superdex 200 and SDS-PAGE. Ig $\kappa$  was detected by goat anti-mouse Ig $\kappa$  antibody. Mass spectrometry analysis of a ~26 kDa band, which can be detected by Western blot, revealed several peptide that match mice Ig $\kappa$  light chain. (E) One peptide of Ig $\lambda$  light chain protein was detected in the mass spectrometry analysis of a ~26 kDa band.



**Figure 4.** IgM and IgG were secreted from sorted liver epithelial cells. (A) CK18 was chosen as the sorting marker for liver epithelial cells. P1 represented the larger cells in liver. CK18<sup>+</sup> cells in P1 gate, which were P2 cells, were harvested. (B) Secreted IgM and IgG of liver epithelial cells sorted by CK18 from WT mice and  $\mu$ MT mice were detected by ELISPOT. WT spleen cells were used as positive control.


## Discussion

In this study, by using B cell-deficient  $\mu$ MT mice, we further provided evidence that Ig can be expressed by non-B cells under physiological conditions. We found that liver epithelial cells produced multiple classes of Ig, including IgG, IgM, IgA and IgD. These liver epithelial cell-derived Ig had distinctive characteristics in their variable regions.

In past 10 years, many studies have shown that non-B cell cancer cells, including epithelial cancer cells, and other types of malignant cells can produce Ig, especially IgG, which is involved in cell survival, proliferation, tumorigenesis and metastasis<sup>35–39</sup>. Ig expression in non-B cells was thought to be unique to malignant cells due to the genomic instability in these cells. However, in recent years, growing evidence has indicated that normal non-B cells, such as epithelial cells, endothelial cells, neurons, and even germ cells, can also express Ig. The expression of Ig in these non-B cells is of unknown significance, and its function remains poorly understood. Because these findings challenge the important classical theory that B cells are the only source of Ig, some immunologists are still questioning the possibility that non-B cells produce Ig. To address this issue, we avoided the contamination of B cells by using  $\mu$ MT mice (which have a deletion in the transmembrane domain of the Ig  $\mu$  chain, thus blocking B cell development after the pro-B stage) as a model to determine whether Ig can be produced in non-B cells.

First, we found that mature B cells are not present in peripheral blood and lymphoid organs of  $\mu$ MT mice; furthermore, these mice have very low levels (~100 times lower than WT mice) of Ig in the bloodstream. Subsequently, using a series of techniques, including immunohistochemistry, western blotting, ELISPOT, MS, RT-PCR and DNA sequencing, we found that different classes of Ig could be expressed and secreted from non-B cells (such as liver epithelial cells) at both the mRNA and protein level. Moreover, the levels of Ig in non-immune organs in  $\mu$ MT mice was similar to those in WT mice, which suggests that non-B cells can spontaneously express Ig that is retained locally, whereas the Ig in the bloodstream may be derived mainly from B cells.

Hepatocytes are epithelial cells that are involved in synthesizing protein, cholesterol, bile salts, fibrinogen, phospholipids and glycoproteins. So far, there are no reports indicating that hepatocytes can express Ig or perform Ig-related functions. However, mouse liver tissue was used as a negative control by Tonegawa (1987 Nobel laureate in Physiology for discovery of Ig gene rearrangement) when he confirmed that Ig genes are rearranged in Ig-producing cells. In 1976, he used Southern blotting to show that a mouse myeloma cell line, but not liver tissue,



**Figure 5. Ig gene rearrangement and transcription in  $\mu$ MT mice liver epithelial cells.** (A) Northern blot analysis. Total RNA from liver cells of three  $\mu$ MT mice and WT spleen cells (positive control) were transferred to nylon membrane, and probed with probes to constant region of  $Ig\mu$  and  $Ig\kappa$ ; 28 s and 18 s RNA were shown by DNA agarose gel electrophoresis. (B) *In situ* hybridization analysis was shown for detection of  $Ig\mu$  and  $Ig\kappa$  transcripts in  $\mu$ MT mice with antisense or sense probes for C region of  $\mu$  chain or J-C region of  $\kappa$  chain. (C) a It was shown for the primer design of  $Ig\mu$  variable region,  $Ig\kappa$  variable region and  $Ig\lambda$ ; b Sorted liver epithelial cells from two  $\mu$ MT mice were analyzed for Ig rearrangement and transcription by RT-PCR. WT spleen cells were used as the positive control. Variable region and constant region of heavy chain ( $\mu$ ,  $\alpha$ ,  $\delta$ ,  $\gamma$ ,  $\epsilon$ ) and light chain ( $\kappa$ ,  $\lambda$ ), CD20 and GAPDH were amplified; “1, 2” represented two samples of sorted  $\mu$ MT liver epithelial cells.

was able to bind to a probe made from the mouse myeloma cell line's mRNA fragment of an Ig variable region<sup>40–42</sup>. It was concluded that Ig gene rearrangement occurred in the myeloma cells but not in liver tissue. This conclusion further strengthened the classical concept that only B cells can produce Ig. However, in the current study, we found that Ig gene rearrangement also occurs in  $\mu$ MT mouse liver epithelial cells sorted by FACS with the specific liver epithelial cell marker CK18. In light of the current knowledge of Ig gene rearrangement, it is likely that the conclusion made by Tonegawa did not take substantial differences in Ig gene rearrangement among Ig-producing cells into account. Different conclusion could have been reached if Tonegawa had used a liver cell-derived Ig variable region mRNA fragment as a probe in his experiments. In addition, our results reveal a novel Ig source in  $\mu$ MT mice because Ig can be produced from many non-B cells, such as liver epithelial cells, but is not limited to z “a small population of B cells” as described previously<sup>32–34</sup>. In fact, we did not find any B cells in peripheral blood or mature B cells in BM in the  $\mu$ MT mice we used.

V(D)J gene rearrangement and transcription provide crucial evidence for Ig expression. We detected rearranged transcripts of Ig genes in sorted liver epithelial cells. To exclude the possible contamination of any other types of cells, we gated on the larger liver cells when CK18<sup>+</sup> liver cells were sorted. As expected, rearranged functional heavy chain and light chain transcripts of IgG, IgM, IgD and IgA, but not IgE, have been found in the sorted liver epithelial cells. Sequencing analysis revealed that similar to B cells, the liver epithelial cells derived V(D)J or VJ showed classical Ig gene rearrangement patterns and had typical N/P additions among the V, D and J segments. Moreover, liver epithelial cell-derived IgA and IgG, but not IgM and IgD, also tended to show hypermutation. However, in the same liver epithelial cell population of  $\mu$ MT mice, each Ig heavy chain, including  $\mu$ ,  $\delta$ ,  $\gamma$ ,  $\alpha$ , showed distinctive variable region sequences, but not the same VDJ sequences. This finding suggests that class switching does not occur in liver epithelial cells, unlike the classical concept of class switching when IgM switches to IgA, IgG or IgE, only the constant region (not the variable region containing the VDJ sequences) is replaced. Indeed, we also found that non-B cells lack class switch mechanisms in our previous studies<sup>6</sup>. This finding suggests that the mechanism of Ig production in non-B cells is different from that in B cells, although the mechanism remains unknown.

Based on our current and previous results, non-B cell are able to express Ig; however, its function remains unclear. Our previous studies suggested that cervical cancer cell-produced IgM have natural IgM activity that could bind to ssDNA, dsDNA and bacteria<sup>22</sup>. Jiang *et al.* found that normal epidermal cells can express and secrete IgG and IgA, which can bind to different strains of bacteria<sup>9</sup>. We have preliminary results showing that IgM extracted from liver tissue of both WT and  $\mu$ MT mice possesses natural IgM activity (data not shown). This

	Ig $\mu$			Ig $\alpha$			Ig $\gamma$			Ig $\delta$			Ig $\kappa$		
	V <sub>H</sub> DJ <sub>H</sub> recombination	No. of clones	Identity with germline (%)	V <sub>H</sub> DJ <sub>H</sub> recombination	No. of clones	Identity with germline (%)	V <sub>H</sub> DJ <sub>H</sub> recombination	No. of clones	Identity with germline (%)	V <sub>H</sub> DJ <sub>H</sub> recombination	No. of clones	Identity with germline (%)	VJ recombination	No. of clones	Identity with germline (%)
No. 1	IGHV1-55*01/ IGHD2-1*01/ IGHJ4*01	7/8	97	IGHV1-80*01/ IGHD2-4*01/ IGHJ4*01	5/9	97.6	IGHV1-53*01/ IGHD2-5*01/ IGHJ2*01	5/7	99.6	IGHV17*01/ IGHD2-4*01/ IGHJ4*01	3/8	99.6	IGKV14-111*01/ IGKJ2*01	7/8	96
				IGHV1-72*01/ IGHD2-1*01/ IGHJ2*01	2/9	97.3				IGHV1-39*01/ IGHD2-3*01/ IGHJ2*01	2/8	100			
	IGHV1-37*01/ IGHD2-13*01/ IGHJ2*01	1/8	98.6	IGHV1-64*01/ IGHD2-9*01/ IGHJ2*01	1/9	98	IGHV1-39*01/ IGHD2-1*01/ IGHJ3*01	2/7	95.1	IGHV1-55*01/ IGHD1-1*01/ IGHJ3*01	1/8	100	IGKV9-120*01/ IGKJ2*01	1/8	100
				IGHV1-7*02/ IGHD2-1*01/ IGHJ2*01	1/9	99.6				IGHV1-18*01/ IGHD2-3*01/ IGHJ4*01	1/8	99.3			
No. 2	IGHV14-1*01/ IGHD3-2*01/ IGHJ2*01	4/6	98.6	IGHV1-82*01/ IGHD3-3*01/ IGHJ2*01	3/6	97	IGHV1-69*02/ IGHD1-1*01/ IGHJ1*03	5/7	97	IGHV1-7*01/ IGHD2-9*01/ IGHJ4*01	3/8	100	IGKV9-120*01/ IGKJ2*01	5/8	97.5
				IGHV1-81*01/ IGHD5-8*01/ IGHJ4*01	1/6	97				IGHV1-76*01/ IGHD2-4*01/ IGHJ4*01	1/8	100			
	IGHV1-82*01/ IGHD1-1*01/ IGHJ1*03	1/6	95.2	IGHV1-82*01/ IGHD2-4*01/ IGHJ3*01	1/6	97.6	IGHV1-78*01/ IGHD2-3*01/ IGHJ3*01	2/7	97.9	IGHV1-39*01/ IGHD1-1*01/ IGHJ4*01	1/8	100	IGKV9-120*01/ IGKJ2*01	2/8	99.6
				IGHV1-18*01/ IGHD2-5*01/ IGHJ2*01	1/6	96.3				IGHV1-20*01/ IGHD2-3*01/ IGHJ3*01	1/8	99.3			
IGHV1-55*01/ IGHD1-2*01/ IGHJ4*01	1/6	97							IGHV1-4*01/ IGHD2-1*01/ IGHJ4*01	1/8	100	IGKV14-111*01/ IGKJ2*01	1/8	98.9	
									IGHV1-4*01/ IGHD2-9*01/ IGHJ4*01	1/8	99.3				
No. 3	IGHV6-6*01/ IGHD1-2*01/ IGHJ4*01	5/8	99.7										IGKV9-120*01/ IGKJ5*01	2/4	99.6
	IGH-V1S16*01/ IGHD2-10*02/ IGHJ2*01	1/8	96										IGKV9-120*01/ IGKJ2*01	1/4	99.6
	IGHV1-47*01/ IGHD1-1*01/ IGHJ2*01	1/8	100										IGKV9-120*01/ IGKJ4*02	1/4	99.6
	IGHV1-7*01/ IGHD4-1*02/ IGHJ2*01	1/8	100												
No. 4	IGHV1-39*01/ IGHD2-3*01/ IGHJ4*01	5/9	100										IGKV17-121*01/ IGKJ2*01	6/8	99.6
	IGHV14-3*02/ IGHD5-8*01/ IGHJ3*01	4/9	100										IGKV17-121*01/ IGKJ5*01	2/8	99.3
No. 5	IGHV1-67*01/ IGHD2-3*01/ IGHJ2*01	5/5	96										IGKV4-50*01/ IGKJ2*01	5/7	99.6
														IGKV4-50*01/ IGKJ5*01	2/7

**Table 1. VDJ and VJ recombinations in liver epithelial cells of  $\mu$ MT mice.**

finding suggests that Ig from non-B cells, especially epithelial cells, may serve natural antibody functions and may be involved in natural immunity in local tissues. Our focus will be on the function of these Ig.



WT-Spleen	No. of clones	Recombination	V segment identity with germline (%)
V $\mu$ DJ $\mu$ recombination	2/8	IGHV1-80*01/IGHD2-4*01/IGHJ3*01	100
	1/8	IGHV1-47*01/IGHD2-5*01/IGHJ4*01	97.6
	1/8	IGHV1-82*01/IGHD2-9*01/IGHJ1*03	100
	1/8	IGHV1-31*01/IGHD2-3*01/IGHJ3*01	99.2
	1/8	IGHV1-7*01/IGHD2-5*01/IGHJ2*01	99.2
	1/8	IGHV1-78*01/IGHD2-14*01/IGHJ1*03	98.8
	1/8	IGHV1-76*01/IGHD2-3*01/IGHJ2*01	100
V $\kappa$ J $\kappa$ recombination	1/7	IGKV9-120*01/IGKJ2*01	99.6
	1/7	IGKV9-120*01/IGKJ2*01	99.3
	1/7	IGKV14-111*01/IGKJ2*01	100
	1/7	IGKV14-126*01/IGKJ2*01	100
	1/7	IGKV17-127*01/IGKJ2*01	100
	1/7	IGKV17-127*01/IGKJ2*01	97.9
	1/7	IGKV17-127*01/IGKJ5*01	100

**Table 2.** V(D)J recombination of Ig gene in WT spleen cells.

## Materials and Methods

**Mice.** BALB/c mice were purchased from Charles River Laboratories (Beijing, China).  $\mu$ MT/BALB/c mice were housed in Institute of Biophysics, Chinese Academy of Science. All these mice used in this study were housed in a specific pathogen free environment at the Peking University Health Science Center. And procedures about animal studies were performed in accordance with the guidelines of the People's Republic of China Ministry of Health and approved by the Animal Care and Use Committee of Peking University Health Science Center.

**Immunohistochemistry.** At first, tissues were sectioned and fixed in 10% formalin, and then was embedded in paraffin. The paraffin sections were rehydrated by ethanol with the concentrations of 75%, 80%, 90%, 95% and 100% after deparaffinization. Antigen retrieval was performed in tris-EDTA buffer (pH 9.0) at 90 °C for 5-min in a microwave oven. To block the endogenous peroxidase activity, the slide was treated with 0.3% hydrogen peroxide for 5 min, and washed in PBS. Then, the sections were blocked with 10% normal goat serum for 10 min. Slides were incubated with primary antibody, goat anti-mouse IgM (VECTOR LAB, CA, USA) or goat anti-mouse Ig $\kappa$  (Southern Biotech, UBA, USA), for 60 min at 37 °C in a humidified chamber. After washed, the sections were incubated with the HRP-conjugated second antibodies (rabbit anti-goat IgG) at 37 °C for 40 min. After washing with PBS, the signal was detected using DAB (Dako, CA, USA). Sections stained without primary antibody were used as negative controls.


**Protein extraction.** Heart perfusion was performed to the mice anaesthetized by chloral hydrate before sacrifice. The tissues were cut into small pieces and lysed in TSD buffer (50 mM Tris-HCL, 0.5% sodium dodecyl sulfate, 5 mM DTT) followed by ultrasonication. Lysate were centrifuged at 13,000 rpm for 20 min at 4 °C, and the supernatants were collected for ELISA or Western blot.

**Rapid Mouse Ig Isotyping Array.** As mentioned above, tissues lysate was prepared. Serum from mice was diluted for 100 times. Then the Ig was detected according to the instructions of Rapid Mouse Ig Isotyping Array (Ray Biotech, Norcross GA, USA).

**ELISA.** To detect the concentration of IgM and IgG in liver lysate of mice liver and serum, we used Mouse IgM ELISA Quantitation Set and Mouse IgG ELISA Quantitation Set (Bethyl Laboratories, USA). 80  $\mu$ g liver lysate or 500 times diluted serum was performed ELISA analysis by following the standard protocol. OD450 was measured with a microplate reader (Bio-Tek, Winooski, VT).

**SDS-PAGE, Western Blot and MS.** The reduced (with  $\beta$ -mercaptoethanol) tissue lysate samples were separated by SDS-PAGE. For Western blot, the separate proteins were transferred to nitrocellulose membranes. After blocked with tris-buffered saline containing and 5% defatted milk for 2 h at room temperature, membranes were incubated with the appropriate primary antibodies: Biotin-conjugated goat anti-mouse IgM ( $\mu$  chain specific) antibody (VECTOR LAB, CA, USA) and goat anti-mouse Ig $\kappa$  antibody (Southern Biotech, UBA, USA) overnight at 4 °C. Following the wash with TBS, membranes were incubated with the appropriate IRDye 800 or IRDye 700-conjugated secondary antibodies (LI-COR Bioscience Inc., Lincoln, NE, USA) in the dark. The Odyssey Imaging System (LI-COR Bioscience Inc., Lincoln, NE, USA) was used to detect the signals. For MS, the SDS-PAGE bands at specific molecular weight were cut to perform MS analysis.

**Isolation of mice liver epithelial cells.** Heart perfusion was performed to the mice anaesthetized by chloral hydrate before sacrifice. The livers were cut into little pieces and digested with 150 U/ml DNase I (Sigma, St. Louis, Mo, USA) and 500 U/ml collagenase IV (Sigma, St. Louis, Mo, USA) at 37 °C. Then the liver single-cell



**Figure 6. Characters of liver epithelial cell-derived Ig transcripts in μMT mice. (A)** The mutation points of VDJ recombinases and VJ recombinases were shown, which were preponderant rearrangement pattern in μMT mice. “.” represents the matched nucleotides of liver epithelial cell-derived Ig transcripts compared with the Ig germline genes. “?” represents the missing nucleotides of junctions compared with the Ig germline genes. Each point represents one nucleotide. **(C)** The frequency of JH and Jκ used by Ig derived from liver epithelial cells of μMT mice.

suspensions was obtained after filtering by mesh. The liver cells were stained with rabbit anti-mouse CK 18 (Santa Cruz, San Diego, CA, USA), and then were further labeled with secondary antibody (FITC-conjugated anti-rabbit IgG). The CK18<sup>+</sup> liver epithelial cells were sorted by FACSaria™ II Flow Cytometer (Becton Dickinson, San Diego, CA, USA). The sorted liver epithelial cells were used for RT-PCR, ELISPOT and WB analysis.

**Flow cytometry.** To analyze B cells in mice, blood cells and bone marrow cells were harvest and washed with PBS. Before staining, cells were blocked with 5% FBS in PBS for 30 min at 4 °C. And then cells were stained with membrane marker antibodies: PE-CD43, FITC-IgM, PE-B220 and PE/cy7-B220 purchased from eBioscience (USA). The control was performed that cells were stained with corresponding fluorescein conjugated isotype control antibody.

**ELISPOT.** To detect the secretion of IgG and IgM in liver epithelial cells, ELISPOT was performed. 1 × 10<sup>5</sup> sorted liver epithelial cells or 1 × 10<sup>5</sup> spleen cells were used for the IgM or IgG ELISPOT. MultiScreen filter plates for ELISPOT were coated with goat anti-mouse IgM (Bethyl Laboratories, USA) or goat anti-mouse IgG (Bethyl

Laboratories, USA) for overnight at 4 °C. Following the blocking with 10% FBS, cells were incubated with plates for 24 h at 37 °C. After washed with PBS, HRP conjugated detection antibody were incubated for 1 h at room temperature. Then AEC coloration was performed and analyzed by ELISPOT reader.

**Northern blot analysis.** Total RNA was extracted from liver or spleen cells with TRIzol reagent. A total of 10 µg liver-derived RNA or 5 µg spleen-derived RNA was separated in 1% agarose gel with formaldehyde, and then transferred to a nylon membrane. The sequence of probe specific to I $\mu$  constant region gene was: ttcatctctg-cgacagctggaatgggacacat, and the probe specific to I $\kappa$  constant region was: cgccatttgtcttcactgcatcaatcttc. The probes were conjugated with biotin. The membrane was hybridized with 60 ng/ml denatured probe in hybridization buffer (Thermo) after prehybridization at 56 °C for 12–16 h. And the signal was detected with North2South<sup>®</sup> Chemiluminescent Hybridization and Detection Kit (Thermo) according to the manufacturer's instructions.

**In situ hybridization.** *In situ* hybridization was performed on 6 µm serial sections of paraffin-embedded tissue sections. Plasmids inserted with constant region fragments of the I $\mu$  and I $\kappa$ , which were obtained by PCR with I $\mu$  CH3-CH4 primers and Jk-Ck primers (Table S1), were linearized. The RNA probes were labeled with digoxigenin (DIG) and transcribed by T7 RNA polymerase (for the antisense probe) or SP6 RNA polymerase (for the sense probe) by DIG northern starter kit (Roche, Rotkreuz, Switzerland). Paraffin-embedded tissue sections were performed deparaffinized and dehydrate. Then sections were treated with Proteinase K, fixed with paraformaldehyde, prehybridized at 42 °C for 2 h, and hybridized with the DIG-labeled RNA probe (5 µg/mL) at 42 °C overnight. After hybridization, the sections were washed in 2 × SSC and 0.1 × SSC at 37 °C, respectively, and then treated with RNase A. The samples were incubated with alkaline phosphatase-conjugated antidigoxigenin antibody (1:250; Roche, Rotkreuz, Switzerland). BCIP/NBT (Sigma, Saint Louis, USA) was used for the color reaction. Corresponding sense probes were used as controls.

**RNA extraction and RT-PCR.** Total RNA in spleen was extracted using TRIzol reagent (Invitrogen, Carlsbad, USA). For the sorted liver epithelial cells, RNA was extracted with RNeasy Mini Kit (Qiagen, Hilden, Germany) according to the standard protocol. Reverse transcription (RT) was carried out with RevertAid First Strand cDNA Synthesis Kit (Fermentas, Glen Burnie, USA) by following the manufacturer's instructions. And then the cDNA was prepared for PCR.

The primers used in this study were shown in Table S1. The Ig constant region genes were amplified at the annealing temperature of 56 °C for 35 cycles. Semi-nested PCR was performed for Ig V region genes. cDNA derived from BALB/c mice spleen cells was used as positive control. PCR products were separated with 1% agarose gel by electrophoresis, and then were stained with ethidium bromide. The PCR products were cloned to pGEM-T Easy Vector (Promega) to be sequenced by ABI 3730XL Genetic Analyzer.

**Sequence analysis.** In order to analyze the usage and junctions of Ig variable region genes, all sequence derived from PCR products were submitted to the IMGT V-QUEST program. Alignments were performed with Lasergene software (DNASar) or with BLAST in NCBI to compare with published sequences.

**Statistical analysis.** All statistical calculations were performed using the statistical software program (GraphPad Prism 5.0 software). Differences between different groups were evaluated by the Student's *t* test. Differences were considered to be statistically significance when *P* was <0.05.

## References

1. Qiu, X. *et al.* Human epithelial cancers secrete immunoglobulin g with unidentified specificity to promote growth and survival of tumor cells. *Cancer Res* **63**, 6488–6495 (2003).
2. Deng, Y. Q. *et al.* [Immunoglobulin expression in colon cancer cell line HT-29 and its biological activities]. *Zhonghua Zhong Liu Za Zhi* **28**, 88–91 (2006).
3. Zhu, X. *et al.* Immunoglobulin mRNA and protein expression in human oral epithelial tumor cells. *Applied immunohistochemistry & molecular morphology: AIMM/official publication of the Society for Applied Immunohistochemistry* **16**, 232–238 (2008).
4. Huang, J., Zhang, L., Ma, T., Zhang, P. & Qiu, X. Expression of immunoglobulin gene with classical V-(D)-J rearrangement in mouse testis and epididymis. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society* **57**, 339–349 (2009).
5. Huang, J. *et al.* Expression of immunoglobulin gene with classical V-(D)-J rearrangement in mouse brain neurons. *Int J Biochem Cell Biol* **40**, 1604–1615 (2008).
6. Zheng, J. *et al.* Immunoglobulin gene transcripts have distinct VHDJH recombination characteristics in human epithelial cancer cells. *J Biol Chem* **284**, 13610–13619 (2009).
7. Zhang, S. *et al.* Immunoglobulin gene locus events in epithelial cells of lactating mouse mammary glands. *Cell Mol Life Sci* **67**, 985–994 (2010).
8. Huang, J. *et al.* Rearrangement and expression of the immunoglobulin mu-chain gene in human myeloid cells. *Cell Mol Immunol* **11**, 94–104 (2014).
9. Jiang, D. *et al.* IgG and IgA with potential microbial-binding activity are expressed by normal human skin epidermal cells. *Int J Mol Sci* **16**, 2574–2590 (2015).
10. Zheng, H. *et al.* Immunoglobulin alpha heavy chain derived from human epithelial cancer cells promotes the access of S phase and growth of cancer cells. *Cell biology international* **31**, 82–87 (2007).
11. Kimoto, Y. Expression of heavy-chain constant region of immunoglobulin and T-cell receptor gene transcripts in human non-hematopoietic tumor cell lines. *Genes Chromosomes Cancer* **22**, 83–86 (1998).
12. Li, J. *et al.* Proteomic detection of changes in protein synthesis induced by NGX6 transfected in human nasopharyngeal carcinoma cells. *Journal of protein chemistry* **20**, 265–271 (2001).
13. Li, M. *et al.* Expression of immunoglobulin kappa light chain constant region in abnormal human cervical epithelial cells. *Int J Biochem Cell Biol* **36**, 2250–2257 (2004).
14. Babbage, G., Ottensmeier, C. H., Blaydes, J., Stevenson, F. K. & Sahota, S. S. Immunoglobulin heavy chain locus events and expression of activation-induced cytidine deaminase in epithelial breast cancer cell lines. *Cancer Res* **66**, 3996–4000 (2006).

15. Chen, Z. & Gu, J. Immunoglobulin G expression in carcinomas and cancer cell lines. *FASEB J* **21**, 2931–2938 (2007).
16. Liu, H. D. *et al.* Upregulated expression of kappa light chain by Epstein-Barr virus encoded latent membrane protein 1 in nasopharyngeal carcinoma cells via NF- $\kappa$ B and AP-1 pathways. *Cell Signal* **19**, 419–427 (2007).
17. Zheng, H. *et al.* Expression and secretion of immunoglobulin alpha heavy chain with diverse VDJ recombinations by human epithelial cancer cells. *Mol Immunol* **44**, 2221–2227 (2007).
18. Lee, G., Laflamme, E., Chien, C. H. & Ting, H. H. Molecular identity of a pan cancer marker, CA215. *Cancer Biol Ther* **7**, 2007–2014 (2008).
19. Lee, G. Cancer cell-expressed immunoglobulins: CA215 as a pan cancer marker and its diagnostic applications. *Cancer Biomark* **5**, 137–142 (2009).
20. Qiu, X. *et al.* Immunoglobulin gamma heavy chain gene with somatic hypermutation is frequently expressed in acute myeloid leukemia. *Leukemia* **27**, 92–99 (2013).
21. Zhao, Y., Liu, Y., Chen, Z., Korteweg, C. & Gu, J. Immunoglobulin g (IgG) expression in human umbilical cord endothelial cells. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society* **59**, 474–488 (2011).
22. Hu, F. *et al.* Spontaneous production of immunoglobulin M in human epithelial cancer cells. *PLoS One* **7**, e51423 (2012).
23. Nemazee, D. Receptor selection in B and T lymphocytes. *Annual review of immunology* **18**, 19–51 (2000).
24. Kitamura, D., Roes, J., Kuhn, R. & Rajewsky, K. A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin mu chain gene. *Nature* **350**, 423–426 (1991).
25. Qin, Z. *et al.* B cells inhibit induction of T cell-dependent tumor immunity. *Nat Med* **4**, 627–630 (1998).
26. Smelt, S. C., Cotterell, S. E., Engwerda, C. R. & Kaye, P. M. B cell-deficient mice are highly resistant to Leishmania donovani infection, but develop neutrophil-mediated tissue pathology. *J Immunol* **164**, 3681–3688 (2000).
27. Dittel, B. N., Urbana, T. H. & Janeway, C. A. Jr. Relapsing and remitting experimental autoimmune encephalomyelitis in B cell deficient mice. *J Autoimmun* **14**, 311–318 (2000).
28. Li, H. *et al.* Experimental autoimmune myasthenia gravis induction in B cell-deficient mice. *Int Immunol* **10**, 1359–1365 (1998).
29. Homann, D. *et al.* Evidence for an underlying CD4 helper and CD8 T-cell defect in B-cell-deficient mice: failure to clear persistent virus infection after adoptive immunotherapy with virus-specific memory cells from muMT/muMT mice. *J Virol* **72**, 9208–9216 (1998).
30. Langhorne, J., Cross, C., Seixas, E., Li, C. & von der Weid, T. A role for B cells in the development of T cell helper function in a malaria infection in mice. *Proc Natl Acad Sci USA* **95**, 1730–1734 (1998).
31. Macpherson, A. J. *et al.* IgA production without mu or delta chain expression in developing B cells. *Nat Immunol* **2**, 625–631 (2001).
32. Lagerquist, M. K. *et al.* 17Beta-estradiol expands IgA-producing B cells in mice deficient for the mu chain. *Scand J Immunol* **67**, 12–17 (2008).
33. Orinska, Z. *et al.* Novel B cell population producing functional IgG in the absence of membrane IgM expression. *Eur J Immunol* **32**, 3472–3480 (2002).
34. Ghosh, S., Hoselton, S. A. & Schuh, J. M. mu-chain-deficient mice possess B-1 cells and produce IgG and IgE, but not IgA, following systemic sensitization and inhalational challenge in a fungal asthma model. *J Immunol* **189**, 1322–1329 (2012).
35. Qiu, Y. *et al.* Immunoglobulin G expression and its colocalization with complement proteins in papillary thyroid cancer. *Mod Pathol* **25**, 36–45 (2012).
36. Ma, C. *et al.* Immunoglobulin G expression and its potential role in primary and metastatic breast cancers. *Current molecular medicine* **13**, 429–437 (2013).
37. Niu, N. *et al.* IgG expression in human colorectal cancer and its relationship to cancer cell behaviors. *PLoS One* **7**, e47362 (2012).
38. Celis, J. E. *et al.* Proteomic characterization of the interstitial fluid perfusing the breast tumor microenvironment: a novel resource for biomarker and therapeutic target discovery. *Molecular & cellular proteomics: MCP* **3**, 327–344 (2004).
39. Liu, Y. *et al.* Binding of the mAb RP215 to IgG in metastatic lung adenocarcinomas is correlated with poor prognosis. *Histopathology* **67**, 645–653 (2015).
40. Hozumi, N. & Tonegawa, S. Evidence for somatic rearrangement of immunoglobulin genes coding for variable and constant regions. 1976 [classical article]. *J Immunol* **173**, 4260–4264 (2004).
41. Brack, C. & Tonegawa, S. Variable and constant parts of the immunoglobulin light chain gene of a mouse myeloma cell are 1250 nontranslated bases apart. *Proc Natl Acad Sci USA* **74**, 5652–5656 (1977).
42. Bernard, O., Hozumi, N. & Tonegawa, S. Sequences of mouse immunoglobulin light chain genes before and after somatic changes. *Cell* **15**, 1133–1144 (1978).

## Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (No. 81272237, No. 91229102, No. 81320108020).

## Author Contributions

W.S. and X.Q. designed the experiment; W.S., J.M., Z.Z., X.G. and Z.Q. led the sample collection; Z.Q. supplied mice used in this study; W.S., E.L., L.Z. and C.Z. did the other experiments; W.S. and L.Z. carried out the data analysis; W.S., Z.Q. and X.Q. wrote the paper; all authors reviewed the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Shao, W. *et al.* Identification of Liver Epithelial Cell-derived Ig Expression in  $\mu$  chain-deficient mice. *Sci. Rep.* **6**, 23669; doi: 10.1038/srep23669 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>