

Nicotinamide phosphoribosyltransferase inhibits receptor activator of nuclear factor- κ B ligand-induced osteoclast differentiation *in vitro*

JONG MIN BAEK^{1*}, SUNG-JUN AHN^{1*}, YOON-HEE CHEON^{1,2},
MYEUNG SU LEE³⁻⁵, JAEMIN OH^{1,4,5} and JU-YOUNG KIM⁴

¹Department of Anatomy, School of Medicine; ²Center for Metabolic Function Regulation; ³Division of Rheumatology, Department of Internal Medicine; ⁴Imaging Science-Based Lung and Bone Diseases Research Center; ⁵Institute for Skeletal Disease, Wonkwang University, Iksan, Jeonbuk 570-749, Republic of Korea

Received November 19, 2015; Accepted November 22, 2016

DOI: 10.3892/mmr.2016.6069

Abstract. The adipokine nicotinamide phosphoribosyltransferase (*Nampt*), also known as pre-B-cell colony-enhancing factor or the insulin-mimetic hormone visfatin, has a crucial role in the conversion of nicotinamide to nicotinamide mononucleotide during biosynthesis of the coenzyme nicotinamide adenine dinucleotide. Previous reports have demonstrated the inhibitory effects of *Nampt* on osteoclast formation from human peripheral blood mononuclear cells and CD14⁺ monocytes. However, the effects of *Nampt* on bone marrow macrophage (BMM)-derived osteoclastogenesis and its precise role in the process remain unclear. The present *in vitro* study used recombinant *Nampt* and BMMs as osteoclast precursors demonstrated that *Nampt* suppresses receptor activator of nuclear factor- κ B ligand (RANKL)-induced osteoclastogenesis by decreasing the phosphorylation of various early signal transducers, including c-Jun N-terminal kinase, Akt, glycogen synthase kinase-3 β , Bruton's tyrosine kinase and phospholipase C γ -2. In addition, western blotting and reverse transcription-quantitative polymerase chain reaction analysis indicated that *Nampt* downregulates the mRNA and protein expression levels of c-Fos and nuclear factor of

activated T cells, cytoplasmic 1, leading to a decrease in the expression of osteoclast-specific genes including tartrate-resistant acid phosphatase, osteoclast-associated receptor and cathepsin K. However, the bone-resorbing activity of mature osteoclasts treated with *Nampt* was similar to untreated control osteoclasts. This finding indicates that *Nampt* exerts its anti-osteoclastogenic activity by targeting osteoclast precursor cells rather than mature osteoclasts. Consequently, the present study demonstrated that *Nampt* acts as a negative regulator of RANKL-mediated differentiation of BMMs into osteoclasts, suggesting the potential therapeutic targets to treat bone-related disorders such as osteoporosis.

Introduction

Nicotinamide phosphoribosyltransferase (*Nampt*) is a novel adipokine, which has been reported to be expressed in adipose tissue, chondrocytes in the articular cartilage matrix and peripheral blood mononuclear cells (PBMCs) (1-4). It has been revealed that *Nampt* is closely associated with various biological processes, including nicotinamide adenine dinucleotide (NAD) biosynthesis, cellular metabolism and immunomodulatory responses. In the process of NAD biosynthesis, *Nampt* regulates the activity of the NAD-dependent deacetylase silent information regulator 2 (Sir2) through increasing the cellular level of NAD, and subsequently promoting Sir2 transcriptional activity in mammalian cells (5). The potent *Nampt* inhibitor FK866 negatively regulates glycolysis by altering the initial steps in glucose oxidation and leads to changes in carbohydrate metabolism in cancer cells (6). Furthermore, *Nampt* is an essential catabolic mediator of osteoarthritis, which is the most common form of inflammatory arthritis, and regulates hypoxia-inducible factor 2 α -mediated matrix metalloproteinase (*MMP*) expression in chondrocytes, leading to the destruction of osteoarthritic cartilage (7).

For the treatment of metabolic bone diseases, including osteoporosis, adipokines are considered to be a therapeutic target via their effects on two types of bone cell, osteoclasts and osteoblasts (8). Osteoclasts are well-characterized cells that are

Correspondence to: Dr Jaemin Oh, Department of Anatomy, School of Medicine, Wonkwang University, 344-2 Sinyong-dong, Iksan, Jeonbuk 570-749, Republic of Korea
E-mail: jmoh@wku.ac.kr

Dr Ju-Young Kim, Imaging Science-Based Lung and Bone Diseases Research Center, Wonkwang University, 344-2 Sinyong-dong, Iksan, Jeonbuk 570-749, Republic of Korea
E-mail: kimjy1014@gmail.com

*Contributed equally

Key words: nicotinamide phosphoribosyltransferase, adipokine, osteoclast, osteoporosis

required for bone resorption and excessive osteoclast differentiation is a predominant indicator of osteoporosis. Osteoblasts are responsible for bone formation. A previous study indicated that osteoblast proliferation and differentiation are enhanced *in vitro*, and that acceleration of bone formation and mineral apposition rates are observed *in vivo* in the absence of the adipokine apelin, which is a ligand of the Gi-G protein-coupled receptor APJ. These data suggest a crucial role of apelin in bone homeostasis as a physiological antianabolic factor (9). Another adipokine, visceral adipose tissue-derived serine protease inhibitor (*vaspin*), has been reported to suppress receptor activator of nuclear factor- κ B ligand (RANKL)-mediated differentiation of RAW264.7 cells and bone marrow cells (BMCs) into mature osteoclasts by reducing the expression of nuclear factor of activated T cells, cytoplasmic 1 (*NFATc1*) and the subsequent induction of osteoclast-specific gene markers, such as *MMP-9* and cathepsin K (10). Adiponectin is an important adipokine that regulates energy homeostasis, which also inhibits RANKL-induced osteoclastogenesis by decreasing the expression of several osteoclastogenic factors, including *NFATc1*, tumor necrosis factor receptor-associated factor 6, cathepsin K and tartrate-resistant acid phosphatase (*TRAP*), and induces apoptosis in mature osteoclasts (11). It has previously been demonstrated that *Nampt* attenuates osteoclast differentiation derived from PBMCs in patients with multiple myeloma and human CD14⁺ monocytes; however, the role of *Nampt* in the differentiation of murine bone marrow macrophages (BMMs) into osteoclasts and its underlying mechanisms have not yet been revealed (12,13).

The present study investigated the effects of *Nampt* on RANKL-mediated osteoclast differentiation and functional bone-resorbing activity. In addition, the present study determined whether *Nampt* is involved in RANKL-dependent intracellular signaling pathways and the expression of osteoclast-specific gene markers.

Materials and methods

Preparation of *Nampt* and reagents. Recombinant mouse *Nampt* (visfatin/pre-B-cell colony-enhancing factor) was purchased from Adipogen International, Inc. (San Diego, CA, USA). Recombinant soluble human macrophage colony-stimulating factor (M-CSF) and human RANKL were obtained from PeproTech EC Ltd. (London, UK). Anti-p38 (cat. no. 9212), anti-phosphorylated (p)-p38 (cat. no. 9211), anti-extracellular signal-regulated protein kinases (ERK) 1/2 (cat. no. 9102), anti-p-ERK 1/2 (cat. no. 9101), anti-c-Jun N-terminal kinase (JNK; cat. no. 9252), anti-p-JNK (cat. no. 9251), anti-Akt (cat. no. 9272), anti-p-Akt (cat. no. 9271), anti-glycogen synthase kinase-3 β (GSK3 β ; cat. no. 9315), anti-p-GSK3 β (cat. no. 9323) and anti-Bruton's tyrosine kinase (Btk; cat. no. 3533) antibodies were purchased from Cell Signaling Technology, Inc. (Beverly, MA, USA). Anti-c-Fos (cat. no. sc-7202), anti-NFATc1 (cat. no. sc-7294), anti-phospholipase C γ -2 (PLC γ 2; cat. no. sc-5283) and anti-p-PLC γ 2 (cat. no. sc-101785) antibodies were purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA). Anti-p-Btk (cat. no. GTX61792) and monoclonal anti- β -actin (cat. no. GTX109639) antibodies were obtained from GeneTex, Inc. (Irvine, CA, USA) and Sigma-Aldrich (Merck

Millipore; Darmstadt, Germany), respectively. Fetal bovine serum (FBS), α -minimum essential medium (α -MEM) and penicillin/streptomycin were purchased from Gibco (Thermo Fisher Scientific, Inc., Waltham, MA, USA). All other chemicals were of analytical grade or complied with the standards required for cell culture experiments.

Mouse BMM preparation and osteoclast differentiation. A total of 10 male ICR strain mice (age, 5 weeks; weight, 30 \pm 2 g) were purchased from Samtako (Osan, Korea). During the experimental period, the mice were maintained in a temperature- and humidity-controlled environment at 22–24°C and 55–60% humidity, with a 12-h light/dark cycle and access to sterilized water and standard rodent chow (Samtako) *ad libitum*. All experiments were conducted according to the guidelines of the Institutional Animal Care and Use Committee of Wonkwang University (WKU-14-23; Iksan, Korea). BMMs from mice were cultured as described previously (14). Briefly, to obtain BMMs, BMCs were cultured in α -MEM supplemented with 10% FBS and M-CSF (10 ng/ml) for 1 day. Non-adherent cells were further cultured in the presence of M-CSF (30 ng/ml) for 3 days. Subsequently, the adherent cells were used as BMMs. BMMs were cultured in 48-well plates at 37°C in 5% CO₂ for 4 days in the condition of M-CSF (30 ng/ml) and RANKL (100 ng/ml), and pretreated with *Nampt* (100, 250 or 500 ng/ml). The cells were fixed in 3.7% formalin, permeabilized with 0.1% Triton X-100, and stained with TRAP solution. The stained multinucleated cells (MNCs) with >5 nuclei were counted to determine the level of osteoclast differentiation.

Cell viability assay, western blotting, reverse transcription-quantitative polymerase chain reaction (RT-qPCR) analysis and bone resorption assay. The XTT cell viability assay, western blot analysis, RT-qPCR analysis and the resorption pit assay were performed as described previously (14). Resorption pits were imaged and analyzed using Image Pro-Plus version 4.5 (Media Cybernetics, Inc., Rockville, MD, USA). Primers used for PCR are summarized in Table I. The western blots were analyzed using ImageJ (imagej.nih.gov/).

Retroviral gene transfection. Packaging of the retroviral vectors pMX-IRES-EGFP, pMX-cFos-IRES-EGFP and pMX-NFATc1-IRES-EGFP was performed using transient transfection of these pMX vectors (Cell Biolabs, Inc., San Diego, CA, USA) into platinum-E (plat-E) retroviral packaging cells (Cell Biolabs, Inc.) using X-tremeGENE 9 (Roche, Nutley, NJ, USA) according to the manufacturer's protocol. Following incubation at 37°C in fresh medium for 2 days, the culture supernatants of the retrovirus-producing cells were collected. For retroviral infection, non-adherent BMCs were cultured in M-CSF (30 ng/ml) for 2 days. The BMMs were incubated with viral supernatant medium of pMX-IRES-EGFP, pMX-cFos-IRES-EGFP and pMX-NFATc1-IRES-EGFP virus-producing plat-E cells together with polybrene (10 ng/ml) and M-CSF (30 ng/ml) for 6 h. The infection efficiency of the retrovirus was determined by green fluorescent protein (GFP) expression and was always >80%. Post-infection, the BMMs were induced to differentiate in the presence of M-CSF (30 ng/ml) and RANKL (100 ng/ml) for 4 days. The expression of each construct was detected using a fluorescence

Table I. Primer sequences used for reverse transcription-quantitative polymerase chain reaction analysis.

Gene	Primer sequence (5'-3')
<i>GAPDH</i>	
Forward	5'-TCAAGAAGGTGGTGAAGCAG-3'
Reverse	5'-AGTGGGAGTTGCTGTTGAAGT-3'
<i>c-Fos</i>	
Forward	5'-GGTGAAGACCGTGTCCAGGAG-3'
Reverse	5'-TATTCGGTTCCTTCGGATT-3'
<i>NFATc1</i>	
Forward	5'-GAGTACACCTTCCAGCACCTT-3'
Reverse	5'-TATGATGTCGGGAAAGAGA-3'
<i>TRAP</i>	
Forward	5'-TCATGGGTGGTGTCTGCT-3'
Reverse	5'-GCCACAGCCACAAATCT-3'
<i>OSCAR</i>	
Forward	5'-GGAATGGTCCTCATCTCCTT-3'
Reverse	5'-TCCAGGCAGTCTCTCAGTTT-3'
<i>DC-STAMP</i>	
Forward	5'-TCCTCCATGAACAAACAGTTCCA-3'
Reverse	5'-AGACGTGGTTTAGGAATGCAGCTC-3'
<i>Atp6vOd2</i>	
Forward	5'-GACCCTGTGGCACTTTTTGT-3'
Reverse	5'-GTGTTTGAGCTTGGGGAGAA-3'
<i>Cathepsin K</i>	
Forward	5'-CCAGTGGGAGCTATGGAAGA-3'
Reverse	5'-CTCCAGGTTATGGGCAGAGA-3'
α v-integrin	
Forward	5'-ACAAGCTCACTCCCATCACC-3'
Reverse	5'-ATATGAGCCTGCCGACTGAC-3'
β 3-integrin	
Forward	5'-GGAGTGGCTGATCCAGATGT-3'
Reverse	5'-TCTGACCATCTTCCCTGTCC-3'
<i>CTR</i>	
Forward	5'-TCCAACAAGGTGCTTGGGAA-3'
Reverse	5'-CTTGAAGTGCCTCCACTGGC-3'
<i>Nampt</i>	
Forward	5'-ATCCAGGAGGCCAAAGAAGT-3'
Reverse	5'-CGGGAGATGACCATCGTATT-3'

CTR, calcitonin receptor; *DC-STAMP*, dendritic cell-specific transmembrane protein; *Nampt*, nicotinamide phosphoribosyltransferase; *NFATc1*, nuclear factor of activated T cells, cytoplasmic 1; *OSCAR*, osteoclast-associated receptor; *TRAP*, tartrate-resistant acid phosphatase.

microscope and osteoclast formation was determined by fixing in 3.7% formalin, permeabilizing with 0.1% Triton X-100, and staining with TRAP solution.

Statistical analysis. Each experiment was performed at least three times and all quantitative data are presented as

the mean \pm standard deviation. All statistical analyses were performed using SPSS (Korean version 14.0; SPSS, Inc., Chicago, IL, USA). Student's t-test was used to compare the parameters between two groups, whereas the one-way analysis of variance test, followed by the Tukey post hoc test, was used to compare the parameters among three groups. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

***Nampt* inhibits RANKL-mediated osteoclast formation in a dose-dependent manner with no cytotoxicity.** The present study analyzed the expression of *Nampt* in BMM cultures treated with M-CSF (30 ng/ml) and RANKL (100 ng/ml). As shown in Fig. 1A, the mRNA expression levels of *Nampt* were reduced in the presence of RANKL. To validate the effects of *Nampt* on osteoclast differentiation, mouse primary BMMs were treated with M-CSF (30 ng/ml) and RANKL (100 ng/ml) in the presence or absence of various concentrations of *Nampt*. As expected, the control untreated group generated TRAP-positive (TRAP⁺) osteoclasts. However, the presence of *Nampt* suppressed the formation of TRAP⁺ multinucleated osteoclasts in a dose-dependent manner (Fig. 1B and C). Subsequently, XTT cell viability assays were conducted to ascertain whether *Nampt* induced cytotoxicity during RANKL-induced osteoclast differentiation. The addition of *Nampt* did not affect cell viability at any of the concentrations used in the present study (Fig. 1D).

***Nampt* regulates osteoclastogenesis via mediating RANKL-dependent early signaling pathways.** To elucidate a molecular mechanism that underlies the inhibitory effects of *Nampt* on osteoclastogenesis, *Nampt* was added to BMM cultures treated with M-CSF (30 ng/ml) and RANKL (100 ng/ml), at three different time points after RANKL treatment. The results indicated that *Nampt* (500 ng/ml) significantly blocked osteoclast differentiation when the cells were exposed on days 0-1 after RANKL treatment but not on days 1-2 or 2-3 (Fig. 2A and B). As shown in Fig. 2C and D, *Nampt* negatively affected the phosphorylation of JNK, Akt and GSK3 β . In addition, *Nampt* downregulated the phosphorylation of Btk and PLC γ 2, which are required for calcium signaling during osteoclast differentiation (Fig. 2E and F). These results indicated that *Nampt* is involved in the early stages of osteoclast differentiation by inducing dephosphorylation of JNK, Akt and its downstream target GSK3 β , Btk and PLC γ 2.

***Nampt* downregulates the expression levels of *c-Fos*, *NFATc1* and osteoclast-specific marker genes.** To examine whether *Nampt* regulates RANKL-induced osteoclast differentiation by downregulating the activation of *c-Fos* and *NFATc1*, the present study evaluated the effects of *Nampt* on RANKL-induced *c-Fos* and *NFATc1* expression. When BMMs were stimulated for 12-48 h with RANKL (100 ng/ml), the mRNA expression levels of *c-Fos* and *NFATc1* were increased in the control group, whereas *Nampt* treatment reduced their expression (Fig. 3A). Similarly, western blot analysis demonstrated that *Nampt* significantly reduced the protein levels of *c-Fos* and *NFATc1* (Fig. 3B). Subsequently, the present study examined




Figure 1. *Nampt* attenuates TRAP-positive osteoclast formation without cytotoxicity. (A) BMMs were cultured in the presence of M-CSF (30 ng/ml) and then stimulated with RANKL (100 ng/ml) for the indicated durations. Total RNA was isolated from cells using QIAzol reagent and *Nampt* mRNA levels were evaluated by reverse transcription-quantitative polymerase chain reaction. *** $P < 0.001$ vs. control group. (B) BMMs were cultured for 3 days in the presence of M-CSF (30 ng/ml) and RANKL (100 ng/ml), with or without the indicated concentrations of *Nampt*. Cells were fixed, permeabilized and stained with TRAP solution. Images of TRAP⁺ cells were captured under a light microscope (magnification, $\times 5$). (C) BMMs were seeded into a 96-well plate and cultured for 3 days in the presence of M-CSF (30 ng/ml) with the indicated concentrations of *Nampt* (100, 250 or 500 ng/ml). After 3 days, cell viability was analyzed by the XTT assay. (D) TRAP⁺ MNCs with > 5 nuclei were counted as osteoclasts. *** $P < 0.001$ vs. control group. BMMs, bone marrow macrophages; M-CSF, macrophage colony-stimulating factor; MNCs, mononucleated cells; N, nuclei; *Nampt*, nicotinamide phosphoribosyltransferase; RANKL, receptor activator of nuclear factor- κ B ligand; TRAP, tartrate-resistant acid phosphatase.

whether ectopic expression of *c-Fos* or *NFATc1* is sufficient to rescue the inhibitory effects of *Nampt* on osteoclastogenesis using a retroviral system. BMMs were infected with *c-Fos* or *NFATc1*-encoding retroviruses and cultured with M-CSF (30 ng/ml) and RANKL (100 ng/ml) in the presence or absence of *Nampt* (500 ng/ml). Indeed, the overexpression of *c-Fos* or *NFATc1* rescued the anti-osteoclastogenic effect of *Nampt* (Fig. 3C). In addition, it was determined whether *Nampt* regulates the mRNA expression of various osteoclast-specific transcription factors, including *TRAP*, osteoclast associated receptor (*OSCAR*), cathepsin K, calcitonin receptor (*CTR*), *Atp6v0d2*, dendritic cell-specific transmembrane protein (*DC-STAMP*), α v-integrin and β 3-integrin. These genes are associated with osteoclast formation and function during RANKL-induced

osteoclast differentiation. The mRNA expression levels of *TRAP*, *OSCAR*, cathepsin K, *CTR*, *Atp6v0d2*, *DC-STAMP*, α v-integrin and β 3-integrin were significantly decreased by *Nampt* (Fig. 4). These results suggested that *Nampt* efficiently inhibits *c-Fos* and *NFATc1* activation, leading to the downregulation of osteoclast marker gene expression during RANKL-mediated osteoclast differentiation.

Nampt is not associated with the bone-resorbing activity of mature osteoclasts. The present study subsequently examined whether *Nampt* regulates osteoclastic bone-resorptive functions. To investigate this role, mature osteoclasts were seeded onto the top of hydroxyapatite-coated plates in the presence or absence of *Nampt* (500 ng/ml). However, the number and area




Figure 2. *Nampt* inhibits the early stages of osteoclastogenesis by downregulating RANKL-dependent early signaling pathways. (A) BMMs were cultured for 3 days in the presence of M-CSF (30 ng/ml) and RANKL (100 ng/ml) with or without *Nampt* (500 ng/ml) for the indicated durations. Cells were then stained with TRAP solution and images of TRAP⁺ cells were captured under a light microscope (magnification, 5x). The diagram shows the time period of *Nampt* treatment. (B) TRAP⁺ MNCs with >5 nuclei were counted as osteoclasts. ****P*<0.001 vs. control group. (C) BMMs were pretreated with or without *Nampt* (500 ng/ml) for 1 h in the presence of M-CSF (30 ng/ml) prior to RANKL (100 ng/ml) stimulation at the indicated time points. Whole-cell lysates were analyzed by western blotting with the indicated antibodies. β -actin was used as the internal control. (D) Semi-quantification of blots was performed using ImageJ. ****P*<0.001 vs. the control group. (E) BMMs were pretreated with or without *Nampt* (500 ng/ml) for 1 h in the presence of M-CSF (30 ng/ml) prior to RANKL (100 ng/ml) stimulation at the indicated time points. Whole-cell lysates were analyzed by western blotting with the indicated antibodies. β -actin was used as the internal control. (F) Semi-quantification of western blot bands was performed using ImageJ. ****P*<0.001 vs. the control group. BMMs, bone marrow macrophages; Btk, Bruton's tyrosine kinase; ERK, extracellular signal-regulated protein kinases; GSK3 β , glycogen synthase kinase-3 β ; JNK, c-Jun N-terminal kinase; M-CSF, macrophage colony-stimulating factor; MNCs, mononucleated cells; N, nuclei; *Nampt*, nicotinamide phosphoribosyltransferase; p-, phosphorylated; PLC γ 2, phospholipase C γ -2 RANKL, receptor activator of nuclear factor- κ B ligand; TRAP, tartrate-resistant acid phosphatase.




Figure 3. *Nampt* reduces the expression of *c-Fos* and *NFATc1*. (A) BMMs were pretreated with or without *Nampt* (500 ng/ml) for 1 h in the presence of M-CSF (30 ng/ml), and then stimulated with RANKL (100 ng/ml) for the indicated times. The mRNA expression levels of *c-Fos* and *NFATc1* were analyzed by reverse transcription-quantitative polymerase chain reaction. * $P < 0.05$, *** $P < 0.001$ vs. control group at the indicated time points. (B) The effects of *Nampt* on protein levels of *c-Fos* and *NFATc1* were evaluated by western blot analysis with the indicated antibodies. β -actin was used as the internal control. (C) BMMs were infected with retroviruses expressing pMX-IRES-EGFP (pMX), pMX-*c-Fos*-IRES-EGFP (*c-Fos*) or pMX-*NFATc1*-IRES-EGFP (*NFATc1*). Infected BMMs were cultured with or without *Nampt* (500 ng/ml) in the presence of M-CSF (30 ng/ml) and RANKL (100 ng/ml) for 4 days. After culturing, the cells were stained with TRAP solution. Images of TRAP⁺ cells were captured under a light microscope (magnification, 5x). TRAP⁺ MNCs with >5 nuclei were counted as osteoclasts. *** $P < 0.001$ vs. control group and ### $P < 0.001$ vs. *Nampt* group. BMMs, bone marrow macrophages; M-CSF, macrophage colony-stimulating factor; MNCs, mononucleated cells; N, nuclei; *Nampt*, nicotinamide phosphoribosyltransferase; *NFATc1*, nuclear factor of activated T cells, cytoplasmic 1; RANKL, receptor activator of nuclear factor- κ B ligand; TRAP, tartrate-resistant acid phosphatase.




Figure 4. *Nampt* reduces the expression of osteoclast-specific genes. BMMs were pretreated with or without *Nampt* (500 ng/ml) for 1 h in the presence of M-CSF (30 ng/ml) and then stimulated with RANKL (100 ng/ml) for the indicated times. Total RNA was isolated from cells using QIAzol reagent and the mRNA expression levels of *TRAP*, *OSCAR*, *cathepsin K*, *CTR*, *Atp6v0d2*, *DC-STAMP*, β 3-integrin and α v-integrin were evaluated by reverse transcription-quantitative polymerase chain reaction. *** $P < 0.001$ vs. control group at the indicated time. BMMs, bone marrow macrophages; *CTR*, calcitonin receptor; *DC-STAMP*, dendritic cell-specific transmembrane protein; M-CSF, macrophage colony-stimulating factor; *Nampt*, nicotinamide phosphoribosyltransferase; *OSCAR*, osteoclast-associated receptor; RANKL, receptor activator of nuclear factor- κ B ligand; *TRAP*, tartrate-resistant acid phosphatase.

of resorption pits were unaffected by *Nampt* treatment (Fig. 4), suggesting that *Nampt* does not have a role in the resorbing activity of mature osteoclasts.

Discussion

The present study demonstrated that *Nampt* attenuated RANKL-mediated differentiation of primary mouse BMMs

into TRAP⁺ MNCs in a dose-dependent manner without cytotoxic effects. During this process, *Nampt* decreased the phosphorylation of various early signal transducers, including JNK and Akt, and its downstream target, GSK3 β , as well as calcium-dependent signaling pathways, including PLC γ 2 and Btk. Furthermore, the mRNA and protein expression levels of two master regulators of osteoclastogenesis, *c-Fos* and *NFATc1*, were significantly decreased by *Nampt*




Figure 5. *Nampt* does not affect the bone-resorbing activity of mature osteoclasts. Mature osteoclasts were seeded on hydroxyapatite-coated plates and treated for 24 h with *Nampt* (500 ng/ml). Attached cells were removed and images were captured under a light microscope (magnification, x5). Pit areas on the hydroxyapatite plate were quantified using the Image Pro-Plus (version 4.5) software. *Nampt*, nicotinamide phosphoribosyltransferase.

treatment, leading to decreased expression levels of various key transcription factors in osteoclast differentiation including *TRAP*, *OSCAR*, cathepsin K, *CTR*, *Atp6vOd2*, *DC-STAMP*, β 3- and α v-integrin.

The differentiation of monocyte/macrophage lineage precursors into bone-resorbing osteoclasts is initiated in response to two important cytokines, M-CSF and RANKL, resulting in the activation of early downstream pathways (15). During this process, the phosphorylation of numerous signal transducers, including mitogen-activated protein kinases (MAPKs), which are comprised of p38, ERK and JNK; nuclear factor- κ B; phosphatidylinositol 3-kinase/Akt; PLC γ 2 and Btk occurs (16-19). RANKL-mediated activation of JNK is known to have an anti-apoptotic function in osteoclastogenesis, and Akt is a potent inducer of osteoclast differentiation by promoting the formation of an inactive form of GSK3 β (p-GSK3 β) and the nuclear translocation of NFATc1 (20,21). In addition, it has been well established that calcium signaling is crucial for RANKL-dependent osteoclastogenesis. PLC γ 2 activation requires phosphorylation of its tyrosine residues to induce calcium oscillations and the translocation of NFATc1 by forming a complex with regulatory adapter molecule GRB2-associated binding protein 2 and modulating its recruitment to RANK (22,23). PLC γ 2 is regulated by the upstream tyrosine kinase Btk, which is involved in osteoclast differentiation by linking RANK and immunoreceptor tyrosine-based activation motif signaling, with subsequent regulation of the formation of Btk/BLNK-containing complex and activation of PLC γ 2-dependent calcium signaling (19). The

results of the present study revealed that *Nampt* suppressed RANKL-induced osteoclast differentiation by interfering with survival-related signaling pathways that contain JNK and Akt, as well as Btk-PLC γ 2-dependent intracellular calcium signaling. Since *Nampt* affected several signals associated with the early stages of osteoclastogenesis, the present study examined whether *Nampt* is involved in the expression of late-stage transcription factors, *c-Fos* and *NFATc1*. A previous report indicated that *c-Fos* knock-out mice exhibit morphological characteristics of osteopetrosis, owing to osteoclast malfunction, whereas impaired osteoclastogenesis in murine BMMs is completely rescued by exogenous overexpression of *c-Fos* (24,25). In response to the activation of *c-Fos*, another master regulator, *NFATc1*, serves a crucial role in osteoclast differentiation. *NFATc1* inhibition in embryonic stem cells suppresses their ability to differentiate into normal osteoclasts, and this phenomenon is reversed by ectopic expression of *NFATc1* even in the absence of RANKL (26,27). This c-Fos-NFATc1 activation cascade leads to the elevated expression of osteoclast-specific gene markers, such as *TRAP*, *OSCAR*, *CTR*, cathepsin K, *DC-STAMP* and β 3-integrin. The present data revealed that *Nampt* expression significantly decreased mRNA and protein levels of *c-Fos* and *NFATc1* compared with the control, resulting in the downregulation of various transcription factors associated with osteoclast formation and function.

Although the relationship between bone regulation and *Nampt* has previously been reported, the present study is the first, to the best of our knowledge, to identify the effects of

Nampt on mouse BMM-derived osteoclastogenesis and its molecular mechanisms. In conclusion, the present study demonstrated that the adipokine *Nampt* effectively interferes with RANKL-mediated osteoclast differentiation by inactivating several early signal transducers. These signaling molecules include JNK, Akt and GSK3 β , as well as Btk-PLC γ 2-calcium signaling. Furthermore, *Nampt* treatment decreases c-Fos and NFATc1 mRNA and protein levels, resulting in the downregulation of various target gene mRNA levels. *Nampt* does not influence the bone-resorbing activity of mature osteoclasts; instead, *Nampt* exerts its anti-osteoclastogenic effects by targeting osteoclast precursors rather than mature multinucleated osteoclasts. Although further studies are required to reveal the restorative effect of *Nampt* on osteoporotic bone loss in mouse models, it may be suggested that increased *Nampt* is a potential target for the treatment of metabolic bone diseases, such as osteoporosis, by suppressing osteoclast differentiation and function.

Acknowledgments

The present study was supported by a grant from the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (grant no. NRF-2016R1D1A1B03933712).

References

- Samal B, Sun Y, Steams G, Xie C, Suggs S and McNiece I: Cloning and characterization of the cDNA encoding a novel human pre-B-cell colony-enhancing factor. *Mol Cell Biol* 14: 1431-1437, 1994.
- Luk T, Malam Z and Marshall JC: Pre-B cell colony-enhancing factor (PBEF)/visfatin: A novel mediator of innate immunity. *J Leukoc Biol* 83: 804-816, 2006.
- Gosset M, Berenbaum F, Salvat C, Sautet A, Pigenet A, Tahiri K and Jacques C: Crucial role of visfatin/pre-B cell colony-enhancing factor in matrix degradation and prostaglandin E2 synthesis in chondrocytes: Possible influence on osteoarthritis. *Arthritis Rheum* 58: 1399-1409, 2008.
- Moschen AR, Kaser A, Enrich B, Mosheimer B, Theurl M, Niederegger H and Tilg H: Visfatin, an adipocytokine with proinflammatory and immunomodulating properties. *J Immunol* 178: 1748-1758, 2007.
- Revollo JR, Grimm AA and Imai S: The NAD biosynthesis pathway mediated by nicotinamide phosphoribosyltransferase regulates Sir2 activity in mammalian cells. *J Biol Chem* 279: 50754-50763, 2004.
- Tan B, Dong S, Shepard RL, Kays L, Roth KD, Geeganage S, Kuo MS and Zhao G: Inhibition of nicotinamide phosphoribosyltransferase (NAMPT), an enzyme essential for NAD⁺ biosynthesis, leads to altered carbohydrate metabolism in cancer cells. *J Biol Chem* 290: 15812-15824, 2015.
- Yang S, Ryu JH, Oh H, Jeon J, Kwak JS, Kim JH, Kim HA, Chun CH and Chun JS: NAMPT (visfatin), a direct target of hypoxia-inducible factor-2 α , is an essential catabolic regulator of osteoarthritis. *Ann Rheum Dis* 74: 595-602, 2015.
- Liu Y, Song CY, Wu SS, Liang QH, Yuan LQ and Liao EY: Novel adipokine and bone metabolism. *Int J Endocrinol* 2013: 895045, 2013.
- Wattanachanya L, Lu WD, Kundu RK, Wang L, Abbott MJ, O'Carroll D, Quertermous T and Nissenson RA: Increased bone mass in mice lacking the adipokine apelin. *Endocrinology* 154: 2069-2080, 2013.
- Kamio N, Kawato T, Tanabe N, Kitami S, Morita T, Ochiai K and Maeno M: Vaspin attenuates RANKL-induced osteoclast formation in RAW264.7 cells. *Connect Tissue Res* 54: 147-152, 2013.
- Tu Q, Zhang J, Dong LQ, Saunders E, Luo E, Tang J and Chen J: Adiponectin inhibits osteoclastogenesis and bone resorption via APPL1-mediated suppression of Akt1. *J Biol Chem* 286: 12542-12553, 2011.
- Venkateshaiah SU, Khan S, Ling W, Bam R, Li X, van Rhee F, Usmani S, Barlogie B, Epstein J and Yaccoby S: NAMPT/PBEF1 enzymatic activity is indispensable for myeloma cell growth and osteoclast activity. *Exp Hematol* 41: 547.e2-557.e2, 2013.
- Moschen AR, Geiger S, Germer R and Tilg H: Pre-B cell colony enhancing factor/NAMPT/visfatin and its role in inflammation-related bone disease. *Mutat Res* 690: 95-101, 2010.
- Kim JY, Cheon YH, Oh HM, Rho MC, Erkhembaatar M, Kim MS, Lee CH, Kim JJ, Choi MK, Yoon KH, *et al*: Oleonic acid acetate inhibits osteoclast differentiation by downregulating PLC γ 2-Ca(2+)-NFATc1 signaling, and suppresses bone loss in mice. *Bone* 60: 104-111, 2014.
- Glantschnig H, Fisher JE, Wesolowski G, Rodan GA and Reszka AA: M-CSF, TNF α and RANK ligand promote osteoclast survival by signaling through mTOR/S6 kinase. *Cell Death Differ* 10: 1165-1177, 2003.
- Li X, Udagawa N, Itoh K, Suda K, Murase Y, Nishihara T, Suda T and Takahashi N: p38 MAPK-mediated signals are required for inducing osteoclast differentiation but not for osteoclast function. *Endocrinology* 143: 3105-3113, 2002.
- Kobayashi N, Kadono Y, Naito A, Matsumoto K, Yamamoto T, Tanaka S and Inoue J: Segregation of TRAF6-mediated signaling pathways clarifies its role in osteoclastogenesis. *EMBO J* 20: 1271-1280, 2001.
- Gingery A, Bradley E, Shaw A and Oursler MJ: Phosphatidylinositol 3-kinase coordinately activates the MEK/ERK and AKT/NF κ B pathways to maintain osteoclast survival. *J Cell Biochem* 89: 165-179, 2003.
- Shinohara M, Koga T, Okamoto K, Sakaguchi S, Arai K, Yasuda H, Takai T, Kodama T, Morio T, Geha RS, *et al*: Tyrosine kinases Btk and Tec regulate osteoclast differentiation by linking RANK and ITAM signals. *Cell* 132: 794-806, 2008.
- Ikeda F, Matsubara T, Tsurukai T, Hata K, Nishimura R and Yoneda T: JNK/c-Jun signaling mediates an anti-apoptotic effect of RANKL in osteoclasts. *J Bone Miner Res* 23: 907-914, 2008.
- Moon JB, Kim JH, Kim K, Youn BU, Ko A, Lee SY and Kim N: Akt induces osteoclast differentiation through regulating the GSK3 β /NFATc1 signaling cascade. *J Immunol* 188: 163-169, 2012.
- Wilde JI and Watson SP: Regulation of phospholipase C gamma isoforms in haematopoietic cells: Why one, not the other? *Cell Signal* 13: 691-701, 2001.
- Mao D, Eppele H, Uthgenannt B, Novack DV and Faccio R: PLC γ 2 regulates osteoclastogenesis via its interaction with ITAM proteins and GAB2. *J Clin Invest* 116: 2869-2879, 2006.
- Wisdon R and Verma IM: Transformation by Fos proteins requires a C-terminal transactivation domain. *Mol Cell Biol* 13: 7429-7438, 1993.
- Grigoriadis AE, Wang ZQ, Cecchini MG, Hofstetter W, Felix R, Fleisch HA and Wagner EF: c-Fos: A key regulator of osteoclast-macrophage lineage determination and bone remodeling. *Science* 266: 443-448, 1994.
- Takayanagi H: The role of NFAT in osteoclast formation. *Ann N Y Acad Sci* 1116: 227-237, 2007.
- Takayanagi H, Kim S, Koga T, Nishina H, Isshiki M, Yoshida H, Saiura A, Isobe M, Yokochi T, Inoue J, *et al*: Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell* 3: 889-901, 2002.