



# New primers for promising single-copy genes in fungal phylogenetics and systematics

I. Schmitt<sup>1</sup>, A. Crespo<sup>2</sup>, P.K. Divakar<sup>2</sup>, J.D. Fankhauser<sup>1</sup>, E. Herman-Sackett<sup>3</sup>,  
K. Kalb<sup>4</sup>, M.P. Nelsen<sup>3,5</sup>, N.A. Nelson<sup>1,6</sup>, E. Rivas-Plata<sup>3,7</sup>, A.D. Shimp<sup>1</sup>,  
T. Widhelm<sup>3,7</sup>, H.T. Lumbsch<sup>3</sup>

## Key words

*Ascomycota*  
DNA replication licensing factor  
evolution  
lichenised fungi  
*Mcm7*  
MS277  
MS456  
phylogeny  
pre-rRNA processing protein  
protein-coding  
*Tsr1*

**Abstract** Developing powerful phylogenetic markers is a key concern in fungal phylogenetics. Here we report degenerate primers that amplify the single-copy genes *Mcm7* (MS456) and *Tsr1* (MS277) across a wide range of *Peizizomycotina* (*Ascomycota*). Phylogenetic analyses of 59 taxa belonging to the *Eurotiomycetes*, *Lecanoromycetes*, *Leotiomycetes*, *Lichinomycetes* and *Sordariomycetes*, indicate the utility of these loci for fungal phylogenetics at taxonomic levels ranging from genus to class. We also tested the new primers in silico using sequences of *Saccharomycotina*, *Taphrinomycotina* and *Basidiomycota* to predict their potential of amplifying widely across the *Fungi*. The analyses suggest that the new primers will need no, or only minor sequence modifications to amplify *Saccharomycotina*, *Taphrinomycotina* and *Basidiomycota*.

**Article info** Received: 5 May 2009; Accepted: 1 July 2009; Published: 4 August 2009.

## INTRODUCTION

Molecular systematics has revolutionised our view of fungal evolution. Recent large scale sequencing efforts resulted in comprehensive multi-locus phylogenies, which have significantly improved our understanding of phylogenetic relationships within fungi (Binder & Hibbett 2002, Lumbsch et al. 2004, Lutzoni et al. 2004, James et al. 2006). These data led to the first phylogenetic classification of the *Fungi* (Hibbett et al. 2007). However, early events in fungal evolution still remain uncertain because of missing support and resolution at the backbone of the phylogeny. We lack information, for example, about the relationships of the different ascomycete classes to one another, or the evolution within major lineages, such as the lichenised *Lecanoromycetes*, or the basidiomycete clade *Agaricomycetes*. Robust and well-supported phylogenies are essential for a better understanding of fungal evolution, and a prerequisite for studies aiming at reconstructing the evolution of non-molecular characters on the background of a molecular phylogeny.

Commonly used molecular loci in fungal phylogenetics include nuclear and mitochondrial ribosomal rDNA (18S, 28S, ITS, IGS, mtSSU, mtLSU), as well as protein-coding genes, such

as RNA polymerases (*RPB1* and *RPB2*),  $\beta$ -tubulin,  $\gamma$ -actin, ATP synthase (*ATP6*), and elongation factor EF-1 $\alpha$  (*TEF1* $\alpha$ ). Some single-copy protein-coding genes such as *RPB1* and *RPB2* are promising for yielding well resolved and highly supported phylogenies (Liu & Hall 2004, Reeb et al. 2004, Crespo et al. 2007, Lumbsch et al. 2007). Other protein-coding genes, such as the tubulins, are present in the genome in multiple copies and thus have the potential of being phylogenetically misleading (Landvik et al. 2001). Generally, slow evolving loci are more suitable for reconstruction of deep phylogenetic relationships, while loci with high rates of evolution are better for the reconstruction of more recent evolutionary events. Ribosomal loci with high and heterogeneous rates of change, such as ITS, IGS and mtSSU rDNA, can be used to distinguish taxa at the genus and species level. However, the non-coding regions of these loci are prone to significant length variation, making alignment of distantly related taxa problematic. Fast evolving ribosomal genes are therefore less useful in large scale concatenated analyses involving higher-level phylogenetic relationships. Molecular systematists are constantly searching for loci that are conserved enough to produce reliable alignments, and at the same time have sufficient variability to yield well resolved and well supported phylogenies. Analysing phylogenetic relationships at lower and higher taxonomic levels simultaneously, while using only a few loci, is desirable, because sequencing entire genomes or even multiple loci is not feasible for many phylogenetically interesting taxa. Fungal material suitable for molecular study is often limited, and culturing of many species impossible.

In a recent study Aguilera et al. (2008) used a bioinformatics approach to assess the performance of single-copy protein-coding genes for fungal phylogenetics. Their analyses of 30 published fungal genomes revealed two loci, MS277 and MS456, which outperformed all other single-copy genes in phylogenetic utility. MS277 corresponds to the gene *Tsr1*, required for rRNA

<sup>1</sup> Department of Plant Biology and Bell Museum of Natural History, University of Minnesota, 1445 Gortner Ave, St. Paul, MN 55108, USA; corresponding author e-mail: schm2109@umn.edu.

<sup>2</sup> Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense de Madrid, Madrid 28040, Spain.

<sup>3</sup> Department of Botany, The Field Museum, 1400 S. Lake Shore Dr., Chicago, IL 60605, USA.

<sup>4</sup> Im Tal 12, D-92318 Neumarkt, Germany.

<sup>5</sup> Committee on Evolutionary Biology, University of Chicago, 1025 E. 57th Street, Chicago, IL 60637, USA.

<sup>6</sup> Augsburg College, 2211 Riverside Avenue South, Minneapolis, MN 55454, USA.

<sup>7</sup> Biological Sciences Department, University of Illinois at Chicago, 845 W. Taylor St., Chicago, IL 60607, USA.

**Table 1** Material and DNA sequences used in this study.

Species	Order	Class	Source	GenBank accession	
				<i>Mcm7</i> (MS456)	<i>Tsr1</i> (MS277)
<i>Ajellomyces capsulatus</i>	Onygenales	Eurotiomycetes	–	XM_001538714	XM_001541629
<i>Arctomia delicatula</i>	<i>incertae sedis</i>	Lecanoromycetes	Sweden, 2002, <i>Palice s.n.</i> (F)	GQ272388	GQ272430
<i>Arctomia teretiuscula</i>	<i>incertae sedis</i>	Lecanoromycetes	China (GZU – holotype)	GQ272389	GQ272431
<i>Aspergillus clavatus</i>	Eurotiales	Eurotiomycetes	–	XM_001275314	XM_001275562
<i>Aspergillus fumigatus</i>	Eurotiales	Eurotiomycetes	–	XM_750254	XM_750526
<i>Aspergillus nidulans</i>	Eurotiales	Eurotiomycetes	–	XM_658504	XM_658778
<i>Aspergillus nidulans</i>	Eurotiales	Eurotiomycetes	–	XM_001213626	XM_001208611
<i>Aspergillus niger</i>	Eurotiales	Eurotiomycetes	–	XM_001397760	XM_001399262
<i>Aspergillus oryzae</i>	Eurotiales	Eurotiomycetes	–	XM_001826176	XM_001821764
<i>Aspicilia caesiocinerea</i>	Pertusariales	Lecanoromycetes	USA, <i>Lumbsch 19277e</i> (F)	GQ272390	GQ272432
<i>Aspicilia cinerea</i>	Pertusariales	Lecanoromycetes	USA, <i>Lumbsch 19190c</i> (F)	GQ272391	GQ272433
<i>Botryotinia fuckeliana</i>	Helotiales	Leotiomycetes	–	XM_001556412	XM_001554531
<i>Bulbothrix apophysata</i>	Lecanorales	Lecanoromycetes	Costa Rica, <i>Lücking 16650btu</i> (F)	GQ272392	GQ272434
<i>Cetrariastrum andense</i>	Lecanorales	Lecanoromycetes	Peru, <i>Lumbsch 19334</i> (MAF)	GQ272429	GQ272471
<i>Cetrariastrum dubitans</i>	Lecanorales	Lecanoromycetes	Peru, <i>Lumbsch 19366</i> (MAF)	GQ272427	GQ272470
<i>Chaetomium globosum</i>	Sordariales	Sordariomycetes	–	XM_001220296	XM_001225626
<i>Coccidioides immitis</i>	Onygenales	Eurotiomycetes	–	XM_001240385	XM_001245725
<i>Dermatocarpon intestininforme</i>	Verrucariales	Eurotiomycetes	Turkey, 27.7.1997, <i>John</i> (F)	GQ272393	GQ272435
<i>Dermatocarpon miniatum</i>	Verrucariales	Eurotiomycetes	Germany, 17.10.2001, <i>Zimmermann</i> (F)	GQ272394	GQ272436
<i>Everniastrum lipidiferum</i>	Lecanorales	Lecanoromycetes	Peru, <i>Lumbsch 19309b</i> (MAF)	GQ272395	GQ272437
<i>Everniopsis trulla</i>	Lecanorales	Lecanoromycetes	Peru, <i>Lumbsch 19309b</i> (F)	GQ272396	GQ272438
<i>Flavoparmelia marchantii</i>	Lecanorales	Lecanoromycetes	Australia, (MAF-Lich 10492)	GQ272420	GQ272463
<i>Gibberella zeae</i>	Hypocreales	Sordariomycetes	–	XM_387281	XM_384579
<i>Lecanora allophana</i>	Lecanorales	Lecanoromycetes	Turkey, <i>Lumbsch 19618d</i> (F)	GQ272399	GQ272444
<i>Lecanora carpinea</i>	Lecanorales	Lecanoromycetes	Turkey, <i>Lumbsch 19611m</i> (F)	GQ272400	GQ272443
<i>Lecanora chlorotera</i>	Lecanorales	Lecanoromycetes	Turkey, <i>Lumbsch 19622e</i> (F)	GQ272398	GQ272440
<i>Lecanora margarodes</i>	Lecanorales	Lecanoromycetes	Australia, <i>Lumbsch 19086b</i> (F)	GQ272401	GQ272439
<i>Lecanora pulicaris</i>	Lecanorales	Lecanoromycetes	Turkey, <i>Lumbsch 19627c</i> (F)	GQ272419	GQ272441
<i>Lecanora subcarpinea</i>	Lecanorales	Lecanoromycetes	Turkey, <i>Lumbsch 19622a</i> (F)	GQ272428	GQ272442
<i>Lobothallia radiosa</i>	Pertusariales	Lecanoromycetes	Switzerland, 9.8.2004, <i>Lumbsch</i> (F)	GQ272397	GQ272445
<i>Magnaporthe grisea</i>	Sordariales	Sordariomycetes	–	XM_364455	XM_368157
<i>Malcolmiella psychotrioides</i>	Ostropales	Lecanoromycetes	Costa Rica, <i>Lücking s.n.</i> (F)	GQ272412	GQ272456
<i>Malcolmiella sp. 1</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37092</i> (hb. Kalb)	GQ272402	GQ272447
<i>Malcolmiella sp. 2</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 36969</i> (hb. Kalb)	GQ272411	GQ272455
<i>Malcolmiella sp. 3</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37093</i> (hb. Kalb)	GQ272405	GQ272450
<i>Malcolmiella sp. 4</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 36858</i> (hb. Kalb)	GQ272403	GQ272448
<i>Malcolmiella sp. 5</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37060</i> (hb. Kalb)	GQ272407	GQ272446
<i>Malcolmiella sp. 6</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37072</i> (hb. Kalb)	GQ272408	GQ272452
<i>Malcolmiella sp. 7</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 36832</i> (hb. Kalb)	GQ272406	GQ272451
<i>Malcolmiella sp. 8</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37005</i> (hb. Kalb)	GQ272409	GQ272453
<i>Malcolmiella sp. 9</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 36963</i> (hb. Kalb)	GQ272404	GQ272449
<i>Malcolmiella sp. 10</i>	Ostropales	Lecanoromycetes	Thailand, <i>Kalb 37086</i> (hb. Kalb)	GQ272410	GQ272454
<i>Neosartorya fischeri</i>	Eurotiales	Eurotiomycetes	–	XM_001260497	XM_001260746
<i>Neurospora crassa</i>	Sordariales	Sordariomycetes	–	XM_958785	XM_951859
<i>Ochrolechia parella</i>	Pertusariales	Lecanoromycetes	Turkey, <i>Lumbsch 19625g</i> (MIN)	GQ272421	GQ272464
<i>Ochrolechia subpallascens</i>	Pertusariales	Lecanoromycetes	USA, <i>Lumbsch 19900a</i> & <i>Schmitt</i> (MIN)	GQ272422	GQ272465
<i>Parmeliopsis hyperopta</i>	Lecanorales	Lecanoromycetes	Spain (MAF-Lich 10181)	GQ272426	GQ272468
<i>Peltula euploca</i>	Lichinales	Lichinomycetes	USA, <i>Lumbsch 19923b</i> & <i>Schmitt</i> (MIN)	GQ272424	GQ272467
<i>Penicillium marneffeii</i>	Eurotiales	Eurotiomycetes	–	XM_002146315	XM_002148793
<i>Pertusaria amara</i>	Pertusariales	Lecanoromycetes	USA, <i>Lumbsch 19925a</i> & <i>Schmitt</i> (MIN)	GQ272423	GQ272466
<i>Pertusaria velata</i>	Pertusariales	Lecanoromycetes	USA, <i>Lumbsch 19913c</i> & <i>Schmitt</i> (MIN)	GQ272425	GQ272469
<i>Podospora anserina</i>	Sordariales	Sordariomycetes	–	XM_001912857	XM_001909251
<i>Psiloparmelia denotata</i>	Lecanorales	Lecanoromycetes	Peru, <i>Lumbsch 19302g</i> (F)	GQ272413	GQ272457
<i>Pyrenula subpraelucida</i>	Pyrenulales	Eurotiomycetes	Costa Rica, <i>Lücking 17550f</i> (F)	GQ272414	GQ272459
<i>Pyrgillus javanicus</i>	Pyrenulales	Eurotiomycetes	Australia, <i>Lumbsch 19115e</i> (F)	GQ272415	GQ272458
<i>Sclerotinia sclerotiorum</i>	Helotiales	Leotiomycetes	–	XM_001586126	XM_001593622
<i>Umbilicaria leprosa</i>	<i>incertae sedis</i>	Lecanoromycetes	Peru, <i>Lumbsch 19355a</i> (F)	GQ272416	GQ272460
<i>Usnea endochrysa</i>	Lecanorales	Lecanoromycetes	USA, <i>Buck 51175</i> (hb. Lendemer)	GQ272417	GQ272461
<i>Verrucaria muralis</i>	Verrucariales	Eurotiomycetes	Czech Republic, <i>Palice 6011</i> (hb. Palice)	GQ272418	GQ272462

**Table 2** Primers developed in the current study.

Primer Name	Direction	Sequence (5'-3')	Position in <i>A. nidulans</i> mRNA (XM_658504 and XM_658778)	Corresponding amino acid sequence in <i>A. nidulans</i> (AN5992 and AN6266)	Length	Degeneracy
<i>Mcm7</i> -709for	For	ACI MGI GTI TCV GAY GTH AAR CC	709	TRVSDVKP	23 bp	32
<i>Mcm7</i> -1348rev	Rev	GAY TTD GCI ACI CCI GGR TCW CCC AT	1348	MGDPGVAKS	26 bp	16
<i>Mcm7</i> -1447rev	Rev	C ATI ACI GCI GCI GTR AGR CC	1447	GLTAAVM	21 bp	4
<i>Tsr1</i> -1453for	For	GAR TTC CCI GAY GAR ATY GAR CT	1453	EFPDEIEL	23 bp	32
<i>Tsr1</i> -1459for	For	CCI GAY GAR ATY GAR CTI CAY CC	1459	PDEIELHP	23 bp	32
<i>Tsr1</i> -2308rev	Rev	CTT RAA RTA ICC RTG IGT ICC	2308	GTHGYFK	21 bp	8

**Table 3** Annealing conditions and PCR success rates for primers used in this study.

Gene	Primer combination	Approximate fragment length	Annealing temp.	PCR success (% of attempts)
<i>Mcm7</i> (MS456)	<i>Mcm7</i> -709for/ <i>Mcm7</i> -1348rev	640 bp	56 °C	80 %
<i>Mcm7</i> (MS456)	<i>Mcm7</i> -709for/ <i>Mcm7</i> -1447rev	740 bp	56 °C	50 %
<i>Tsr1</i> (MS277)	<i>Tsr1</i> -1459for/ <i>Tsr1</i> -2308rev	750 bp	49 °C	40 %
<i>Tsr1</i> (MS277)	<i>Tsr1</i> -1453for/ <i>Tsr1</i> -2308rev	750 bp	49 °C	40 %

accumulation during biogenesis of the ribosome (Gelperin et al. 2001), while MS456 corresponds to the gene *Mcm7*, a DNA replication licensing factor required for DNA replication initiation and cell proliferation (Moir et al. 1982, Kearsley & Labib 1998). Alignments based on these two loci alone recovered phylogenies that had the same topology, resolution power, and branch support as phylogenies based on a concatenated analysis of all 135 orthologous single-copy genes identified from fungal genomes (Aguileta et al. 2008). Strikingly, the authors report that most protein-coding genes commonly used in fungal systematics, such as *RPB1*, *RPB2*, *TEF1 $\alpha$* ,  $\beta$ -tubulin, and  $\gamma$ -actin are not found among the best performing genes.

In the current study we designed degenerate primers to amplify a 600–800 bp fragment of each, MS277 and MS456, over a wide range of *Peizizomycotina*. We tested variability and phylogenetic utility of these loci at taxonomic levels ranging from genus to class. Our analyses include in silico comparisons of the new primers to sequences of *Saccharomycotina* and *Basidiomycota* to predict primer utility in these phylogenetic groups.

## MATERIALS AND METHODS

Material and GenBank sequences used in the current study are listed in Table 1. We designed new degenerate primers based on amino acid alignments of *Mcm7* (MS456) and *Tsr1* (MS277) of euascomycete sequences available in GenBank. These alignments included members of *Dothideomycetes*, *Eurotiomycetes*, *Leotiomycetes* and *Sordariomycetes*. Primer sequences and annealing conditions are reported in Table 2 and 3. The locations of the fragments amplified by the new primers are indicated in Fig. 1. We used *Aspergillus nidulans* mRNA sequences of *Mcm7* and *Tsr1* as reference sequences (GenBank accession numbers XM\_658504 and XM\_658778). *Saccharomycotina*, *Taphrinomycotina* and *Basidiomycota* used for in silico analysis of primer fit are listed in Table 4.

### Molecular procedures


We extracted total genomic DNA from our samples using the Qiagen Plant Mini Kit (Qiagen). PCR reactions (25  $\mu$ L) contained PuReTaq Ready-To-Go PCR beads (GE Healthcare), 1.25  $\mu$ L of each primer (10 mM), 19.5  $\mu$ L H<sub>2</sub>O, and 3  $\mu$ L DNA template. Alternatively we used 0.125  $\mu$ L AmpliTaq Gold Taq (Applied Biosystems), 2.5  $\mu$ L buffer, 2  $\mu$ L dNTPs, 2.5–4  $\mu$ L

MgCl (20 mM), 0–5  $\mu$ L BSA, 1.25  $\mu$ L of each primer, and 3  $\mu$ L DNA template. We found that increasing the amount of forward primer *Tsr1*-1459for to 2.5  $\mu$ L, as well as adding 2  $\mu$ L MgCl (20 mM) to PCR reactions involving PCR beads often improved PCR results. PCR cycling conditions for *Mcm7*-709for/*Mcm7*-1447rev and *Mcm7*-709for/*Mcm7*-1348rev (MS456) were: initial denaturation 94 °C for 10 min, followed by 38 cycles of 94 °C for 45 s, 56 °C for 50 s, 72 °C for 1 min, and final elongation 72 °C for 5 min. PCR cycling conditions for *Tsr1*-1459for/*Tsr1*-2308rev (MS277) were the same as above except with 49 °C annealing temperature. Amplification products were stained with EZ-Vision DNA dye (Amresco) and viewed on 1 % low melt agarose gels. We excised bands of the expected length from the gel and purified them using GELase (Epicentre). Alternatively, PCR products were cleaned using the Bioclean Columns kit (Biotools, Madrid) according to the manufacturer's instructions. We sequenced the fragments using Big Dye v3.1 chemistry (Applied Biosystems) and the same primers as for PCR. Cycle sequencing was executed with the following program: initial denaturation for 1 min at 96 °C followed by 32 cycles of 96 °C for 15 s, 50 °C for 10 s, 60 °C for 4 min. Sequenced products were precipitated with 25  $\mu$ L of 100 % EtOH mixed with 1  $\mu$ L of 3 M NaOAc, and 1  $\mu$ L of EDTA, before they were loaded on an ABI PRISM™ 3730 DNA Analyser (Applied Biosystems). We assembled partial sequences using SeqMan v4.03 (Lasergene) and edited conflicts manually. We aligned the sequences based on amino acid sequence using ClustalW as implemented in the program BioEdit v7.0.9 (Hall 1999) and subsequently translated them back to nucleotides.

### Phylogenetic analyses

We assembled two alignments including the same 59 taxa each. For phylogenetic analysis we used a maximum parsimony (MP), maximum likelihood (ML) and a Bayesian approach (B/MCMC) (Larget & Simon 1999, Huelsenbeck et al. 2001). We performed all analyses on the single gene alignments as well as on a combined alignment. We tested for potential conflict between individual datasets by comparing the 75 % MP bootstrap consensus trees.

We used PAUP v4.0 (Swofford 2003), GARLI v0.96 (Zwickl 2006) and MrBayes v3.1.2. (Huelsenbeck & Ronquist 2001) to analyse the alignments. MP analyses included 100 replicates with random sequence additions and TBR branch swapping in effect. MP bootstrapping (Felsenstein 1985) was performed based on 2 000 replicates with the same settings as for the



**Fig. 1** Locations of the new primers for *Mcm7* and *Tsr1* using *Aspergillus nidulans* mRNA (XM\_658504 and XM\_658778) as reference sequence. Shaded areas in *Tsr1* indicate regions of high sequence variability.

**Table 4** Taxa used to test the fit of the new primers in silico.

Taxon	<i>Mcm7</i>	<i>Tsr1</i>
<i>Saccharomycotina</i>		
<i>Ashbya gossypii</i>	NP_984137	NP_984911
<i>Kluyveromyces lactis</i>	XP_454998	XP_454177
<i>Saccharomyces cerevisiae</i>	NP_009761	NP_010223
<i>Yarrowia lipolytica</i>	XP_501070	XP_500653
<i>Taphrinomycotina</i>		
<i>Schizosaccharomyces pombe</i>	NP_596545	NP_593391
<i>Basidiomycota</i>		
<i>Coprinopsis cinerea</i>	EAU88865	EAU91047
<i>Cryptococcus neoformans</i>	XP_571487	XP_570891
<i>Ustilago maydis</i>	EAK87259	EAK85759




MP search. Likelihood analyses were run using the GTR+I+G model and default settings in GARLI. For Bayesian analyses we partitioned the dataset into three parts (each codon position) and each partition was allowed to have its own parameter values (Nylander et al. 2004). No molecular clock was assumed, and no interpartition rate heterogeneity was allowed. Heating of the chains was set to 0.2. A run with 3 000 000 generations starting with a random tree and employing 4 simultaneous chains was executed for the individual datasets. Every 100th tree was saved into a file. The first 300 000 generations (i.e. the first 3 000 trees) were deleted as the 'burn in' of the chain. For the combined alignment dataset we executed a run with 6 000 000 generations and deleted the initial 600 000 generations (i.e. the first 6 000 trees). We plotted the log-likelihood scores of sample points against generation time using TRACER v1.0 (<http://tree.bio.ed.ac.uk/software/tracer/>) to ensure that stationarity was achieved after the first 300 000 (600 000 for the combined alignment dataset) generations by checking whether the log-likelihood values of the sample points reached a stable equilibrium value (Huelsenbeck & Ronquist 2001).

Additionally, we used AWTY (Nylander et al. 2008) to compare splits frequencies in the different runs and to plot cumulative split frequencies to ensure that stationarity was reached. We calculated a majority rule consensus tree with average branch lengths of the remaining 54 000 trees (27 000 from each of the parallel runs) using the sumt option of MrBayes. For the combined alignment dataset the majority rule consensus tree consisted of 108 000 ( $2 \times 54\,000$ ) trees from the stationarity phase. Posterior probabilities were obtained for each clade. Clades with posterior probabilities  $\geq 0.95$  were considered as strongly supported. Phylogenetic trees were visualised using the program Treeview (Page 1996).

## RESULTS

We report 84 new sequences of *Mcm7* (MS456) and *Tsr1* (MS277) for 42 lichenised ascomycetes belonging to the classes *Eurotiomycetes*, *Lecanoromycetes* and *Lichinomycetes* (Table 1). PCR success rates for our newly developed primers were highest for the primer combination *Mcm7*-709for/*Mcm7*-



**Fig. 2** Phylogeny of *Pezizomycotina* (*Ascomycota*) based on a combined alignment of *Mcm7* (MS456) and *Tsr1* (MS277) sequences. Total alignment length is 1203 bp. This is a 50 % majority rule consensus tree based on a sampling of 108 000 B/MCMC trees. **Bold** branches indicate posterior probabilities  $\geq 0.95$ . Numbers above branches are maximum parsimony bootstrap support values  $\geq 70$  based on 2 000 random addition replicates.

**Table 5** *Mcm7* and *Tsr1* sequence and alignment properties.

	<i>Mcm7</i> (MS456)	<i>Tsr1</i> (MS277)
Introns	None	some (length: 189–272 bp)
Total alignment length (bp)	573	827
Hypervariable (excluded) sites	None	198
Variable sites	357/573 (62.3 %)	489/629 (77.7 %)
Constant sites	216/573 (37.7 %)	140/629 (22.3 %)
Within-genus sequence variation (p-distances) excluding hypervariable sites:		
<i>Malcolmiella</i> (11 OTUs)	0.0055–0.2227	0.0332–0.2193
<i>Aspergillus</i> (7 OTUs)	0.0230–0.2307	0.0357–0.3076
<i>Lecanora</i> (6 OTUs)	0.0377–0.2756	0.0226–0.4148

1348rev ( $\pm 80\%$ ), while *Mcm7*-709for/*Mcm7*-1447rev worked in  $\pm 50\%$  of the attempted PCRs, and the *Tsr1* primers in  $\pm 40\%$ . Multiple bands were sometimes present when we used the primer combinations *Mcm7*-709for/*Mcm7*-1447rev and *Tsr1*-1459for/*Tsr1*-2308rev. *Tsr1*-1453for is a modification of *Tsr1*-1459for that we used under the same annealing conditions. We used the *Aspergillus nidulans* mRNA sequences of *Mcm7* (XM\_658504) and *Tsr1* (XM\_658778) as references for the locations of our primers. The full length genomic DNA sequences of *Aspergillus nidulans* *Mcm7* and *Tsr1* contain 1–2 introns of  $\pm 60$  bp length, which, however, do not overlap with the sequence fragments amplified by primers developed in this study. We found introns (length: 189–272 bp) with characteristic GT-intron-AG splice sites near the reverse primer (*Tsr1*-2308rev) in *Tsr1* in three *Lecanora* species. Two hypervariable regions containing many gaps (*Tsr1*: positions 198–221 and 518–628) were excluded from the phylogenetic analysis. The *Mcm7* alignment contained no gaps and no ambiguously aligned regions. Properties of the sequences and alignments are summarized in Table 5. We performed parsimony bootstrap analyses on each individual dataset, and examined 75 % bootstrap consensus trees for conflict (Lutzoni et al. 2004). We used the program Modeltest v3.7 (Posada &

**Table 6** Comparison of phylogenetic analyses (MP, ML, B/MCMC) between single and combined datasets.

	<i>Mcm7</i> (MS456)	<i>Tsr1</i> (MS277)	Combined
MP tree length	3537	4606	8200
Number of MP trees	1	12	8
Consistency Index (CI) excluding uninformative sites	0.195	0.216	0.205
# of nodes supported by bootstrap $\geq 70$ in MP analyses (based on 2 000 replicates)	23	30	37
ML score using GTR+I+G (GARLI)	-13732	-18424	-32262
# of nodes supported by PP $\geq 95$ in B/MCMC analyses	36	38	44

Crandall 1998) to determine the nucleotide substitution model that best fit our data. For both datasets the program selected the GTR+I+G model.

The tree topologies obtained from the single gene datasets resulting from MP, ML and Bayesian analyses did not show any strongly supported conflicts. Thus, we present only the B/MCMC tree of the combined analysis (Fig. 2). Statistical values and number of supported nodes obtained by MP, ML and Bayesian analyses of single and combined datasets are summarised in Table 6. The *Sordariomycetes* were used as outgroup. The classes *Sordariomycetes*, *Leotiomycetes*, *Eurotiomycetes* and *Lecanoromycetes* are monophyletic and highly supported (PP  $\geq 95$ ). *Lichinomycetes* is only represented by a single species, *Peltula euploca*. The phylogenetic estimate obtained from the combined analysis of *Mcm7* and *Tsr1* agrees with previously published phylogenies (Gargas et al. 1995, James et al. 2006). *Lecanoromycetes* form a supported sister group relationship with *Eurotiomycetes*. Basal to this are *Lichinomycetes* and *Leotiomycetes*. Within *Lecanoromycetes*, the subclasses *Lecanoromycetidae* and *Ostropomycetidae* form supported groups, while the genus *Umbilicaria* is in an unsupported position at the base of *Lecanoromycetes*. Within

<i>Mcm7</i> -709for	ACIMGIGTITTCVGYGTHAARCC	<i>Mcm7</i> -1348rev	ATGGGWGAYCCIGGIGTIGCHAARTC
Ashbya	ACTAGGATATCTGACGTTAAGCC	Ashbya	ATGGGTGATCCTGGTGTGGCCAAGTC
Kluyveromyces	ACCAGAGTCTCTGATGTGAAGCC	Kluyveromyces	ATGGGTGATCCTGGTGTGGCTAAATC
Saccharomyces	ACCAGAGTTCTGATGTCAAACC	Saccharomyces	ATGGGTGATCCTGGTGTGGCCAATC
Yarrowia	ACACGAGTTTCTGATGTCAAAGCC	Yarrowia	ATGGAGATCCAGGTGTGGCCAAGTC
Schizosacch.	ACTCGTACAGTATGTAAAGCC	Schizosacch.	ACTGGTATCCTGGTGTGGCAAATC
Coprinopsis	ACGCGCGTCTCAGAAGTCAAACC	Coprinopsis	ATGGGTGATCCTGGTGTGGCCAATC
Cryptococcus	ACCCGTGTTTCTGAAGTCAAAGCC	Cryptococcus	ATGGGTGATCCTGGTGTGGCCAATC
Ustilago	ACGCGCGTCTCCGAGGTAAAGCC	Ustilago	ATGGGGATCCTGGTGTGGGAAATC
<i>Mcm7</i> -1447rev	GGYCTYACIGCIGCIGTIATG	<i>Tsr1</i> -1453for	GARTTCCCIGAYGARATYGARCT
Ashbya	GGGTTGACGGCCGCGTTATG	Ashbya	GAGTTCCTCAGATGAGATTGAACT
Kluyveromyces	GGTCTAACAGCCGCGTTATG	Kluyveromyces	GAATTTCTGACGAAATGAACT
Saccharomyces	GGTCTGACCCGCTGCCGTCATG	Saccharomyces	GAGTTCCTCAGATGAGATTGAACT
Yarrowia	GGTCTTACAGCAGCTGTGATG	Yarrowia	GATTTCCCTGATGAACTGCAACT
Schizosacch.	GGTTTAACTGCTGCTGTAATG	Schizosacch.	GAATTTCCCTGATGAGATTGAACT
Coprinopsis	GGACTCACTGCTGCTGTCATG	Coprinopsis	GGTTTCCCTGATGAACTGCAACT
Cryptococcus	GGTTTACAGCCGCGTTATG	Cryptococcus	ATGTTCCCTGATGAACTGCAACT
Ustilago	GGGTTGACGGCAGCAGTATG	Ustilago	GAGTTCCTGAGCAGATTGACAC
<i>Tsr1</i> -1459for	CCIGAYGARATYGARCTICAYCC	<i>Tsr1</i> -2308rev	GGIACICAYGGITAYTTYAAG
Ashbya	CCAGATGAGATTGAACTGGATCC	Ashbya	GGTACTCATGGTACTTCAAG
Kluyveromyces	CCTGACGAAATTGAACTTGACCC	Kluyveromyces	GGTACTCATGGTACTTCAAG
Saccharomyces	CCCGATGAGATCGAACTAGAGCC	Saccharomyces	GGTACTCATGGTACTTCAAG
Yarrowia	CCTGATGAAGTCAAGTCAAGCC	Yarrowia	GGTACTCATGGTACTTCAAG
Schizosacch.	CCCGATGAGGTAGAGCTTCAACC	Schizosacch.	GGTACTCATGGTACTTCAAG
Coprinopsis	CCGATGAAGTCAAGTCAAGCC	Coprinopsis	GGTACTCATGGTACTTCAAG
Cryptococcus	CCGACGAAAGTCAAGTCAAGCC	Cryptococcus	GGTACTCATGGTACTTCAAG
Ustilago	CCGACGAAAGTCAAGTCAAGCC	Ustilago	GGTACTCATGGTACTTCAAG

**Fig. 3** Comparison of the new primers to homologous sequences in *Saccharomycotina* (*Ashbya*, *Kluyveromyces*, *Saccharomyces*, *Yarrowia*), *Taphrinomycotina* (*Schizosaccharomyces*) and *Basidiomycota* (*Coprinopsis*, *Cryptococcus*, *Ustilago*). 100 % matches between primer sequence and gene sequences studied are indicated by grey shading. High sequence similarities indicate that the new primers are likely to fit in members of the analysed groups. Some primer sequences may require slight modifications.

*Eurotiomycetes*, *Eurotiomycetidae* and *Chaetothyriomycetidae* form supported clades. We included multiple species/strains of the genera *Aspergillus* (7), *Lecanora* (6), and *Malcolmiella* (11) to assess within-genus variation of the analysed loci, as well as resolution power at low taxonomic levels. Genetic distances within *Aspergillus*, *Lecanora* and *Malcolmiella* are reported in Table 5. Each of these genera forms a supported monophyletic clade with high internal resolution and support (Fig. 2).

We aligned selected members of *Saccharomycotina*, *Taphrinomycotina* and *Basidiomycota* (Table 4) with our datasets and compared the new primer sequences to the corresponding positions in these taxa. The low number of mismatches suggests that the new primers will need no adjustments or only minor modifications to also fit these phylogenetic groups (Fig. 3).

## DISCUSSION

We developed new degenerate primers, which amplify fragments of the single-copy protein-coding genes *Mcm7* and *Tsr1* in *Pezizomycotina*. Our study confirms that *Mcm7* and *Tsr1* are suitable loci for the reconstruction of phylogenetic relationships among fungi (Aguileta et al. 2008). We were able to obtain sequences from representatives of 5 classes and 11 orders of euascomycetes, demonstrating the ability of the primers to amplify a wide range of unrelated taxa. Additionally we tested primer fit in silico using members of *Saccharomycotina*, *Taphrinomycotina* and *Basidiomycota* and found that the new primers can be used for these groups as well, possibly with slight sequence modifications.

Our analyses within *Pezizomycotina* show that *Mcm7* and *Tsr1* are able to resolve large scale as well as fine scale phylogenetic relationships. The sequences are alignable across a wide range of unrelated taxa and at the same time have sufficient variability to resolve within-genus relationships (Table 5). This property sets the new loci apart from commonly used ribosomal markers, such as ITS or mtSSU, which also have the power to resolve lower level phylogenetic relationships, but may yield ambiguous and saturated alignments, when used to compare distantly related taxa. We predict that *Mcm7* and *Tsr1* have an even higher potential to resolve phylogenetic relationships between fungi when analyzed in combination with other routinely used datasets, such as 18S, 28S, *RPB1* and *RPB2*.

*Mcm7* and *Tsr1* are two relatively long (~ 2.5 kb) single-copy genes which can be aligned across major fungal lineages, such as *Ascomycota* and *Basidiomycota* (Aguileta et al. 2008). The fact that *Homo sapiens* sequences can be used as outgroups (Aguileta et al. 2008, www.systematicbiology.org, online Appendix 5) indicates that these loci might also be useful for phylogenetic studies involving fungi as well as non-fungal organisms.

**Acknowledgements** We thank Fabian Ernemann (Chicago) and Paul Nelson (St. Paul) for support with lab work and sequence editing. This study was supported by start-up funds to I.S. from the University of Minnesota, Student Research Funding to NAN from Augsburg College, NSF grants DEB-0516116 (PI: HTL) and DEB-0715660 (PI: Robert Lücking) to The Field Museum, and the Spanish Ministry of Science and Innovation through a Ramon y Cajal grant (RYC02007-01576) to PKD. We wish to thank James Lendemer (New York), Robert Lücking (Chicago), and Zdenek Palice (Praha) for allowing us to use their collections for DNA isolation. Several of the new sequences were generated in the Pritzker Laboratory at the Field Museum.

## REFERENCES

Aguileta G, Marthey S, Chiapello H, Lebrun MH, Rodolphe F, Fournier E, Gendraud-Jacquemard A, Giraud T. 2008. Assessing the performance of single-copy genes for recovering robust phylogenies. *Systematic Biology* 57: 613–627.

Binder M, Hibbett DS. 2002. Higher-level phylogenetic relationships of homobasidiomycetes (mushroom-forming fungi) inferred from four rDNA regions. *Molecular Phylogenetics and Evolution* 22: 76–90.

Crespo A, Lumbsch HT, Mattsson JE, Blanco O, Divakar PK, Articus K, Wiklund E, Bawingan PA, Wedin M. 2007. Testing morphology-based hypotheses of phylogenetic relationships in Parmeliaceae (Ascomycota) using three ribosomal markers and the nuclear *RPB1* gene. *Molecular Phylogenetics and Evolution* 44: 812–824.

Felsenstein J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39: 783–791.

Gargas A, Depriest PT, Grube M, Tehler A. 1995. Multiple origins of lichen symbioses in Fungi suggested by SSU rDNA phylogeny. *Science* 268: 1492–1495.

Gelperin D, Horton L, Beckman J, Hensold J, Lemmon SK. 2001. *Bms1p*, a novel GTP-binding protein, and the related *Tsr1p* are required for distinct steps of 40S ribosome biogenesis in yeast. *Rna* a Publication of the Rna Society 7: 1268–1283.

Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.

Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, et al. 2007. A higher-level phylogenetic classification of the Fungi. *Mycological Research* 111: 509–547.

Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755.

Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP. 2001. Evolution – Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* 294: 2310–2314.

James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, et al. 2006. Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature* 443: 818–822.

Kearsey SE, Labib K. 1998. MCM proteins: evolution, properties, and role in DNA replication. *Biochimica et Biophysica Acta* 1398: 113–136.

Landvik S, Eriksson OE, Berbee ML. 2001. Neolecta – a fungal dinosaur? Evidence from beta-tubulin amino acid sequences. *Mycologia* 93: 1151–1163.

Larget B, Simon DL. 1999. Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Molecular Biology and Evolution* 16: 750–759.

Liu YJ, Hall BD. 2004. Body plan evolution of ascomycetes, as inferred from an RNA polymerase II, phylogeny. *Proceedings of the National Academy of Sciences* 101: 4507–4512.

Lumbsch HT, Schmitt I, Mangold A, Wedin M. 2007. Ascus types are phylogenetically misleading in Trapeliaceae and Agyriaceae (Ostropomycetidae, Ascomycota). *Mycological Research* 111: 1133–1141.

Lumbsch HT, Schmitt I, Palice Z, Wiklund E, Ekman S, Wedin M. 2004. Supraordinal phylogenetic relationships of Lecanoromycetes based on a Bayesian analysis of combined nuclear and mitochondrial sequences. *Molecular Phylogenetics and Evolution* 31: 822–832.

Lutzoni F, Kauff F, Cox CJ, McLaughlin D, Celio G, et al. 2004. Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *American Journal of Botany* 91: 1446–1480.

Moir D, Stewart SE, Osmond BC, Botstein D. 1982. Cold-sensitive cell-division-cycle mutants of yeast: Isolation, properties, and pseudoreversion studies. *Genetics* 100: 547–563.

Nylander JAA, Ronquist F, Huelsenbeck JP, Nieves-Aldrey JL. 2004. Bayesian phylogenetic analysis of combined data. *Systematic Biology* 53: 47–67.

Nylander JAA, Wilgenbusch JC, Warren DL, Swofford DL. 2008. AWTY (are we there yet?): a system for graphical exploration of MCMC convergence in Bayesian phylogenetics. *Bioinformatics* 24: 581–583.

Page RDM. 1996. TreeView: An application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357–358.

Posada D, Crandall KA. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14: 817–818.

Reeb V, Lutzoni F, Roux C. 2004. Contribution of *RPB2* to multilocus phylogenetic studies of the euascomycetes (*Pezizomycotina*, Fungi) with special emphasis on the lichen-forming *Acarosporaceae* and evolution of polyspory. *Molecular Phylogenetics and Evolution* 32: 1036–1060.

Swofford DL. 2003. PAUP\* phylogenetic analysis using parsimony and other methods, v4.0b10. Sinauer Associates, Sunderland, Massachusetts, USA.

Zwickl DJ. 2006. Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. PhD dissertation The University of Texas at Austin, USA.