

REVIEW ARTICLE OPEN



The impact of the lung environment on macrophage development, activation and function: diversity in the face of adversity

Calum C. Bain^{1,3} and Andrew S. MacDonald^{2,3}

© The Author(s) 2022

The last decade has been somewhat of a renaissance period for the field of macrophage biology. This renewed interest, combined with the advent of new technologies and development of novel model systems to assess different facets of macrophage biology, has led to major advances in our understanding of the diverse roles macrophages play in health, inflammation, infection and repair, and the dominance of tissue environments in influencing all of these areas. Here, we discuss recent developments in our understanding of lung macrophage heterogeneity, ontogeny, metabolism and function in the context of health and disease, and highlight core conceptual advances and key unanswered questions that we believe should be focus of work in the coming years.

Mucosal Immunology (2022) 15:223–234; <https://doi.org/10.1038/s41385-021-00480-w>

INTRODUCTION

Macrophages are largely sessile, tissue dwelling phagocytes that are present in every organ of the body. The biology of macrophages has been studied for well over one hundred years since being described by Elie Metchnikoff at the end of the nineteenth century. However, arguably, it is the last 10–15 years that has seen major revisions of key concepts in macrophage biology, from nomenclature of subsets and activation states to the ontogeny of these cells. For instance, while macrophages have historically been described as M1 (“classically-activated”) or M2 (“alternatively-activated”) largely based on *in vitro* culture systems¹, it is now beyond doubt that a binary classification such as this is inadequate to capture the complexity of macrophage plasticity and activation states, particularly *in vivo*². Technological advances have revealed tremendous diversity and heterogeneity between macrophages from different tissues and even within different niches of the same tissue in terms of phenotype, transcriptome and metabolome^{3,4}. While macrophages generally excel at the ‘silent’ clearance of debris, apoptotic host cells and the capture and destruction of microbial intruders, it is now clear they also play far broader roles and are exquisitely tailored to meet the demands of the local tissue microenvironment in which they reside. Indeed, the *in vivo* tissue environment may be the major determinant governing macrophage development, recruitment, activation and function, highlighting the importance of careful consideration of the distinctive properties that different tissues possess, in steady state and during inflammation, to fully understand the role of macrophages in different locations throughout the body.

In this article, we review recent developments in the understanding of macrophage heterogeneity, ontogeny and function in

lung health and during inflammation, immunity and tissue repair. Using pulmonary fibrosis as an example, we discuss how dysregulated macrophage behaviour can contribute to lung pathology and how mouse models have revealed the complexity of the macrophage response to lung injury. Along the way we identify key areas that we believe warrant further investigation.

DEFINING PULMONARY MACROPHAGES IN THE HEALTHY LUNG

Macrophages are distributed throughout the lung and can broadly be divided into those present in the airways/alveoli and those in the tissue interstitium/parenchyma. While it has been clear for decades that pulmonary macrophages exist in both airways and tissues, it is only recently that we have come to appreciate the degree of heterogeneity and diversity between different macrophage subsets in each location. In particular, the advent of single cell technologies, such as single cell RNA sequencing (scRNA-seq), has allowed tissue macrophage heterogeneity to be assessed in a completely unbiased manner across several species^{5–11}.

In mice, alveolar macrophages (AlvMφs; see Box 1) in the healthy lung are defined by their high and uniform expression of CD11c, SiglecF and CD169 (Siglec1; sialoadhesin), and lack of CD11b expression^{12–14}. Their residence in the airways can be confirmed by performing bronchoalveolar lavage (BAL) where they are the sole macrophage population in health¹⁵, although it is important to note that this method only retrieves a fraction of the macrophages resident in the airways. Murine interstitial macrophages (IntMφs) express high levels of CD11b, but lack expression of SiglecF^{12–14}. Compared with AlvMφs, considerably less is known about the IntMφ compartment, likely reflecting the fact that these macrophages are

¹The University of Edinburgh Centre for Inflammation Research, Queen’s Medical Research Institute, Edinburgh Bioquarter, Edinburgh EH16 4TJ, UK. ²Lydia Becker Institute of Immunology and Inflammation, University of Manchester, Manchester M13 9NT, UK. ³These authors contributed equally: Calum C. Bain, Andrew S. MacDonald.

✉email: calum.bain@ed.ac.uk; andrew.macdonald@manchester.ac.uk

Received: 19 October 2021 Revised: 4 December 2021 Accepted: 18 December 2021

Published online: 11 January 2022

Box 1. What's in a name?

In order to discuss pulmonary macrophage subsets, some clarity on nomenclature is needed. Historically, macrophages found in the airways/bronchoalveolar lavage fluid (BALF) of naïve mice have been termed 'alveolar macrophages', which express high levels of CD11c, SiglecF and CD169 while lacking CD11b expression. These airway/alveolar macrophages are phenotypically and transcriptionally distinct from those found in the lung tissue, which express CD11b and variable levels of CD11c, but lack SiglecF expression and are often termed as 'lung tissue', 'interstitial' or 'parenchymal' macrophages. It is now clear that, during inflammation, macrophages with a tissue/interstitial phenotype can be readily found in the airways/BALF. Thus, for clarity, in this review we will use 'alveolar' only to refer to airway macrophages (AMφs) that are CD11c^{hi}SiglecF⁺ (AlvMφs), and 'interstitial' to refer to CD11b⁺SiglecF⁻ lung tissue macrophages (IntMφs). We will use the term monocyte-derived AMφs to refer to macrophages in the BALF that have been shown to arise in the context of inflammation and exhibit some, but not all, features of bona fide AlvMφs. We will use the term 'ex-interstitial' (ex-IntMφ) to try to make clear when we are referring to cases where macrophages with this phenotype are found in the airways. Since it is currently impossible to distinguish tissue vs airway origins of human pulmonary macrophages, we will simply refer to cells obtained by BAL as airway macrophages (AMφs) and those obtained from human lung tissue as tissue macrophages (tissue Mφs).

difficult to isolate from lung tissue using standard enzymatic protocols. Indeed, flow cytometric analysis of whole mouse lung digests suggests that IntMφs form a small fraction of the overall macrophage compartment in health, with AlvMφs outnumbering them by 5–10-fold^{15,16}. Whether this is also the case in humans is currently hard to determine with certainty, as no comparable set of surface markers has yet been identified to unequivocally distinguish human AlvMφs from IntMφs in BAL, sputum or in lung tissue digests. Nevertheless, limited fluorescence microscopy of mouse and human lung, which circumvents the need to dissociate solid tissue, suggests that tissue (IntMφs) are more abundant than suggested by flow cytometry^{17–19}. Thus, the notion that these cells represent a minor macrophage population should be reconsidered (Box 2).

It is also important to note that due to the overlapping expression of CD11c, CD11b and SiglecF by other myeloid cells, in particular CD11c⁺ dendritic cells (DCs) and SiglecF⁺CD11b⁺ eosinophils, expression of these surface markers alone is not sufficient for the characterisation of murine lung macrophages. Instead, a more rigorous approach is needed to define bona fide murine pulmonary macrophage subsets, with a growing consensus that the optimal strategy is by their co-expression of the high affinity FcγR1 (CD64) and Mer-tyrosine kinase (MerTK), a key efferocytic receptor^{12,15,20}. In health, AlvMφs and IntMφs can then be identified amongst the CD64⁺MerTK⁺ fraction by their distinct CD11c/CD11b profiles and other phenotypic traits (Fig. 1). As discussed below, distinction between these anatomically distinct cells becomes less apparent when homeostasis is perturbed. Moreover, recent work has shown that some DCs can acquire expression of CD64 in certain contexts²¹, emphasising the need for multi-parameter analysis when characterising these cells. Notably, neither AlvMφs nor IntMφs can be defined using the M1/M2 nomenclature system. Indeed, in the healthy lung both populations co-express markers historically considered "M1" and "M2" specific^{14,15}. For instance, murine AlvMφs constitutively co-express CD11c and Ym-1 (encoded by *Chil3*) which have been used by some as defining features of so-called "M1" and "M2" macrophages^{22,23}. However, Ym-1 expression by homeostatic AlvMφs is independent of IL-4–IL-4R signalling¹⁵, the axis controlling "alternative" activation of macrophages, and CD11c expression is independent of exposure to microbial products or inflammatory cytokines thought to drive so-called "M1" polarisation^{24,25}. This highlights the inadequate nature of the M1/M2 nomenclature in defining macrophages in vivo and that as a field we need to abandon using it²⁶.

Recent studies employing scRNA-seq have begun to reveal additional heterogeneity in the murine pulmonary macrophage compartment. Whereas AlvMφs appear to be relatively

Box 2. Key Challenges

1. Establish "signatures" to faithfully distinguish human AlvMφs from IntMφs and ex-IntMφs
2. Develop methods of lineage tracing in human lung macrophage subsets
3. Reduce reliance on in vitro functional assays of questionable relevance and develop models/systems/approaches that better represent the lung environment
4. Reach consensus on the heterogeneity, function and spatial distribution of IntMφs
5. Develop novel transgenic murine models to selectively target AlvMφs vs IntMφs vs ex-IntMφs and understand the molecular levers that control their differentiation
6. Transition to using mouse models with microbial exposure/status more aligned with that encountered by humans.

homogeneous⁶, the IntMφ compartment harbours at least two distinct subsets defined by their expression of CD206 (mannose receptor)^{14,18,27}. The anatomical locale in which each of these subsets are found remains contentious. CD206⁻ IntMφs that mostly express MHCII⁺ have been suggested to be enriched in the interstitial tissue adjacent to the alveoli, whereas their CD206⁺ MHCII⁻ counterparts appear more numerous in the interstitium surrounding the bronchi¹⁸. Multi-parameter fluorescence microscopy has also started to elucidate the nature of the niche in which these subsets may exist. For instance, CD206⁺MHCII⁺ IntMφs can be found to interact with nerves, whereas CD206⁺MHCII⁻ cells may occupy a perivascular niche²⁷. Transcriptional profiling has revealed other useful markers for the identification of these subsets, including Lyve-1, folate receptor beta (FRβ) and CD36, which appear to be more highly expressed by perivascular CD206⁺ MHCII⁻ macrophages. Work by the Khanna group identified a population of CD169-expressing IntMφs²⁸, although it seems these largely overlap transcriptionally with the CD206⁻MHCII⁺ IntMφs identified by others. CX3CR1 has also been suggested as a defining marker of these subsets²⁷, although other studies have shown that CX3CR1 is expressed equally across these subsets^{14,18,29}. Thus, while it is clear that the murine IntMφ compartment is heterogeneous, consensus is yet to be reached on the best strategy to define these cells.

The identity of macrophage subsets in the human lung has also become clearer recently. Human AMφs express high levels of HLA-DR, CD11b and CD206 together with CD169 and MARCO^{7,13,30}. CD163 appears to define two subsets of AMφ in lung tissue from humans and non-human primates. However, whereas CD163^{hi} AMφs are abundant in BALF, CD163^{lo} AMφs are relatively rare¹³, questioning whether they truly represent AlvMφs. Importantly, while SiglecF has become somewhat of a de facto marker of murine AlvMφs, its human paralog, Siglec8, is absent from AMφs in man¹³. Despite only partial conservation in terms of phenotype between mice and humans, recent transcriptional profiling suggests that these cells share a core gene signature, including expression of *PPARG*, *FABP4*, *FFAR4*, *FN1*^{7,31–34} (Fig. 1). The phenotypic and transcriptional identity of human IntMφs during health is limited by access to healthy lung tissue. Nevertheless, they appear to be defined as HLA-DR⁺CD11b⁺CD36⁺ cells lacking CD169 expression¹³, a phenotype that has recently been confirmed using humanised mice³⁵. Moreover, heterogeneity similar to that seen in mice is likely to exist amongst human IntMφs²⁷, with one study suggesting the intensity of HLA-DR and CD36 a defining characteristic⁷.

MACROPHAGE FUNCTION IN HEALTH

As immunologists, we tend to consider the primary function of macrophages is to act as the first line of defence against pathogens. Indeed, the positioning of macrophages in the airways




Fig. 1 Heterogeneity, phenotypic profiles and functions of macrophages in the healthy lung. The lung macrophage compartment is heterogeneous, with at least two populations occupying distinct anatomical niches in the healthy lung. Macrophages are present in the bronchoalveolar space, including the alveoli where gaseous exchange occurs. Alveolar macrophages (AlvMφs) are defined by their expression of CD11c, MARCO and CD169 in both mice and humans, although additional species-specific markers must be used to define them accurately. AlvMφs are crucial for regulating surfactant produced by the respiratory epithelium as well as maintaining epithelial integrity and responsiveness. Their high phagocytic capability allows them to clear apoptotic/senescent cells and inhaled particles efficiently. They also act as the first line of defence against air-borne pathogens, although the relative role of resident AlvMφs versus elicited, monocyte-derived macrophages in immune protection varies depending on the nature of the insult (see text and Fig. 3). Macrophages are also found in the interstitial space between the alveoli and the capillary beds, as well as surrounding larger airways (bronchi). These interstitial macrophages (IntMφs) are phenotypically distinct from AlvMφs and at least two subsets exist in mouse and man defined by differential expression of MHCII (HLA-DR), Lyve-1 and/or CD36. IntMφs may act as a second line of defence under the epithelial barrier and basement membrane. In health, they may support the stromal/structural compartment through growth factor supply, as well as maintaining T cells and acting a rich source of IL-10. Although nerve- and blood vessel (BV)-associated IntMφs have been described, whether these represent obligate niches is under debate.

means that they will be one of the first cells to encounter airborne pathogens and altering the ability of airway macrophages to detect, engulf and kill respiratory pathogens leads to increased susceptibility to a variety of bacteria, viruses and fungi^{36–41}. However, as described below, to avoid excessive and potentially harmful pro-inflammatory responses being mounted against environmental antigens encountered by airway macrophages, these cells are held in a state of relative hyporesponsiveness via a range of powerful inhibitory mechanisms⁴². In the absence of infection, the principal role of airway macrophages is the regulation of pulmonary surfactant, the lipid-protein complex produced by the respiratory epithelium to lubricate the lungs and allow frictionless expansion/contraction. Consistently, the AlvMφ transcriptional signature is dominated by genes associated with lipid metabolism in mouse and man^{7,43,44}. Macrophages are indispensable for this function, as spontaneous pulmonary alveolar proteinosis (PAP) develops in mice and humans with absent or dysfunctional AlvMφs^{45–50}. Likewise, dead, dying and senescent cells accumulate in the absence of functional airway macrophages, demonstrating their key efferocytic role⁴³. Airway macrophages may also maintain the integrity and responsiveness of the respiratory epithelium. For instance, production of immunoregulatory cytokines, such as TGFβ and IL-10, may modify epithelial cell function through regulation of ion and fluid transport^{42,51} and AlvMφ-derived fibronectin may act as a proliferative factor for airway epithelial cells⁵². Suppressor of cytokine signalling (SOCS)-containing vesicles released from AlvMφs may regulate the responsiveness of the epithelium to e.g., TLR ligands^{53,54}.

Compared with AlvMφs, the mechanisms of regulation and homeostatic functions of IntMφs are poorly understood⁵⁵, although

it is clear that all murine IntMφ subsets are avidly phagocytic and can capture *E.coli* bioparticles *in vivo*^{14,18}, suggesting they may act as a second line of defence should the epithelial barrier be breached. Their constitutive production of IL-10 under normal physiological conditions, in both mouse and man^{18,29,56–58}, suggests an immunoregulatory role. It is likely this involves supporting regulatory T cells locally in the lung parenchyma, given that IntMφs are thought to be non-migratory⁵⁹. However, IntMφs may also alter T cell responses indirectly through IL-10-dependent modulation of DC migration and priming activity⁵⁶.

Exposure to bacterial CpG DNA leads to IntMφ expansion and augmentation of IL-10 production²⁹, suggesting these cells may be inherently anti-inflammatory. Indeed, *Cx3cr1*-mediated deletion of *Il10* leads to increases susceptibility to allergic asthma in mice⁶⁰. Production of platelet-derived growth factor (PDGF) by IntMφs suggests they may also support fibroblast and epithelial homeostasis⁶¹. Finally, given their occupation of distinct anatomical niches, it is intriguing to speculate they may differentially contribute to nerve and vascular endothelial cell homeostasis, similar to their counterparts in the gut wall⁶².

LUNG MACROPHAGE ONTOGENY

Macrophage origins in health

Historically, macrophages were thought to be part of a linear mononuclear phagocyte system where tissue macrophages were continually replaced by blood monocytes, which themselves are replaced by dedicated bone marrow progenitors⁶³. However, over the last 10 years there has been a conceptual revolution in our understanding of macrophage ontogeny with the discovery that many tissue macrophages derive from embryonic progenitors and




Fig. 2 Pulmonary macrophage ontogeny during health. The contribution of distinct progenitors to the pulmonary macrophage compartments is highly dynamic and alters with age. During embryonic development (in mice) yolk sac-derived macrophages colonise the lung and these remain present at birth. However, these are outnumbered by foetal liver-derived progenitors that enter the lung prior to birth, some of which move into the airways upon alveolarization within the first days of life. During the neonatal period, where there is massive tissue growth, all macrophages show high levels of proliferation to occupy the newly created niches. This is sufficient to expand the AlvMφ compartment with little, if any, contribution from bone marrow-derived monocytes. However, recent work has suggested that during adulthood under homeostatic conditions AlvMφs are replenished, albeit at low rates, by bone marrow-derived, CCR2-dependent monocytes. These monocytes replace IntMφs at a higher rate, although in the unperturbed lung, the IntMφ compartment likely contains macrophages derived from the yolk sac, foetal liver and bone marrow, with the latter dominating numerically.

maintain themselves autonomously through in situ self-renewal^{64–74}. While some older studies had demonstrated the ability of macrophages to self-renew⁷⁵, it is the development of elegant lineage tracing models that has led to major advances in our understanding of macrophage origins. For instance, genetic fate mapping using mice with tamoxifen-inducible Cre recombinase under the control of the *Csf1r*, *Runx1*, *Cx3cr1* or *Tie2* promoters has shown that brain microglia derive from yolk sac progenitors and require little, if any, contribution from blood monocytes across the life of an animal^{64,65,67,69,76}. Using similar systems, it was shown that yolk sac progenitors contribute minimally to lung AlvMφs⁷⁷. Instead, tracing of foetal and adult haematopoiesis using *Flt3*^{Cre} mice shows haematopoietic stem cell (HSC)-derived cells make a major contribution to AlvMφs^{65,68,74}. This, combined with the fact that AlvMφs develop within the first few days of life in mice and humans in parallel with alveolarisation of the lung^{13,78,79}, led to the idea that they derive predominantly from foetal monocytes. That AlvMφs are unaffected in adult monocytopenic *Ccr2*^{-/-} mice and show little exchange in the context of parabiosis or tissue protected bone marrow chimeric mice supported the notion that these cells self-maintain throughout adult life in the absence of inflammation or infection^{68,78}. Moreover, analysis of AMφ longevity in the human context supported these observations in mice. By analysing macrophages obtained by transbronchial biopsies of recipients of sex mismatched lung transplants in a longitudinal manner, Eguiluz-Garcia *et al.* showed that the majority of AMφs remain of donor origin in this setting, suggesting human AMφs maintain themselves autonomously in situ⁸⁰, a finding supported by an independent study analysing AMφs obtained by bronchoalveolar lavage⁸¹.

However, several recent studies have started to challenge this model. First, longitudinal analysis of *Flt3*^{Cre}-*Rosa26*^{LSL-YFP} reporter mice showed increases in labelling of AlvMφs over time, indicative of age-dependent contribution of HSC-derived cells to the AlvMφ pool, a phenomenon not seen in brain microglia⁶⁵. This highlights the need for longitudinal analysis when considering macrophage dynamics, something that was not always performed in early lineage tracing studies^{64,74}. Indeed, longitudinal analysis of *Ms4a3*^{Cre} reporter mice, which allow tracing of all cells deriving from bone marrow granulocyte-monocyte progenitors (GMPs),

supports the idea that AlvMφs require replenishment from bone marrow-derived monocytes over the life course⁷¹ (Fig. 2). These data are consistent with recent work assessing AlvMφ turnover using so-called 'MISTRG' humanised mice, which have genes encoding human M-CSF (also known as CSF-1), GM-CSF (also known as CSF-2), IL-3 and thrombopoietin 'knocked-in' to their respective mouse loci to support human myeloid cell development, as well as a transgene encoding human SIRPα to prevent engulfment and destruction of human cells^{35,82}. Moreover, the idea of AlvMφ replenishment by monocytes is supported by a recent study using scRNA-seq to determine AMφ longevity in the context of sex-mismatch lung transplants where the majority of donor AMφs appear to be replaced by recipient cells⁸³. Why different studies using transplanted tissue reach discordant conclusions is unclear, but could reflect differences in methodologies used, for instance scRNA-seq versus fluorescence in situ hybridisation (FISH) for X/Y chromosomes, or the degree of injury caused by transplantation-related ischaemia and reperfusion. Clearly further work is warranted to clarify the dominant replenishment mechanisms underlying the homeostatic maintenance of AlvMφs.

The developmental origin of IntMφs has started to be unravelled in the past few years, although these cells have attracted much less attention than their AlvMφ counterparts. Genetic fate mapping indicates an initial contribution of yolk sac progenitors to the IntMφ pool^{65,77}, but these appear to be largely replaced, first by foetal liver-derived macrophages and then by HSC-derived macrophages during the early post-natal period^{27,65} (Fig. 2). CCR2-dependent bone marrow-derived cells continue to replenish IntMφs during adulthood, albeit at a low rate, and despite the heterogeneity described above, IntMφ subsets appear to display similar replenishment kinetics^{27,84}. Notably, despite clear evidence of progressive replenishment by monocytes, intact *Ccr2*^{-/-} mice have normal numbers of IntMφs^{18,29}, serving as a cautionary note that *Ccr2*^{-/-} mice cannot be used in isolation to determine the contribution of monocytes to tissue macrophage pools, at least in health. This suggests potential redundancy between chemokine receptors involved in monocyte navigation⁸⁵ and/or that compensatory mechanisms maintain macrophages in the context of monocytopenia. Indeed, both AlvMφs and IntMφs




Fig. 3 Pulmonary macrophage dynamics during inflammation and resolution. In inflammation caused by agents such as pathogens, pollutants or allergens, most resident AlvMφs are lost and replaced by monocyte-derived Mφs and perhaps ex-IntMφs. This occurs in parallel to accumulation of other inflammatory cells such as neutrophils and eosinophils, recruitment of which to the airways is facilitated by chemokines and disrupted barrier integrity. During resolution of the damage caused by acute inflammation, and/or in the face of chronic low-level inflammation, residual AlvMφs can self-renew through proliferation, clear up dying or dysfunctional cells in the airways, as well as be replenished through conversion of monocyte-derived macrophages and ex-IntMφs which are transcriptionally, epigenetically and functionally conditioned by the airway environment to take on AlvMφ identity.

can and do proliferate under normal physiological conditions^{15,68}. While this appears to be insufficient to maintain these populations, it may be sufficient in the absence of monocytes. It is also important to appreciate that self-renewal and derivation from monocytes are not mutually exclusive mechanisms of macrophage maintenance and, in certain contexts, monocyte-derived macrophages proliferate more readily than their embryonic counterparts^{73,86,87}. Replenishment by monocytes has been proposed to arise in response to niche availability⁸⁸, although the factors that govern 'availability' remain poorly understood and it is unclear if this differs in distinct subanatomical niches within the lung. Alternatively, there is evidence that different precursors have distinct metabolic states that may determine their long-term persistence in the lung, suggesting that cell-intrinsic properties may also govern replenishment kinetics⁸⁹.

Macrophage origin following injury, inflammation or infections

The mechanisms that govern macrophage expansion or repopulation following injury, infection or inflammation appear to be dependent on the nature of the insult. Acute inflammation or severe infection often leads to loss of tissue resident macrophages, a phenomenon described as the 'macrophage disappearance reaction' (Fig. 3). For instance, sterile inflammation induced by instillation of LPS in mice leads to a transient loss of AlvMφs and expansion of IntMφs in the tissue (unpublished observations). IntMφ expansion is impeded by *Ccr2* deficiency²⁹, suggesting a major role for monocyte recruitment in this process. In contrast, AlvMφ repopulation during inflammation resolution appears to rely exclusively on local proliferation⁷¹. In contrast, following a more substantial inflammatory insult, such as that induced by administration of bleomycin or silica to model lung fibrosis or infection with influenza, leads to replacement of resident AlvMφs with monocyte-derived cells^{90,91}. Consistent with this, monocyte-derived cells come to dominate the AMφ compartment in individuals with severe Sars-CoV-2 infection⁵. Whether this reflects direct effects of severe inflammation on the

self-renewal capacity of AlvMφs, or if severe inflammation leads to structural alterations, such as breakdown of basement membrane and epithelial integrity, is unknown. While classical monocytes can enter the airways in response to injury or infection, there is evidence that monocyte-derived, elicited macrophages in the interstitium may also transition to the airways. Whether these alternative differentiation routes influence the fate and function of these cells is currently unclear.

ENVIRONMENTAL IMPRINTING OF LUNG MACROPHAGES

The diversity of macrophages within the lung results from their plasticity and ability to respond to local environmental cues. In this regard, the pulmonary environment is unusual, even in comparison to other barrier sites, in terms of the wide range of environmental features it presents that can influence immune cell recruitment, activation and function, and that will have a particular impact on lumen-dwelling AlvMφs. These include the unique makeup of the airway fluids (predominantly composed of surfactant and mucins), commensals (bacteria, viruses and fungi) and nutrient levels (e.g., both host and microbial derived metabolites), all of which can change markedly during inflammation.

As well as acting as a lubricant, pulmonary surfactant constituents can also influence AlvMφ behaviour. For instance, in the absence of surfactant protein D (SP-D) AlvMφs display an unusual CD11b^{hi} phenotype and constitutively produce TNFα⁹². While phenotypic alterations are not evident in AlvMφs from naïve SP-A deficient mice⁹³, SP-A may amplify IL-4Rα-mediated AlvMφ activation while regulating their responsiveness to exogenous stimulation through direct interactions with Toll-like receptor 4 (TLR4) and MD-2⁹³⁻⁹⁵. As collectins, surfactant conformation can dictate their function in a context-dependent manner, with structural changes altering their binding to targets and in doing so conferring pro- vs anti-inflammatory ability⁹⁶.

The other major constituent of pulmonary fluids is mucus, as vital for lubricating the airways as surfactant, and also for entrapment

and ciliary clearance of inhaled particles and microbes⁹⁷. Additionally, mucus can be important for control of bacterial infection, regulation of hydration, resolution of inflammation, and modulation of immune and epithelial cell function^{98,99}. The main mucins expressed in the lung are Muc5b and Muc5ac, with Muc5b being essential for maintenance of healthy airways^{97,99}, while Muc5ac is up-regulated during inflammation^{100–102}. Each of these mucins has different properties that determine their dominant function. Mucus also provides a home for commensal lung microbes⁹⁷, which can exert a dramatic effect over airway function, for example through invasion/colonisation, consumption of nutrients, and production of metabolites – all of which can influence pulmonary inflammation and macrophage activation and function. Although most is currently known about bacteria and their products in this context, more widespread metagenomic approaches will increase our understanding of the abundance and diversity of bacterial, viral and fungal commensals in the lung, and how this changes during inflammation and disease.

In terms of nutrients, the airways present AlV ϕ s with one of the lowest glucose environments in the body, a tissue adaptation that has likely evolved to prevent outgrowth of glucose-hungry opportunistic bacteria^{103,104}. Indeed, epithelial cells lining the lung express high levels of glucose transporters apically¹⁰⁵, with which they can rapidly and efficiently reduce glucose levels in the airways, with the blood and tissues underlying the epithelium possessing over twelve times the levels of glucose than are found in the airway fluids^{103,105}. This low-glucose environment, coupled with abundant lipid-rich surfactant, likely plays a central role in governing AlV ϕ metabolism and function. Indeed, it is now clear that metabolism is a central factor in governing macrophage activation and function, with the general principle being that glycolysis may be more associated with 'type 1' macrophages, while lipid metabolism tends to be more typical of 'type 2' macrophages (reviewed by^{106,107}). However, the majority of the work that has established this paradigm has relied on *in vitro* assessment of bone marrow or monocyte-derived macrophages, with much less understanding of how tissue environments influence macrophage metabolism *in vivo*. In the context of the lung, we and others have shown that AlV ϕ s display a distinctive metabolic profile, expressing elevated levels of genes associated with lipid metabolism and peroxisome proliferator-activated receptor gamma (PPAR γ), and reduced expression of genes associated with glycolysis^{15,20}. Further, we have shown that glycolysis is a key determinant of AlV ϕ activation and function in type 2 inflammation¹⁵, suggesting that glycolytic ability, and availability of glucose, may be centrally involved in enabling AlV ϕ activation and function in the airways.

During inflammation, with compromised epithelial integrity, the balance of airway nutrients can change markedly, with reports of elevated airway glucose evident in a range of disease settings, including COPD, cystic fibrosis and asthma^{108–111}. This may be particularly relevant in chronic conditions where metabolite balance in the airways can be modified long-term, with consequent long-term alteration of AlV ϕ metabolism and function. In more acute settings, such mechanisms may enable a window of opportunity for glycolytic 'boosting' of AlV ϕ function that will reduce as the epithelium heals and metabolite levels recalibrate, in essence providing a metabolic rheostat for fine-tuning of AlV ϕ activation and function directly linked to how local substrate levels change in line with levels of tissue damage vs resolution.

In contrast to AlV ϕ s, less is currently known about metabolic control of IntM ϕ activation and function. IntM ϕ s residing in the more nutrient-rich environment of the lung tissue appear much more glycolytically active than their AlV ϕ counterparts, and are consequently more effective at producing the reactive oxygen species necessary for killing intracellular bacteria such as *Mycobacterium tuberculosis*^{20,112}. In this way, IntM ϕ s may be less susceptible to substrate-related functional control than AlV ϕ s,

though this likely changes if IntM ϕ s migrate into the airways in the context of inflammation as they adapt to this new environment. Indeed, a key outstanding question is to what extent does the metabolic status of monocyte-derived macrophages play in their differentiation to AlV ϕ s during and following an inflammatory insult. It is important, however, to remember that the methods to extract tissue macrophages can have profound effects on their biology, including their transcriptional and metabolic status. Given that IntM ϕ s are embedded in the tissue, it is plausible that some of the differences seen in their metabolic profile may reflect their response to extraction.

As described above, there is continual bi-directional crosstalk between macrophages and the structural/stromal cells that comprise their niche, and this crosstalk leads to niche-specific outcomes for macrophage recruitment, differentiation and function. Removing macrophages from their niche leads to phenotypic and transcriptional changes, directly demonstrating the need for continual crosstalk with structural cells³. In the airways, AlV ϕ s are highly dependent on GM-CSF for their development and maintenance^{78,113–117}, consistent with their high expression of GM-CSF receptor. Consequently, disruption to the GM-CSF-GM-CSFR axis leads to defective AlV ϕ differentiation and the development of PAP in both mice and humans^{45–50}. Recent work using GM-CSF reporter mice and cell-specific deletion of GM-CSF has identified alveolar type 2 epithelial cells as the indispensable source of GM-CSF during the pre- and post-natal stages of AlV ϕ development from foetal progenitors¹¹⁷. Interestingly, although innate lymphoid cells (ILCs) are major sources of GM-CSF in the steady state lung, haematopoietic deletion of GM-CSF does not affect AlV ϕ development¹¹⁷. Moreover, although it has been proposed that GM-CSF may control AlV ϕ s indirectly through induction of *Il6*, *Il13* and *Csf1* expression in lung basophils¹¹⁸, genetic depletion of basophils has little, if any, effect on AlV ϕ number or phenotype¹¹⁷. Indeed, that CD11c-mediated deletion of *Csf2rb*, which encodes one of the GM-CSFR subunits, or *Stat5*, which lies downstream of GM-CSFR, leads to aborted differentiation of AlV ϕ s^{119,120} supports the notion that GM-CSF acts directly on developing AlV ϕ s.

As mentioned above, crosstalk between AlV ϕ s and alveolar epithelial cells also involves the TGF β -TGF β R axis. TGF β is a potent immunomodulatory cytokine which is abundant in the mouse and human lung in health. It is produced in a latent form and must be converted to active TGF β to have biological effects. In the airways, integrin-mediated activation is thought to be the principal mechanism of TGF β activation. Specifically, the α v β 6 integrin, which is expressed by alveolar epithelial cells, is crucial for generating active TGF β ¹²¹. Consequently, genetic disruption of *Itgb6*, which encodes integrin β 6, leads to development of emphysema due to excessive production of MMP12 by dysfunctional AlV ϕ s; a phenotype that can be rescued by constitutive expression of TGF β 1^{121,122}. Human AM ϕ s are known to have a gene signature consistent with TGF β R signalling¹²³ and myeloid-specific deletion of TGF β R in mice leads to aborted AlV ϕ development, demonstrating a need for cell intrinsic TGF β R for this process^{124,125}. Interestingly, although many cells can produce TGF β , macrophages themselves are thought to be an important source^{123,124}. Moreover, AlV ϕ s can facilitate integrin-mediated release of active TGF β through production of amphiregulin, at least in the context of helminth infection¹²⁶. Loss of TGF β R signalling leads to reduced expression of GM-CSFR, suggesting cooperation between these factors¹²⁴. However, the indispensable nature of TGF β in their development makes assessing its role in regulation of AlV ϕ behaviour during homeostasis difficult. Nevertheless, TGF β is thought to upregulate the inhibitory receptor CD200R1, which is known to be key for maintaining the activation threshold of AlV ϕ s¹²⁷. Consistent with this, loss of autocrine TGF β leads to spontaneous production of pro-inflammatory cytokines and chemokines by AlV ϕ s¹²³.

Both GM-CSF and TGF β induce expression of the transcription factor PPAR γ , which is considered the master transcription factor for Alveolar Macrophages (AlvM ϕ s)^{43,124}. Mice with myeloid-specific deletion of *Pparg* also develop PAP, consistent with regulation of molecules involved in lipid catabolism by PPAR γ ^{43,128,129}. Given that PPAR γ is expressed by macrophages in other tissues, including splenic red pulp macrophages and macrophages of the erythroblastic islands in the bone marrow^{129,130}, until recently, it remained unclear how specificity was conferred to AlvM ϕ s. We recently uncovered the transcription factor early growth response 2 (EGR2) as a key evolutionarily conserved regulator of AlvM ϕ differentiation downstream of PPAR γ in the lung but not spleen¹²⁵. Interestingly, mice with *Egr2* deficient AlvM ϕ s and individuals with mutations in *EGR2* do not appear to develop spontaneous PAP^{125,131}, demonstrating that PPAR γ must cooperate with other transcriptional regulators to regulate distinct aspects of AlvM ϕ biology. For instance, EGR2 appears particularly important for regulating expression of adhesion molecules, chemotactic machinery and apparatus for the detection and elimination of respiratory pathogens¹²⁵. EGR2 appears to maintain expression of CCAAT/enhancer-binding protein beta (C/EBP β), which has been implicated in AlvM ϕ differentiation¹³². The transcription factors Bhlhe40 and Bhlhe41 have also been shown to control the phenotypic identity and proliferative capacity of AlvM ϕ s, and seem to rely on TGF β R signalling in a PPAR γ -independent manner¹³³. The histone deacetylase (HDAC) sirtuin 1 (SIRT1) also plays a key role in regulating the proliferative activity of AlvM ϕ s¹³⁴. Finally, Bach2 (B lymphoid transcriptional repressor BTB and CNC homology 2) has been shown to be essential for surfactant regulation by AlvM ϕ s¹³⁵. Thus, while much progress has been made in understanding the transcriptional control of lung macrophages, if and how these transcriptional regulators interact or cooperate to control the discrete molecular programmes required for homeostatic function of AlvM ϕ s is only starting to be understood and warrants further study using state-of-the-art technologies.

The environmental control of IntM ϕ s is much less well understood. Despite high expression of CX3CR1 by at least some IntM ϕ s, their survival, phenotype and proliferative capacity is unaffected by *Cx3cr1* deficiency¹⁴. Unlike their alveolar counterparts, IntM ϕ s rely on signalling through CSF1R for their development and maintenance as evidenced by their depletion with anti-CSF1R antibody treatment²⁸ and failure to develop from *Csf1r*^{-/-} precursors in a competitive bone marrow chimera setting¹³⁶. The relative role of the ligands for the CSF1R, M-CSF and IL-34, has not been examined exhaustively. For instance, although cDC2s are reported to be affected in *Il34*^{LacZ/LacZ} mice¹³⁷, these cells were simply defined as CD11c⁺CD11b⁺ non-AlvM ϕ s and it is highly likely this compartment contains both IntM ϕ s and cDC2s. Similarly, analysis of *Csf1*^{op/op} mice, which have a naturally occurring inactivating mutation in the *Csf1* gene, has shown an effect on the abundance of CD169⁺ but not CD169⁻ IntM ϕ s²⁸, suggesting differential reliance on M-CSF by discrete IntM ϕ subsets. Application of novel reporter and conditional "KO" mice, such as those used to identify the cellular sources of M-CSF in the lymph node and spleen^{138,139}, should help discern the relative roles and cellular origin of M-CSF and IL-34 in regulating survival and differentiation of IntM ϕ subsets.

The downstream molecular pathways that govern IntM ϕ differentiation remain largely elusive. Although expression of *Maf*, *Mafb*, *Irf5*, *Jun* and *Atf3* have been identified through scRNA-seq studies as highly expressed by murine IntM ϕ s^{6,125}, if and how these transcription factors control their differentiation remains unexplored. Furthermore, given that some of these (e.g., IRF5) have been implicated in AlvM ϕ homeostasis¹⁴⁰, high expression does not always equate to specificity. Importantly, although dispensable for the phenotypic identity and survival of IntM ϕ s, β -catenin signalling has recently been implicated in the control of the metabolic profile of IntM ϕ s, in response to the Wnt family

molecule R-spondin3 derived from pulmonary endothelial cells¹⁴¹. IntM ϕ s are intimately associated with extracellular matrix and interaction with collagen via the collagen receptor, LAIR1, appears to alter the composition of the IntM ϕ pool. Notably, LAIR1 appears to regulate CSF1R expression¹⁴² and therefore interactions with the ECM may regulate macrophage longevity, although this remains to be tested experimentally.

Thus, it is clear that the lung environment exerts multiple layers of control over macrophage development, activation and function in health and during inflammation. This highlights the current over-reliance on in vitro methods to research lung macrophages, particularly for human research, which are likely of questionable relevance. Innovative new approaches are needed that better reflect the lung environment, such as 'lung on a chip'¹⁴³⁻¹⁴⁶, organoids^{147,148}, and maintenance of whole lung tissue ex vivo.

MACROPHAGES IN PULMONARY FIBROSIS

Despite their key roles in lung homeostasis, macrophages are implicated in the pathogenesis of many chronic lung pathologies, including pulmonary fibrosis (PF). PF is a common feature of a group of conditions known as interstitial lung diseases (ILDs), where excessive ECM deposition leads to irreversible scarring of the lung (reviewed by^{149,150}). In many cases the cause of pulmonary fibrosis is not identified (idiopathic pulmonary fibrosis (IPF)), whereas in others it can be attributed to exposure to certain occupational substances (e.g., asbestos, silica) or drugs (e.g., bleomycin, methotrexate). Moreover, there are indications that following severe coronavirus disease 19 (COVID-19), certain individuals develop pulmonary fibrosis¹⁵¹, although whether this results in permanent, irreversible scarring is still being understood.

While the prevailing school of thought is that PF arises from ineffective repair of airway epithelium following repetitive injury, there is now compelling evidence that macrophages contribute to PF pathology¹⁵² and that targeting macrophages could be beneficial in human disease¹⁵³. First, there is vast macrophage accumulation in the lung parenchyma during PF and experimental fibrosis where they co-localise with collagen-producing myofibroblasts and support their proliferation and function through production of PDGF α , PDGF β , TGF β 1 and Galectin-3^{30,154-157}. Macrophages have also been shown to be rich sources of osteopontin (encoded by *Spp1/SPP1*)^{32,90,155,158}, which has long been established as a pro-fibrotic mediator, in part through activation of TGF β 1¹⁵⁸. Osteopontin-producing macrophages are found in the airways and parenchyma of IPF individuals¹⁵⁹, although highest expression is attributed to *MAFB*⁺*PPARG*⁻ macrophages, which most likely represent IntM ϕ s³². Moreover, high expression of inhibitors of collagenolytic enzymes, including tissue inhibitor of metalloproteinases 1 (TIMP1) and TIMP2 is a feature of fibrosis-associated macrophages¹⁵⁴. Studies in mice have shown macrophages to be able to produce certain collagens (e.g., collagen VI), and collagen VI deficiency limited to the haematopoietic compartment ameliorates experimental bleomycin-induced fibrosis¹⁶⁰. Interestingly, however, elevated expression of matrix metalloproteinases, such as MMP-9, MMP-12 and MMP-14, also defines fibrosis-associated macrophages across species. Thus, the relative contribution of collagen production versus collagen clearance remains poorly understood.

Macrophage accumulation results, at least in part, from de novo recruitment of CCR2⁺ monocytes in both experimental models and human PF¹⁶¹. The presence of CCR2⁺ monocytes and their macrophage progeny correlates with the presence of fibrotic tissue in mouse and man¹⁶¹, and bleomycin-induced experimental fibrosis can be exacerbated by adoptive transfer of classical monocytes¹⁶². Consistent with this, experimental fibrosis is blunted in monocytopenic *Ccr2* deficient mice^{161,163,164}, by neutralisation of the CSF1-CSF1R axis¹⁵⁴ or by rendering monocyte-derived macrophages susceptible to apoptosis¹⁶⁵.

Moreover, one of the few treatments for PF, pirfenidone, has recently been shown to reduce accumulation of CCR2⁺ monocytes in bleomycin-induced experimental fibrosis¹⁶¹. However, the Akira group has suggested that developmentally distinct, pro-fibrotic monocytes termed “segregated-nucleus-containing atypical monocytes” (SatM) arise in the context of experimental lung fibrosis and are responsible for driving disease¹⁶⁶. So-called ‘SatM’ appear to depend on the transcription factor C/EBP β , but derive from Fc ϵ R1⁺ granulocyte/macrophage progenitors (GMPs) and not macrophage/DC progenitors (MDPs)¹⁶⁶. Indeed, work since has described distinct pathways to generate monocytes from GMPs and MDPs in health and following infection¹⁶⁷. How these SatM relate to the CCR2-dependent monocytes described in other studies remains unclear.

Whether pro-fibrotic macrophages are limited to the lung parenchyma or if monocyte-derived AM ϕ s also contribute to fibrosis is still under debate^{90,125,154,156}. In support of the latter, sustained epithelial injury is a feature of human PF^{155,168} and reducing epithelial damage through administration of a specific inhibitor of sphingosine kinase 1, which is elevated in IPF lungs, reduces experimental fibrosis, at least in part by reducing recruitment of fibrogenic monocytes¹⁶⁹. Attributing key pathogenic roles to macrophage subsets is made difficult by the breakdown in clear phenotypic boundaries between parenchymal and AlV ϕ s in the context of inflammation and fibrosis, and by the fact that, at least some, IntM ϕ s may differentiate into AlV ϕ s during lung repair^{125,156}. PF incidence and severity positively correlates with age^{170–172} and given monocytes may progressively replace embryonically-derived AlV ϕ s with age, it is intriguing to speculate that these two phenomena could be related. In addition to blood-derived monocytes, it is plausible that macrophages in the pleural cavity may contribute to the pro-fibrotic pool of macrophages in the lung. Consistent with this notion, fibrosis in IPF patients is often concentrated in the subpleural region¹⁵⁰ and there is transcriptional similarity between pleural and MHCII⁻ IntM ϕ s in mice^{27,173}. Moreover, it’s been suggested serous cavity macrophages may contribute to tissue repair in neighbouring solid organs following injury in mice^{174,175}. However, elegant intersectoral genetics and a combination of injury models have shown that, while pleural GATA6⁺ macrophages may accumulate on the pleural membrane, they do not migrate deep into the lung parenchyma nor are they essential for fibrogenesis or resolution¹⁷⁶.

Why macrophages become excessively pro-fibrotic is only starting to be understood, although, again, this probably reflects their plasticity. IL-4, IL-13, IL-33 and TGF β have all been implicated in altering macrophage behaviour in PF (reviewed by¹⁵²), but the relative and combinatorial roles of these factors is poorly understood. Recently, overactive Notch signalling has been implicated in the pro-fibrotic behaviour of macrophages, as deletion of RBPJ reduces fibrosis in mice through abrogating TGF β production by Ly6C^{hi}MHCII⁺ monocyte-derived macrophages¹⁷⁷. Whether these effects are attributable to airway or IntM ϕ s is difficult to discern in this study¹⁷⁷. TGF β is of particular interest given its long-standing role in tissue fibrosis¹⁷⁸. Recent work has uncovered discrete functions of TGF β isoforms in the fibrotic process¹⁷⁹. However, if and how excessive TGF β isoforms influence macrophage function in the fibrotic niche in vivo has not been tested directly.

Like in many pathologies, the role of macrophages in lung fibrosis is very much context dependent. The self-resolving nature of some experimental models allows macrophage dynamics and behaviour to be assessed during fibrosis regression and resolution, something that cannot be gleaned from human disease. This has revealed that severe lung injury leads to almost complete replacement of embryonically derived AlV ϕ s with monocyte-derived cells^{90,125}. We have recently shown that this process is highly dependent on the transcription factor EGR2 and that

EGR2-dependent monocyte-derived AM ϕ s are indispensable for resolution of fibrosis and restoration of airway homeostasis¹²⁵, findings consistent with older non-specific depletion studies in mice¹⁶². There is evidence this may involve direct clearance of collagen by (monocyte-derived) AM ϕ s. For instance, genetic ablation of milk fate globule epidermal growth factor 8 (Mfge8), a receptor typically associated with efferocytosis, leads to failed collagen clearance following bleomycin-induced injury¹⁸⁰. Moreover, macrophage-derived ApoE may facilitate binding and targeting of type I collagen for phagocytosis via the low-density lipoprotein receptor-related protein 1 (LRP1)¹⁸¹. Pro-resolution roles of monocyte-derived AM ϕ s are also seen following influenza infection and their absence can result in the development of fibrosis, at least in mice¹⁸². Such functions may include metabolic rewiring of monocyte-derived macrophages as deletion of *Acod1*, the enzyme required for generation of the metabolite itaconate, leads to persistent fibrosis¹⁸³. Thus, while generally considered as key pro-fibrotic cellular players, monocyte-derived AM ϕ s appear to have a crucial role in re-establishing lung homeostasis and may hold great promise for therapeutic targeting to promote fibrosis regression and lung repair.

CONCLUSION

The past few years have seen a leap forwards in our understanding of pulmonary macrophage development, heterogeneity and function, and how environmental features of the lung can exert a dramatic influence over these processes in both disease and in health. A major challenge for the coming years is to develop much greater clarity on how different inflammatory conditions alter pulmonary macrophage subset diversity and function, in particular in the context of human disease, to identify core mechanisms that might enable development of the targeted therapeutics of the future.

REFERENCES

- Mills, C. D., Kincaid, K., Alt, J. M., Heilman, M. J. & Hill, A. M. M-1/M-2 macrophages and the Th1/Th2 paradigm. *J. Immunol.* **164**, 6166–6173 (2000).
- Murray, P. J. et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* **41**, 14–20 (2014).
- Lavin, Y. et al. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. *Cell* **159**, 1312–1326 (2014).
- Mulder, K. et al. Cross-tissue single-cell landscape of human monocytes and macrophages in health and disease. *Immunity* **54**, 1883–1900.e5 (2021).
- Liao, M. et al. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat. Med.* **26**, 842–844 (2020).
- Sajti, E. et al. Transcriptomic and epigenetic mechanisms underlying myeloid diversity in the lung. *Nat. Immunol.* **21**, 221–231 (2020).
- Leach, S. M. et al. Human and mouse transcriptome profiling identifies cross-species homology in pulmonary and lymph node mononuclear phagocytes. *Cell Rep.* **33**, 108337 (2020).
- Lavin, Y. et al. Innate immune landscape in early lung adenocarcinoma by paired single-cell analyses. *Cell* **169**, 750–765.e17 (2017).
- Zilionis, R. et al. Single-cell transcriptomics of human and mouse lung cancers reveals conserved myeloid populations across individuals and species. *Immunity* **50**, 1317–1334.e10 (2019).
- Raredon, M. S. B. et al. Single-cell connectomic analysis of adult mammalian lungs. *Sci. Adv.* **5**, eaaw3851 (2019).
- Travaglini, K. J. et al. A molecular cell atlas of the human lung from single-cell RNA sequencing. *Nature* **587**, 619–625 (2020).
- Gautier, E. L. et al. Gene-expression profiles and transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue macrophages. *Nat. Immunol.* **13**, 1118–1128 (2012).
- Bharat, A. et al. Flow cytometry reveals similarities between lung macrophages in humans and mice. *Am. J. Respir. Cell Mol. Biol.* **54**, 147–149 (2016).
- Gibbins, S. L. et al. Three unique interstitial macrophages in the murine lung at steady state. *Am. J. Respir. Cell Mol. Biol.* **57**, 66–76 (2017).
- Svedberg, F. R. et al. The lung environment controls alveolar macrophage metabolism and responsiveness in type 2 inflammation. *Nat. Immunol.* **20**, 571–580 (2019).

16. Becher, B. et al. High-dimensional analysis of the murine myeloid cell system. *Nat. Immunol.* **15**, 1181–1189 (2014).
17. Hoffmann, F. M. et al. Distribution and interaction of murine pulmonary phagocytes in the naive and allergic lung. *Front Immunol.* **9**, 1046 (2018).
18. Schyns, J. et al. Non-classical tissue monocytes and two functionally distinct populations of interstitial macrophages populate the mouse lung. *Nat. Commun.* **10**, 3964 (2019).
19. Hume, P. S. et al. Localization of macrophages in the human lung via design-based stereology. *Am J. Respir. Crit. Care Med.* **201**, 1209–1217 (2020).
20. Huang, L., Nazarova, E. V., Tan, S., Liu, Y. & Russell, D. G. Growth of mycobacterium tuberculosis in vivo segregates with host macrophage metabolism and ontogeny. *J. Exp. Med.* **215**, 1135–1152 (2018).
21. Bosteels, C. et al. Inflammatory type 2 cDCs acquire features of cDC1s and macrophages to orchestrate immunity to respiratory virus infection. *Immunity* **52**, 1039–1056.e9 (2020).
22. Ono, Y. et al. CD11c+ M1-like macrophages (MΦs) but not CD206+ M2-like MΦ are involved in folliculogenesis in mice ovary. *Sci. Rep.* **8**, 8171–10 (2018).
23. Röszer, T. Understanding the mysterious M2 macrophage through activation markers and effector mechanisms. *Mediators Inflamm.* **2015**, 816460 (2015).
24. Saini, Y. et al. Gene expression in whole lung and pulmonary macrophages reflects the dynamic pathology associated with airway surface dehydration. *BMC Genomics* **15**, 726 (2014).
25. Lai, J. F. et al. Critical role of macrophages and their activation via MyD88-NFκB signaling in lung innate immunity to *Mycoplasma pneumoniae*. *PLoS One* **5**, e14417 (2010).
26. Nahrendorf, M. & Swirski, F. K. Abandoning M1/M2 for a network model of macrophage function. *Circ. Res.* **119**, 414–417 (2016).
27. Chakarov, S. et al. Two distinct interstitial macrophage populations coexist across tissues in specific subtissular niches. *Science* **363**, eaau0964 (2019).
28. Ural, B. B. et al. Identification of a nerve-associated, lung-resident interstitial macrophage subset with distinct localization and immunoregulatory properties. *Sci. Immunol.* **5**, eaax8756 (2020).
29. Sabatel, C. et al. Exposure to bacterial CpG DNA protects from airway allergic inflammation by expanding regulatory lung interstitial macrophages. *Immunity* **46**, 457–473 (2017).
30. Yu, Y.-R. A. et al. Flow cytometric analysis of myeloid cells in human blood, bronchoalveolar lavage, and lung tissues. *Am. J. Respir. Cell Mol. Biol.* **54**, 13–24 (2016).
31. McQuattie-Pimentel, A. C. et al. The lung microenvironment shapes a dysfunctional response of alveolar macrophages in aging. *J. Clin. Invest.* **131**, e140299 (2021).
32. Morse, C. et al. Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. *Eur. Respir. J.* **54**, 1802441 (2019).
33. Vieira Braga, F. A. et al. A cellular census of human lungs identifies novel cell states in health and in asthma. *Nat. Med.* **25**, 1153–1163 (2019).
34. Mould, K. J. et al. Airspace macrophages and monocytes exist in transcriptionally distinct subsets in healthy adults. *Am J. Respir. Crit. Care Med.* **203**, 946–956 (2021).
35. Evren, E. et al. Distinct developmental pathways from blood monocytes generate human lung macrophage diversity. *Immunity* **54**, 259–275.e7 (2021).
36. Aberdein, J. D., Cole, J., Bewley, M. A., Marriott, H. M. & Dockrell, D. H. Alveolar macrophages in pulmonary host defence the unrecognized role of apoptosis as a mechanism of intracellular bacterial killing. *Clin. Exp. Immunol.* **174**, 193–202 (2013).
37. Preston, J. A. et al. Alveolar macrophage apoptosis-associated bacterial killing helps prevent murine pneumonia. *Am. J. Respir. Crit. Care Med.* **200**, 84–97 (2019).
38. Schneider, C. et al. Alveolar macrophages are essential for protection from respiratory failure and associated morbidity following influenza virus infection. *PLoS Pathog.* **10**, e1004053 (2014).
39. Xu, J. et al. Scavenger receptor MARCO orchestrates early defenses and contributes to fungal containment during cryptococcal infection. *J. Immunol.* **198**, 3548–3557 (2017).
40. Dorrington, M. G. et al. MARCO is required for TLR2- and Nod2-mediated responses to *Streptococcus pneumoniae* and clearance of pneumococcal colonization in the murine nasopharynx. *J. Immunol.* **190**, 250–258 (2013).
41. Arredouani, M. et al. The scavenger receptor MARCO is required for lung defense against pneumococcal pneumonia and inhaled particles. *J. Exp. Med.* **200**, 267–272 (2004).
42. Hussell, T. & Bell, T. J. Alveolar macrophages: plasticity in a tissue-specific context. *Nat. Rev.* **14**, 81–93 (2014).
43. Schneider, C. et al. Induction of the nuclear receptor PPAR-γ by the cytokine GM-CSF is critical for the differentiation of fetal monocytes into alveolar macrophages. *Nat. Immunol.* **15**, 1026–1037 (2014).
44. Gibbings, S. L. et al. Transcriptome analysis highlights the conserved difference between embryonic and postnatal-derived alveolar macrophages. *Blood* **126**, 1357–1366 (2015).
45. Nishinakamura, R. et al. Mice deficient for the IL-3/GM-CSF/IL-5 beta c receptor exhibit lung pathology and impaired immune response, while beta IL3 receptor-deficient mice are normal. *Immunity* **2**, 211–222 (1995).
46. Robb, L. et al. Hematopoietic and lung abnormalities in mice with a null mutation of the common beta subunit of the receptors for granulocyte-macrophage colony-stimulating factor and interleukins 3 and 5. *Proc. Natl Acad. Sci. U.S.A.* **92**, 9565–9569 (1995).
47. Kitamura, T. et al. Idiopathic pulmonary alveolar proteinosis as an autoimmune disease with neutralizing antibody against granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* **190**, 875–880 (1999).
48. Suzuki, T. et al. Familial pulmonary alveolar proteinosis caused by mutations in CSF2RA. *J. Exp. Med.* **205**, 2703–2710 (2008).
49. van de Laar, L. et al. Yolk sac macrophages, fetal liver, and adult monocytes can colonize an empty niche and develop into functional tissue-resident macrophages. *Immunity* <https://doi.org/10.1016/j.immuni.2016.02.017> (2016).
50. Stanley E. et al. Granulocyte/macrophage colony-stimulating factor-deficient mice show no major perturbation of hematopoiesis but develop a characteristic pulmonary pathology. *Proc Natl Acad Sci USA.* **91**, (1994).
51. Peters, D. M. et al. TGF-β directs trafficking of the epithelial sodium channel ENaC which has implications for ion and fluid transport in acute lung injury. *Proc. Natl Acad. Sci. U.S.A.* **111**, E374–E383 (2014).
52. Han, S. W. & Roman, J. Fibronectin induces cell proliferation and inhibits apoptosis in human bronchial epithelial cells: pro-oncogenic effects mediated by PI3-kinase and NF-κB. *Oncogene* **25**, 4341–4349 (2006).
53. Bourdonnay, E. et al. Transcellular delivery of vesicular SOCS proteins from macrophages to epithelial cells blunts inflammatory signaling. *J. Exp. Med.* **212**, 729–742 (2015).
54. Draijer, C. et al. Resident alveolar macrophage-derived vesicular SOCS3 dampens allergic airway inflammation. *FASEB J.* **34**, 4718–4731 (2020).
55. Schyns, J., Bureau, F. & Marichal, T. Lung interstitial macrophages: past, present, and future. *J. Immunol. Res.* **2018**, 5160794 (2018).
56. Bedoret, D. et al. Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. *J. Clin. Invest.* **119**, 3723–3738 (2009).
57. Kawano, H. et al. IL-10-producing lung interstitial macrophages prevent neurophilic asthma. *Int. Immunol.* **28**, 489–501 (2016).
58. Hoppstädter, J. et al. Differential cell reaction upon toll-like receptor 4 and 9 activation in human alveolar and lung interstitial macrophages. *Respir. Res.* **11**, 124–15 (2010).
59. Plantinga, M. et al. Conventional and monocyte-derived CD11b(+) dendritic cells initiate and maintain T helper 2 cell-mediated immunity to house dust mite allergen. *Immunity* **38**, 322–335 (2013).
60. Ren, J. et al. Intranasal delivery of MSC-derived exosomes attenuates allergic asthma via expanding IL-10 producing lung interstitial macrophages in mice. *Int Immunopharmacol.* **91**, 107288 (2021).
61. Brody, A. et al. Interstitial pulmonary macrophages produce platelet-derived growth factor that stimulates rat lung fibroblast proliferation in vitro. *J. Leukoc. Biol.* **6**, 640–648 (1992).
62. De Schepper, S. et al. Self-maintaining gut macrophages are essential for intestinal homeostasis. *Cell* **175**, 400–415.e13 (2018).
63. van Furth, R. et al. The mononuclear phagocyte system: a new classification of macrophages, monocytes, and their precursor cells. *Bull. World Health Organ* **46**, 845–852 (1972).
64. Schulz, C. et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science* **336**, 86–90 (2012).
65. Gomez Perdiguero, E. et al. Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid progenitors. *Nature* **518**, 547–551 (2015).
66. Hoeffel, G. et al. Adult Langerhans cells derive predominantly from embryonic fetal liver monocytes with a minor contribution of yolk sac-derived macrophages. *J. Exp. Med.* **209**, 1167–1181 (2012).
67. Ginhoux, F. et al. Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science* **330**, 841–845 (2010).
68. Hashimoto, D. et al. Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity* **38**, 792–804 (2013).
69. Hoeffel, G. et al. C-Myb(+) erythro-myeloid progenitor-derived fetal monocytes give rise to adult tissue-resident macrophages. *Immunity* **42**, 665–678 (2015).
70. Sheng, J., Ruedl, C. & Karjalainen, K. Most tissue-resident macrophages except microglia are derived from fetal hematopoietic stem cells. *Immunity* **43**, 382–393 (2015).
71. Liu, Z. et al. Fate mapping via Ms4a3-expression history traces monocyte-derived. *Cells Cell* **178**, 1509–1525.e19 (2019).

72. Shaw, T. N. et al. Tissue-resident macrophages in the intestine are long lived and defined by Tim-4 and CD4 expression. *J. Exp. Med.* **215**, 1507–1518 (2018).
73. Bain, C. C. et al. Long-lived self-renewing bone marrow-derived macrophages displace embryo-derived cells to inhabit adult serous cavities. *Nat. Commun.* **7**, ncomms11852 (2016).
74. Epelman, S. et al. Embryonic and adult-derived resident cardiac macrophages are maintained through distinct mechanisms at steady state and during inflammation. *Immunity* **40**, 91–104 (2014).
75. Tarling, J. D., Lin, H. S. & Hsu, S. Self-renewal of pulmonary alveolar macrophages: evidence from radiation chimera studies. *J. Leukoc. Biol.* **42**, 443–446 (1987).
76. Goldmann, T. et al. A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nat. Neurosci.* **16**, 1618–1626 (2013).
77. Tan, S. Y. S. & Krasnow, M. A. Developmental origin of lung macrophage diversity. *Development* **143**, 1318–1327 (2016).
78. Guillems, M. et al. Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. *J. Exp. Med.* **210**, 1977–1992 (2013).
79. Alenghat, E. & Esterly, J. R. Alveolar macrophages in perinatal infants. *Pediatrics* **74**, 221–223 (1984).
80. Eguiluz-Gracia, I. et al. Long-term persistence of human donor alveolar macrophages in lung transplant recipients. *Thorax* <https://doi.org/10.1136/thoraxjnl-2016-208292> (2016).
81. Nayak, D. K. et al. Long-term persistence of donor alveolar macrophages in human lung transplant recipients that influences donor-specific immune responses. *Am. J. Transpl.* **16**, 2300–2311 (2016).
82. Rongvaux, A. et al. Development and function of human innate immune cells in a humanized mouse model. *Nat. Biotechnol.* **32**, 364–372 (2014).
83. Byrne, A. J. et al. Dynamics of human monocytes and airway macrophages during healthy aging and after transplant. *J. Exp. Med.* **217**, e20191236 (2020).
84. Hawley, C. A. et al. Csf1r-mApple transgene expression and ligand binding in vivo reveal dynamics of CSF1R expression within the mononuclear phagocyte system. *J. Immunol.* **200**, 2209–2223 (2018).
85. Dyer, D. P. et al. Chemokine receptor redundancy and specificity are context dependent. *Immunity* **50**, 378–389.e5 (2019).
86. Jenkins, S. J. et al. IL-4 directly signals tissue-resident macrophages to proliferate beyond homeostatic levels controlled by CSF-1. *J. Exp. Med.* **210**, 2477–2491 (2013).
87. Davies, L. C. et al. Distinct bone marrow-derived and tissue-resident macrophage lineages proliferate at key stages during inflammation. *Nat. Commun.* **4**, 1886 (2013).
88. Guillems, M. & Scott, C. L. Does niche competition determine the origin of tissue-resident macrophages? *Nat. Rev.* **17**, 451–460 (2017).
89. Li, F., Okreglicka, K. M., Pohlmeier, L. M., Schneider, C. & Kopf, M. Fetal monocytes possess increased metabolic capacity and replace primitive macrophages in tissue macrophage development. *EMBO J.* **39**, e103205 (2020).
90. Misharin, A. V. et al. Monocyte-derived alveolar macrophages drive lung fibrosis and persist in the lung over the life span. *J. Exp. Med.* **214**, 2387–2404 (2017).
91. Aegerter, H. et al. Influenza-induced monocyte-derived alveolar macrophages confer prolonged antibacterial protection. *Nat. Immunol.* **21**, 145–157 (2020).
92. Hortobágyi, L. et al. Surfactant protein D inhibits TNF- α production by macrophages and dendritic cells in mice. *J. Allergy Clin. Immunol.* **122**, 521–528 (2008).
93. Minutti, C. M. et al. Local amplifiers of IL-4Ra-mediated macrophage activation promote repair in lung and liver. *Science* **356**, 1076–1080 (2017).
94. Yamada, C. et al. Surfactant protein A directly interacts with TLR4 and MD-2 and regulates inflammatory cellular response. Importance of supratrimeric oligomerization. *J. Biol. Chem.* **281**, 21771–21780 (2006).
95. Sano, H. et al. Pulmonary surfactant protein A modulates the cellular response to smooth and rough lipopolysaccharides by interaction with CD14. *J. Immunol.* **163**, 387–395 (1999).
96. Murugaiah, V., Tsolaki, A. G. & Kishore, U. Collectins: innate immune pattern recognition molecules. *Adv. Exp. Med. Biol.* **1204**, 75–127 (2020).
97. McShane, A. et al. Mucus. *Curr. Biol.* **31**, R938–R945 (2021).
98. Ridley, C. & Thornton, D. J. Mucins: the frontline defence of the lung. *Biochem Soc. Trans.* **46**, 1099–1106 (2018).
99. Roy, M. G. et al. Muc5b is required for airway defence. *Nature* **505**, 412–416 (2014).
100. Barbier, D. et al. Influenza A induces the major secreted airway mucin MUC5AC in a protease-EGFR-extracellular regulated kinase-Sp1-dependent pathway. *Am. J. Respir. Cell Mol. Biol.* **47**, 149–157 (2012).
101. Ehre, C. et al. Overexpressing mouse model demonstrates the protective role of Muc5ac in the lungs. *Proc. Natl Acad. Sci. USA.* **109**, 16528–16533 (2012).
102. Evans, C. M. et al. The polymeric mucin Muc5ac is required for allergic airway hyperreactivity. *Nat. Commun.* **6**, 6281 (2015).
103. Gill, S. K. et al. Increased airway glucose increases airway bacterial load in hyperglycaemia. *Sci. Rep.* **6**, 27636 (2016).
104. Baker, E. H. & Baines, D. L. Airway glucose homeostasis: a new target in the prevention and treatment of pulmonary infection. *Chest* **153**, 507–514 (2018).
105. Baker, E. H. et al. Hyperglycemia and cystic fibrosis alter respiratory fluid glucose concentrations estimated by breath condensate analysis. *J. Appl. Physiol.* (1985) **102**, 1969–1975 (2007).
106. O'Neill, L. A. J. & Pearce, E. J. Immunometabolism governs dendritic cell and macrophage function. *J. Exp. Med.* **213**, 15–23 (2016).
107. Van den Bossche, J., O'Neill, L. A. & Menon, D. Macrophage immunometabolism: where are we (going)? *Trends Immunol.* **38**, 395–406 (2017).
108. Mallia, P. et al. Role of airway glucose in bacterial infections in patients with chronic obstructive pulmonary disease. *J. Allergy Clin. Immunol.* **142**, 815–823.e6 (2018).
109. Garnett, J. P. et al. Hyperglycaemia and *Pseudomonas aeruginosa* acidify cystic fibrosis airway surface liquid by elevating epithelial monocarboxylate transporter 2 dependent lactate-H⁺ secretion. *Sci. Rep.* **6**, 37955 (2016).
110. Ho, W. E. et al. Metabolomics reveals altered metabolic pathways in experimental asthma. *Am. J. Respir. Cell Mol. Biol.* **48**, 204–211 (2013).
111. Ogger, P. P. & Byrne, A. J. Macrophage metabolic reprogramming during chronic lung disease. *Mucosal Immunol.* **14**, 282–295 (2021).
112. Pisu, D., Huang, L., Grenier, J. K. & Russell, D. G. Dual RNA-Seq of Mtb-infected macrophages in vivo reveals ontologically distinct host-pathogen interactions. *Cell Rep.* **30**, 335–350.e4 (2020).
113. Martinez-Moczygemba, M. et al. Pulmonary alveolar proteinosis caused by deletion of the GM-CSFR α gene in the X chromosome pseudoautosomal region 1. *J. Exp. Med.* **205**, 2711–2716 (2008).
114. Shibata, Y. et al. GM-CSF regulates alveolar macrophage differentiation and innate immunity in the lung through PU.1. *Immunity* **15**, 557–567 (2001).
115. Suzuki, T. et al. Pulmonary macrophage transplantation therapy. *Nature* **514**, 450–454 (2014).
116. Willinger, T. et al. Human IL-3/GM-CSF knock-in mice support human alveolar macrophage development and human immune responses in the lung. *Proc. Natl Acad. Sci. USA.* **108**, 2390–2395 (2011).
117. Gschwend, J. et al. Alveolar macrophages rely on GM-CSF from alveolar epithelial type 2 cells before and after birth. *J. Exp. Med.* **218**, e20210745 (2021).
118. Cohen, M. et al. Lung single-cell signaling interaction map reveals basophil role in macrophage imprinting. *Cell* **175**, 1031–1044.e18 (2018).
119. Croxford, A. L. et al. The Cytokine GM-CSF drives the inflammatory signature of CCR2⁺ monocytes and licenses autoimmunity. *Immunity* **43**, 502–514 (2015).
120. Eddy, W. E. et al. Stat5 is required for cd103⁺ dendritic cell and alveolar macrophage development and protection from lung injury. *J. Immunol.* **198**, 4813–4822 (2017).
121. Morris, D. G. et al. Loss of integrin α (v) β 6-mediated TGF- β activation causes Mmp12-dependent emphysema. *Nature* **422**, 169–173 (2003).
122. Koth, L. L. et al. Integrin β 6 mediates phospholipid and collectin homeostasis by activation of latent TGF- β 1. *Am. J. Respir. Cell Mol. Biol.* **37**, 651–659 (2007).
123. Branchett, W. J. et al. Airway macrophage-intrinsic TGF- β 1 regulates pulmonary immunity during early-life allergen exposure. *J. Allergy Clin. Immunol.* **147**, 1892–1906 (2021).
124. Yu, X. et al. The cytokine TGF- β promotes the development and homeostasis of alveolar macrophages. *Immunity* **47**, 903–912.e4 (2017).
125. McCowan, J. et al. The transcription factor EGR2 is indispensable for tissue-specific imprinting of alveolar macrophages in health and tissue repair. *Sci. Immunol.* **6**, eabj2132 (2021).
126. Minutti, C. M. et al. A macrophage-pericyte axis directs tissue restoration via amphiregulin-induced transforming growth factor β activation. *Immunity* **50**, 645–654.e6 (2019).
127. Snelgrove, R. J. et al. A critical function for CD200 in lung immune homeostasis and the severity of influenza infection. *Nat. Immunol.* **9**, 1074–1083 (2008).
128. Baker, A. D. et al. PPAR γ regulates the expression of cholesterol metabolism genes in alveolar macrophages. *Biochem Biophys. Res Commun.* **393**, 682–687 (2010).
129. Gautier, E. L. et al. Systemic analysis of PPAR γ in mouse macrophage populations reveals marked diversity in expression with critical roles in resolution of inflammation and airway immunity. *J. Immunol.* **189**, 2614–2624 (2012).
130. Okreglicka, K. et al. PPAR γ is essential for the development of bone marrow erythroblastic island macrophages and splenic red pulp macrophages. *J. Exp. Med.* **218**, e20191314 (2021).
131. Szigeti, K. et al. Functional, histopathologic and natural history study of neuropathy associated with EGR2 mutations. *Neurogenetics* **8**, 257–262 (2007).
132. Cain, D. W. et al. Identification of a tissue-specific, C/EBP β -dependent pathway of differentiation for murine peritoneal macrophages. *J. Immunol.* **191**, 4665–4675 (2013).

133. Rauschmeier, R. et al. Bhlhe40 and Bhlhe41 transcription factors regulate alveolar macrophage self-renewal and identity. *EMBO J.* **38**, e101233 (2019).
134. Imperatore, F. et al. SIRT1 regulates macrophage self-renewal. *EMBO J.* <https://doi.org/10.15252/emj.201695737> (2017).
135. Nakamura, A. et al. Transcription repressor Bach2 is required for pulmonary surfactant homeostasis and alveolar macrophage function. *J. Exp. Med.* **210**, 2191–2204 (2013).
136. Schlitzer, A. et al. IRF4 transcription factor-dependent CD11b+ dendritic cells in human and mouse control mucosal IL-17 cytokine responses. *Immunity* **38**, 970–983 (2013).
137. Wang, Y. et al. IL-34 is a tissue-restricted ligand of CSF1R required for the development of Langerhans cells and microglia. *Nat. Immunol.* **13**, 753–760 (2012).
138. Bellomo, A. et al. Reticular fibroblasts expressing the transcription factor WT1 define a stromal niche that maintains and replenishes splenic red pulp macrophages. *Immunity* **53**, 127–142.e7 (2020).
139. Mondor, I. et al. Lymphatic endothelial cells are essential components of the subcapsular sinus macrophage niche. *Immunity* **50**, 1453–1466.e4 (2019).
140. Albers, G. J. et al. IRF5 regulates airway macrophage metabolic responses. *Clin. Exp. Immunol.* **204**, 134–143 (2021).
141. Zhou, B. et al. The angiocrine Rspodin3 instructs interstitial macrophage transition via metabolic-epigenetic reprogramming and resolves inflammatory injury. *Nat. Immunol.* **21**, 1430–1443 (2020).
142. Keerthivasan, S. et al. Homeostatic functions of monocytes and interstitial lung macrophages are regulated via collagen domain-binding receptor LAIR1. *Immunity* **54**, 1511–1526.e8 (2021).
143. Huh, D., Torisawa, Y.-S., Hamilton, G. A., Kim, H. J. & Ingber, D. E. Microengineered physiological biomimicry: organs-on-chips. *Lab Chip* **12**, 2156–2164 (2012).
144. Bhatia, S. N. & Ingber, D. E. Microfluidic organs-on-chips. *Nat. Biotechnol.* **32**, 760–772 (2014).
145. Ahadian, S. et al. Organ-on-a-chip platforms: a convergence of advanced materials, cells, and microscale technologies. *Adv. Healthc. Mater.* **7**, <https://doi.org/10.1002/adhm.201700506> (2018).
146. Zamprogno, P. et al. Second-generation lung-on-a-chip with an array of stretchable alveoli made with a biological membrane. *Commun. Biol.* **4**, 168–10 (2021).
147. Leibel, S. L., McVicar, R. N., Winquist, A. M. & Snyder, E. Y. Generation of 3D whole lung organoids from induced pluripotent stem cells for modeling lung developmental biology and disease. *J. Vis. Exp.* e62456 <https://doi.org/10.3791/62456> (2021).
148. Miller, A. J. et al. Generation of lung organoids from human pluripotent stem cells in vitro. *Nat. Protoc.* **14**, 518–540 (2019).
149. Coward, W. R., Saini, G. & Jenkins, G. The pathogenesis of idiopathic pulmonary fibrosis. *Ther. Adv. Respir. Dis.* **4**, 367–388 (2010).
150. Martinez, F. J. et al. Idiopathic pulmonary fibrosis. *Nat. Rev. Dis. Prim.* **3**, 17074–19 (2017).
151. Vijayakumar, B. et al. CT Lung abnormalities after COVID-19 at 3 months and 1 year after hospital discharge. *Radiology* 211746 <https://doi.org/10.1148/radiol.2021211746> (2021).
152. Byrne, A. J., Maher, T. M. & Lloyd, C. M. Pulmonary macrophages: a new therapeutic pathway in fibrosing lung disease? *Trends Mol. Med.* <https://doi.org/10.1016/j.molmed.2016.02.004> (2016).
153. Hirani, N. et al. Target inhibition of galectin-3 by inhaled TD139 in patients with idiopathic pulmonary fibrosis. *Eur. Respir. J.* **57**, (2021).
154. Joshi, N. et al. A spatially restricted fibrotic niche in pulmonary fibrosis is sustained by M-CSF/M-CSFR signalling in monocyte-derived alveolar macrophages. *Eur. Respir. J.* **55**, 1900646 (2020).
155. Reyfman, P. A. et al. Single-cell transcriptomic analysis of human lung provides insights into the pathobiology of pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.* **199**, 1517–1536 (2019).
156. Aran, D. et al. Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).
157. Humphries, D. C. et al. Selective myeloid depletion of galectin-3 offers protection against acute and chronic lung injury. *Front. Pharm.* **12**, 715986 (2021).
158. Berman, J. S. et al. Altered bleomycin-induced lung fibrosis in osteopontin-deficient mice. *Am. J. Physiol. Lung Cell Mol. Physiol.* **286**, L1311–L1318 (2004).
159. Adams, T. S. et al. Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. *Sci. Adv.* **6**, eaba1983 (2020).
160. Uceroglu, A. C. et al. Fra-2-expressing macrophages promote lung fibrosis in mice. *J. Clin. Invest.* **129**, 3293–3309 (2019).
161. Brody, S. L. et al. Chemokine receptor 2-targeted molecular imaging in pulmonary fibrosis. A clinical trial. *Am. J. Respir. Crit. Care Med.* **203**, 78–89 (2021).
162. Gibbons, M. A. et al. Ly6Chi monocytes direct alternatively activated profibrotic macrophage regulation of lung fibrosis. *Am. J. Respir. Crit. Care Med.* **184**, 569–581 (2011).
163. Osterholzer, J. J. et al. Implicating exudate macrophages and Ly-6C(high) monocytes in CCR2-dependent lung fibrosis following gene-targeted alveolar injury. *J. Immunol.* **190**, 3447–3457 (2013).
164. Okuma, T. et al. C-C chemokine receptor 2 (CCR2) deficiency improves bleomycin-induced pulmonary fibrosis by attenuation of both macrophage infiltration and production of macrophage-derived matrix metalloproteinases. *J. Pathol.* **204**, 594–604 (2004).
165. McCubrey, A. L. et al. Deletion of c-FLIP from CD11bhi macrophages prevents development of bleomycin-induced lung fibrosis. *Am. J. Respir. Cell Mol. Biol.* **58**, 66–78 (2018).
166. Satoh, T. et al. Identification of an atypical monocyte and committed progenitor involved in fibrosis. *Nature* **541**, 96–101 (2017).
167. Yáñez, A. et al. Granulocyte-monocyte progenitors and monocyte-dendritic cell progenitors independently produce functionally distinct monocytes. *Immunity* **47**, 890–902.e4 (2017).
168. Habermann, A. C. et al. Single-cell RNA sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Sci. Adv.* **6**, eaba1972 (2020).
169. Chereshe, P. et al. The Sphingosine Kinase 1 Inhibitor, PF543, Mitigates pulmonary fibrosis by reducing lung epithelial cell mtDNA damage and recruitment of fibrogenic monocytes. *Int. J. Mol. Sci.* **21**, 5595, <https://doi.org/10.3390/ijms21165595> (2020).
170. Thannickal, V. J. et al. Blue journal conference. Aging and susceptibility to lung disease. *Am. J. Respir. Crit. Care Med.* **191**, 261–269 (2015).
171. Ragu, G., Weycker, D., Edelsberg, J., Bradford, W. Z. & Oster, G. Incidence and prevalence of idiopathic pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.* **174**, 810–816 (2006).
172. Watanabe, S. et al. Resetting proteostasis with ISRIB promotes epithelial differentiation to attenuate pulmonary fibrosis. *Proc. Natl. Acad. Sci. USA* **118**, e2101100118 (2021).
173. Bain, C. C. et al. Rate of replenishment and microenvironment contribute to the sexually dimorphic phenotype and function of peritoneal macrophages. *Sci. Immunol.* **5**, eabc4466 (2020).
174. Wang, J. & Kubes, P. A reservoir of mature cavity macrophages that can rapidly invade visceral organs to affect tissue repair. *Cell* **165**, 668–678 (2016).
175. Deniset, J. F. et al. Gata6+ pericardial cavity macrophages relocate to the injured heart and prevent cardiac fibrosis. *Immunity* **51**, 131–140.e5 (2019).
176. Jin, H. et al. Genetic fate-mapping reveals surface accumulation but not deep organ invasion of pleural and peritoneal cavity macrophages following injury. *Nat. Commun.* **12**, 2863 (2021).
177. Zhang, N. et al. Myeloid-specific blockade of Notch signaling alleviates murine pulmonary fibrosis through regulating monocyte-derived Ly6clo MHCIIhi alveolar macrophages recruitment and TGF- β secretion. *FASEB J.* **34**, 11168–11184 (2020).
178. Frangogiannis, N. Transforming growth factor- β in tissue fibrosis. *J. Exp. Med.* **217**, e20190103 (2020).
179. Sun, T. et al. TGF β 2 and TGF β 3 isoforms drive fibrotic disease pathogenesis. *Sci. Transl. Med.* **13**, eabe0407 (2021).
180. Atabai, K. et al. Mfge8 diminishes the severity of tissue fibrosis in mice by binding and targeting collagen for uptake by macrophages. *J. Clin. Invest.* **119**, 3713–3722 (2009).
181. Cui, H. et al. Monocyte-derived alveolar macrophage apolipoprotein E participates in pulmonary fibrosis resolution. *JCI Insight* **5**, e134539 (2020).
182. Huang, S. et al. Macrophage PPAR- γ suppresses long-term lung fibrotic sequelae following acute influenza infection. *PLoS ONE* **14**, e0223430 (2019).
183. Ogger, P. P. et al. Itaconate controls the severity of pulmonary fibrosis. *Sci. Immunol.* **5**, eabc1884 (2020).

ACKNOWLEDGEMENTS

CCB is funded by a Sir Henry Dale Fellowship from the Wellcome Trust/Royal Society (206234/Z/17/Z). ASM is funded by the Medical Research Council UK. Images from Servier Medical Art (smart.servier.com) were used in some of the figures.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Calum C. Bain or Andrew S. MacDonald.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the

article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022