

RESEARCH

Open Access

Good performance in the management of acute heart failure in cardiogeriatric departments: the ICREX-94 experience



Emmanuelle Berthelot^{1,2*}, Amaury Broussier^{3,4}, Thibaud Damy^{3,5}, Cristiano Donadio^{1,2,6}, Stephane Cosson⁷, Xavier Rovani⁷, Emmanuel Salengro⁸, Gilles Billebeau⁸, Richard Megbemado⁹, Noomen Rezik⁹, Christian Godreuil¹⁰, Kevin Richard¹¹, Jason Shourick¹², Patrick Assayag^{1,2}, Joel Belmin⁶, Jean Philippe David^{3,13}, Luc Hittinger^{3,5} and for the FINC-94 network

Abstract

Context: A growing number of elderly patients hospitalized for Acute Heart Failure (AHF) are being managed in cardiogeriatrics departments, but their characteristics and prognosis are poorly known. This study aimed to investigate the profile and outcome (rehospitalization at 90 days) of patients hospitalized for AHF in cardiogeriatrics departments in the Val-de-Marne area in the suburbs of Paris, and to compare them to AHF patients hospitalized in cardiology departments in the same area.

Methods: Observational study, ICREX-94, conducted in seven cardiology departments in France and three specific cardiogeriatrics departments in Val-de-Marne.

Results: A total of 308 patients were hospitalized for AHF between October 2017 and January 2019. During the 90 days following discharge, 29.6% patients were readmitted to the hospital. Compared with patients hospitalized in cardiology departments, patients in cardiogeriatrics departments were older ($p < 0.001$), less independent (living more often alone or in an institution) ($p < 0.001$), more often depressed ($p < 0.001$), had more often major neurocognitive disorder ($p < 0.001$), had a higher Human Development Index (HDI, $p < 0.001$), and were less often diagnosed with amyloidosis ($p < 0.001$). There was no difference in outcome whether patients were discharged from cardiology or cardiogeriatrics departments. The most frequent precipitating factors underlying AHF decompensation between the first and second hospitalization were arrhythmia and infection.

Conclusion: AHF patients discharged from cardiogeriatrics departments, compared to cardiology departments, showed clinical differences but had the same prognosis regarding AHF rehospitalization at 90 days.

Keywords: Heart failure, Cardiogeriatrics, Prognosis, Profile, Risk factors

* Correspondence: emmanuelle.berthelot@gmail.com

¹Université Paris Sud, Le Kremlin-Bicêtre, France

²APHP, Department of Cardiology, Hôpital Bicêtre, 78, rue du général Leclerc, 94270 Le Kremlin Bicêtre, France

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Heart failure is a disease that affects about 2% of the population in western countries [1, 2]. With the aging population, the prevalence of acute heart failure (AHF), measured as the annual incidence of hospitalizations for AHF, rises to more than 10% in patients over 70 years of age [3].

After discharge from the hospital for AHF, readmissions are frequent [4, 5]. Although overall hospitalization rates for heart failure have decreased, unplanned readmissions continue to be a common occurrence, with nearly 30% of patients being readmitted within 90 days of discharge [6, 7]. Among older patients, hospitalization is associated with markedly adverse outcomes, including increased mortality, morbidity, and health care expenditures [8].

Often, heart failure patients are older than 75 years and have other common geriatric conditions including frailty, depression, cognitive impairment, malnutrition, disabilities, and chronic diseases other than a heart condition [9, 10]. The management of these patients depends on geriatrics and cardiology particularities. Cardiologists' and geriatricians' awareness and perception of heart failure, comorbidities and functional status can be different and complementary [11]. Indeed, the intervention of cardiologists in the course of care for elderly patients has been shown to improve short-term mortality and readmission outcomes [12, 13]. Calls have been made for a new paradigm in cardiac care for older adults or for closer collaboration between the two specialties [14, 15]. No study has yet established the contribution to HF care of geriatricians with expertise in cardiology.

Some studies have focused on the profile of patients at risk of being rehospitalized [16, 17], and other studies have looked at precipitating factors that trigger acute heart failure [18, 19]. In the context of hospitalization in cardiology and cardiogeriatrics units, the various factors influencing rehospitalization are not clearly established.

We hypothesized that the geriatric conditions of patients hospitalized for AHF could impact the rate and precipitating factors of rehospitalization.

The objectives of this study were (i) to determine the profile and outcome (rehospitalization at 90 days) of patients admitted to Val-de-Marne hospitals for HF, depending on whether they were hospitalized in a cardiology or a specialized cardiogeriatrics department and (ii) to analyze modes and precipitating factors of rehospitalization in the two types of departments.

Methods

The ICRES-94 research was a prospective, non-interventive, observational, transversal, multicentric registry conducted in seven cardiology units and three

cardiogeriatrics units in the Val-de-Marne department (zip code 94) in France.

The present study was conducted in the Val-de-Marne department: 245 sq. km, 1.4 million inhabitants, a mix of residential cities and low-incomes cities (mean HDI 0.58, max 0.78, min 0.35). The Val-de-Marne health system comprises 48 hospitals, with a total capacity of 9500 beds. In 2018, in the 10 hospitals participating in this study, 2393 heart failure admissions were recorded by the "*Caisse Primaire d'Assurance Maladie*" (French health insurance fund), representing 85% of all AHF admissions in Val-de-Marne. In 2016, ten Val-de-Marne cardiology and cardiogeriatrics departments, academic and non-academic, public and private, large and small, interested in HF care, decided to create a network (FINC94) and to collaborate, in order to share their experiences, train healthcare professionals and conduct clinical studies such as this one [20].

There were seven classical cardiology departments, found in both teaching and nonteaching hospitals, public and private, and three cardiogeriatrics departments specialized in heart failure management, with geriatricians who had received specific academic and practical training in cardiovascular medicine. In addition to their geriatrics background, the geriatricians in these units trained for several months in cardiology departments specialized in HF, and have university diplomas in echocardiography and cardiovascular disease of elderly patients. Therefore, they work in close cooperation with the HF team of their departments. There was no specific protocol when patients were hospitalized in cardiology or cardiogeriatrics departments, except to follow 2016 ESC guidelines on HF. Upon admission to cardiogeriatrics departments, an individual and multidisciplinary approach (by geriatricians, physiotherapists, dietitians and social workers) was established, focused on stabilization of comorbidities, return to self-sufficiency and renutrition in addition to specific cardiology follow-up.

There were no specific guidelines to direct patients to a cardiology or cardiogeriatrics department.

Consecutive patients over 18 years of age, hospitalized for acute HF and alive at discharge were eligible for the study. Diagnosis of AHF was based on signs and symptoms of HF— clinical point of view, BNP at admission > 100 pg/ml and heart structure suggesting HF on echocardiograms, as recommended (ESC guidelines). Patients who did not understand the French language were excluded. The study was compliant with Helsinki rules and was approved by the local ethics committee (Commission éthique et déontologie de la Faculté de Médecine Paris-Saclay #20181128163709). All patients gave their informed consent. Informed consent was obtained for all the participants.

Table 1 Baseline characteristics of all patients depending on department (cardiology or cardiogeriatrics)

Department	Overall	Cardiogeriatrics	Cardiology	p
N (%)	308	66 (21.4)	242 (78.6)	
Age, yrs.	75.8 ± 13.5	82.4 ± 9.2	74.0 ± 13.8	< 0.001
Age classes, n(%)				< 0.001
< 45 yrs.	9 (2.9)	0 (0)	9 (3.7)	
45–74 yrs.	125 (40.6)	12 (16.7)	114 (47.1)	
75–85 yrs.	98 (31.8)	26 (39.4)	72 (29.7)	
> 85 yrs.	76 (24.6)	28 (42.4)	48 (19.8)	
Female Gender, n(%)	133 (43.3)	32 (49.2)	101 (41.7)	
Live alone, n(%)	54 (17.5)	24 (36.4)	30 (12.4)	< 0.001
Married, n(%)	70 (22.7)	14 (21.2)	56 (23.1)	0.869
Live with family, n(%)	31 (10.1)	6 (9.1)	25 (10.3)	1
Live in institution, n(%)	3 (1)	3 (4.5)	0	0.009
IDH	0.6 ± 0.1	0.6 ± 0.1	0.54 ± 0.1	< 0.001
Coronary heart disease, n(%)	115 (37.8)	23 (35.4)	92 (38.5)	0.668
Amyloidosis, n(%)	28 (9.2)	1 (1.5)	27 (11.2)	0.014
Hypertension, n(%)	207 (67.9)	44 (67.7)	163 (67.9)	1
Diabetes, n(%)	112 (36.7)	17 (26.2)	95 (39.6)	0.059
Smoker, n(%)	38 (12.5)	8 (12.3)	30 (12.5)	1
Chronic alcohol intake, n(%)	23 (7.5)	5 (7.7)	18 (7.5)	1
Chronic kidney disease, n(%)	141 (46.2)	27 (41.5)	114 (47.5)	0.404
COPD, n(%)	65 (21.3)	14 (21.5)	51 (21.2)	1
History of heart failure, n(%)	203 (66.8)	43 (66.1)	160 (66.9)	1
Stroke, n(%)	47 (15.4)	13 (20)	34 (14.2)	0.25
Atrial fibrillation, n(%)	181 (59.3)	38 (58.5)	143 (59.6)	0.888
Dementia, n(%)	24 (7.9)	20 (30.6)	11 (4.68)	< 0.001
Depression, n(%)	15 (4.9)	9 (13.6)	6 (2.5)	0.001
BMI, kg/m²	27.1 ± 7.2	26.2 ± 6.6	27.2 ± 7.3	0.03
SBP, mmHg	122 ± 20	122 ± 18	123 ± 22	0.855
LVEF classes, n(%)				0.584
> 50%	69 (22.4)	13 (19.7)	56 (23.1)	
40–50%	53 (17.2)	14 (21.2)	39 (16.1)	
< 40%	186 (60.4)	39 (59.19)	147 (60.7)	
BNP at discharge, pg/ml	554 ± 745	618 ± 823	300 ± 112	0.001
Length of stay, days	12.7 ± 10.1	19.3 ± 15.4	11.2 ± 7.3	< 0.001
Creatinemia at discharge, days	134 ± 63	118 ± 38	137 ± 67	0.18
Prescription at discharge, n(%)	292 (95.74)	63 (95.45)	229 (95.82)	0.254
ACEi, n(%)	133 (43.9)	34 (53.1)	99 (41.4)	0.118
ACEisup50, n(%)	53 (17.5)	12 (18.7)	41 (17.1)	0.853
ARNi, n(%)	25 (8.2)	5 (7.8)	20 (8.3)	1
ARNisup50, n(%)	13 (4.3)	5 (7.8)	8 (3.3)	0.157
Diuretic, n(%)	259 (85.2)	51 (79.7)	208 (86.7)	0.169
Dose Diuretic, n(%)	118 ± 190	71 ± 117	131 ± 204	0.002
Beta blocker, n(%)	208 (68.4)	46 (71.9)	162 (67.5)	0.548
Beta blockersup50, n(%)	83 (27.3)	14 (21.9)	69 (28.7)	0.344

Table 1 Baseline characteristics of all patients depending on department (cardiology or cardiogeriatrics) (Continued)

Department	Overall	Cardiogeriatrics	Cardiology	p
MRA, n(%)	61 (20.1)	7 (10.9)	54 (22.5)	0.052
MRAsup50, n(%)	45 (14.8)	6 (9.4)	39 (16.2)	0.234
Ivabradine, n(%)	7 (2.3)	2 (3.1)	5 (2.1)	0.641
LCZ696, n(%)	25 (8.2)	5 (7.8)	20 (8.3)	1
Pacemaker, n(%)	41 (13.4)	16 (24.2)	25 (10.4)	0.007
ICD, n(%)	43 (14.1)	2 (3)	41 (17.1)	0.002
Well Treated, n(%)	24 (7.9)	36 (56.2)	118 (49.2)	0.328

Note. The data are quoted as the number (%), or mean ± sd, "well treated" was defined when patients were treated with BB+ACEi or ARNi
 Abbreviations: ACEi Angiotensin-Converting Enzyme inhibitors, ARNi Angiotensin Receptor Nephilysin inhibitor, BMI Body Mass Index, COPD Chronic Obstructive Pulmonary Disease, HDI Human Development Index, HF Heart Failure, HR Heart Rate, ICD Implantable Cardioverter Defibrillator, MRA Mineralocorticoid Receptor Antagonists, SBP Systolic Blood Pressure, LVEF Left Ventricular Ejection Fraction

Baseline data collection

The following data was collected at inclusion and if patients were rehospitalized: HF type (i.e., right, left, global), etiology of HF, date of diagnosis of HF, clinical characteristics including geriatric comorbidities like dementia and depression, ECG data (sinus rhythm, atrial fibrillation), and biological data such as BNP, haemoglobinemia, and serum creatinine. In addition, the human development index (HDI), which evaluates the progress of a country or a region in the long term, adapted to the Ile-de-France region, was determined by the town of residency. The HDI takes into account three basic dimensions of human development: a long and healthy life (life expectancy), access to knowledge (education) and a decent standard of living (income) [21]. We recorded echocardiographic characteristics, such as left ventricular ejection fraction (LVEF), and medical treatments with

respective doses and whether the patient had a multi-site and/or defibrillator pacemaker. We defined patients as "well-treated" when they had received more than 50% of the target dose of the treatment by ARB and beta blockers.

Follow-up data collection

Patients were followed over 90 days after discharge from hospital by direct phone calls and correspondence. If the patient did not answer, we called the patient's family, caregiver, general practitioner or cardiologist. Rehospitalizations within the 90 days were recorded, with medical reports and the same clinical, ECG and biological variables as on first admission. Cause and mode of rehospitalization were analyzed by Clinical Endpoints Committees (CEC) set up to review all medical reports of rehospitalized patients. Each CEC included one




Fig. 1 Flow chart

cardiologist and one geriatrician trained in endpoint adjudication. All events were reviewed independently by each CEC. Any disagreement between CECs was resolved by a third physician as CEC chairman (EB, KR, LH, CD, TD).

CECs divided hospital admissions in four classes: AHF Planned Rehospitalization, AHF Unplanned Rehospitalization, Non-AHF Planned Rehospitalization, and Non-AHF Unplanned Rehospitalization. For AHF readmissions, the underlying causes were classified by CECs as follows: infection, unstable hypertension, arrhythmia, medical treatment modification, non-adherence, anaemia, myocardial infarction, pulmonary embolism, acute renal failure, very severe chronic heart failure (i.e. “frequent flyer” patients with ≥ 3 hospitalizations in the year or with NTproBNP > 5000 pg/ml).

Statistical analysis

Continuous variables are expressed as median [interquartile range (IQR)], and categorical variables are expressed as number or frequency (percentage). Differences in patient clinical characteristics between cardiology and cardiogeriatrics departments were tested by the χ^2 or fisher test for categorical data and by the Wilcoxon test for continuous data.

Differences in clinical characteristics between patients hospitalized for acute heart failure and non hospitalized patients were obtained with univariate logistic regression and the Wald test.

Finally we produced a Kaplan-Meier curve of readmission for acute heart failure within 90 days depending on

the type of department and did a survival analysis using a univariate cox regression.

A two-sided p -value < 0.05 was considered statistically significant. All statistical analyses were performed using R version 4.

Results

Characteristics of the study population

Three hundred and eight patients were included between October 2017 and January 2019. Patient characteristics are presented in Table 1. Patients were 75.8 ± 13.5 years old, and 133 (43.3%) were women. Regarding comorbidities, 203 (66.8%) patients had a history of HF, 115 (37.8%) of coronary artery disease, 112 (36.7%) of diabetes, 181 (59.3%) were in atrial fibrillation and 28 (9.2%) had been diagnosed with amyloidosis. Sixty-nine (22.4%) had LVEF > 50% and 186 (60.4%) LVEF < 40%. Mean LVEF was $39 \pm 19\%$. For HF treatment, 259 (85.2%) patients took diuretics, 208 (68.4%) beta-blockers, 183 (59.4%) ACE inhibitors or ARNi or LCZ696, and 60 (20.1%) MRA.

Characteristics of patients hospitalized depending on the department

The clinical profiles of AHF hospitalized patients depending on the department (cardiogeriatrics or cardiology) are shown in Table 1. In cardiogeriatrics units, they were older ($p < 0.001$) and less often diagnosed with amyloidosis ($p < 0.001$), and had higher BNP levels at discharge ($p < 0.001$) and longer stays ($p < 0.001$). They more frequently lived alone ($p < 0.001$) or in institutions ($p < 0.001$), with a major neurocognitive disorder ($p <$





Fig. 2 Patients’ outcome at 90 days according to the type of department, cardiologic or cardiogeriatric: AHF planned or unplanned



0.001) or depression ($p < 0.001$). However, their HDI was higher ($p < 0.001$). LVEF was higher in cardiogeriatrics departments ($43 \pm 14\%$ vs $36 \pm 19\%$). Regarding treatment, patients from cardiogeriatrics departments took lower doses of diuretics ($p = 0.002$), and more frequently

had an implanted pacemaker ($p = 0.007$) and less frequently an Implantable Cardioverter-Defibrillator (ICD) ($p = 0.002$) (Table 1). There was no difference between cardiogeriatrics and cardiology units in the prescription of angiotensin-converting enzyme inhibitors (ACEI),

Table 2 Events in the population following discharge, depending on department

Type of Center	Overall	Cardio-geriatrics	Cardiology	p
N	308	66	242	
Reasons and mode of rehospitalization, n(%)				0.53
AHF Unplanned, n(%)	41 (13.3)	8 (12.1)	33 (13.6)	
AHF Planned, n(%)	11 (3.6)	2 (3.0)	9 (3.7)	
Non-AHF Planned, n(%)	25 (8.1)	3 (4.5)	22 (9.1)	
Non-AHF Unplanned, n(%)	14 (4.5)	3 (4.5)	11 (4.5)	
Number of events in the first 90 days, n(%)	91 (29.5)	16 (24.2)	75 (31)	0.11

Note. The data are quoted as the number (%)
 Abbreviations: AHF Acute Heart Failure

angiotensin II receptor blockers (ARBs) or beta-blockers (BB-).

Patient outcomes at 90 days

At 90 days, of the 308 patients discharged from hospital, 52 (17%) were readmitted for AHF and 39 (12.6%) for non-cardiovascular causes (Fig. 1). Of the 52 readmitted for AHF, 41 (13.3%) were unplanned and 11 (3.6%) planned readmissions. Of the 39 readmitted for other reasons, 25 (8.1%) were planned, and 14 (4.5%) unplanned non-AHF rehospitalization. Regarding the department of discharge, cardiogeriatrics or cardiology, there was no statistical difference in the primary endpoint “90 days rehospitalization for AHF” (Fig. 2). Regarding death at 90 days, there were 22 deaths in patients discharged from cardiologic department and 2 deaths in patients discharged from cardiogeriatric department, with no statistical difference ($p = 0.146$).

Compared with cardiology departments, when patients had been initially hospitalized in cardiogeriatrics departments, there was no statistical difference regarding the type of rehospitalization (AHF unplanned, AHF planned, non-AHF planned, non-AHF unplanned) (Fig. 3 and Table 2).

Risk of AHF rehospitalization according to causes

In univariate analysis, after a first AHF rehospitalization, the profiles of patients at risk of being rehospitalized for AHF in our study were analyzed. A past history of HF, amyloidosis, high Heart Rate (HR) at discharge and

intracardiac defibrillator usage was associated with a higher risk of AHF rehospitalization at 90 days (Fig. 4).

Causes of hospitalization in HF patients by department: first and second hospitalization in the 90-day follow-up

The most frequent precipitating factors for AHF decompensation between the first and second hospitalizations (H1 and H2) were arrhythmia (42.3%), infection (30.8%) and very severe symptoms (17.3%) (Table 3 and Fig. 5). When analyzing differences between cardiogeriatrics or cardiology admissions, we found no difference in the causes for hospitalization, which were predominantly arrhythmia and infection.

Discussion

In a multicentric study in the Val-de-Marne area south-east of Paris, we prospectively conducted a comprehensive assessment of AHF patient profiles and the modes and causes of rehospitalization within 90 days, depending on whether patients were hospitalized in cardiogeriatrics or cardiology departments.

To our knowledge, this is the first study comparing patients hospitalized in cardiology vs. cardiogeriatrics with specific geriatrician training.

Our study had two main findings:

- While AHF patients in cardiogeriatrics were older, less independent, less often diagnosed with amyloidosis, more often living alone, more often with major neurocognitive disorder or depression,




Fig. 4 Forest plot of different parameters influencing AHF hospitalization in the overall population

Table 3 Causes of rehospitalization of HF patients depending on department: first rehospitalization (H1) and second rehospitalization (H2) during 90-day follow-up

Type of Center	Overall	Cardio-geriatrics	Cardiology	p
H1: First hospitalization				
Arrhythmias, n(%)	88 (28.6)	15 (22.7)	73 (30.2)	0.283
Infection, n(%)	76 (24.7)	17 (25.8)	59 (24.4)	0.872
Poor adherence to TT, n(%)	17 (5.5)	0 (0)	17 (7.0)	0.029
Medical TT modification, n(%)	2 (0.6)	1 (1.5)	1 (0.4)	0.383
Poor diet observance, n(%)	26 (8.4)	3 (4.5)	23 (9.5)	0.316
Anemia, n(%)	18 (5.8)	7 (10.6)	11 (4.5)	0.076
Myocardial infarction, n(%)	22 (7.1)	6 (9.1)	16 (6.6)	0.589
Pulmonary Embolism, n(%)	2 (0.6)	0 (0)	2 (0.8)	1
Acute Renal Failure, n(%)	23 (7.5)	2 (3.0)	21 (8.7)	0.184
Very Severe HF, n(%)	8 (2.6)	1 (1.2)	7 (2.9)	1
Unstable hypertension, n(%)	6 (1.9)	2 (3.0)	4 (1.6)	0.612
Other, n(%)	60 (19.5)	15 (22.7)	45 (18.6)	0.484
H2: Second hospitalization				
Arrhythmias, n(%)	28 (21.9)	4 (23.5)	24 (21.6)	1
Infections, n(%)	28 (21.9)	2 (11.7)	26 (23.4)	0.36
Poor adherence to treatment, n(%)	8 (6.2)	0 (0)	8 (7.2)	0.596
Medical TT modification, n(%)	2 (1.6)	0 (0)	2 (1.8)	1
Poor regimen adherence, n(%)	6 (4.7)	1 (5.9)	5 (4.5)	0.583
Anemia, n(%)	5 (3.9)	2 (11.8)	3 (2.7)	0.131
Myocardial infarction, n(%)	3 (2.3)	0 (0)	3 (2.7)	1
Pulmonary Embolism, n(%)	1 (0.8)	0 (0)	1 (0.9)	1
Acute Renal Failure, n(%)	7 (5.5)	1 (5.9)	6 (5.4)	1
Very Severe HF, n(%)	2 (0.6)	0 (0)	2 (0.8)	1
Other n(%)	38 (29.7)	8 (47.1)	30 (27)	0.151

Note. The data are quoted as the number (%)
Abbreviations: TT Treatment, HF Heart Failure

but with higher HDIs, there was no statistical difference in the primary endpoint “hospitalization for AHF” depending on the specialty department of discharge.

- The most frequent precipitating factors underlying AHF decompensation between the first and second hospitalization were arrhythmia, infection or very severe symptoms, and it made no difference whether patients were discharged from cardiogeriatrics or cardiology units.


Readmission for HF

The characteristics of our population are similar to those in previous reports on the general population in terms of age, gender, risk factors, coronary artery disease, diabetes mellitus, atrial fibrillation, prescription of diuretics, BB-, ACE-I/ARB/ARNi and MRA [19, 22].

In our study, the readmission rate at 90 days was 29.6%, comparable to rates previously reported for the

same timeframe [6, 7]. We chose to present data at 90 days because the restricted 30-day window has been questioned. Readmissions for HF are a real problem. Strategies intended to reduce rates of premature admission have been developed. Good stabilization of HF can reduce the occurrence of readmission [6, 23]. In addition, data comparing the relative utility of a 30-day window versus other post-discharge timeframes showed limited differences in assessing overall hospital performance [24].

While previous studies demonstrated worse outcomes in elderly patients [24], in the present study at 90 days there was no difference in the rate of readmission for AHF nor of death (Fig. 3). In our study, cardiogeriatrics patients were older, had higher rates of depression and neurocognitive disorders, and lived more frequently alone or in an institution. They had higher BNP levels. Post discharge, they received a lower dosage of diuretics and were more frequently implanted with a



pacemaker and less frequently with an ICD. The absence of difference in the rehospitalization rate of patients from cardiogeriatrics departments vs. cardiology departments may be partly due to the similarity in maintenance therapy (BB-, ACE, i/ARNi), and cardiogeriatrics patients' higher HDI, which may counterbalance the effects of age, dementia or depression on the rate of readmission [25].

Management of elderly HF patients

A recent study concluded that a “cardiogeriatrics model” of managing HF did not improve the prognosis of HF patients at 30 days [26]. There is a room for innovative care for elderly HF patients [15, 16]. Though cardiogeriatrics patients were older and more socially isolated and dependent, with more mood disorders and major neurocognitive disorders, there was no significant difference in rates of readmission for AHF at 90 days compared to cardiology patients. This similarity in prognosis may be linked to a similar efficacy in therapeutic management, a comprehensive and specific multidisciplinary approach and to a longer stay in cardiogeriatrics departments allowing for better stabilization of comorbidities that may lead to rehospitalization [27]. It is also possible that the recruitment and care of our elderly patients through cardiogeriatrics departments may differ from usual geriatrics

departments. Indeed, the cardiogeriatrics departments are characterized by geriatricians with specific competences in cardiology, with easy access to echocardiography and BNP measurement, and in close contact with cardiologists in the cardiology departments of our area.

Comprehensive patient management seems essential to reduce readmissions and thus improve the quality of life of these patients. Achieving this outcome will require training cardiologists to manage multiple morbidities and frailty, and improving the skills of geriatricians in HF management [15, 16]. The use of a frailty score accessible to cardiologists will facilitate the collaboration between cardiologists and geriatricians within the heart team serving the patient [27].

Risk of acute heart failure hospitalization according to profile and causes

In our population, the factors associated with AHF readmissions were: previous AHF history, higher HR at discharge, cardiac amyloidosis, and intracardiac defibrillator use. These factors have previously been shown to influence HF readmissions [17]. Precipitating factors and their contribution to hospitalization of patients with HF have been previously described [7, 18, 19]. In the present study, according to those reports, factors that influenced the most readmissions were infection, cardiac arrhythmia and

severity of heart failure. Interestingly, when precipitating factors were analyzed for the second readmissions, factors remained similar for some patients, but differed for others, showing the complexity and heterogeneity of the heart failure process in different patients. Moreover, the same frequent causes were found in both cardiology and cardiogeriatrics departments.

Limitations

The present study has several limitations.

The study was performed in the Val-de-Marne department of France, and therefore may have limited implications for other territories with different environments or healthcare systems. The numbers of patients recruited in cardiology and cardiogeriatrics departments were not evenly balanced, thus limiting the strength of our results; however, the data was recorded prospectively over the same period of time. The percentage of patients with preserved ejection fraction appears lower than usually observed in elderly patient studies [26, 28]. Due to the mode of recruitment, our study includes relatively few patients and may lack power. In a future work, the patient cohort could be larger, better distributed between the admissions departments. Results could include mortality, and more diverse geriatric outcomes, such as functional outcomes, necessity of (nursing) home care after first admission or patient satisfaction data.

Some elements that could explain rehospitalizations were not noted (multidomain assessment of frailty) and could be the subject of future work.

Conclusion

While clinically different, AHF patients discharged from cardiogeriatrics compared to cardiology departments, had similar prognosis regarding rehospitalization for AHF at 90 days. Among other possibilities, care provided in cardiogeriatrics departments by geriatricians with cardiology/HF expertise may have played a role, suggesting the effectiveness of innovative management of elderly HF patients. The main precipitating factors underlying AHF decompensation for the first rehospitalization were arrhythmia, infection, and very severe symptoms, in both cardiology and cardiogeriatrics departments, and remained proportionally similar during the second hospitalization. Further studies are needed to confirm these conclusions.

Abbreviations

ACEi: Angiotensin-Converting-Enzyme inhibitors; ARNI: Angiotensin Receptor Neprilysin inhibitor; BMI: Body Mass Index; COPD: Chronic Obstructive Pulmonary Disease; HDI: Human Development Index; HF: Heart Failure; HR: Heart Rate; ICD: Implantable Cardioverter-Defibrillator; MRA: Mineralocorticoid Receptor Antagonists; SBP: Systolic Blood Pressure; LVEF: Left Ventricular Ejection Fraction

Acknowledgements

We are indebted to the following persons and agencies, without support this study could have not been successfully completed: Sophie Bataille &

Charlotte Rodier Daval from the Agence Regionale de Sante Ile de France; David Cheutin, Minh-Thu Vu-Nguyen from the Caisse Primaire d'Assurance Maladie Val de Marne; Michel Marty from the Caisse primaire d'Assurance Maladie Ile de France.

Authors' contributions

E.B. and T. D and L. H and A.B. wrote the main manuscript. J.S. did the statistical analysis and the figures T.D. and E.B. Did the tables and figures EB, TD, CD, SC, XR, ES, GB, RM, NR, CG, KR, JS, PA, JB, JPD, LH, collected data. E. B, T.D., L.H., K.R. reviewed all the data and were the clinical adjudication committee. The authors read and approved the final manuscript.

Funding

The study was funded by Agence régionale de santé Ile de France and the Association pour la Recherche MultiDisciplinaire en Cardiologie-CHU Henri Mondor - 94000 Créteil.

Availability of data and materials

Data and materials are available: Raw data are available on the following link: https://drive.google.com/file/d/1bHDRR_ZOPDXPoulMZG08E-9AtFKM7ryR/view?usp=sharing

Declarations

Ethics approval and consent to participate

Ethics approval and consent to participate were obtained: The study was compliant with Helsinki rules and was approved by the local ethics committee (commission éthique et déontologie de la Faculté de Médecine Paris-Saclay #20181128163709). Informed consent was obtained for all the participants.

Consent for publication

Consent for publication is approved by all the Authors.

Competing interests

There was no Competing interests.

Author details

¹Université Paris Sud, Le Kremlin-Bicêtre, France. ²APHP, Department of Cardiology, Hopital Bicêtre, 78, rue du général Leclerc, 94270 Le Kremlin Bicêtre, France. ³Université Paris Est, Créteil, INSERM, IMRB, Equipe CEpiA, F-94010 Créteil, France. ⁴Department of Geriatrics, AP-HP, Henri-Mondor/Emile-Roux hospitals, F-94456 Limeil-Brevannes, France. ⁵Department of Cardiology, heart failure and amyloidosis unit, Referral Center For Cardiac Amyloidosis, AP-HP, Henri-Mondor/Albert-Chenevier hospitals, F-94010 Créteil, France. ⁶Department of geriatrics, AP-HP, Hôpital Charles Foix and Sorbonne Université, F-94200 Ivry-sur-Seine, France. ⁷Hôpital privé Paul Déginge, 4 avenue Marx Dormoy, F-94500 Champigny-sur-Marne, France. ⁸Centre Hospitalier de Villeneuve St Georges, 40 allée de la Source, F-94190 Villeneuve-Saint-Georges, France. ⁹Hôpital Sainte Camille, 2 rue des Pères Camilliens, F-94360 Bry-sur-Marne, France. ¹⁰Hôpital d'Instruction des Armées Bégin, 69 avenue de Paris, F-94160 Saint-Mandé, France. ¹¹AP-HP Centre hospitalier Chenevier, 40 rue de Mesly, F-94000 Créteil, France. ¹²AP-HP, BioStatisticien, Paris Sud, Paris, France. ¹³Department of Internal Medicine and Geriatrics, AP-HP, Henri Mondor hospitals, F-94010 Créteil, France.

Received: 28 December 2020 Accepted: 30 March 2021

Published online: 01 May 2021

References

- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, et al. Authors/Task Force Members. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2016;37:2129–200.
- Maggioni AP, Dahlstrom U, Filippatos G, Chioncel O, Crespo Leiro M, Drozd J, et al. Heart failure association (HFA) of the European Society of Cardiology. EURObservational research Programme: regional differences and 1-year follow-up results of the heart failure pilot survey (ESC-HF pilot). *Eur J Heart Fail*. 2013;15(7):808–17. <https://doi.org/10.1093/eurjhf/hft050>.

3. Mozaffarian D, Benjamin EJ, Go AS. Heart disease and stroke Statistics-2016 update: a Report from the American Heart Association. *Circulation*. 2016;133:e38–48.
4. Tuppin P, Cuerq A, de Peretti C, Fagot-Campagna A, Danchin N, Juillièrre Y, et al. Two-year outcome of patients after a first hospitalization for heart failure: a national observational study. *Arch Cardiovasc*. 2014;107:165–8.
5. Carson PE, Anand IS, Win S, Rector T, Haass M, Lopez-Sendon J, et al. The hospitalization burden and post-hospitalization mortality risk in heart failure with preserved ejection fraction: results from the I-PRESERVE trial (Irbesartan in heart failure and preserved ejection fraction). *JACC Heart Fail*. 2015;3(6):429–41. <https://doi.org/10.1016/j.jchf.2014.12.017>.
6. Fonarow GC, Abraham WT, Albert NM, Stough WG, Gheorghiane M, Greenberg BH, et al. Association between performance measures and clinical outcomes for patients hospitalized with heart failure. *JAMA*. 2007;297(1):61–70. <https://doi.org/10.1001/jama.297.1.61>.
7. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the medicare fee-for-service program. *N Engl J Med*. 2009;360(14):1418–28. <https://doi.org/10.1056/NEJMsa0803563>.
8. Shah RU, Tsai V, Klein L, Heidenreich PA. Characteristics and outcomes of very elderly patients after first hospitalization for heart failure. *Circ Heart Fail*. 2011;4(3):301–7. <https://doi.org/10.1161/CIRCHEARTFAILURE.110.959114>.
9. Cannon JA, McMurray JJ, Quinn TJ. "Hearts and minds": association, causation and implication of cognitive impairment in heart failure. *Alzheimers Res Ther*. 2015;7(1):22. <https://doi.org/10.1186/s13195-015-0106-5>.
10. Chaudhry SI, Mc Avay G, Chen S, Whitson H, Newman AB, Krumholz HM, et al. Risk factors for hospital admission among older persons with newly diagnosed heart failure: findings from the cardiovascular health study. *J Am Coll Cardiol*. 2013;6:635–42.
11. Remme WJ, McMurray JJ, Hobbs FD, Cohen-Solal A, Lopez-Sendon J, Boccanelli A, et al. Awareness and perception of heart failure among European cardiologists, internists, geriatricians, and primary care physicians. *Eur Heart J*. 2008;29(14):1739–52. <https://doi.org/10.1093/eurheartj/ehn196>.
12. Selim AM, Mazurek JA, Iqbal M, Wang D, Negassa A, Zolty R, et al. Mortality and readmission rates in patients hospitalized for acute decompensated heart failure: a comparison between cardiology and general-medicine service outcomes in an underserved population. *Clin Cardiol*. 2015;38(3):131–8. <https://doi.org/10.1002/clc.22372>.
13. Tabit CE, Coplan MJ, Spencer KT, Alcaïn CF, Spiegel T, Vohra AS, et al. Cardiology Consultation in the Emergency Department Reduces Re-hospitalizations for Low-Socioeconomic Patients with Acute Decompensated Heart Failure. *Am J Med*. 2017;130:1112 e17–1112.e31.
14. Dewan P, Jackson A, Jhund PS, et al. The prevalence and importance of frailty in heart failure with reduced ejection fraction — an analysis of PARA DIGM-HF and ATMOSPHERE [published online ahead of print, Apr 30, 2020]. *Eur J Heart Fail*. 2020;22(11):2123–33.
15. Azad NA, Mielniczuk L. A Call for Collaboration: Improving Cardiogeriatric Care. *Can J Cardiol*. 2016;32(9):1041–4. <https://doi.org/10.1016/j.cjca.2016.01.025>.
16. Vader JM, LaRue SJ, Stevens SR, Mentz RJ, DeVore AD, Lala A, et al. Timing and causes of readmission after acute heart failure hospitalization—insights from the heart failure network trials. *J Card Fail*. 2016;22(11):875–83. <https://doi.org/10.1016/j.cardfail.2016.04.014>.
17. Quock TP, Yan T, Tieu R, D'Souza A, Broder MS. Untangling the clinical and economic burden of hospitalization for cardiac amyloidosis in the United States. *Clinicoecon Outcomes Res*. 2019;11:431–9. <https://doi.org/10.2147/CEOR.S207127>.
18. Fonarow GC, Abraham WT, Albert NM, Stough WG, Gheorghiane M, Greenberg BH, et al. Factors identified as precipitating hospital admissions for heart failure and clinical outcomes: findings from OPTIMIZE-HF. *Arch Intern Med*. 2008;168(8):847–54. <https://doi.org/10.1001/archinte.168.8.847>.
19. Opasich C, Febo O, Riccardi PG, Traversi E, Forni G, Pinna G, et al. Concomitant factors of decompensation in chronic heart failure. *Am J Cardiol*. 1996;78(3):354–7. [https://doi.org/10.1016/S0002-9149\(96\)00294-9](https://doi.org/10.1016/S0002-9149(96)00294-9).
20. Galinier M, Roubille F, Berdague P, et al. Telemonitoring versus standard of care in heart failure: a randomised multicentre trial [published online ahead of print, 2020 May 21]. *Eur J Heart Fail*. 2020. <https://doi.org/10.1002/ehf.1906>.
21. United Nations Development Programme (UNDP). Human Development Report 2016. New York; 2016b.
22. Logeart D, Isnard R, Resche-Rigon M, Seronde MF, de Groote P, Jondeau G, et al. Heart failure of the French Society of Cardiology. Current aspects of the spectrum of acute heart failure syndromes in a real-life. OFICA study. *Eur J Heart Fail*. 2013;15(4):465–76. <https://doi.org/10.1093/eurjhf/hfs189>.
23. Vaduganathan M, Bonow RO, Gheorghiane M. Thirty-day readmissions: the clock is ticking. *JAMA*. 2013;309(4):345–6. <https://doi.org/10.1001/jama.2012.05110>.
24. Sato T, Yamauchi H, Kanno Y, Suzuki S, Yoshihisa A, Yamaki T, et al. Comparisons of prognostic factors between Young and elderly patients with chronic heart failure. *Geriatr Gerontol Int*. 2015;15(4):435–42. <https://doi.org/10.1111/ggi.12293>.
25. Arbaje AI, Wolff JL, Yu Q, Powe NR, Anderson GF, Boulton C. Post-discharge environmental and socioeconomic factors and the likelihood of early hospital readmission among community-dwelling Medicare beneficiaries. *Gerontologist*. 2008;48(4):495–504. <https://doi.org/10.1093/geront/48.4.495>.
26. Mulligan A, Seman M, Abramowski R, Falls R, Scholes E, Vogrin S, et al. Cardio-geriatric model of care in acute heart failure: initial experience of a multidisciplinary approach in complex elderly patients *Intern. Med J*. 2020;50(4):488–92. <https://doi.org/10.1111/imj.14793>.
27. Vitale C, Jankowska E, Hill L, Piepoli M, Doehner W, Anker SD, et al. Heart Failure Association/European Society of Cardiology position paper on frailty in patients with heart failure. *Eur J Heart Fail*. 2019;21:1299–305.
28. Garnick DW, DeLong ER, Luft HS. Measuring hospital mortality rates: are 30-day data enough? *Ischemic Heart Disease Patient Outcomes Research Team. Health Serv Res*. 1995;29(6):679–95.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

