

ORAL PRESENTATION

Open Access

Familial chilblain lupus caused by an activating mutation in STING

N König¹, C Fiehn², H-M Lorenz^{2,3}, MA Lee-Kirsch^{1*}

From 8th International Congress of Familial Mediterranean Fever and Systemic Autoinflammatory Diseases Dresden, Germany. 30 September - 3 October 2015

Familial chilblain lupus is a monogenic form of cutaneous lupus erythematosus characterized by cold-induced cutaneous lesions at acral location. It is caused by loss-of-function mutations in the nucleic acid metabolizing enzymes TREX1 or SAMHD1. Gain-of-function mutations in STING (stimulator of Interferon genes) have been described in an infancy-onset autoinflammatory syndrome with fever, inflammatory cutaneous lesions and interstitial lung disease.

Here we report on a family with dominant chilblain lupus over 4 generations. Affected family members presented with acral inflammatory and partially necrotizing lesions beginning in early childhood. In some cases, low-titered ANAs and immune complexes were detectable. The family tested negative for TREX1 or SAMHD1 mutations. Exome sequencing revealed a heterozygous STING mutation segregating with chilblain lupus in the family. The mutation affects a highly conserved residue within the STING dimer interface and is predicted to be pathogenic. Quantitative RT-PCR analysis showed an increased expression of IFN-stimulated genes in blood cells of affected family members suggesting that the identified mutation has an activating effect on type I IFN signaling. Taken together, our findings demonstrate that gain-of-function mutations in STING can cause familial chilblain lupus and expand the spectrum of type I interferonopathies.

Authors' details

¹Technische Universität Dresden, Molekulare Pädiatrie, Klinik für Kinder- und Jugendmedizin, Dresden, Germany. ²ACURA-Rheumazentrum, Baden-Baden, Germany. ³Universität Heidelberg, Sektion Rheumatologie, Medizinische Klinik V, Heidelberg, Germany.

Published: 28 September 2015

¹Technische Universität Dresden, Molekulare Pädiatrie, Klinik für Kinder- und Jugendmedizin, Dresden, Germany

Full list of author information is available at the end of the article

doi:10.1186/1546-0096-13-S1-O62

Cite this article as: König et al.: Familial chilblain lupus caused by an activating mutation in STING. *Pediatric Rheumatology* 2015 13(Suppl 1): O62.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

