

Corrigendum

Corrigendum to “Periorbital facial rejuvenation; applied anatomy and pre-operative assessment” [J. Curr. Ophthalmol. 29 (3), (September 2017) 154–168]

Mohsen Bahmani Kashkouli*, Parya Abdolalizadeh, Navid Abolfathzadeh, Hamed Sianati, Maria Sharepour, Yasaman Hadi

Eye Research Center, Oculo-Facial Plastic Surgery, Rassoul Akram Hospital, Iran University of Medical Sciences, Tehran, Iran

Received 28 November 2017; accepted 29 November 2017

Available online 6 March 2018

The authors regret to inform a few displaced figures in our manuscript. While text and figure legends are correct, the following 4 figures should be re-placed.

Figure 15 should be figure 12.

Figure 12 should be figure 13.

Figure 13 should be figure 14.

Figure 14 should be figure 15.

The correctly renumbered figures are also provided below.



Fig. 12. The two most common causes of lateral hooding are lateral eyebrow ptosis (left, arrows) and lacrimal gland prolapse (right, arrow).

DOI of original article: <https://doi.org/10.1016/j.joco.2017.04.001>.

* Corresponding author. Rassoul Akram Hospital, Sattarkhan-Niyesh St., Tehran, 14455-364, Iran.

E-mail addresses: info@eyeplasticsurgeries.com, mkashkouli2@gmail.com (M.B. Kashkouli).

Peer review under responsibility of the Iranian Society of Ophthalmology.

Authors obtain inform consents from the patients to publish their photoes.

<https://doi.org/10.1016/j.joco.2017.11.013>

2452-2325/Copyright © 2017, Iranian Society of Ophthalmology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Fig. 13. Eyelid-globe vector assessment. In the lateral view, a line dropped from the supraorbital rim to the infraorbital rim. Positive vector is when the cornea is posterior to this line (left), neutral vector is when they touch (middle), and negative vector is when the corneal apex is anterior to the line (right).



Fig. 14. Asymmetric eyebrow ptosis and higher tarsal plate show on the left side point out the possibility of left upper blepharoptosis.



Fig. 15. While patients with lower hair line (left) are good candidates for endoscopic upper face lifting, other forehead and eyebrow procedures should be sought for patients with high hairline (right).

The authors would like to apologize for any inconvenience caused.