

## Review Article

# Effect of Tai Chi on Cognitive Function among Older Adults with Cognitive Impairment: A Systematic Review and Meta-Analysis

Renjun Gu <sup>1</sup>, Yujia Gao,<sup>2</sup> Chunbing Zhang,<sup>3</sup> Xiaojuan Liu,<sup>4</sup> and Zhiguang Sun <sup>1</sup>

<sup>1</sup>Jiangsu Provincial Second Chinese Medicine Hospital, The Second Affiliated Hospital of Nanjing University of Chinese Medicine, Nanjing 210017, China

<sup>2</sup>Rehabilitation Medical Center, Jiangsu Province Hospital, Nanjing 210029, Jiangsu, China

<sup>3</sup>Jiangsu Hospital of Chinese Medicine, Nanjing 210046, Jiangsu, China

<sup>4</sup>The First School of Clinical Medicine, Nanjing University of Chinese Medicine, Nanjing 210023, Jiangsu, China

Correspondence should be addressed to Zhiguang Sun; zhiguangsun2020@163.com

Received 28 December 2020; Revised 20 July 2021; Accepted 22 July 2021; Published 5 August 2021

Academic Editor: Yingchun Zeng

Copyright © 2021 Renjun Gu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Cognitive decline occurs in all persons during the aging process and drugs can only alleviate symptoms and are expensive. Some researches demonstrated that Tai Chi had potential in preventing cognitive decline while others' results showed Tai Chi had no influence on cognitive impairment. Therefore, we conduct a systematic review and meta-analysis to assess the efficacy and safety of cognitive impairment patients practicing Tai Chi. **Methods.** A comprehensive literature search was carried out in multiple databases, including PubMed, Cochrane, MEDLINE (Ovid), Web of Science, Embase, Scopus, PsycInfo (Ovid), CKNL, Wan Fang, VIP, SinoMed, and ClinicalTrials, from their inception to 1 July 2020 to collect randomized controlled trials about practicing Tai Chi for patients with cognitive impairment. Primary outcomes included changes of cognitive function and secondary outcomes included changes of memory functions. Data were extracted by two independent individuals and Cochrane Risk of Bias tool version 2.0 was applied for the included studies. Systematic review and meta-analysis were performed by RevMan 5.3 software. **Results.** The results included 827 cases in 9 studies, of which 375 were in the experimental group and 452 were in the control group. Meta-analysis showed that Mini-Mental State Examination WMD = 1.52, 95% CI [0.90, 2.14]; Montreal Cognitive Assessment WMD = 3.5, 95% CI [0.76, 6.24]; Clinical Dementia Rating WMD = -0.55, 95% CI [-0.80, -0.29]; logical memory delayed recall WMD = 1.1, 95% CI [0.04, 2.16]; digit span forward WMD = 0.53, 95% CI [-0.65, 1.71]; and digit span backward WMD = -0.1, 95% CI [-0.38, 0.19]. No adverse events were reported in the included articles. **Conclusion.** There is limited evidence to support that practicing Tai Chi is effective for older adults with cognitive impairment. Tai Chi seems to be a safe exercise, which can bring better changes in cognitive function score.

## 1. Introduction

While social aging is a trend, cognitive decline can occur for everyone. Eventually, this may result in mild cognitive impairment and dementia [1]. Mild cognitive impairment occurs along a continuum from normal cognition to dementia [2]. It is widely recognized as the intermediate stage of cognitive impairment between the changes seen in normal cognitive aging and dementia [3]. At present, drugs for cognitive impairment can only alleviate the symptoms of cognitive disorders and their price is usually high. Therefore, complementary and alternative therapies have become a hot

research topic for improving cognitive impairment in recent years [4].

Tai Chi has a long history and culture. Participants take deep breathing and mental concentration in order to carrying out smooth and continuous body movements [5]. It combines Chinese martial arts and meditative movements that promote balance of mind and body for healing [6]. Physical exercise and fitness have been proposed as potential factors that may promote healthy cognitive aging [7] and aerobic exercise was proven to improve cognitive function in adults with neurological disorders [8]. In recent years, long-term cognitive training and physical exercise had been

confirmed its benefits for delaying the cognitive decline for the elderly [9, 10]. Tai Chi might improve memory and executive function in older adults with amnesic-mild cognitive impairment, possibly via an upregulation of brain-derived neurotrophic factor [11, 12]. The studies suggested Tai Chi has impacts on global cognitive functions, visuospatial skills, semantic memory, verbal learning memory, and self-perception of memory [13]. It may also have direct benefits on enhancing attention and executive functions [14].

At present, a growing body of evidence supports that Tai Chi may help improve cognitive function and mental well-being for older adults with mild dementia [15, 16]. It also proved that Tai Chi has psychophysiological benefits for motor coordination and memory [17–20]. However, some studies reported no significant differences in assessment of cognitive function [21]. Previous meta-analysis revealed that Tai Chi had no influence on individuals with cognitive impairment [22]. In addition, previous meta-analysis only searched the English databases [23] while Tai Chi is most practiced in China. Therefore, we will conduct a meta-analysis and systematic review without language limitation to assess the effect of Tai Chi on cognitive function among older adults with cognitive impairment.

## 2. Information and Methods

**2.1. Study Protocol.** This systematic review and meta-analysis followed the preferred reporting items for systematic reviews and meta-analyses (PRISMA) of 2015 guideline [24]. The protocol was registered at PROSPERO (<http://www.crd.york.ac.uk/PROSPERO>), registration number: CRD42020171559.

**2.2. Search Strategy.** Electronic literature searches were performed in the database of PubMed, Cochrane, MEDLINE (Ovid), Web of Science, Embase, Scopus, PsycInfo (Ovid), CKNI, Wan Fang, VIP, SinoMed, and ClinicalTrials from inception to July 2020. Search strategy of PubMed is shown in Figure 1.

**2.3. Inclusion Criteria.** (a) Published literature; (b) RCTs; (c) inclusion of people with cognitive impairment; (d) people over 65 years old according to classification for older adults by World Health Organization in 2020 [25]; (e) practicing Tai Chi for more than one month but no more than one year; (f) interventions using Tai Chi as a main treatment; the combination therapy of Tai Chi and other interventions compared with the same other interventions alone were also included; and (g) reporting more than one of the following primary or secondary outcomes.

**2.4. Exclusion Criteria.** (a) Nonclinical studies (experimental and basic studies); (b) observational or retrospective studies; (c) lack of sufficient information on baseline or primary or secondary outcome data; (d) narrative reviews, systematic

reviews, case reports, letters, editorials, clinical guidelines, and commentaries.

**2.5. Primary Outcome.** Any change in cognitive function (such as MMSE, MoCA, and CDR).

**2.6. Secondary Outcome.** Any change in memory function (such as LMD, DSF, and DSB).

**2.7. Patient and Public Involvement.** Neither patients nor public were involved in the design of this study. This systematic review and meta-analysis did not recruit any patients.

**2.8. Data Collection.** Data were extracted by two independent reviewers (RG and YG) using a standardized form including study demographics, baseline characteristics, study design, intervention methods, outcome measures, and results. We resolved any disagreement through discussion and we consulted a third review author (CZ or XL).

**2.9. Bias Risk Assessment.** According to the risk of bias assessment tool from the Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019), two authors independently assessed the risk of bias of the included study, and any conflicts were resolved through consensus. Bias risk assessment was evaluated from the following seven items: random sequence generation, assignment concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other bias. These items are described as green, yellow, and red colors and “+”, “-”, and “?”. The symbols indicate “low”, “high”, and “unclear” risk of bias.

**2.10. Statistical Analysis.** The statistical analyses were performed by using Review Manager software (RevMan version 5.3, Cochrane Collaboration, Oxford, UK). Weighted mean difference (WMD) and 95% CI were used as the effect quantity to merge the continuous variables included in the study.  $P$  value and  $I^2$  statistic were used to test heterogeneity between trial results. When more than two articles were included, heterogeneity was considered. If the  $I^2$  was  $>50\%$ , the random effect model was applied according to the clinical heterogeneity. Subgroup analysis was used to evaluate the source of heterogeneity. The statistical calculation process was completed by RevMan 5.3 software.

## 3. Results

**3.1. Literature Search.** Initial searches generated 1316 related literatures. According to the inclusion criteria and exclusion criteria, 9 literatures were included [16, 21, 26–32] (see Figure 2).

- #1. Search "Tai Ji"[Mesh]
- #2. Search ((((((((((Tai-ji) OR Tai Chi) OR Chi, Tai) OR Tai Ji Quan) OR Ji Quan, Tai) OR Quan, Tai Ji) OR Taiji) OR Taijiquan) OR T'ai Chi) OR Tai Chi Chuan
- #3. #1 OR #2
- #4. Search "Cognitive Dysfunction"[Mesh]
- #5. Search (((((((((((((((((((Cognitive Dysfunctions) OR Dysfunction, Cognitive) OR Dysfunctions, Cognitive) OR Cognitive Impairments) OR Cognitive Impairment) OR Impairment, Cognitive) OR Impairments, Cognitive) OR Mild Cognitive Impairment) OR Cognitive Impairment, Mild) OR Cognitive Impairments, Mild) OR Impairment, Mild Cognitive) OR Impairments, Mild Cognitive) OR Mild Cognitive Impairments) OR Mild Neurocognitive Disorder) OR Disorder, Mild Neurocognitive) OR Disorders, Mild Neurocognitive) OR Mild Neurocognitive Disorders) OR Neurocognitive Disorder, Mild) OR Neurocognitive Disorders, Mild) OR Cognitive Decline) OR Cognitive Declines) OR Decline, Cognitive) OR Declines, Cognitive) OR Mental Deterioration) OR Deterioration, Mental) OR Deteriorations, Mental) OR Mental Deteriorations
- #6. #4 OR #5
- #7. #3 AND #6

FIGURE 1: Literature search strategy.




FIGURE 2: Flowchart of study selection.

3.2. *Characteristics of the Study.* Nine articles [16, 21, 26–32] contained 375 cases in the experimental group and 452 cases in the control group (see Tables 1 and 2).

3.3. *Risk of Bias.* The results of the risk of bias assessment of the 9 studies [16, 21, 26–32] are summarized in Figure 3. Three literatures [28, 29, 31] were scored as high risk without

TABLE 1: Characteristics of the literatures.

Study	Exp. average age/range	Exp. group number	Con. average age/range	Con. group number	Exp. group method	Con. group method	Duration of Tai Chi	Country	Measure	Research designs
Kasal 2010 [26]	73.54	13	74.54	13	Tai Chi	N/A	6 months	Brazil	5.6	RCT
Lam 2012 [27]	77.2	92	78.3	169	Tai Chi + stretching and relaxation exercises	Stretching and relaxation exercises	12 months	China	1.3.4.5.6	RCT
Li 2014 [28]	75	22	77	24	Tai Chi	N/A	14 weeks	China	1	RCT
Tai 2016 [21]	70.21	14	76.3	10	Tai Chi	Nonhealth-related social activities	6 weeks	China	1.3	RCT
Sungkarat 2017 [29]	68.3	33	67.5	33	Tai Chi + education	Education	15 weeks	Thailand	4	RCT
Siu 2018 [30]	—	80	—	80	Tai Chi	N/A	16 weeks	China	1	RCT
Huang 2019 [16]	81.9	36	81.9	38	Tai Chi + routine treatments	Routine treatments	10 months	China	1.2	RCT
Wang 2019 [31]	65–69 (3) 70–79 (32) 80–85 (11)	54	65–69 (2) 70–79 (30) 80–85 (16)	54	Tai Chi	N/A	6 months	China	1.4	RCT
Bao 2019 [32]	65.62	31	68.22	31	Tai Chi + health education	Health education	6 months	China	1.2	RCT

Measure: 1, Mini-Mental State Examination; 2, Montreal Cognitive Assessment; 3, Clinical Dementia Rating; 4, logical memory delayed recall score; 5, digit span forward; 6, digit span backward.

TABLE 2: Characteristics of the literature’s inclusion criteria of cognitive impairment.

Study	Inclusion criteria of cognitive impairment	Style of Tai Chi
Kasal 2010 [26]	(i) Memory complaint offered by the patient or by family members over the previous year; (ii) screening score of the Rivermead Behavioral Memory Test lower than 10; (iii) Mini-Mental State Examination (MMSE) within normality, corrected by educational level	Yang style
Lam 2012 [27]	(i) CDR of 0.5 or (ii) neuropsychological criteria for amnesic-mild cognitive impairment (MCI) with subjective cognitive complaints [21]; objective memory impairment with reference to delayed recall of list learning test at greater than or equal to 1.5 SD below education- and age-matched subjects with CDR 0; (iii) no previous regular practice of Tai Chi or other mind-body exercise for more than 6 months	Yang 24-form style
Li 2014 [28]	(i) Having MMSE scores between 20 and 30	N/A
Tai 2016 [21]	(i) Alzheimer with a Clinical Dementia Rating (CDR) score of 0.5–1; (ii) upper limb mobility sufficient to perform requisite finger-pointing tasks, such as flexing and extending the shoulder, elbow, wrist, and fingers	Yang style
Sungkarat 2017 [29]	(i) Petersen’s criteria for diagnosing amnesic multiple-domain MCI (a-MCI) had scores of 24 or greater on the Mini-Mental State Examination (MMSE) and less than 26 on the Montreal Cognitive Assessment (MoCA), had adequate memory if cued, and comprehended instructions required for study participation	10-form style
Siu 2018 [30]	(i) The CMMSE screening score ranging from 19 to 28, which was corrected based on educational level ( $\geq 18$ for illiterate respondents and $\geq 22$ for those having received more than two years of schooling)	Yang style
Huang 2019 [16]	(i) Diagnosed with dementia based on the diagnostic criteria 128 of the Diagnostic and Statistical Manual of Mental 129 Disorders, 4th edition; (ii) a clinical dementia 130 rating score $< 2$	N/A
Wang 2019 [31]	(i) According to the diagnostic criteria set by the National Institute on Aging and the Alzheimer’s Association (NIA, AA), the patients were screened as MCI, i.e., subjective cognitive function: the patients who complained or knew about cognitive impairment; (ii) objective cognitive function: according to the Peking Union Medical College, version of the total score of MoCA-p is 25 for the elderly aged 65–79 and 21–24 for the elderly aged 80–85; (iii) the total score of activities of daily living (ADL) is $\leq 26$ , and the complex engineering daily living (ADL) is $\geq 10$	8-form style
Bao 2019 [32]	(i) Having memory decline; (ii) the course of disease was more than 3 months; Global Deterioration Scale (GDS) was 2–3, Clinical Dementia Rating Scale was 0.5, memory test score was below 1.5 standard deviation of age and education matched control group, MMSE score met illiteracy (18–21), primary school culture (21–24), secondary school culture (25–27), and daily life ability score was lower than 26; (iii) memory impairment and other aspects of cognitive function retention	Yang style

using random sequence generation. All literatures did not describe detection bias. One article [27] was scored as high risk with incomplete outcome data. All trials measured outcomes listed in their studies and reported on all expected outcome measures of interest (low risk of bias).

**3.4. Mini-Mental State Examination.** Six literatures [16, 21, 27, 28, 30, 32] reported the MMSE. Subgroup analysis was carried to analyze heterogeneity. American Academy of Neurology guideline reported that short-term exercise training (6 months) is likely to improve cognitive measure [35]. Therefore, we divide MMSE into different time periods in order to assessing short-term (less than 6 months) and long-term (more than 6 months) effects. In terms of practicing Tai Chi for less than 6 months (including 6 months), the combined effect is  $WMD = 1.81$ , 95% CI [1.32, 2.30],  $P < 0.05$ . The data were statistically significant. On the other hand, the result of practicing Tai Chi for more than 6 months is  $WMD = 0.61$ , 95% CI [-0.16, 1.38],  $P = 0.12$ , which was not statistically significant. The total combined effect is  $WMD = 1.52$ , 95% CI [0.90, 2.14],  $I^2 = 63%$ , and the data were statistically significant. It indicated that practicing Tai Chi less than 6 months (including 6 months) might improve MMSE for patients with cognitive impairment (see Figure 4).

**3.5. Montreal Cognitive Assessment.** Three literatures described the MoCA [16, 31, 32]. The combined effect was  $WMD = 3.5$ , 95% CI [0.76, 6.24],  $P < 0.05$ . The data was statistically significant (see Figure 5).

**3.6. Clinical Dementia Rating.** Two literatures reported CDR [21, 27]. The combined effect was  $WMD = -0.55$ , 95% CI [-0.80, -0.29],  $P < 0.05$ , which was statistically significant (see Figure 6).

**3.7. Logical Memory Delayed Recall Score.** Three literatures reported LMD [27, 29, 31]. One article [27] described practicing Tai Chi for more than 6 months, and the MD = 0.4, 95% CI [-0.19, 0.99],  $P = 0.18$ , which had no statistical significance. In terms of practicing Tai Chi for less than 6 months,  $WMD = 1.53$ , 95% CI [0.99, 2.08],  $P < 0.05$ . The result was statistically significant, and it indicated that practicing Tai Chi less than 6 months might improve LMD (see Figure 7).

**3.8. Digit Span Forward.** Two experiments [26, 27] mentioned DSF, and the combined effect amount was  $WMD = 0.53$ , 95% CI [-0.65, 1.71],  $P = 0.38$ . It included 105 cases in the experimental group and 182 cases in the control group. The comparison results were not statistically significant and the evidence level was low (see Figure 8).

**3.9. Digit Span Backward.** DSB was included in two articles [26, 27]. The combined effect amount was  $WMD = -0.1$ , 95% CI [-0.38, 0.19],  $P = 0.5$ . It had no statistical significance (see Figure 9).

**3.10. Adverse Events.** No article reported the adverse events of Tai Chi.

## 4. Discussion

**4.1. Summary of Main Findings.** The objective of this review was to summarize and evaluate the effectiveness of Tai Chi for cognitive impairment. Nine researches including 827 patients were carried out in China, Brazil, and Thailand. The evidence shows that Tai Chi is more likely to improve cognitive impairment comparing to control group. We found statistically significant benefits of Tai Chi as follows: MMSE  $WMD = 1.52$ , 95% CI [0.90, 2.14]; MoCA  $WMD = 3.5$ , 95% CI [0.76, 6.24]; CDR  $WMD = -0.55$ , 95% CI [-0.80, -0.29]; LMD  $WMD = 1.1$ , 95% CI [0.04, 2.16]. However, DSF  $WMD = 0.53$ , 95% CI [-0.65, 1.71], and DSB  $WMD = -0.1$ , 95% CI [-0.38, 0.19], were not statistically significant. In addition, there is no research reporting the adverse events of Tai Chi and it may be a safe exercise for people with cognitive impairment.

**4.2. Applicability of the Current Evidence.** Previous Cochrane Review showed Tai Chi did significantly reduce risk of falling [4], but it did not analyze any influence on cognitive impairment. It had been affirmed that exercising activities play a positive role in declining risk of the elderly [14, 33], but the effect of Tai Chi was uncertain. Other correlative meta-analyses only focused on English databases [14, 36]. We added studies from Chinese databases in our systematic review. Therefore, this study is an update meta-analysis which evaluates the role of Tai Chi in the prevention of cognitive impairment. Cognitive impairment is a long-term process [2] and it is often considered irreversible [37]. In recent years, performing Tai Chi has been stressed in the process of preventing cognitive impairment [34]. Our research finds that practicing Tai Chi is helpful for cognitive function, but it seems to have no effect on logical memory. Considering that the test results are not necessarily positively correlated with clinical symptoms and conditions, this study only provides reference for clinical practice.

**4.3. Limitations of This Review.** Although new published literatures were added to this systematic review, the risk of this limitation has not been avoided. First, according to Cochrane's bias risk assessment tool, 5 of the 9 included studies are considered to have a high bias risk due to the lack of randomization, blind method implementation, and distribution concealment. The quality of evidence is not good enough because most researches did not describe the detailed methods in their research. Second, the type of Tai Chi was not assessed in this study according to lack of related RCTs. Third, different severity of cognitive impairment may lead to different effects by performing Tai Chi, but no article mentioned that.

**4.4. Implications for Further Studies.** This study provides a certain value for the future research [38]. The current data show that the Tai Chi group has better cognitive function






FIGURE 3: Quality assessment of the included studies.




FIGURE 4: Forest plot of Mini-Mental State Examination.

score than the control group and Tai Chi is a safe exercise. In terms of clinical rehabilitation, more long-term follow-ups are needed to provide more RCTs and mechanical researches. In addition, most of the included studies either inadequately reported or did not clearly report methods

related to important biases such as randomization/allocation concealment and blinding methods. Future trials should be improved for their reporting quality by following the Consolidated Standards of Reporting Trials (CONSORT) statement [39].




FIGURE 5: Forest plot of Montreal Cognitive Assessment.




FIGURE 6: Forest plot of Clinical Dementia Rating.




FIGURE 7: Forest plot of logical memory delayed recall score.




FIGURE 8: Forest plot of digit span forward.




FIGURE 9: Forest plot of digit span backward.

## 5. Conclusion

There is limited evidence supporting that practicing Tai Chi can bring better changes in cognitive function score (MMSE, MoCA, CDR, and LMD). However, there is no influence on DSF and DSB. Current evidence indicates that Tai Chi is a safe exercise for people with cognitive impairment. There is still a need for increasing RCTs to address whether practicing Tai Chi is effective for patients with cognitive impairment.

## Abbreviations

MMSE: Mini-Mental State Examination  
 MoCA: Montreal Cognitive Assessment  
 CDR: Clinical Dementia Rating  
 LMD: Logical memory delayed recall  
 DSF: Digit span forward  
 DSB: Digit span backward  
 RCT: Randomized controlled trial.

## Data Availability

No primary data were used in this article.

## Conflicts of Interest

All authors of this study have declared that they have no conflicts of interest.

## Authors' Contributions

Gu Renjun and Gao Yujia worked equally in this article. Sun Zhiguang and Gu Renjun conceived and designed the analysis. Gu Renjun, Gao Yujia, Zhang Chunbing, and Liu Xiaojuan completed the data retrieval and analyzed the data. Gu Renjun wrote the paper. All authors read and approved the final manuscript.

## Acknowledgments

This study was supported by Jiangsu Leading Talents Project of Traditional Chinese Medicine (SLJ0226).

## Supplementary Materials

PRISMA 2009 checklist. (*Supplementary Materials*)

## References

- [1] J. E. Morley, "An overview of cognitive impairment," *Clinics in Geriatric Medicine*, vol. 34, no. 4, pp. 505–513, 2018.
- [2] A. M. Sanford, "Mild cognitive impairment," *Clinics in Geriatric Medicine*, vol. 33, no. 3, pp. 325–337, 2017.
- [3] J. N. Vega and P. A. Newhouse, "Mild cognitive impairment: diagnosis, longitudinal course, and emerging treatments," *Current Psychiatry Reports*, vol. 16, no. 10, p. 490, 2014.
- [4] D. Forbes, S. Forbes, D. G. Morgan, M. Markle-Reid, J. Wood, and I. Culum, "Physical activity programs for persons with dementia," *Cochrane Database System Reviews*, vol. 3, Article ID CD006489, 2008.
- [5] S. R. Nyman, W. Ingram, J. Sanders et al., "Randomised controlled trial of the effect of tai chi on postural balance of people with dementia," *Clinical Interventions in Aging*, vol. 14, pp. 2017–2029, 2019.
- [6] F. Wang, E.-K. O. Lee, T. Wu et al., "The effects of tai chi on depression, anxiety, and psychological well-being: a systematic review and meta-analysis," *International Journal of Behavioral Medicine*, vol. 21, no. 4, pp. 605–617, 2014.
- [7] D. I. Miller, V. Taler, P. S. Davidson, and C. Messier, "Measuring the impact of exercise on cognitive aging: methodological issues," *Neurobiology of Aging*, vol. 33, no. 3, p. 622, 2012.
- [8] M. N. McDonnell, A. E. Smith, and S. F. Mackintosh, "Aerobic exercise to improve cognitive function in adults with neurological disorders: a systematic review," *Archives of Physical Medicine and Rehabilitation*, vol. 92, no. 7, pp. 1044–1052, 2011.
- [9] R. C. Petersen, B. Caracciolo, C. Brayne, S. Gauthier, V. Jelic, and L. Fratiglioni, "Mild cognitive impairment: a concept in evolution," *Journal of Internal Medicine*, vol. 275, no. 3, pp. 214–228, 2014.
- [10] J. Lee, "Effects of aerobic and resistance exercise interventions on cognitive and physiologic adaptations for older adults with mild cognitive impairment: a systematic review and meta-analysis of randomized control trials," *International Journal of Environmental Research and Public Health*, vol. 17, no. 24, p. 9216, 2020.
- [11] P. W. H. Peng, "Tai chi and chronic pain," *Regional Anesthesia and Pain Medicine*, vol. 37, no. 4, pp. 372–382, 2012.
- [12] S. Sungkarat, S. Boripuntakul, S. Kumfu, S. R. Lord, and N. Chattipakorn, "Tai chi improves cognition and plasma BDNF in older adults with mild cognitive impairment: a randomized controlled trial," *Neurorehabilitation and Neural Repair*, vol. 32, no. 2, pp. 142–149, 2018.
- [13] K. Lim, A. Pysklywec, M. Plante, and L. Demers, "The effectiveness of Tai Chi for short-term cognitive function improvement in the early stages of dementia in the elderly: a systematic literature review," *Clinical Interventions in Aging*, vol. 14, pp. 827–839, 2019.
- [14] P. M. Wayne, J. N. Walsh, R. E. Taylor-Piliae et al., "Effect of tai chi on cognitive performance in older adults: systematic review and meta-analysis," *Journal of the American Geriatrics Society*, vol. 62, no. 1, pp. 25–39, 2014.
- [15] P. J. Klein, J. Baumgarten, and R. Schneider, "Qigong and tai chi as therapeutic exercise: survey of systematic reviews and meta-analyses addressing physical health conditions," *Alternative Therapies in Health and Medicine*, vol. 25, no. 5, pp. 48–53, 2019.
- [16] N. Huang, W. Li, X. Rong et al., "Effects of a modified tai chi program on older people with mild dementia: a randomized controlled trial," *Journal of Alzheimer's Disease*, vol. 72, no. 3, pp. 947–956, 2019.
- [17] P. Huston and B. McFarlane, "Health benefits of tai chi: what is the evidence?" *Can Fam Physician*, vol. 62, no. 11, pp. 881–890, 2016.
- [18] S. A. Saeed, D. J. Antonacci, and R. M. Bloch, "Exercise, yoga, and meditation for depressive and anxiety disorders," *American Family Physician*, vol. 81, no. 8, pp. 981–986, 2010.
- [19] R. T. Ho, T. C. Fong, A. H. Wan et al., "A randomized controlled trial on the psychophysiological effects of physical exercise and Tai-chi in patients with chronic schizophrenia," *Schizophrenia Research*, vol. 171, no. 1–3, pp. 42–49, 2016.
- [20] J. Tao, X. Chen, N. Egorova et al., "Tai Chi Chuan and baduanjin practice modulates functional connectivity of the cognitive control network in older adults," *Scientific Reports*, vol. 7, no. 1, Article ID 41581, 2017.



- [21] S.-Y. Tai, C.-L. Hsu, S.-W. Huang, T.-C. Ma, W.-C. Hsieh, and Y.-H. Yang, "Effects of multiple training modalities in patients with Alzheimer's disease: a pilot study," *Neuropsychiatric Disease and Treatment*, vol. 12, pp. 2843–2849, 2016.
- [22] S. Wang, H. Yin, Y. Jia, L. Zhao, L. Wang, and L. Chen, "Effects of mind-body exercise on cognitive function in older adults with cognitive impairment," *Journal of Nervous & Mental Disease*, vol. 206, no. 12, pp. 913–924, 2018.
- [23] M.-L. Chen, S. B. Wotiz, S. M. Banks, S. A. Connors, and Y. Shi, "Dose-response association of tai chi and cognition among community-dwelling older adults: a systematic review and meta-analysis," *International Journal of Environmental Research and Public Health*, vol. 18, no. 6, p. 3179, 2021.
- [24] L. Shamseer, D. Moher, M. Clarke et al., "Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation," *BMJ*, vol. 354, p. i4086, 2016.
- [25] F. C. Bull, S. S. Al-Ansari, S. Biddle et al., "World Health Organization 2020 guidelines on physical activity and sedentary behaviour," *British Journal of Sports Medicine*, vol. 54, no. 24, pp. 1451–1462, 2020.
- [26] J. Y. T. Kasai, A. L. Busse, R. M. Magaldi et al., "Effects of Tai Chi Chuan on cognition of elderly women with mild cognitive impairment," *Einstein (São Paulo)*, vol. 8, no. 1, pp. 40–45, 2010.
- [27] L. C. Lam, R. C. Chau, B. M. Wong et al., "A 1-year randomized controlled trial comparing mind body exercise (Tai Chi) with stretching and toning exercise on cognitive function in older Chinese adults at risk of cognitive decline," *Journal of the American Medical Directors Association*, vol. 13, no. 6, p. 568, 2012.
- [28] F. Li, P. Harmer, Y. Liu, and L.-S. Chou, "Tai Ji Quan and global cognitive function in older adults with cognitive impairment: a pilot study," *Archives of Gerontology and Geriatrics*, vol. 58, no. 3, pp. 434–439, 2014.
- [29] S. Sungkarat, S. Boripuntakul, N. Chattipakorn, K. Watcharasaksilp, and S. R. Lord, "Effects of tai chi on cognition and fall risk in older adults with mild cognitive impairment: a randomized controlled trial," *Journal of the American Geriatrics Society*, vol. 65, no. 4, pp. 721–727, 2017.
- [30] M.-Y. Siu and D. T. F. Lee, "Effects of tai chi on cognition and instrumental activities of daily living in community dwelling older people with mild cognitive impairment," *BMC Geriatrics*, vol. 18, no. 1, p. 37, 2018.
- [31] Q. Wang and Y. Sheng, "Time effect analysis of Taijiquan on cognitive function of patients with mild cognitive impairment," *Journal of Nursing Management*, vol. 19, no. 2, pp. 141–145, 2019.
- [32] N. Bao and C. Liu, "Study on the influence of Taijiquan on the cognitive function of patients with amnesic mild cognitive impairment," *Medical Information*, vol. 32, no. 2, pp. 115–117, 2019.
- [33] J. M. Northey, N. Cherbuin, K. L. Pumpa, D. J. Smees, and B. Rattray, "Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis," *British Journal of Sports Medicine*, vol. 52, no. 3, pp. 154–160, 2018.
- [34] Y. Zeng, X. Xie, and A. S. K. Cheng, "Qigong or tai chi in cancer care: an updated systematic review and meta-analysis," *Current Oncology Reports*, vol. 21, no. 6, p. 48, 2019.
- [35] R. C. Petersen, O. Lopez, M. J. Armstrong et al., "Practice guideline update summary: mild cognitive impairment," *Neurology*, vol. 90, no. 3, pp. 126–135, 2018.
- [36] M. E. Kelly, D. Loughrey, B. A. Lawlor, I. H. Robertson, C. Walsh, and S. Brennan, "The impact of exercise on the cognitive functioning of healthy older adults: a systematic review and meta-analysis," *Ageing Research Reviews*, vol. 16, pp. 12–31, 2014.
- [37] R. C. Petersen, "Mild cognitive impairment," *Continuum*, vol. 22, pp. 404–418, 2016.
- [38] G. Zheng, F. Liu, S. Li, M. Huang, J. Tao, and L. Chen, "Tai chi and the protection of cognitive ability," *American Journal of Preventive Medicine*, vol. 49, no. 1, pp. 89–97, 2015.
- [39] K. F. Schulz, D. G. Altman, D. G. Altman, and D. Moher, "CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials," *BMC Medicine*, vol. 8, no. 1, p. 18, 2010.