

Molecular tracking of individual host use in the Shiny Cowbird – a generalist brood parasite

Ma Alicia de la Colina¹, Mark E. Hauber^{2,3}, Bill M. Strausberger⁴, Juan Carlos Reboreda¹ & Bettina Mahler¹

¹Departamento de Ecología, Genética y Evolución, and IEGEBA-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

²Department of Psychology, Hunter College and the Graduate Center of the City University of New York, 695 Park Avenue, New York, New York 10065

³School of Biological Sciences, University of Auckland, 3A Symonds Street, PB 92019, Auckland, New Zealand

⁴Pritzker Laboratory for Molecular Systematics and Evolution, Field Museum of Natural History, 1400 S. Lake Shore Drive, Chicago, Illinois 60605

Keywords

Host preference, laying patterns, microsatellites, *Molothrus bonariensis*, mtDNA, nest-use strategies.

Correspondence

Ma Alicia de la Colina, IEGEBA-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Intendente Güiraldes 2160, lab. 40, 4to piso, Pabellón 2. Ciudad Universitaria C1428EGA. Buenos Aires, Argentina.

Tel: ++54 11 4576-3300 int. 200;

Fax: +54-11-45763384;

E-mail: madelacolina@ege.fcen.uba.ar

Funding Information

Royal Society of New Zealand's ISAT grant program; Fondo para la Investigación Científica y Tecnológica; Universidad de Buenos Aires; Consejo Nacional de Investigaciones Científicas y Técnicas

Received: 22 December 2015; Revised: 13 May 2016; Accepted: 16 May 2016

Ecology and Evolution 2016; 6(14): 4684–4696

doi: 10.1002/ece3.2234

Introduction

Brood parasitic species exploit the parental care of other host species (Friedmann 1964, Payne 1977; Rothstein 1990; Davies 2000; Schulze-Hagen et al. 2009). In order to reproduce successfully, among the most important decisions a brood parasitic female has to make include where, when and how many eggs to lay. Females of generalist brood parasitic species (i.e., that parasitize many hosts species) must make decisions to select both suitable host

Abstract

Generalist parasites exploit multiple host species at the population level, but the individual parasite's strategy may be either itself a generalist or a specialist pattern of host species use. Here, we studied the relationship between host availability and host use in the individual parasitism patterns of the Shiny Cowbird *Molothrus bonariensis*, a generalist avian obligate brood parasite that parasitizes an extreme range of hosts. Using five microsatellite markers and an 1120-bp fragment of the mtDNA control region, we reconstructed full-sibling groups from 359 cowbird eggs and chicks found in nests of the two most frequent hosts in our study area, the Chalk-browed Mockingbird *Mimus saturninus* and the House Wren *Troglodytes aedon*. We were able to infer the laying behavior of 17 different females a posteriori and found that they were mostly faithful to a particular laying area and host species along the entire reproductive season and did not avoid using previously parasitized nests (multiple parasitism) even when other nests were available for parasitism. Moreover, we found females using the same host nest more than once (repeated parasitism), which had not been previously reported for this species. We also found few females parasitizing more than one host species. The use of an alternative host was not related to the main hosts' nest availability. Overall, female shiny cowbirds use a spatially structured and host species specific approach for parasitism, but they do so nonexclusively, resulting in both detectable levels of multiple parasitism and generalism at the level of individual parasites.

species (Teuschl et al. 1998; Hahn et al. 1999; Strausberger and Ashley 2003; Langmore and Kilner 2007; Strausberger and Rothstein 2009) and lay in nests of available individual hosts among suitable breeders (Soler et al. 1995; Hauber 2001; Polačiková et al. 2008; Fiorini et al. 2009a; Soler and Pérez-Contreras 2012). Additionally, parasitism must be synchronized with the host's laying cycle to maximize incubation schedules, timing of hatching, and subsequent chick survival (Davies and Brooke 1988; Hauber 2003; Ellison et al. 2006; Moskat et al. 2006; Fiorini et al. 2009).

Individual host-use strategies differ among generalist brood parasitic species. While in the common cuckoo *Cuculus canorus* each female parasitizes predominantly one host species (Marchetti *et al.* 1998; Skjelseth *et al.* 2004; Fossøy *et al.* 2011, 2016), evidence indicates there exist both host-specialist and host-generalist individuals within populations of both the brown-headed cowbird *Molothrus ater* (Alderson *et al.* 1999; Woolfenden *et al.* 2003; Strausberger and Ashley 2005) and the bronzed cowbird *Molothrus aeneus* (Ellison *et al.* 2006). In these species, egg-laying decisions are sometimes flexible and related to the availability and/or quality of hosts, being thus variable among and within breeding seasons (Woolfenden *et al.* 2003; Strausberger and Ashley 2005). Parasites should avoid the use of a host that has been already parasitized to prevent the competition with other parasitic chicks (Strausberger 1998; Hahn *et al.* 1999; Trine 2000; Hoover 2003; McLaren *et al.* 2003; Moskat *et al.* 2006; Goguen *et al.* 2011). However, if the host is able to raise more than one brood parasite, this intraspecific competition might be less costly than investing time and energy in searching for a different nest (Martínez *et al.* 1998). Multiple parasitism, whereby different females lay eggs in the same nest, is widespread among parasitic *Molothrus* cowbirds. Several studies have shown that although females defend territories (Hauber 2001), multiple parasitism is frequent in the brown-headed cowbird with different females parasitizing the same nest (also called superparasitism) and/or the same parasitic female using one nest more than once (*i.e.*, repeated parasitism; Alderson *et al.* 1999; Hahn *et al.* 1999; McLaren *et al.* 2003; Ellison *et al.* 2006; Hauber *et al.* 2012). Previous studies showed that multiple parasitism increases with density of parasitic females and reduced availability of host nests across different avian host–parasite systems (Strausberger 1998; Moskat *et al.* 2006; Rivers *et al.* 2012). In the shiny cowbird *Molothrus bonariensis* (Fig. 1), multiple parasitism is common (Lyon 1997; Ortega 1998; Gloag *et al.* 2012) and probably consequence of an absence of territoriality and defensive

behaviors by parasitic females (Mermoz and Reboreda 2003; Scardamaglia and Reboreda 2014). However, no events of repeated parasitism have been reported yet for this species (Kattan 1997; Lyon 1997; Gloag *et al.* 2014).

The aim of this study was to analyze individual nest-use strategies of the shiny cowbird and test whether parasitism strategies are related to hosts' nests availability. This obligate brood parasite is extremely generalist at population level, known to use the nests of more than 260 species (Friedmann and Kiff 1985; Lowther 2014). Indirect, genetic evidence based on population parameters suggests that females do however not lay their eggs randomly but preferentially parasitize certain host species and nest location types (López-Ortiz *et al.* 2006; Mahler *et al.* 2007), which vary according to the particular host community that is being parasitized (De Mársico *et al.* 2010). Nonrandom host use by shiny cowbird females was supported by other findings, based on morphological differentiation between cowbird eggs laid in different colonies (Lyon 1997) or in different host nests (de la Colina *et al.* 2010; Tuero *et al.* 2012) and implied from behaviors like synchronization of laying and egg-puncturing that vary according to the host (Fiorini *et al.* 2009). Here, for the first time, we studied the individual laying behavior of shiny cowbird females (Fig. 1B) and investigate host preference, nest selection, and temporal laying patterns using genetic tools. We analyze individual female's nest use and explore whether it is related to nest availability. We also analyze temporal laying ranges of individual females and examine whether individuals continue to lay throughout the breeding season or different females sequentially replace each other during the reproductive season. Knowing individual host-use strategies in generalist brood parasites is crucial to understand coevolutionary interactions between parasites and hosts and the impact of parasitism on hosts' reproductive success and the viability of their populations. In the case of generalist females, parasitism can constitute a serious threat to less abundant or preferred hosts as parasite's population




Figure 1. Male (A) and female (B) of Shiny Cowbird (Photos: F. Furiolo).

dynamics will mainly depend on the availability of abundant hosts and will not be affected by the contraction of less abundant ones. Therefore, these hosts can experience increases in parasitism frequencies although their populations are declining. On the contrary, if females are specialists, population dynamics of each “host-specific race” will be associated in a density-dependent way to this host and parasite’s population number will be regulated by the host it uses (May and Robinson 1985; Ney-Nifle et al. 2005). Discerning individual host-use strategies will also improve the comprehension of coevolutionary interactions with host species and the selective pressures operating on them.

Methods

Study system

We conducted the study at the private reserve “El Destino” (35°8’S, 57°23’W) near the town of Magdalena, Buenos Aires Province, Argentina. The study site is nearly flat, marshy grassland with interspersed woodland patches dominated by *Celtis ehrenbergiana* and *Scutia buxifolia*. We collected shiny cowbird eggs found in hosts’ nests in an area of approximately 580 ha within the reserve. This area was not continuous but was subdivided into three plots separated by forest patches of exotic species (*Ligustrum* spp. and *Eucalyptus* spp.) where abundance of cowbird hosts is very low and one area used for other experiments by other members of our research group (Fig. 2). While

there are various host species in this area (Mason 1986), two species are the main hosts: the chalk-browed mockingbird *Mimus saturninus* (hereafter: mockingbird) and the house wren *Troglodytes aedon* (hereafter: wren) (De Mársico et al. 2010), with parasitism frequencies of 89% (Gloag et al. 2012) and 60% (Tuero et al. 2007), respectively. During their breeding seasons (October–January) of 2008–2009, 2009–2010, and 2010–2011, we systematically searched for the nests of these main hosts in the study area and collected cowbird samples (shiny cowbird eggs or blood samples of cowbird chicks). Other host species are also present and breeding in the study area, but they experience considerably lower parasitism frequencies (*Zonotrichia capensis* 25%, *Furnarius rufus* 20%, *Sicalis flaveola* 8%, *Agelaioides badius* 20%, Mason 1986); in the case of the two latter, they are also less abundant than the two main host species (Mason 1985), with only a couple of nests found within the patches used for sample collection. We nonetheless collected cowbird offspring samples from parasitized nests of these hosts that were found occasionally during nest searching and monitoring of mockingbird and wren breeding sites.

Mockingbirds lay three to five eggs, incubation starting with the laying of the penultimate egg and lasting ~14 days. Chicks leave the nest when they are 12–14 days old (Fiorini 2007). We searched for nests by focusing on adult mockingbird activity and then inspecting potential nesting sites within the territory of each breeding pair. Adults are extremely territorial and maintain their territories all year round (Fraga 1985) often using the same



Figure 2. Study site with pinpointed geographical locations of all nests found in the year 2008; black pins correspond to nests of chalk-browed mockingbirds ($N = 77$), white pins to house wren nests ($N = 35$). Polygons show maximal laying areas of the 10 females with full-sibling offspring groups estimated for 2008 (see text). The black polygon shows the complete searching area.

shrubs for nesting. It was possible to follow all breeding attempts of mockingbird pairs within our study area. We monitored nests daily during the periods of advanced construction and laying in order to detect parasitism events. Mockingbirds are partial accepters and only reject immaculate cowbird eggs within the first 24 h after parasitism (Fraga 1985; Mason 1986; Sackmann and Rebores 2003, de la Colina *et al.* 2012). The immaculate eggs represent a low proportion of cowbird eggs in the study area (7%, Gloag *et al.* 2014). To reduce the probability of losing these cowbird eggs, mockingbird's nests were monitored mainly in the first hours of the morning.

Wrens have an average clutch size of five eggs, incubation starts with the laying of the penultimate egg and lasts ~15 days, nestling period is 16 days (Skutch 1953; Tuero *et al.* 2007). To facilitate the monitoring of wren nests, we placed 54 wooden nest boxes in the collection area with holes large enough to be accessed by cowbirds. These nest boxes were within or nearby mockingbird territories. Nest boxes are readily used by wrens and mostly preferred over natural cavities (Llambías and Fernández 2009). We monitored all nests constructed in nest boxes every 2 days during advanced construction and daily during laying. Although some wren breeding pairs nesting in natural cavities may have gone undetected, we are confident that we monitored the majority of the breeding pairs in the area as wrens prefer newly placed nest boxes over natural cavities in 95% of the cases (Llambías and Fernández 2009). Wrens accept all cowbird eggs (Mason 1986).

With the aim of studying the relationship between host availability and host use, for each nest, we recorded the starting date of the laying stage (appearance of the first host egg). To facilitate proper monitoring of the nest and determine the start of incubation stage, we also recorded the date of laying of the remaining host eggs, as well as the laying date of all parasite's eggs collected for molecular analyses (parasite eggs were marked with a permanent marker for later individual identification). In the cases where the nest was found with host eggs without incubation, we waited until the end of laying to calculate the starting date, assuming a clutch of four eggs for mockingbirds and five eggs for wrens. In this way, estimations of the starting date of each nest are not affected by host's egg losses. In the cases where the nest was found during the incubation stage (warm eggs), we determined the starting date from the hatching date of the first egg, by accounting for the average incubation period of each species (above). The exact locations of all nests were marked on a satellite image with Google Earth[®] software (Google Inc. Mountain View, CA) (Fig. 2). For nests belonging to other hosts that we located occasionally, we simply recorded host species, geographical location and whether it was found during laying or incubation.

Sample collection

Parasite's freshly laid eggs were artificially incubated (Yonar[®], model 50/E) at $37.5 \pm 1^\circ\text{C}$ for 48 h to obtain adequate embryonic development and then frozen until processed (Strausberger and Ashley 2001), while eggs found with some degree of incubation (warm at touch, confirmed later with ovoscopy) were directly frozen (total eggs = 359). In cases where cowbird chicks were found in nests of the host species ($N = 11$), blood samples were taken via wing venipuncture and stored in lysis buffer (100 mmol/L Tris pH 8, 10 mmol/L NaCl, 100 mmol/L EDTA, 2% SDS). During the first two breeding seasons, we also captured 32 adults with walk-in traps baited with millet (2008: six females and 10 males, 2009: seven females and nine males; sexed by plumage). The traps were placed in the same three plots used for egg collection. We took blood samples using the same procedure as in chicks and ringed individuals before release.

Analysis of genetic data

For genetic analyses, embryonic tissue was extracted from the eggs and stored in DMSO buffer (20% v/v DMSO, 250 mmol/L EDTA, NaCl). DNA extraction of embryonic samples was performed following a standard protocol of dehydration and precipitation with ethanol and NaCl (Miller *et al.* 1988). Seven microsatellite loci designed for brown-headed cowbirds were amplified using two multiplex-touchdown polymerase chain reaction (PCRs): (1) CB1, CB12, and CB15 (Longmire *et al.* 2001), and (2) Ma μ 20 (Gibbs *et al.* 1997), Ma μ 25, Ma μ 29, and Dp μ 15b (Alderson *et al.* 1999). PCR amplifications for both sets of primers were performed in 10- μL reaction volumes using 20–60 ng of DNA template, 0.2 $\mu\text{mol/L}$ forward and reverse primers, 0.25 $\mu\text{mol/L}$ dNTPs, 2.5 mmol/L MgCl₂, and 0.25 U Taq polymerase. Cycling temperatures were 95°C for 4 min then 10 cycles of 94°C for 30 sec, $55\text{--}53^\circ\text{C}$ for 45 sec, and 72°C for 45 sec, then 35 cycles of 94°C for 30 sec, 53°C for 45 sec, and 72°C for 45 sec, finishing with 72°C for 40 min. The forward primer for each locus was fluorescently labeled and analyzed on an Applied Biosystems Model 3130xl Genetic Analyzer. Genotypes were assigned using Peak Scanner[™] v.1.0 (Applied Biosystems, Foster City, CA).

Genotypes of adult individuals were used to estimate population genetic parameters. The calculation of observed and expected heterozygosities and tests for departures from Hardy–Weinberg and linkage equilibrium were conducted with Genepop v. 4.0 (Raymond and Rousset 1995; Rousset 2008). Loci were checked for null alleles with Micro-checker v. 2.2.3 (Van Oosterhout *et al.* 2004).

In order to analyze an individual female's laying behavior, we assigned eggs to particular females creating full-sibling groups. To assess the statistical confidence for individual identification with our set of markers, we calculated the probability of identity $PI_{(ID)}$ and the probability of identity between siblings $PI_{(SIB)}$ (the probability that two individuals drawn at random from a population will have the same genotype at multiple loci, Waits *et al.* 2001) using the software Gimlet v. 1.2.3 (Valière 2002). Sibling group reconstruction was used to assemble sets of offspring that belonged to individual females which had also been fertilized by the same male, thus including only full-siblings. Relatedness coefficients (r) among shiny cowbird samples were calculated using ML-Relate (Kalinowski *et al.* 2006). Full-siblings were identified for r -values above the empirical cutoff value calculated with the software iRel v. 1.0 (Gonçalves da Silva and Russello 2011) following the procedure of Russello and Amato (2004). For egg pairs with full-sibling r -values, we tested specific hypotheses of full-siblings versus half-siblings and unrelated samples to discard full-sibling relationships assigned by chance. Hypothesized relationships of full-siblings were accepted for $P > 0.99$ (Marshall *et al.* 1998; Goodnight and Queller 1999; McPeck and Sun 2000; Kalinowski *et al.* 2006).

To confirm female identity, we also sequenced an 1120-bp fragment of the mtDNA control region (Mahler *et al.* 2007) for individuals assigned to full-sibling groups. Amplified products were sequenced on an Applied Biosystems Model 3130xl Genetic Analyzer and manually edited using CodonCode Aligner v. 5.0.1 (CodonCode Corporation, Centerville, MA). As full-siblings must share the maternally inherited mitochondrial haplotype, we excluded those individuals that showed a different haplotype from the rest of the full-sibling group. Full-sibling groups composed of two members with different haplotypes were excluded from the analyses.

Analysis of parasitism strategies

To analyze the spatial distribution of nests parasitized by the same female, we obtained coordinates of the nests from which her offspring was collected. Using the package "Geosphere" (Hijmans and Williams 2012) for R software (R Core Team, 2013), we calculated the distances between each pair of nests with one female's offspring. We then compared the distribution of distance data of parasitized nests with related eggs (for 17 sibling groups) and the distribution of distance data of parasitized nests with unrelated eggs using a nonparametric Kolmogorov–Smirnov test with InfoStat v. 2015 software (Di Rienzo *et al.* 2015). For the first group, 62 samples assigned to 17 females were included, while for the second group, 111 samples were included. This generated 100

distance values for samples of the first group and 5460 distance values for the second group.

For each cowbird female, we identified the host species utilized, and the breeding attempt at which they were parasitized. In cases of multiple and repeated parasitism, we determined the availability of nests appropriate for parasitism at that time, considering as "suitable" those nest at laying or early incubation stage (Fiorini *et al.* 2009). We also analyzed temporal laying patterns and calculated the maximum laying range for cowbird females in the study area (in number of days).

Results

Data collection and genetic data

In total, 316 nests were monitored during the three breeding seasons, including 220 mockingbird nests and 96 wren nests belonging to an average of 20 and 28 reproductive couples per season, respectively (Table 1). Mockingbird reproductive pairs had an average (\pm SD, standard deviation) of 2.8 (\pm 0.3) breeding attempts per season versus 1.2 (\pm 0.1) in wrens. Mean frequency of parasitism was 61.3% (\pm 3.8) and 40.1% (\pm 2.7) nests parasitized, and the intensity of parasitism was 2.2 and 1.2 eggs per nest, respectively (Table 1). Although climatic conditions differed among years, with very dry (2008) and very rainy (2009) seasons, parasitism frequencies and intensities per host remained fairly constant. We collected 402 individual cowbird samples during the three breeding seasons, corresponding to 311 offspring samples taken from mockingbird nests, 53 from wren nests, 6 from rufous-collared sparrows *Z. capensis* nests, and 32 adult cowbird samples.

We were able to genotype 198 offspring samples and the 32 adult samples. Unfortunately, for the remaining eggs, we either could not ensure embryonic development despite pre-incubation for 48 h, or the extracted DNA was not sufficient for successful amplification or genotyping. Genetic variability data for the adults' samples are shown in Table 2. Locus *Dpμ* 15b presented many amplification problems and was subsequently eliminated for genotyping. The locus *CB1* showed deviation from Hardy–Weinberg equilibrium and evidence of null alleles and was thus excluded from kinship analyses. Probabilities of identity for the remaining five loci were $PI_{(ID)} = 3.2E^{-08}$ and $PI_{(sib)} = 4.2E^{-03}$, below the thresholds suggested by Waits *et al.* (2001; $PI_{(ID)} < 0.001$ and $PI_{(sib)} < 0.05$).

Kinship analysis

Allele frequencies used for kinship analysis were calculated from the adult samples. To determine the empirical cutoff value with software iRel, r_{QG} showed greater discriminatory

Table 1. Details of nests monitored in the three breeding seasons (2008–2010) by host species.

	2008					2009					2010				
	Bp	N	N _p	I	R	Bp	N	N _p	I	R	Bp	N	N _p	I	R
Chalk-browed mockingbird	24	77	50	2.3 ± 1.8	1–11	22	68	42	2.5 ± 1.6	1–6	22	75	43	1.8 ± 1.0	1–5
House wren	25	35	14	1.2 ± 0.4	1–2	41	48	15	1.2 ± 0.4	1–2	20	21	9	1.2 ± 0.4	1–2
Rufous-collared sparrow	2	2	2	1.5 ± 0.0	1–2	2	2	2	1.5 ± 0.0	1–2					
Number of collected samples	134					148					88				

Bp: number of breeding pairs; N: number of nests found; N_p: number of parasitized nests; I: intensity of parasitism (mean number ± standard error of parasite eggs per parasitized nests); R: minimum and maximum number of cowbird eggs found in a nest.

Table 2. Characteristics of the six microsatellite loci genotyped from a sample of 32 adult shiny cowbirds.

	K	Range (pb)	H _o	H _e	P-value _{EHW}
CB. 1	12	203–267	0.688	0.884	0.0003
CB. 12	14	152–248	0.955	0.905	0.7709
CB. 15	8	241–273	0.938	0.844	0.9831
Maμ 20	14	120–172	0.903	0.840	0.2274
Maμ 25	10	122–154	0.917	0.850	0.2567
Maμ 29	11	115–179	0.708	0.811	0.1053

K: number of alleles and its molecular weight range in base pairs (pb); H_o: observed heterozygosity; H_e: estimated heterozygosity; P-value_{EHW}: departure from Hardy–Weinberg equilibrium. Bold value indicates statistical significance.

power between adjacent categories of kinship and the allocation of full-siblings was above $r = 0.363$. Using this cut-off value, pairs of full-siblings were identified and subject to hypothesis testing. There were 87 eggs assigned to 26 groups of full-siblings, with an average of 3.3 eggs per group (minimum 2–maximum 7). After mtDNA analysis of these individuals, we found 8 of the haplotypes found by Mahler *et al.* (2007). Then, 25 samples had to be excluded, leaving 17 groups of full-siblings (2008: $N = 10$, 2009: $N = 5$, 2010: $N = 2$). No full-sibling groups containing members from different years were established, so each group was considered to be independent (i.e., from a different female). Remaining offspring ($N = 136$) could not be assigned to full-sibling groups and thus were not further considered for analyses on laying strategies.

Spatial and temporal nest use

The spatial distribution analysis showed that eggs belonging to full-sibling groups were found in nests that were closer than nests with unrelated eggs (Kolmogorov–Smirnov, $KS = 0.6$, $P < 0.01$, Fig. 3). Nests with related eggs were found at a median distance of 440 m mostly within the same collection area (Fig. 2). In contrast, eggs laid by different females showed a bimodal distribution, corresponding to eggs laid within the same area and in different collection areas (Fig. 2). In our study area, the laying

season lasted 86 ± 9.5 days, beginning the first days of October and finishing by mid-January. Laying periods among identified females mostly overlapped during the breeding season. We found a maximum individual laying period of 65 days (Fig. 4) and a maximum laying rate of 0.5 eggs per day, corresponding to three eggs found for a female in a time range of 6 days (Fig. 4). This value is somewhat lower than the one documented by Kattan (1993) of 0.66 eggs per day.

Host selection

Of the 17 different full-sibling groups, 15 were composed of eggs found in nests of a single host species (13 in mockingbird nests and two in wren nests, Fig. 5). The remaining two full-sibling groups were composed of eggs from nests of more than one host species: one with eggs from mockingbird and wren nests and the other one from mockingbird and rufous-collared sparrow nests (Fig. 5).

Considering that full-sibling groups represent different females, we studied individual parasite females laying behavior analyzing host selection and nest use. When analyzing nest availability for the two females that used nests of more than one host species (female 3 and 12, Fig. 5), we found that nests of both hosts were available during laying and early incubation for each parasitism event, thus indicating that use of another host was not related to an absence of nests of the other one.

For eight females, eggs were found in nests of different breeding pairs (Fig. 6). For nine females, some eggs were found in nests of the same breeding pair. Four cowbird females parasitized only different breeding attempts, which were successive in some cases and interspersed in others, while for five females eggs were found in the same nest (Fig. 6). It is noteworthy that one female laid twice in the same nest of two different breeding pairs. This indicates that at least five of 17 females engaged in repeated parasitism. Use of the same nest was made with an interval of 1–4 days. A detailed evaluation of nest availability at the date when the female used the same nest for the second time showed that there were other




Figure 3. Spatial distribution of nests parasitized with eggs of the same female (black) and with unrelated eggs (gray). Kolmogorov–Smirnov, $KS = 0.6$, $P < 0.01$.




Figure 4. The number of eggs in relation to seasonal laying range (range of days between the date of the first and the last egg assigned to each female) for the 17 females studied (Simple linear regression: $R^2 = 0.23$, $N = 17$, $P = 0.48$, $P = 0.049$).

options for laying: At least seven available nests were in the proximity of the nest used by female 7, four nests for females 8 and 10, and five nests for female 9. In the case of female 15, that showed repeated parasitism twice, there were three and one other suitable host nests available, respectively, at the time of laying repeatedly.

Discussion

Our results show that the 17 studied laying Shiny Cowbird females were mostly faithful to a particular laying area and showed a preference for one of the monitored host species. Also, they did not avoid using nests previously parasitized by themselves or by another female. Nests parasitized by the same female were within a limited area, showing individual laying site fidelity. Eggs laid by the same female were found <1 km apart, with a mean distance of 488 m. These results are consistent with the findings of Scardamaglia and Reboreda (2014) who studied

the movements of parasitic females during the reproductive season using radio telemetry and found that the area used by a female was on average 25 ha. Taken together, these findings show that cowbird females use relatively constant areas for nest searching and parasitism and that they maintain this area throughout the breeding season.

Use of the same breeding area by several females makes multiple parasitism very common at this study site (Gloag et al. 2012; also see Stevens et al. 2013). We found a high proportion of parasitized nests with more than one parasitic egg (57% in mockingbirds and 21% in wrens). The use of the same area during the entire reproductive season led five parasitic females to parasitize the same breeding pair several times at different reproductive attempts (Fig. 6). Moreover, we found five females that engaged in repeated parasitism. This laying strategy has not been previously reported for this parasite species (Kattan 1997; Lyon 1997; Gloag et al. 2014). Multiple parasitism imposes a cost to parasitic females, because it increases




Figure 5. Detail of the hosts used by each of the 17 females studied: chalk-browed mockingbird (black), house wren (gray), and rufous-collared sparrow (white).




Figure 6. Detail of the nests used by each of the 17 females studied: different breeding pair (black), same breeding pair – different nest (dark gray) and same breeding pair – same nest (light gray).

predation risk, egg puncture by other cowbird females and the competition among gregarious nestlings. Costs from this last reason will be even higher when multiple parasitism involves the offspring of the same female. However, this strategy might be beneficial when the costs of multiple parasitism are lower than those used to locate and monitor additional nests (Gloag *et al.* 2014) or exceeded by the benefits of choosing a host that can successfully raise multiple parasitic chicks (Martínez *et al.* 1998), especially when host availability is low (Lyon 1997). On the one hand, both host species have been shown to raise more than one parasite chick successfully

(Fraga 1985; Fiorini 2007; Tuero *et al.* 2007), although mockingbirds have higher success than wrens in doing so (Fraga 1985; Fiorini 2007) as mortality rates are higher in the latter when more than one parasite is in the nest (Kattan 1997; Tuero *et al.* 2007). On the other hand, the presence of additional nest mates might enhance feeding rates of the foster parents delivered to their own genetic chicks, too (Kilner *et al.* 2004; Gloag *et al.* 2012). We found that multiple parasitism was higher in mockingbirds than in wrens (57% vs. 21%), although it is not clear from our data whether this is adaptive host use or a consequence of greater nest detectability in the

mockingbirds (Strausberger 1998; Strausberger and Ashley 2003; Rivers et al. 2012).

In the case of repeated parasitism, we found that when the female used the same nest that she parasitized before, there were three to five other nests of the host available at the time in five of the six cases. Just for one event of repeated parasitism only one other nest was available. Although we cannot discount that some available nests were not found by parasitic females, evidence suggests that females do not actively avoid parasitizing the same nest. One strategy could be random use among available nests within the laying area. If this was the case, it might be possible that repeated parasitism is a consequence of lower nest density, where lower nest densities increase the probability of using the same nest. Accordingly, in comparisons with the study site of Gloag et al. (2014), who did not find evidence of repeated parasitism, our site had about half the nest density (approx. 0.18 nest/ha vs. 0.08 nest/ha, respectively).

The maximal daily laying rate found at our study site was 0.5 eggs/day, somewhat less than the laying rate found for a population of this species in Central America, which was estimated in 0.6 eggs per day (Kattan 1993; Rueda-Cediel et al. 2008), but quite similar to populations of brown-headed cowbirds of Texas and Florida (0.57 and 0.56 eggs per day, respectively; Reetz 2008). Considering geographical differences in breeding season duration, we expected higher laying rates at our more temperate study site (Reetz 2008). Similar values might be related to physiological constraints or to a dependency on nutritional factors that might account for regional differences in reproductive values (Payne 1965; Scott and Ankney 1980; Fleischer 1985).

We also found that females parasitize the hosts within their laying area along the entire breeding season, but we were unable to determine whether laying is constant throughout the season or whether females lay their eggs in bouts with resting periods in-between, as well as whether the laying pattern is generalized or varies among females. Although we found that three females laid throughout the season with longer intervals (approx. every 10–13 days) and other females laid every other day, we cannot discard that these differences are related to incomplete sampling.

Along three breeding seasons, we observed that parasitism frequencies and intensities were higher in mockingbird than in wren nests. These values did not change even in the 2009 season, when we found many more house wren nests and reproductive couples (Table 1). This suggests a preference of shiny cowbird females for mockingbird nests, supporting the hypothesis that females prefer certain hosts at population level (Mason 1986; De Marsico et al. 2010) and only parasitize a small fraction of the available hosts in

a community, not parasitizing or only infrequently parasitizing a large proportion of available hosts (De Marsico et al. 2010). The majority of the females only used nests of mockingbirds for laying while one only parasitized house wrens, indicating a preference for one host species. Just two females were found to use nests of more than one host. Parasitism of more than one host was also reported by Gloag et al. (2014). Additionally, mtDNA distribution patterns suggested the existence of host preference in the shiny cowbird but also frequent host switches (Mahler et al. 2007; Dominguez et al. 2015). Taken together, these results suggest that the preference for one host might be partial or flexible according to the conditions. Although the use of alternative hosts might be underestimated because some hosts were not sampled in our study area and some eggs might have been laid outside the sampling area or even rejected by mockingbirds (and hence not collected), we are confident that we collected a representative sample of the eggs as females' laying areas are limited and the two hosts that were sampled are among the most abundant (Mason 1985) and have the highest parasitism frequencies. Other abundant host species have considerably lower parasitism frequencies by shiny cowbirds (Mason 1985; Mason 1986) or are even not available at this study plot, like the yellow-winged blackbird *Agelasticus thilius* or the brown-and-yellow marshbird *Pseudoleistes virescens*, which nest in wetlands.

As found for the brown-headed cowbird (Alderson et al. 1999; Woolfenden et al. 2003; Strausberger and Ashley 2005), shiny cowbird females of our population showed both host-specialist and host-generalist laying strategies. Although we cannot be sure that we sampled all eggs from each female, we observed that some of them had a preference for one host species (females 9, 10, 13, 14, and 15, Fig. 5). This host-specialist behavior is associated with a preference for a host which is not expected to be absolute or exclusive (Tversky 1969). For the brown-headed cowbird *M. ater*, several studies showed flexible laying behavior that was related to parasite female density (Alderson et al. 1999; Hahn et al. 1999; McLaren et al. 2003; Ellison et al. 2006; Rivers et al. 2012) or host nest availability (Strausberger and Ashley 2003; Rivers et al. 2012). Also, the Horsfield's bronze cuckoo *Chalcites basalis* shows plasticity in host and habitat preference, allowing the use of secondary hosts when the preferred one is spatially or temporally absent (Langmore and Kilner 2007). In our population, host nest availability does not account for laying decisions: For both females that used more than one host, we assessed the availability of nests of the other species at the day of laying and found that nests of both host species were available at similar proportions in all cases.

One explanation for the use of nests of alternative hosts might be social learning, with females following other females to detect available and suitable nests for parasitism. Radio-tracking studies in our study area (Scardamaglia and Reborada 2014) showed that females roost communally and leave the roosts in groups before dawn. Various females (up to 4) approach host nests simultaneously for laying (Fraga 1985; Gloag *et al.* 2014) which suggests that females might detect an available nest by following other females within their laying territory. Although the mechanism underlying specialist or generalist individual host use is not clear, our results are consistent with the absence of complete maternal lineage sorting of brood parasitism between different host species, as reported by Mahler *et al.* (2007). These results indicate that coevolutionary interactions between shiny cowbirds and their hosts are not restricted to one host-specific group of cowbirds and its particular host but that interactions with hosts rather involve the cowbird population as a whole. This makes monitoring of endangered species used as hosts particularly important (López-Ortiz *et al.* 2006), as parasitism pressure alone or combined with other causes could drive small populations to local extinctions (Domínguez *et al.* 2014). Future work could manipulate host nest availability experimentally to assess the role of individual preferences versus ecological constraints in host species selection and nest use by individual brood parasitic females in this and other species.

Acknowledgments

We thank Fundación Elsa Shaw de Pearson for allowing us to conduct this study at “Reserva El Destino”. We also acknowledge the help provided by Diana Alvarez Prada and Robyn Howitt from Landcare Research with the laboratory work. We thank E. W. Grillo for English language corrections. We are thankful for the funding from Royal Society of New Zealand’s ISAT grant program, Universidad de Buenos Aires, Agencia Nacional de Promoción Científica y Tecnológica de Argentina, and Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina.

Ethical standards

This research was conducted in accordance with relevant Argentinean regulations (Law of Conservation of Wild Fauna), under the permit issued to JCR, University of Buenos Aires.

Conflict of Interest

None declared.

References

- Alderson, G. W., H. L. Gibbs, and S. G. Sealy. 1999. Determining the reproductive behaviour of individual brown-headed cowbirds using and microsatellite DNA markers. *Anim. Behav.* 58:895–905.
- de la Colina, M. A., B. Mahler, and J. C. Reborada. 2010. Differences in morphology and colour pattern of shiny cowbird (*Molothrus bonariensis*) eggs found in nests of two hosts. *Biol. J. Linn. Soc.* 102:838–845.
- de la Colina, M. A., L. Pompilio, M. E. Hauber, J. C. Reborada, and B. Mahler. 2012. Different recognition cues reveal the decision rules used for egg rejection by hosts of a variably mimetic avian brood parasite. *Anim. Cogn.* 15:881–889.
- Davies, N. B. 2000. Cuckoos, cowbirds and other cheats. Oxford Univ. Press, Oxford, U.K.
- Davies, N. B., and L. Brooke. 1988. Cuckoos versus reed warblers: adaptations and counter adaptations. *Anim. Behav.* 36:262–284.
- De Mársico, M. C., B. Mahler, M. Chomnalez, A. G. Di Giacomo, and J. C. Reborada. 2010. Host use by generalist and specialist brood-parasitic cowbirds at population and individual levels. *Adv. Stud. Behav.* 42:83–121.
- Di Rienzo, J. A., F. Casanoves, M. G. Balzarini, L. Gonzalez, M. Tablada, and C. W. Robledo 2015. InfoStat versión 2012. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Córdoba, Argentina. Available at <http://www.infostat.com.ar>.
- Domínguez, M., J. C. Reborada, and B. Mahler. 2014. Impact of Shiny Cowbird and botfly parasitism on the reproductive success of the globally endangered Yellow Cardinal *Gubernatrix cristata*. BCI, Available on CJO 2014. doi:10.1017/S095927091400015X.
- Domínguez, M., M. A. de la Colina, A. Di Giacomo, J. C. Reborada, and B. Mahler. 2015. Host-switching in cowbird brood parasites: how often does it occur? *J. Evol. Biol.* 28:1290–1297.
- Ellison, K., S. G. Sealy, and H. L. Gibbs. 2006. Genetic elucidation of host use by individual sympatric bronzed cowbirds (*Molothrus aeneus*) and brown-headed cowbirds (*M. ater*). *Can. J. Zool.* 84:1269–1280.
- Fiorini, V. D. 2007. Synchronization of parasitism and host selection in a generalist brood parasite, the shiny cowbird *Molothrus bonariensis* (Icterinae, Aves). Dissertation, Buenos Aires University, Buenos Aires, Argentina.
- Fiorini, V. D., D. T. Tuero, and J. C. Reborada. 2009. Shiny cowbirds benefits of synchronizing parasitism and puncturing eggs in large and small hosts. *Anim. Behav.* 77:561–568.
- Fleischer, R. C. 1985. A new technique to identify and assess the dispersion of eggs of individual brood parasites. *Behav. Ecol. Sociobiol.* 17:91–100.
- Fossøy, F., A. Antonov, A. Moksnes, E. Røskaft, J. R. Vikan, A. P. Møller, *et al.* 2011. Genetic differentiation among

- sympatric cuckoo host races: males matter. *Proc. R. Soc. Lond. B* 278:1639–1645.
- Fossøy, F., M. D. Sorenson, W. Liang, T. Ekrem, A. Moksnes, A. P. Møller, *et al.* 2016. Ancient origin and maternal inheritance of blue cuckoo eggs. *Nat. Commun.* 7:10272. doi:10.1038/ncomms10272.
- Fraga, R. M. 1985. Host–parasite interactions between chalk-browed mockingbirds and shiny cowbirds. *Ornithol. Monogr.* 36:829–844.
- Friedmann, H. 1929. The cowbirds. A study in the biology of social parasitism. C. C. Thomas, Springfield, Illinois.
- Friedmann, H., and L. F. Kiff. 1985. The parasitic cowbirds and their hosts. *Proc. West Found. Vert. Zool.* 2:225–302.
- Gibbs, H. L., P. Miller, G. Alderson, and S. G. Sealy. 1997. Genetic analysis of brown-headed cowbirds *Molothrus ater* raised by different hosts: data from mtDNA and microsatellite DNA markers. *Mol. Ecol.* 6:189–193.
- Gloag, R., V. D. Fiorini, J. C. Reboreda, and A. Kacelnik. 2012. Brood parasite eggs enhance egg survivorship in a multiply parasitized host. *Proc. R. Soc. Lond. B* 279:1831–1839.
- Gloag, R., V. D. Fiorini, J. C. Reboreda, and A. Kacelnik. 2014. Shiny cowbirds share foster mothers but not true mothers in multiply parasitized mockingbird nests. *Behav. Ecol. Sociobiol.* 68:681–689.
- Goguen, C. B., D. R. Curson, and N. E. Mathews. 2011. Costs of multiple parasitism for an avian brood parasite, the brown-headed cowbird (*Molothrus ater*). *Can. J. Zool.* 89:1237–1248.
- Gonçalves da Silva, A., and M. A. Russello. 2011. iRel: software for implementing pairwise relatedness estimators and evaluating their performance. *Conserv. Genet. Resour.* 3:69–71. Available at <http://irel.ok.ubc.ca:8080/webMathematica/irel/irel.html>.
- Goodnight, K. F., and D. C. Queller. 1999. Computer software for performing likelihood tests of pedigree relationship using genetic markers. *Mol. Ecol.* 8:1231–1234.
- Hahn, D. C., J. A. Sedgwick, I. S. Painter, and N. J. Casna. 1999. A spatial and genetic analysis of cowbird host selection. *Stud. Avian Biol.* 18:204–217.
- Hauber, M. E. 2001. Site selection and repeatability in brown-headed cowbird (*Molothrus ater*) parasitism of eastern phoebe (*Sayornis phoebe*) nests. *Can. J. Zool.* 79:1518–1523.
- Hauber, M. E. 2003. Hatching asynchrony, nestling competition, and the cost of interspecific brood parasitism. *Behav. Ecol.* 14:224–235.
- Hauber, M. E., B. M. Strausberger, K. A. Feldheim, J. Lock, and P. Cassey. 2012. Indirect estimates of breeding and natal philopatry in an obligate avian brood parasite. *J. Ornithol.* 153:467–475.
- Hijmans, R. J., and E. Williams. 2012. Geosphere: spherical trigonometry. R package version 1.2-28. Available at <http://CRAN.R-project.org/package=geosphere>.
- Hoover, J. P. 2003. Multiple effects of brood parasitism reduce the reproductive success of prothonotary warblers, *Protonotaria citrea*. *Anim. Behav.* 65:923–934.
- Kalinowski, S. T., A. P. Wagner, and M. L. Taper. 2006. ML-Relate: a computer program for maximum likelihood estimation of relatedness and relationship. *Mol. Ecol. Notes* 6:576–579.
- Kattan, G. H. 1993. Reproductive strategy of a generalist brood parasite, the Shiny Cowbird, in the Cauca Valley, Colombia. Ph. D. Thesis, University of Florida, Gainesville, FL.
- Kattan, G. H. 1997. Shiny cowbirds follow the ‘shotgun’ strategy of brood parasitism. *Anim. Behav.* 53:647–654.
- Kilner, R. M., J. R. Madden, and M. E. Hauber. 2004. Brood parasitic cowbird nestlings use host young to procure resources. *Science* 305:877–879.
- Langmore, N. E., and R. M. Kilner. 2007. Breeding site and host selection by Horsfield’s bronze cuckoos, *Chalcites basalys*. *Anim. Behav.* 74:995–1004.
- Llambías, P. E., and G. J. Fernández. 2009. Effects of nest boxes on the breeding biology of Southern house wrens *Troglodytes aedon bonariae* in the southern temperate zone. *The Ibis* 151:113–121.
- Longmire, J. L., J. L. Roach, M. Maltbie, P. S. White, O. L. Tatum, and K. L. Makora. 2001. Tetranucleotide microsatellite markers for the Brown-headed cowbird (*Molothrus ater*). *J. Avian Biol.* 31:76–78.
- López-Ortiz, R., E. A. Ventosa-Febles, K. R. Ramos-Alvarez, R. Medina-Miranda, and A. Cruz. 2006. Reduction in host use suggests host specificity in individual shiny cowbirds (*Molothrus bonariensis*). *Ornitol. Neotrop.* 17:259–269.
- Lowther, P. E. 2014. Lists of victims and hosts of the parasitic cowbirds. The Field Museum, Chicago, IL. Available at http://www.fieldmuseum.org/sites/default/files/plowther/2014/09/24/cowbird_hosts-17sep2014.pdf (accessed 17 September 2014).
- Lyon, B. E. 1997. Spatial patterns of shiny cowbird brood parasitism on chestnut-capped blackbirds. *Anim. Behav.* 54:927–939.
- Mahler, B., V. A. Confalonieri, I. J. Lovette, and J. C. Reboreda. 2007. Partial host fidelity in nest selection by the shiny cowbird (*Molothrus bonariensis*), a highly generalist avian brood parasite. *J. Evol. Biol.* 20:1918–1923.
- Marchetti, K., H. L. Gibbs, and H. Nakamura. 1998. Host-race formation in the common cuckoo. *Science* 282:471–472.
- Marshall, T. C., J. Slate, L. E. B. Kruuk, and J. M. Pemberton. 1998. Statistical confidence for likelihood-based paternity inference in natural populations. *Mol. Ecol.* 7:639–655.
- Martínez, J. G., J. J. Soler, M. Soler, and T. Burke. 1998. Spatial patterns of egg laying and multiple parasitism in a brood parasite: a non-territorial system in the great spotted cuckoo (*Clamator glamdarius*). *Oecologia* 117:286–294.
- Mason, P. 1985. The nesting biology of some passerines of Buenos Aires. Argentina. *Ornithol. Monogr.* 36:954–972.

- Mason, P. 1986. Brood parasitism in a host generalist, the shiny cowbird: II. Host selection. *Auk* 103:61–69.
- May, R. M., and S. K. Robinson. 1985. Population dynamics of avian brood parasitism. *Am. Nat.* 130:161–167.
- McLaren, C. M., S. G. Sealy, B. E. Woolfenden, and H. L. Gibbs. 2003. Genetic and temporal patterns of multiple parasitism by brown-headed cowbirds (*Molothrus ater*) on song sparrows (*Melospiza melodia*). *Can. J. Zool.* 81:281.
- McPeck, M. S., and L. Sun. 2000. Statistical test for detection of misspecified relationships by use of genome-screen data. *Am. J. Hum. Genet.* 66:1076–1094.
- Mermoz, M. E., and J. C. Reboreda. 2003. Reproductive success of shiny cowbird (*Molothrus bonariensis*) parasitizing the larger brown-and-yellow marshbird (*Pseudoleistes virescens*) in Argentina. *Auk* 120:1128–1139.
- Miller, S. A., D. D. Dykes, and H. F. Polesky. 1988. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.* 16:1215.
- Moskat, C., Z. Barta, M. E. Hauber, and M. Honza. 2006. High synchrony of egg laying in common cuckoos (*Cuculus canorus*) and their great reed warbler (*Acrocephalus arundinaceus*) hosts. *Ethol. Ecol. Evol.* 18:159–167.
- Ney-Nifle, M., C. Bernstein, J. C. Reboreda, and A. Kacelnik. 2005. Population dynamics and avian brood parasitism: persistence and invasions in a three species system. *J. Anim. Ecol.* 74:274–284.
- Ortega, C. P. 1998. Cowbirds and other brood parasites. University of Arizona Press, Tucson, AZ.
- Payne, R. B. 1965. Clutch size and numbers of eggs laid by brown-headed cowbirds. *Condor* 67:44–60.
- Payne, R. B. 1977. The ecology of brood parasitism in birds. *Annu. Rev. Ecol. Syst.* 8:1–28.
- Poláčiková, L., P. Procházka, M. I. Cherry, and M. Honza. 2008. Choosing suitable hosts: common cuckoos *Cuculus canorus* parasitize great reed warblers *Acrocephalus arundinaceus* of high quality. *Evol. Ecol.* 23:879–891.
- R Core Team. 2013. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. Available at <http://www.R-project.org/>.
- Raymond, M., and F. Rousset. 1995. GENEPOP version 1.2: population genetics software for exact tests and ecumenicism. *J. Hered.* 86:248–249.
- Reetz, M. J. 2008. Patterns of brown-headed cowbird parasitism in a recently invaded area and potential mechanisms limiting cowbird reproduction. PhD Dissertation, University of Florida, Gainesville, Florida, USA.
- Rivers, J. W., S. Young, E. G. Gonzalez, B. Horton, J. Lock, and R. C. Fleischer. 2012. High levels of relatedness between brown-headed cowbird (*Molothrus ater*) nest mates in a heavily parasitized host community. *Auk* 129:623–631.
- Rothstein, S. I. 1990. A model system for coevolution: avian brood parasitism. *Annu. Rev. Ecol. Evol. Syst.* 21:481–508.
- Rousset, F. 2008. Genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol. Ecol. Resour.* 8:103–106.
- Rueda-Cediel, P., G. Kattan, and M. P. Ramírez-Pinilla. 2008. Ovarian and oviductal morphology of a brood parasitic bird, *Molothrus bonariensis* (Passeriformes, Icteridae). *Acta Zool.* 89:261–276.
- Russello, M. A., and G. Amato. 2004. Ex-situ population management in the absence of pedigree information. *Mol. Ecol.* 13:2829–2840.
- Sackmann, P., and J. C. Reboreda. 2003. A comparative study of Shiny Cowbird parasitism of two large hosts, the Chalk-browed Mockingbird and the Rufous-bellied Thrush. *Condor* 105:728–736.
- Scardamaglia, R. C., and J. C. Reboreda. 2014. Ranging behavior of female and male shiny cowbirds and screaming cowbirds while searching for host nests. *Auk* 131:610–618.
- Schulze-Hagen, K., B. G. Stokke, and T. R. Birkhead. 2009. Reproductive biology of the European cuckoo *Cuculus canorus*: early insights, persistent errors and the acquisition of knowledge. *J. Ornithol.* 150:1–16.
- Scott, D. M., and C. D. Ankney. 1980. Fecundity of the Brown-headed Cowbird in southern Ontario. *Auk* 97:677–683.
- Skjelseth, S., A. Moksnes, E. Røskaft, L. H. Gibbs, M. Taborsky, B. Taborsky, et al. 2004. Parentage and host preference in the common cuckoo *Cuculus canorus*. *J. Avian Biol.* 35:21–24.
- Skutch, A. F. 1953. Life history of the southern house wren. *Condor* 55:121–149.
- Soler, M., and T. Pérez-Contreras. 2012. Location of suitable nests by great spotted cuckoos: an empirical and experimental study. *Behav. Ecol. Sociobiol.* 66:1305–1310.
- Soler, J. J., M. Soler, A. P. Moller, and J. G. Martinez. 1995. Does the great spotted cuckoo choose magpie hosts according to their parenting ability? *Behav. Ecol. Sociobiol.* 36:201–206.
- Stevens, M., J. Troscianko, and C. N. Spottiswoode. 2013. Repeated targeting of the same hosts by a brood parasite compromises host egg rejection. *Nat. Commun.* 4:2475.
- Strausberger, B. M. 1998. Temporal patterns of host availability, brown-headed cowbird parasitism, and parasite egg mass. *Oecologia* 116:267–274.
- Strausberger, B. M., and M. V. Ashley. 2001. Eggs yield nuclear DNA from egg-laying female cowbirds, their embryos and offspring. *Conserv. Genet.* 2:385–390.
- Strausberger, B. M., and M. V. Ashley. 2003. Breeding biology of brood parasitic brown-headed cowbirds (*Molothrus ater*) characterized by parent-offspring and sibling group reconstruction. *Auk* 120:433–445.
- Strausberger, B. M., and M. V. Ashley. 2005. Host use strategies of individual female brown-headed cowbirds *Molothrus ater* in a diverse avian community. *J. Avian Biol.* 36:313–321.
- Strausberger, B. M., and S. I. Rothstein. 2009. Parasitic cowbirds may defeat host defense by causing rejecters to misimprint on cowbird eggs. *Behav. Ecol.* 20:691–699.

- Teuschl, Y., B. Taborsky, and M. Taborsky. 1998. How do cuckoos find their hosts? The role of habitat imprinting. *Anim. Behav.* 56:1425–1443.
- Trine, C. L. 2000. Effects of multiple parasitism on cowbird and wood thrush nesting success. Pp. 135–155 in J. N. M. Smith, T. L. Cook, S. I. Rothstein, G. E. Robinson, S. G. Sealy, eds. *Ecology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX, USA.
- Tuero, D. T., V. D. Fiorini, and J. C. Reboreda. 2007. Effects of shiny cowbird *Molothrus bonariensis* parasitism on different components of house wren *Troglodytes aedon* reproductive success. *The Ibis* 149:521–527.
- Tuero, D. T., V. D. Fiorini, B. Mahler, and J. C. Reboreda. 2012. Shiny cowbird *Molothrus bonariensis* egg size and chick growth vary between two hosts that differ markedly in body size. *J. Avian Biol.* 43:227–233.
- Tversky, A. 1969. The intransitivity of preferences. *Psychol. Rev.* 76:31–48.
- Valière, N. 2002. GIMLET: a computer program for analyzing genetic individual identification data. *Mol. Ecol. Notes* 2:377–379.
- Van Oosterhout, C. V., W. F. Hutchinson, D. P. M. Wills, and P. Shipley. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol. Ecol. Notes* 4:535–538.
- Waits, L. P., G. Luikart, and P. Taberlet. 2001. Estimating the probability of identity among genotypes in natural populations: cautions and guidelines. *Mol. Ecol.* 10: 249–256.
- Woolfenden, B. E., H. L. Gibbs, S. G. Sealy, and D. G. Mc. Master. 2003. Host use and fecundity of individual female brown-headed cowbirds. *Anim. Behav.* 66:95–106.