

ORIGINAL RESEARCH

Responses of macroalgae to CO₂ enrichment cannot be inferred solely from their inorganic carbon uptake strategy

Luna M. van der Loos^{1,2}  | Matthias Schmid¹  | Pablo P. Leal^{1,3}  |
 Christina M. McGraw⁴  | Damon Britton¹  | Andrew T. Revill⁵  | Patti Virtue^{1,5,6}  |
 Peter D. Nichols^{1,5}  | Catriona L. Hurd¹ 

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

²Marine Ecology, University of Groningen, Groningen, The Netherlands

³Instituto de Fomento Pesquero (IFOP), Puerto Montt, Chile

⁴Department of Chemistry, NIWA/University of Otago Research Centre for Oceanography, University of Otago, Dunedin, New Zealand

⁵CSIRO Oceans and Atmosphere, Hobart, Tasmania, Australia

⁶Antarctic Climate and Ecosystems, Cooperative Research Centre, Hobart, Tasmania, Australia

Correspondence

Luna M. van der Loos, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia.
 Email: lunavdloos@gmail.com

Funding information

Holland Scholarship; Groningen University Fund; Deutsche Forschungsgemeinschaft

Abstract

Increased plant biomass is observed in terrestrial systems due to rising levels of atmospheric CO₂, but responses of marine macroalgae to CO₂ enrichment are unclear. The 200% increase in CO₂ by 2100 is predicted to enhance the productivity of fleshy macroalgae that acquire inorganic carbon solely as CO₂ (non-carbon dioxide-concentrating mechanism [CCM] species—i.e., species without a carbon dioxide-concentrating mechanism), whereas those that additionally uptake bicarbonate (CCM species) are predicted to respond neutrally or positively depending on their affinity for bicarbonate. Previous studies, however, show that fleshy macroalgae exhibit a broad variety of responses to CO₂ enrichment and the underlying mechanisms are largely unknown. This physiological study compared the responses of a CCM species (*Lomentaria australis*) with a non-CCM species (*Craspedocarpus ramentaceus*) to CO₂ enrichment with regards to growth, net photosynthesis, and biochemistry. Contrary to expectations, there was no enrichment effect for the non-CCM species, whereas the CCM species had a twofold greater growth rate, likely driven by a downregulation of the energetically costly CCM(s). This saved energy was invested into new growth rather than storage lipids and fatty acids. In addition, we conducted a comprehensive literature synthesis to examine the extent to which the growth and photosynthetic responses of fleshy macroalgae to elevated CO₂ are related to their carbon acquisition strategies. Findings highlight that the responses of macroalgae to CO₂ enrichment cannot be inferred solely from their carbon uptake strategy, and targeted physiological experiments on a wider range of species are needed to better predict responses of macroalgae to future oceanic change.

KEYWORDS

carbon uptake strategy, carbon dioxide-concentrating mechanism, CCM, CO₂ enrichment, macroalgae, non-CCM, ocean acidification, physiology

⁹Holland Scholarship; Groningen University

Fund; M.S. was funded by the Deutsche Forschungsgemeinschaft (DFG, grant ID: SCHM 3335/1-1)

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Since the beginning of the industrial era, atmospheric CO₂ concentrations have increased by 40% (IPCC, 2013). In terrestrial systems, there is evidence of a “CO₂ enrichment” effect that is resulting in a higher plant biomass (Zhu et al., 2016) as the availability of CO₂ for carbon fixation in photosynthetic organisms via the enzyme Rubisco increases. In coastal marine systems, macroalgae fulfill the same functional role as terrestrial plants; they are foundation species that provide food, habitat, and shelter to higher trophic levels (e.g., shellfish and fish), as well as a range of ecosystem services including nutrient cycling. In addition, they have a potential role in the blue carbon economy and climate mitigation (Bennett et al., 2015; Chung, Beardall, Mehta, Sahoo, & Stojkovic, 2011; Tuya, Wernberg, & Thomsen, 2008). By the end of the century, CO₂ concentrations in seawater will increase by 200% under the RCP (*Representative Concentration Pathway*) 6.0 scenario (IPCC, 2013) and, like terrestrial plants, macroalgal productivity is predicted to increase (Koch, Bowes, Ross, & Zhang, 2013; Sunday et al., 2016).

The absorption of CO₂ by seawater and the subsequent chemical reactions cause a suite of changes to the seawater carbonate system, termed ocean acidification: By 2100, the bicarbonate ion will additionally increase by 14% with a concurrent decline in pH of 0.3 units (RCP6.0 scenario; Hurd, Hepburn, Currie, Raven, & Hunter, 2009; IPCC, 2013; The Royal Society, 2005). In addition to taking up CO₂ via diffusion, ~65% of macroalgae can utilize the bicarbonate ion directly from seawater (Kübler & Dudgeon, 2015). Bicarbonate use is an energy-consuming process, and the derivation of CO₂ from bicarbonate (for subsequent carbon fixation) is termed a carbon dioxide-concentrating mechanism (CCM). Species that solely rely on the diffusive uptake of CO₂, termed non-CCM species (mostly Phylum Rhodophyta), were thought to be rare (Kübler & Dudgeon, 2015; Raven, Ball, Beardall, Giordano, & Maberly, 2005). However, Cornwall, Revill, and Hurd (2015) discovered that up to 90% of some populations were non-CCM species in subtidal reef habitats in Tasmania, southern Australia, and on average 35% of macroalgae globally are now considered non-CCM (Kübler & Dudgeon, 2015).

Extensive field surveys have been conducted to study CCM and non-CCM species in temperate (Cornwall et al., 2015; Hepburn et al., 2011) and tropical systems (Diaz-Pulido, Cornwall, Gartrell, Hurd, & Tran, 2016), and along a gradient of CO₂/pH in a volcanic vent in Italy (Cornwall et al., 2017). These surveys indicate that the responses of macroalgae to CO₂ enrichment will depend on their mechanism(s) of inorganic carbon uptake, that is, their carbon uptake strategies (Hepburn et al., 2011). For CCM species, the predominance of the bicarbonate ion in seawater (90% under current pH) may mean that their growth is not limited by dissolved inorganic carbon (DIC) supply and there will be no future CO₂ enrichment effect. However, field surveys have illustrated that within CCM species we can further distinguish between (a) those that are able to use the 200% additional CO₂ predicted for the future and (b) those that cannot (Cornwall et al., 2017; Hepburn et al., 2011; Kübler & Dudgeon, 2015): (a) Species

able to use additional CO₂ either exhibit a low affinity for DIC (carbon-limited CCM species) or have CCMs that can be downregulated (i.e., decreased CCM activity); (b) CCM species that are considered unresponsive to future CO₂ concentrations have either a high affinity for CO₂ or CCMs that cannot be downregulated under elevated CO₂ conditions (Cornwall et al., 2017). In contrast, growth and photosynthesis of non-CCM species are thought to be limited under current CO₂ concentrations and they are predicted to benefit from the future 200% increase in dissolved CO₂ (Cornwall et al., 2015; Hepburn et al., 2011; Kübler & Dudgeon, 2015).

Carbon uptake strategies in macroalgae can be determined by analyzing their carbon stable isotope composition ($\delta^{13}\text{C}$) (Maberly, Raven, & Johnston, 1992): CO₂ is more depleted in ¹³C than HCO₃⁻ and thus has a lower $\delta^{13}\text{C}$ value. Macroalgae that possess a CCM typically have $\delta^{13}\text{C}$ values ranging between -30‰ and -10‰, whereas a lack of a CCM is indicated by $\delta^{13}\text{C}$ < -30‰. A change in the $\delta^{13}\text{C}$ value of the macroalgal tissue indicates a change in carbon uptake strategy and can also indicate biochemical changes in the macroalgae such as the use of sugars and lipid synthesis (Raven et al., 2002). Macroalgae generally exhibit significant changes in both lipids and fatty acids with changing environmental conditions. Such biochemical changes can be observed in the biomass very shortly after exposure (≤ 7 days) as an acclimation mechanism in membrane and/or storage lipids (Al-Hasan, Hantash, & Radwan, 1991; Gosch, Lawton, Paul, Nys, & Magnusson, 2015). Elevated CO₂ concentrations have been shown to either increase or decrease the lipid production of several species of algae, especially when combined with nutrient limitation (Bermúdez et al., 2015; Sun, Chen, & Du, 2016). CO₂ enrichment could therefore not only result in increased growth rates, but also lead to changes in lipid storage. Whether a species will invest in growth or lipid storage may depend on their carbon uptake strategy. Species that are currently saturated for DIC (i.e., CCM species) may use the elevated DIC to invest in lipid storage, since their growth rates are not limited by carbon. Species whose growth rates are currently limited by DIC (i.e., non-CCM species) may increase their growth rates first before investing in storage compounds.

Given that both CO₂ and bicarbonate concentrations are increasing in seawater, and macroalgae have a variety of inorganic carbon uptake strategies (Diaz-Pulido et al., 2016), determining the physiological and growth responses of macroalgae to future CO₂ enrichment is complex. Despite the ecological, economic, and cultural importance of macroalgae, we know very little about their mechanistic physiological responses to future CO₂ enrichment, particularly compared to microalgae and terrestrial plants (Hibberd & Covshoff, 2010; Jungnick et al., 2014; Sandrini, Matthijs, Verspagen, Muyzer, & Huisman, 2014).

While field surveys have proved useful in identifying the general carbon uptake strategies of macroalgae, detailed knowledge of photosynthetic and growth responses to CO₂ enrichment is unknown for most species and mainly focused on CCM species (Britton, Cornwall, Revill, Hurd, & Johnson, 2016; Fernández, Roleda, & Hurd, 2015; Israel & Hophy, 2002) and calcareous species (Hall-Spencer et al., 2008; Jokiel et al., 2008). The effect of

CO₂ enrichment on non-CCM species has been tested only twice, for *Lomentaria articulata* (Kübler, Johnston, & Raven, 1999) and *Amansia rhodantha* (Ho & Carpenter, 2017), with results being inconsistent between studies: *Lomentaria articulata* exhibited an increase in growth, while no response was detected for *Amansia rhodantha*. Likewise, for CCM species, both increased growth (Kim et al., 2016), no response (Fernández et al., 2015; Ho & Carpenter, 2017; Israel & Hophy, 2002; Rautenberger et al., 2015), and decreased growth have been reported (García-Sánchez, Fernández, & Niell, 1994; Kim et al., 2016). Therefore, the question remains as to whether or not carbon uptake strategy can be used as a predictor of macroalgal responses to CO₂ enrichment.

In this study, we compare the effects of CO₂ enrichment on the growth, physiological, and biochemical responses of two temperate red macroalgae: a CCM species, *Lomentaria australis* (Kützting) Levring, and a non-CCM species, *Craspedocarpus ramentaceus* (C. Agardh) Min-Thein & Womersley. Both species are ecological dominants in the subtidal waters of eastern Tasmania, Australia. As this is only the third non-CCM species to be included in a CO₂ enrichment experiment, as well as the first study to compare biochemical responses of a non-CCM species versus a CCM species, we aim to advance our understanding of the mechanisms that underlie macroalgal responses to CO₂ enrichment. In addition, we conduct a comprehensive literature survey, including studies conducted prior to ocean acidification becoming a research field, to better understand the role that carbon uptake strategy plays in effecting the response of fleshy macroalgae to CO₂ enrichment.

We hypothesized that: (a) CO₂ enrichment will have no effect on the growth and photosynthetic rate of *L. australis* (CCM species), if the species used in this study has a high affinity for DIC. However, if the species has a low affinity for DIC, we predict a downregulation of energy-costly uptake of HCO₃⁻, with an increased passive CO₂ uptake resulting in higher rates of growth and photosynthesis. (b) The net photosynthesis and growth of *C. ramentaceus* (non-CCM species) will increase with CO₂ enrichment, based on the likelihood that the species is currently carbon-limited. (c) An elevated DIC availability will be used for the accumulation of storage compounds such as lipids in *L. australis* (CCM species), resulting in composition changes in lipid classes and fatty acids, whereas *C. ramentaceus* (non-CCM species) will primarily invest in growth rather than storage of lipids.

2 | MATERIALS AND METHODS

2.1 | Macroalgal collection and culture maintenance

Twelve individual specimens of *Lomentaria australis* (a CCM species) and *Craspedocarpus ramentaceus* (a non-CCM species) were collected from Tinderbox, Tasmania (S43°03'30.722 E147°19'52.583), on 21 October 2016 at 6 m depth using Scuba and were transported to the laboratory (30 min) in plastic bags containing seawater. Carbon uptake strategy of both species was confirmed by conducting pH-drift experiments following Cornwall et al. (2015) (Supporting information Table S1). In the laboratory, the macroalgae were carefully

rinsed with filtered seawater and visible epiphytes were gently removed. Prior to the experiment, individuals were acclimated to laboratory conditions in 20-L aquaria for 48 hr, with constant aeration and a 12:12 light:dark photoperiod and a photon flux density (PFD) of 18–20 μmol photons m⁻² s⁻¹. During the pretreatment (24 hr) and experiment (7 days), the PFD was raised to 25–30 μmol photons m⁻² s⁻¹ which mimicked the PFD that was measured at the collection site at 6 m using integrating PAR sensors (Odyssey, Dataflow Systems Pty Ltd, Christchurch, New Zealand). In addition, light saturation curves were conducted to ensure that the light levels used in the experiment were sufficient for saturating photosynthesis (Supporting information Figure S1, Table S2). Seawater used in the experiment was collected south of Bruny Island, Tasmania, and was filtered to 1 μm and UV-sterilized (Emperor Aquatics Smart HO UV Sterilizer, 025050-2, 50 W lamp). The ammonium and nitrate concentrations were 0.46 and 0.20 μM, respectively, measured using a QuickChem 8500 Series 2 Automated Ion Analyzer (Lachat Instrument, Loveland, USA).

2.2 | Experimental conditions

For each species, ~1 g of alga was randomly assigned to an individual container (650 ml) of an automated, pH-controlled culture system. For both species, there were two experimental CO₂ treatments: “current” (pH_T = 8.00, pCO₂ = 365.17) and “future” (pH_T = 7.70, pCO₂ = 1,015.24) at 12.5°C (detailed in the Supporting information Table S3). This corresponds to the local ambient pH/CO₂ and the expected future seawater conditions, with *n* = 6 replicates for each species and CO₂ treatment. The future pH was based on the RCP6.0, which is a “business as usual” modeling projection (IPCC, 2013). A magnetic stirrer in each container (set at 650 rpm) mixed the seawater to minimize the diffusion boundary layer around the blades of the macroalgae.

Target CO₂ levels were achieved in each of the 24 culture containers using a system similar to that described in Bockmon, Frieder, Navarro, White-Kershek, and Dickson (2013) with the modifications presented in Reidenbach et al. (2017). Briefly, every 4 hr, seawater in each 650-ml culture container was replaced by flushing the container with ~1 L of seawater using Jebao DP-4 Auto Dosing aquarium pumps. CO₂ was added to this incoming seawater as it entered each culture container using a mixture of air and CO₂, which was briefly in contact with seawater entering each culture container using a membrane contactor (Micromodule, model 0.5 × 1, Membrana, USA) placed near each culture container inlet. The contactors contain a microporous hollow fiber membrane, which allows efficient mixing of liquids and gases. Since the seawater and air/CO₂ were in contact for only a few seconds, full equilibration of the air/CO₂ mixture was not possible and air/CO₂ ratios of approximately 400:1–1,000:1 were needed to achieved target CO₂ levels in the seawater. These ratios were achieved using mass flow controllers (FMA5418A for air, FMA5402A for CO₂; Omega Engineering, USA), whose flow rates were controlled by an analog output module housed in a USB chassis (NI9264 and cDAQ-9174,

National Instruments, USA). To maintain target CO₂ levels, a feedback system monitored pH_T and adjusted the flow rates of the mass flow controllers automatically.

Seawater pH_T was measured using a modified version of the spectrophotometric pH_T system developed by McGraw et al. (2010). Briefly, a syringe pump and rotary valves were used to sample seawater (V6 pump, 24090 and 24493 valves, 23425 valve driver, Norgren, UK) while minimizing gas exchange. Spectra were acquired using an LED light source and a UV-Vis spectrometer (BluLoop and USB2000+, Ocean Optics, USA) with a 1-cm flow-through quartz cuvette. Reference spectra were obtained from a 25 ml seawater sample; sample spectra were obtained by mixing 24.80 ml of seawater and 200 µl of 2 mM metacresol purple dye (857890, Sigma-Aldrich, Australia) within the syringe pump. The temperature of each sample was recorded with a PT100 temperature sensor and a high-precision data logger (PT-104, PICO Technology, UK). All instrument control, spectra manipulations, and pH_T calculations were done using LabVIEW 2014 (National Instruments, USA).

pH_T was calculated from the temperature, salinity, and the absorbance spectra, which was calculated from the individual reference and sample spectra. To improve measurement precision, the absorbance at 434, 578, and 750 nm was determined using the quadratic fits of the absorbance spectra between 429–439 nm and 573–583 nm and a background signal averaged between 750 and 760 nm (McGraw et al., 2010). Each recorded measurement of pH_T was the average of four replicate measurements, which took ~2 min to obtain. The pH system was standardized using certified reference materials provided by Andrew Dickson, Scripps Institute for Oceanography, San Diego, USA (Dickson, Sabine, & Christian, 2007). A 0.03 pH unit offset was added to each measurement based on the difference between the measured pH and that calculated from the known DIC and A_T of the certified reference material. The appropriateness of this correction was verified through pH_T measurements of the Tris buffer.

2.3 | Biotic responses

Growth rate was determined from changes in both thallus length and wet weight. Linear extension (distance from the apices to the main axis) was calculated from photographs of each individual placed on a grid on day 1 and day 7 using the software ImageJ (Schneider, Rasband, & Eliceiri, 2012). Relative growth rate (RGR) on a wet weight basis was calculated for each individual according to Yong, Yong, and Anton (2013): $RGR = \ln(W_t/W_0) \times t^{-1}$, where W_0 is the wet weight at day 1 and W_t is the final wet weight after 7 days. To remove surface water, blades were carefully blotted dry with tissue paper before weighing.

Net photosynthetic rates were measured at the end of the experiment (day 7) at the beginning of the light cycle in each culture tank under experimental conditions using an Orion RDO Probe 087010MD. Initial O₂ measurements were conducted immediately after the culture tanks received a fresh seawater supply, and final O₂ measurements were conducted two hours later. The probe was

calibrated using a standard of 100% air saturation achieved by bubbling with air for 10 min. Oxygen production was calculated as final O₂ measurements – initial O₂ measurements and was standardized to tissue wet weight (g) per hour.

The maximum quantum yield (F_v/F_m) was used as an indicator of the performance of photosystem II (PSII) at the beginning and at the end of the experiment. F_v/F_m was measured using a pulse amplitude modulation (PAM) chlorophyll fluorescence meter (Diving PAM, Walz, Germany) using individuals that had been dark-adapted for 10 min. The gain was set to 2, and F_0 ranged between 200 and 1,000 for each measurement.

The content of photosynthetic pigments (chlorophyll *a* and phycobiliproteins) was analyzed using ~0.2 g (wet weight) of each individual on day 7. Chlorophyll *a* was extracted at room temperature according to Seely, Duncan, and Vidaver (1972) and Schmidt, Maraschin, and Bouzon (2010) using acetone and dimethylsulfoxide (DMSO). Pigment content was quantified spectrophotometrically with a S-22 UV/Vis Spectrophotometer (Boeco, Germany), and concentrations (mg/g of wet weight) were calculated using the equations in Seely et al. (1972). To determine phycobiliprotein content (mg/g of wet weight), samples were flash frozen by immersion in liquid nitrogen and ground into a fine powder with a mortar and pestle. The extraction was carried out at 4°C using a 0.1 M phosphate buffer (pH = 7.2) according to the methods of Schmidt et al. (2010) and Sampath-Wiley and Neefus (2007). All extractions were conducted in the dark.

Tissue samples from the apices were taken during the pre-treatment and at the end of the experiment on day 7 from each individual sample to determine δ¹³C, C:N ratios, and lipid class and fatty acid content and composition. The samples were rinsed in Type I ultrapure water, frozen at –80°C, freeze-dried (Labconco FreeZone 4.5), and ground with a mortar and pestle. δ¹³C, C and N content, and C:N were determined following the methods in Cornwall et al. (2015) using a NA1500 elemental analyzer coupled to a Thermo Scientific Delta V Plus via a ConFlo IV. Combustion and reduction were achieved at 1,020°C and 650°C, respectively. Values were normalized to the VPDB scale (Vienna Pee Dee Belemnite) via a 3-point calibration using certified reference material. Both precision and accuracy were ±0.1 ‰ (1 SD). For the analyses, change in δ¹³C was calculated as the difference in δ¹³C between day 7 and day 1.

Total lipids were extracted following a modified version of Bligh and Dyer (1959) of the dried and weighed (ca. 40 mg) macroalgal tissue. Lipids were extracted overnight using a one-phase methanol (MeOH): dichloromethane (DCM): Milli-Q (2:1:08 v/v/v) solvent mixture. Phase separation was achieved by addition of 10 ml of DCM and 10 ml of Milli-Q water the next day. The lower lipid-containing layer was drained into a round bottom flask. After addition of few drops of MeOH, solvent was removed using a rotary evaporator (ca. 40°C). The lipid extracts were transferred to preweighed vials, and solvents were evaporated under a constant stream of nitrogen gas. The lipid-containing vials were weighed for determination of total lipids. Samples were redissolved in DCM and stored at –20°C until further analysis.

To determine lipid class composition, an aliquot of the total lipid extract was spotted on SIII chromarods (5 μm particle size). Samples were co-eluted with a standard mix to determine the lipid class composition including hydrocarbons (HC), steryl and wax esters (WE), triacylglycerols (TAG), sterols (ST), free fatty acids (FFA), and polar lipids which includes glycolipids and phospholipids (PL). The mobile phase consisted of hexane: diethyl ether (DEE): glacial acetic acid (GAA) (70:10:0.1 v/v/v). Chromarods were developed for 25 min and then dried at 100°C for 10 min. The dried rods were analyzed using an Iatroskan Mark V TH10 thin layer chromatography (TLC) with a flame ionization detector (FID). Peak identification was achieved by comparison with retention times of co-eluted standards. For quantification, the SIC480II Iatroskan™ integrating software (System Instruments, Mitsubishi Chemical Instruments) was used. The integrated areas were transformed to mass per μl spotted using pre-determined linear regression calculations.

To analyze the fatty acids, an aliquot of the total lipid extract was saponified by addition of 2 ml of 5% KOH in 80% MeOH and placed on a heating block (80°C for 3 hr). After cooling, 1 ml of H_2O was added and the mixture was extracted with hexane:DCM (4:1 v/v) to yield a nonsaponifiable neutral lipid fraction in the upper organic layer and free fatty acids in the lower aqueous layer. The aqueous layer was acidified by addition of 0.3 ml concentrated HCl and extracted three times with hexane:DCM (4:1, v/v). Solvents were removed under a stream of inert nitrogen gas, and methylation was performed by addition of MeOH:DCM:conc. HCl (10:1:1, v/v/v) and heating for 1 hr at 80°C. After cooling, 1 ml H_2O was added and the resulting fatty acid methyl esters (FAME) were extracted three times into hexane:DCM (4:1, v/v). Solvents were removed under a stream of nitrogen gas. The samples were analyzed using gas chromatography (GC) using an Agilent Technologies 7890 (Palo Alto, California, USA) GC coupled with a flame ionization detector (FID) analyzer and equipped with a nonpolar Equity™-1 fused silica capillary column (15 m \times 0.1 mm internal diameter, 0.1 μm film thickness). Agilent ChemStation software was used for quantification of FAME peaks. The individual fatty acids were identified using a GC-mass spectrometer (MS) using a Finnigan ThermoQuest GCQ GC-MS System fitted with an on-column injector and using ThermoQuest Xcalibur software.

To determine how macroalgae modify the carbonate chemistry of their local environment, seawater samples (13 ml) were collected from each replicate culture tank immediately after a change in seawater and again four hours later. These samples were collected both at day 1 and day 7. Simultaneously, pH_T was measured in the culture tanks with a Thermo Scientific Orion VERSA STAR 90 meter and pH electrode Orion 8107BNUMD Ross Ultra pH/ATC Triode, calibrated with a Tris buffer. Seawater samples were immediately poisoned with HgCl_2 and stored under constant darkness. DIC concentrations of the water samples were measured using a DIC analyzer (Apollo SciTech DIC Analyzer Model AS-C3) with an inbuilt CO_2 analyzer (LI-COR LI-7000 $\text{CO}_2/\text{H}_2\text{O}$ Analyzer). The CO_2 analyzer was calibrated with a certified reference material provided by Andrew Dickson, Scripps Institute for Oceanography, San Diego,

USA (Dickson et al., 2007). A_T was calculated using the constants of Mehrbach, Culberson, Hawley, and Pytkowicz (1973), refitted by Dickson and Millero (1987).

2.4 | Statistical analyses

All statistical analyses were performed using the software R 3.1.2 (R Core Team, 2016). A multivariate analysis of variance (MANOVA) was used to assess whether there were differences in biological responses (linear extension, wet weight RGR, Chl *a* and phycobiliprotein content, net photosynthesis, lipid content, fatty acid content, change in $\delta^{13}\text{C}$, change in C:N ratios, and change in carbon and nitrogen tissue content) between species and CO_2 treatments. A second MANOVA was performed on the change in carbonate parameters ($[\text{H}^+]$, $[\text{HCO}_3^-]$, $[\text{CO}_3^{2-}]$, $[\text{CO}_2]$, and total DIC). The data passed the tests for MANOVA assumptions (normality and homogeneity of variance). When the interaction or main factor pH was significant (at $\alpha = 0.05$), Tukey's honestly significant difference (THSD) post hoc tests were used to determine which treatments differed from each other (using the `aov()` and `TukeyHSD()` function in R). The Hedge's *g* is used as a measure of effect size (a *g* of 1 indicates that two groups differ by 1 SD). Hedge's *g* is calculated as follows: $(M1-M2)/SD_{\text{pooled}}$, where $M1-M2$ is the difference in means of two groups and SD_{pooled} is the pooled standard deviation. Differences in F_v/F_m values were tested with a generalized linear model using the `lme4` package, with treatment as fixed factor and start/end of the experiment as random factor (Bates, Mächler, Bolker, & Walker, 2015). A correspondence analysis (CA) was performed to detect differences in fatty acid composition between species and treatments using the `Vegan` package v.2.3-5 (Oksanen et al., 2016). In the CA plot, objects (specimens) that are close to one another are likely to have a similar fatty acid composition. Specimens found near the point representing a fatty acid are likely to contain a higher relative level of that fatty acid. For this analysis, only fatty acids that were on average >0.5% of the TFA profile were used. Graphs were customized using the packages `ggplot2` (Wickham, 2011), `plyr` (Wickham, 2011), and `reshape2` (Wickham, 2007).

2.5 | Literature synthesis

To further determine whether or not patterns in the responses of fleshy macroalgae to CO_2 enrichment depend on their carbon uptake strategy, we conducted a comprehensive literature survey. We searched the databases Web of Science and Google Scholar using the keywords "CO2 enrichment", "CO2 fertilization", "ocean acidification", "macroalgae", and "seaweed" in various combinations. Fifty-six studies published between 1989 and 2018 were analyzed, giving a total of 61 species. Studies were only examined if they met specific criteria. (a) The study had to report a growth response to CO_2 enrichment (in addition, photosynthetic response and $\delta^{13}\text{C}$ response were noted if those were reported); (b) the study needed to be a manipulative experiment comparing one or several macroalgal species in different $p\text{CO}_2$ treatments (thus, we excluded field studies

that were conducted in natural $p\text{CO}_2$ gradients); (c) the study had to report whether the species used in the experiment were able to utilize HCO_3^- or not (i.e., if the species possessed a CCM); (d) only studies on marine fleshy macroalgae were considered; (e) the macroalgae used in the experiment had to be identified to species level (i.e., no “turf”-forming mats or community responses were included); (f) studies testing the effect of CO_2 enrichment on microscopic early developmental stages and spore germination were excluded; and (g) the study had to be published in a peer-reviewed journal. Many studies also tested additional experimental conditions (e.g., irradiance or nutrient concentrations), in combination with elevated $p\text{CO}_2$. Those studies were included in our literature synthesis, but care was taken to only take into account the single-term effects of elevated $p\text{CO}_2$.

3 | RESULTS

3.1 | Growth

For *L. australis* (the CCM species), linear extension under the future CO_2 treatment (pH = 7.70) was twofold greater (a Hedge's g effect size of 1.5) than the current conditions treatment (pH = 8.04) (Table 1, Figure 1), whereas for *C. ramentaceus* (the non-CCM species), there was no significant effect of CO_2 enrichment on growth (Table 1, Figure 1). Growth rates of *L. australis* under the future treatment were ca. eightfold and sevenfold greater (a Hedge's g effect size of 2.8) than *C. ramentaceus* under both current and future treatments, respectively ($p < 0.0001$ for both comparisons, Tukey's honestly significant difference tests). Growth rate on a wet weight basis showed a similar CO_2 treatment effect ($p = 0.095$, Table 1; Supporting information Figure S2).

3.2 | Net photosynthesis, pigments, and F_v/F_m

Net photosynthetic rates were similar for *L. australis* and *C. ramentaceus*, ranging from 0.3 to 0.4 $\mu\text{mol hr}^{-1} \text{g}^{-1}$, and were unaffected by the CO_2 treatment (Table 1, Figure 2). *C. ramentaceus* (the non-CCM species) had a higher pigment content of both chlorophyll *a* and phycobiliproteins compared to *L. australis* (the CCM species), but pigment content was not affected by CO_2 treatment (Supporting information Figure S3, Figure S4). The F_v/F_m value ranged between 0.51–0.54 and 0.49–0.58 for *L. australis* and *C. ramentaceus*, respectively, throughout the experiment and was not significantly different between treatments ($p > 0.05$ for all comparisons, linear mixed model Wald chi-square test).

3.3 | Stable isotopes and C:N ratios

L. australis (the CCM species) under the future treatment had a ca. twofold greater decrease (a Hedge's effect size of 1.2) in $\delta^{13}\text{C}$, compared to the individuals of the same species in current conditions (Figure 3, $p < 0.05$, Tukey's honestly significant difference test). CO_2 treatment did not affect the degree of $\delta^{13}\text{C}$ change of *C. ramentaceus* (the non-CCM species) ($p > 0.05$, Tukey's honestly significant difference tests). C:N ratios increased from ~6.5 w/w at the start

to ~8.5 w/w at the end of the experiment for each of the four treatment–species combinations, due to an increase in carbon content of the tissue (Supporting information Table S4). The magnitude of change in C:N ratio and C content did not differ between species and CO_2 treatment (Table 1). The nitrogen content of the tissue remained relatively constant for both *L. australis* and *C. ramentaceus*, with less than 0.39% w/w difference from the start to end of the experiment (Supporting information Table S4).

3.4 | Biochemical responses

L. australis (the CCM species) had a higher total fatty acid content than *C. ramentaceus* (the non-CCM species), but this was not affected by experimental treatment (Table 1; Supporting information Figure S5). Both species had similar total lipid content (Table 1; Supporting information Figure S6). Polar lipids dominated the lipid class composition of both *L. australis* and *C. ramentaceus* (Supporting information Figure S7). There were no differences in lipid class percentage between species and CO_2 treatment (Supporting information Figure S7). A correspondence analysis (CA) showed that fatty acid composition differed significantly between species, but that the composition in each species did not change with CO_2 treatment (Figure 4). The first CA axis explained 98.2% of the variation, and the second axis explained 0.8% of the variation. The difference in fatty acid composition was mainly driven by 16:1n-7, 18:1n-9, 18:0, 20:4n-6, 20:3n-6, and 20:5n-3, where the former four fatty acids were more associated with *C. ramentaceus* and the latter two with *L. australis*.

3.5 | Modification of seawater carbonate chemistry by algae

Over a period of four hours, the algae modified the pH, total DIC, CO_2 , HCO_3^- , and CO_3^{2-} of the water in their culture tank, before they received a fresh supply of seawater (Supporting information Table S5). The $[\text{H}^+]$ decreased (i.e., pH increased), more so in the future treatments than in current treatments, but without differences between species (Table 1). The total DIC, HCO_3^- , and CO_2 decreased, and the CO_3^{2-} increased over the 4 hr. There were no significant differences between species and CO_2 treatments for the increase in CO_3^{2-} and the decrease in HCO_3^- and total DIC. The decrease in pH and CO_2 was greater in future conditions, but did not differ between species (Table 1).

3.6 | Literature synthesis

Approximately 40% of the CCM species that were tested (21 out of 55 species) showed no growth response to CO_2 enrichment, five species had decreased growth rates and 16 species increased growth with elevated CO_2 (Table 2). For 13 CCM species, mixed responses were reported (i.e., several species were tested in multiple studies, and the responses reported were not concordant). Net photosynthetic rates either increased or did not respond to CO_2 enrichment for the majority of the CCM species tested (11 and 10 species, respectively); in four species, photosynthesis decreased; nine species showed mixed

TABLE 1 Multivariate analysis of variance (MANOVA) table displaying *p*-values and *F*-values for all measured responses (linear extension, wet weight relative growth rate, net photosynthesis, chlorophyll a content, phycobiliprotein content, change in $\delta^{13}\text{C}$, change in C:N ratio, change in carbon tissue content, change in nitrogen tissue content, total lipid content, total fatty acid content, change in $[\text{H}^+]$, change in $[\text{HCO}_3^-]$, change in $[\text{CO}_2]$, change in $[\text{CO}_3^{2-}]$, and change in total DIC)

| Response | Interaction | | Factor | | | |
|---|-----------------|-----------------|-----------------|------------------|---------------------------|------------------|
| | | | Species | | CO ₂ treatment | |
| | <i>F</i> -value | <i>p</i> -value | <i>F</i> -value | <i>p</i> -value | <i>F</i> -value | <i>p</i> -value |
| Linear extension | 6.474 | 0.021 | – | – | – | – |
| Wet weight RGR | 3.125 | 0.095 | 15.743 | 0.001 | 2.133 | 0.162 |
| Chlorophyll a content (mg per g wet weight) | 0.509 | 0.485 | 20.563 | <0.001 | 0.169 | 0.686 |
| Phycobiliprotein content (mg per g wet weight) | 1.375 | 0.257 | 4.271 | 0.054 | 0.020 | 0.889 |
| Net photosynthesis ($\mu\text{mol O}_2 \text{ hr}^{-1} \text{ g}^{-1}$) | 1.813 | 0.196 | 1.385 | 0.256 | 0.075 | 0.788 |
| Change in $\delta^{13}\text{C}$ | 1.229 | 0.283 | 8.528 | 0.010 | 7.975 | 0.012 |
| Change in C:N ratio | 0.274 | 0.608 | 1.350 | 0.261 | 0.083 | 0.777 |
| Change in carbon tissue content | 0.012 | 0.914 | 1.051 | 0.320 | 0.110 | 0.744 |
| Change in nitrogen tissue content | 0.031 | 0.861 | 5.813 | 0.028 | 0.140 | 0.713 |
| Total lipid content | 0.134 | 0.719 | 2.458 | 0.135 | 0.121 | 0.733 |
| Total fatty acid content | 0.220 | 0.645 | 35.466 | <0.001 | 3.363 | 0.084 |
| Change in $[\text{H}^+]$ | 0.659 | 0.427 | 0.549 | 0.468 | 133.993 | <0.001 |
| Change in $[\text{HCO}_3^-]$ ($\mu\text{mol/kg}$) | 0.418 | 0.526 | 0.580 | 0.456 | 0.013 | 0.910 |
| Change in $[\text{CO}_2]$ ($\mu\text{mol/kg}$) | 0.501 | 0.488 | 0.395 | 0.537 | 134.789 | <0.001 |
| Change in $[\text{CO}_3^{2-}]$ ($\mu\text{mol/kg}$) | 0.025 | 0.877 | 1.671 | 0.212 | 0.171 | 0.684 |
| Change in total DIC ($\mu\text{mol/kg}$) | 0.754 | 0.396 | 0.119 | 0.734 | 2.572 | 0.125 |

Note. *p*-values and *F*-values of separate factors are not shown when the interaction between factors is significant ($\alpha = 0.05$). *p*-values in bold have a significance of $p < 0.05$.

responses; and for 21 species, photosynthetic responses were not reported (Table 2). Only three non-CCM species (including our work) have been studied in relation to CO₂ enrichment, with contrasting results in terms of both growth and net photosynthesis (Table 2). Overall, $\delta^{13}\text{C}$ values were measured in only a minority of the studies published (10 out of 56 studies) (Table 2). None of the species increased $\delta^{13}\text{C}$ values in response to elevated CO₂, but no clear pattern in relation to growth and photosynthesis can be inferred.

4 | DISCUSSION

Carbon uptake strategy has been a key concept in developing hypotheses related to how fleshy macroalgae will respond to CO₂ enrichment (Gutow et al., 2014; Hepburn et al., 2011; Ho & Carpenter, 2017; Hurd, Lenton, Tilbrook, & Boyd, 2018; Israel & Hophy, 2002; Johnson, Comeau, Lantz, & Smith, 2017; Koch et al., 2013). The aim of this study was to compare the physiological and biochemical responses of *L. australis* (a CCM species) with *C. ramentaceus* (a non-CCM species) to CO₂ enrichment in order to broaden our knowledge on whether or not carbon uptake strategy is a useful predictor of the responses of fleshy macroalgae to CO₂ enrichment. Our results,

along with a careful scrutiny of the literature, reveal that fleshy macroalgal species exhibit a wide array of responses to CO₂ enrichment in terms of both growth and photosynthesis (Table 2). As these responses also vary between species with the same carbon uptake strategy, we suggest that it is not currently possible to predict the responses of macroalgae to a higher CO₂ ocean based solely on the presence/absence or type of CCM: A greater mechanistic understanding is needed before such predictions can be made.

4.1 | The response of CCM species

For CCM species that responded positively to CO₂ enrichment, the range of growth responses varied from a ~ 1.2-fold increase in *Ulva australis* (Kim et al., 2016) to a ~ 4-fold increase in *Gracilaria chilensis* (Gao, Aruga, Asada, & Kiyohara, 1993), fitting well with the twofold response of *Lomentaria australis* (the CCM species in our study). However, it is noteworthy that mixed growth responses to CO₂ enrichment were reported for 13 of the CCM species and mixed photosynthetic responses for nine of the CCM species included in our literature survey (Table 2), as several species have been tested in multiple studies and the results did not always match. This inconsistency is most likely because other environmental factors also affect growth and physiological responses.




FIGURE 1 Linear extension (cm) of an algal species with carbon-concentrating mechanism (*Lomentaria australis*; CCM species) and a species without (*Craspedocarpus ramentaceus*; non-CCM species), with current (8.0) and future (7.7) CO₂ treatment. Data are displayed as mean ± standard error, $n = 6$. Bars sharing a letter are not significantly different (Tukey's honestly significant difference tests, $\alpha = 0.05$)




FIGURE 3 Change in $\delta^{13}\text{C}$ ratios between day 1 (start of the experiment) and day 7 (end of the experiment) of a species with carbon-concentrating mechanism (*Lomentaria australis*; CCM species) and a species without (*Craspedocarpus ramentaceus*; non-CCM species), in current (8.0) and future (7.7) CO₂ treatment. Data are displayed as mean ± standard error, $n = 5-6$ (with $n = 5$ for the ambient CCM treatment, and $n = 6$ for all other treatments). * denotes significantly different treatments (Tukey's honestly significant difference tests, $\alpha = 0.05$)




FIGURE 2 Net photosynthesis rates of a species with carbon-concentrating mechanism (*Lomentaria australis*; CCM species) and a species without (*Craspedocarpus ramentaceus*; non-CCM species), with current (8.0) and future (7.7) CO₂ treatment. Data are displayed as mean ± standard error, $n = 6$. There were no significant differences between species and treatments




FIGURE 4 Correspondence analysis based on fatty acid composition of a species with carbon dioxide-concentrating mechanism (*Lomentaria australis*; CCM species; gray) and a species without (*Craspedocarpus ramentaceus*; non-CCM species; white), under current (circles) and future (squares) conditions. Crosses denote fatty acids. The nine most important fatty acids (the fatty acids that explain most of the variability) have been labeled

TABLE 2 Response to CO₂ enrichment reported in the literature for fleshy macroalgal species, regarding growth, photosynthesis, and δ¹³C values ("I" = increase, "D" = decrease, "NR" = no response, "-" = unreported)

| Phylum and species | Enriched pCO ₂ level (µatm) | Response to CO ₂ enrichment | | | Putative carbon uptake strategy | Reference |
|---|--|--|----------------|-------------------|---------------------------------|---------------|
| | | Growth | Photosynthesis | δ ¹³ C | | |
| Rhodophyta (Red algae) | | | | | | |
| <i>Amansia rhodantha</i> | ~1,000 | NR | - | - | Non-CCM | [1] |
| <i>Chondrus crispus</i> | ~800 | NR & I | NR, D & I | - | CCM | [2] |
| <i>Craspedocarpus ramentaceus</i> | ~1,000 | NR | NR | NR | Non-CCM | current study |
| <i>Gelidium crinale</i> | ~750 | NR | - | - | CCM | [3] |
| <i>Gracilaria chilensis</i> | 650 & 1,250 | I | I | - | CCM | [4] |
| <i>G. conferta</i> | ~750 | NR | - | - | CCM | [3] |
| <i>G. secundata</i> | - | I | - | - | CCM | [5] |
| <i>G. tenuistipitata</i> | - | D | D | - | CCM | [6] |
| <i>G. tikvahiae</i> | (pH = 6.0) | NR | I | - | CCM | [7] |
| <i>Gracilariopsis lemaneiformis</i> * | ~700, ~1,000 & ~1,400 | NR & I | NR & I | - | CCM | [8-12] |
| <i>Grateloupia cornea</i> | 900 & 1,900 | D | D | - | CCM | [13] |
| <i>Hypnea cornuta</i> | ~750 | NR | - | - | CCM | [3] |
| <i>H. musciformis</i> | ~750 | D | - | - | CCM | [3] |
| <i>H. spinella</i> | 700 & 1,600 | I | I | - | CCM | [14] |
| <i>Laurencia intricata</i> | ~1,000 | NR | I | D | CCM | [15] |
| <i>Lomentaria articulata</i> | 700-1,800 (range) | I | - | D | Non-CCM | [16] |
| <i>L. australis</i> | ~1,000 | I | NR | D | CCM | current study |
| <i>Melanothamnus harveyi</i> | ~800 & ~1,500 | NR & I | NR & I | - | CCM | [17] |
| <i>Palmaria palmata</i> | ~1,000 | NR & D | D | - | CCM | [18,19] |
| <i>Phycodrys rubens</i> | ~1,000 | NR | - | - | Possibly non-CCM | [20] |
| <i>Plocamium cartilagineum</i> | 900 | D & I | - | - | CCM | [21] |
| <i>Porphyra linearis</i> | ~750 | D | NR | - | CCM | [22] |
| <i>Ptilota gunneri</i> | ~1,000 | NR | - | - | Possibly non-CCM | [20] |
| <i>Pterocladia capillacea</i> | ~750 | NR | - | - | CCM | [3] |
| <i>Pyropia haitanensis</i> | 1,000 | I | D & I | - | CCM | [23,24] |
| <i>P. leucosticta</i> | - | D | I | - | CCM | [25] |
| <i>P. yezoensis</i> | 1,000 & 1600 | I | I | - | CCM | [26] |
| Chlorophyta (Green algae) | | | | | | |
| <i>Chaetomorpha linum</i> | (pH = 6.73) | I | - | - | CCM | [27] |
| <i>Cladophora coelothrix</i> | (pH = 6.73) | I | - | - | CCM | [27] |
| <i>C. patentiramea</i> | (pH = 6.73) | NR | - | - | CCM | [27] |
| <i>C. vagabunda</i> | (pH = 6.0) | I | I | - | Possibly non-CCM | [7] |
| <i>Codium fragile</i> | 900 & 1,900 | NR | NR | - | CCM | [13] |
| <i>Monostroma grevillei</i> var. <i>arctica</i> | ~1,000 | NR | - | - | CCM | [20] |
| <i>Ulva australis</i> * | ~900 & ~1,000 & ~1,900 | NR & I | NR & I | - | CCM | [13,28-30] |
| <i>U. lactuca</i> | ~700 | NR | NR | - | CCM | [10,31] |
| <i>U. linza</i> | ~750 & ~1,000 | NR | D | - | CCM | [3,32] |
| <i>U. prolifera</i> | ~1,000 | I | NR | - | CCM | [33-35] |
| <i>U. pulchra</i> | - | I | - | - | CCM | [36] |

(Continues)

TABLE 2 (Continued)

| Phylum and species | Enriched $p\text{CO}_2$ level (μatm) | Response to CO_2 enrichment | | | Putative carbon uptake strategy | Reference |
|--|---|--------------------------------------|----------------|-----------------------|---------------------------------|------------------|
| | | Growth | Photosynthesis | $\delta^{13}\text{C}$ | | |
| <i>U. reticulata</i> | – | NR | – | – | CCM | [36] |
| <i>U. rigida</i> * | ~1,200 | NR & I | NR & D | NR | CCM | [36–39] |
| Ochrophyta (Brown algae) | | | | | | |
| <i>Alaria esculenta</i> * | ~1,000 & ~1,300 | NR, D & I | NR | D | CCM | [20,40,41] |
| <i>Chnoospora implexa</i> | ~1,000 | NR | I | D | CCM | [15] |
| <i>Desmarestia aculeata</i> | ~1,000 & ~1,300 | D & I | NR & I | D & NR | CCM | [20,40,42] |
| <i>Dictyopteria undulata</i> | 900 | I | – | – | CCM | [21] |
| <i>Dictyota bartayresiana</i> | ~1,000 | NR | – | – | CCM | [1] |
| <i>Fucus vesiculosus</i> | ~1,200 | NR & D | – | – | CCM | [43–45] |
| <i>F. vesiculosus</i> f. <i>mytili</i> | 1,000 | I | – | – | CCM | [46] |
| <i>Laminaria solidungula</i> | ~1,200 | NR | NR | NR | CCM | [47] |
| <i>Lobophora variegata</i> | ~1,000 | NR | – | – | CCM | [1] |
| <i>Macrocystis pyrifera</i> | ~1,200 | NR | NR | NR | CCM | [48] |
| <i>Nereocystis luetkeana</i> | ~3,000 | I | I | – | CCM | [49,50] |
| <i>Padina pavonica</i> | ~750 | NR | – | – | CCM | [3] |
| <i>Saccharina japonica</i> | ~1800 | NR | I | – | CCM | [51] |
| <i>S. latissima</i> * | ~1,000 & ~1,200 & ~3,000 | NR, D & I | NR & I | NR & D | CCM | [18,41,47,49,52] |
| <i>Saccorhiza dermatodea</i> | ~1,000 | I | – | – | CCM | [20] |
| <i>Sargassum fusiforme</i> * | ~700 & ~1,000 | D & I | NR & I | – | CCM | [53–55] |
| <i>S. horneri</i> | 900 & 1,900 | NR & I | NR | – | CCM | [13,21] |
| <i>S. muticum</i> | 1,000 | I | I | – | CCM | [56] |
| <i>S. thunbergii</i> | 900 & 1,900 | I | I | – | CCM | [13] |
| <i>S. vulgare</i> | ~750 | NR | – | – | CCM | [3] |
| <i>Turbinaria ornata</i> | ~1,000 | NR | NR | NR | CCM | [15] |

Notes. The elevated $p\text{CO}_2$ level (which was compared to ambient control levels in each study), the carbon uptake strategy (non-CCM or CCM), and references are also noted. * indicates species of which detailed physiological and biochemical regulatory mechanisms are known.

References: [1] Ho and Carpenter (2017); [2] Sarker, Bartsch, Olischläger, Gutow, and Wiencke (2013); [3] Israel and Hophy (2002); [4] Gao et al. (1993); [5] Lignell and Pedersén (1989); [6] García-Sánchez et al. (1994); [7] Rivers and Peckol (1995); [8] Chen, Zou, Zhu, and Yang (2017); [9] Xu, Zou, and Gao (2010); [10] Liu, Zou, and Yang (2018); [11] Zou and Gao (2009); [12] Kang, Kambey, Shen, Yang, and Chung (2017); [13] Kim et al. (2016); [14] Suárez-Álvarez, Gómez-Pinchetti, and García-Reina (2012); [15] Bender-Champ, Diaz-Pulido, and Dove (2017); [16] Kübler et al. (1999); [17] Olischläger and Wiencke (2013); [18] Nunes et al. (2016); [19] Sebök, Herppich, and Hanelt (2017); [20] Gordillo, Carmona, Viñegla, Wiencke, and Jiménez (2016); [21] Kram et al. (2016); [22] Israel, Katz, Dubinsky, Merrill, and Friedlander (1999); [23] Liu and Zou (2015a); [24] Xu, Chen, et al. (2017); [25] Mercado, Javier, Gordillo, Xavier Niell, and Figueroa (1999); [26] Gao et al. (1991); [27] de Paula Silva, Paul, Nys, and Mata (2013); [28] Reidenbach et al. (2017); [29] Kang and Chung (2017); [30] Kang and Kim (2016); [31] Liu and Zou (2015b); [32] Gao et al. (2018); [33] Xu and Gao (2012); [34] Li, Xu, and He (2016); [35] Li, Zhong, Zheng, Zhuo, and Xu (2018); [36] Björk, Haglund, Ramazanov, and Pedersén (1993); [37] Gordillo, Niell, and Figueroa (2001); [38] Rautenberger et al. (2015); [39] Gordillo, Figueroa, and Niell (2003); [40] Iñiguez et al. (2016a); [41] Gordillo et al. (2015); [42] Iñiguez, Heinrich, Harms, and Gordillo (2017); [43] Gutow et al. (2014); [44] Kawamitsu and Boyer (1999); [45] Ober and Thornber (2017); [46] Mensch et al. (2016); [47] Iñiguez et al. (2016b); [48] Fernández et al. (2015); [49] Swanson and Fox (2007); [50] Thom (1996); [51] Kang and Chung (2018); [52] Olischläger, Iñiguez, Koch, Wiencke, and Gordillo (2017); [53] Zou (2005); [54] Zou, Gao, and Luo (2011); [55] Jiang, Zou, Lou, and Gong (2018); [56] Xu, Gao, Gao, Xu, and Wu (2017).

For example, the kelp *Saccharina latissima* showed either no response or increased growth rates to elevated $p\text{CO}_2$, depending on levels of UV radiation (Gordillo, Aguilera, Wiencke, & Jiménez, 2015). In another study, *S. latissima* had decreased growth rates in response to CO_2 enrichment (Swanson & Fox, 2007), but the $p\text{CO}_2$ used (~3,000 μatm) was much higher (3 \times) than that typically used in OA experiments and may be unrealistic, as the corresponding decrease in pH is larger than the worst-case IPCC scenario (RCP8.5; IPCC, 2013). Overall, our synthesis

indicates that many (~40%) CCM species are likely to be saturated for inorganic carbon, but that the benefits of CO_2 enrichment for those that are not saturated for inorganic carbon could be substantial in terms of increased growth and productivity.

The reason for the decreased growth rate recorded for five CCM species with CO_2 enrichment is unknown, but may be due to a sensitivity to increasing H^+ concentrations. This sensitivity is known to exist in calcifying organisms (e.g., coccolithophores and




FIGURE 5 Predicted physiological and growth response of fleshy macroalgae to CO_2 enrichment based on their carbon uptake strategy and affinity for DIC. Species with a low affinity for DIC are likely to be limited in DIC under current conditions, and species with a high affinity are likely to be saturated for DIC. There is literature evidence that some CCM species are sensitive to increased H^+ concentrations, illustrated on the right hand side of the figure: Although not illustrated, H^+ sensitivity for non-CCM species may also be possible. This figure builds on that of Cornwall et al. (2017)

zooplankton), as elevated H^+ directly decreases calcification, and in fish, where increasing H^+ causes acidosis and behavioral changes (Hurd et al., 2018). As H^+ is key in regulating cellular homeostasis, changing H^+ concentrations could affect macroalgal metabolic processes and CCM activity. The effect of increasing H^+ concentrations has not been addressed for fleshy macroalgae, with the exception of Roleda, Morris, McGraw, and Hurd (2012), who found an interactive effect of pH (i.e., H^+) and DIC on early microscopic life-history stages of *Macrocystis pyrifera*. As H^+ concentrations will increase by 200% by 2100 (RCP6.0 scenario; IPCC, 2013), this calls for future research on how increasing H^+ concentrations, and the interaction with DIC enrichment, affect macroalgal physiology.

The increase in growth rate of *Lomentaria australis* with elevated CO_2 most likely occurred as a result of the downregulation of an energy-costly CCM. This idea is supported by the greater shift to more negative $\delta^{13}\text{C}$ values under future CO_2 conditions. The increase in tissue carbon content for all treatments and lack of change in lipid accumulation both indicate that the decrease in $\delta^{13}\text{C}$ values is not related to sugar use or lipid accumulation, but was caused by a change in carbon uptake strategy and the relative use of HCO_3^- and CO_2 . As with growth rates and net photosynthesis, mixed results were reported for $\delta^{13}\text{C}$ in the current literature (Table 2). For example, an increased growth rate was combined with decreased $\delta^{13}\text{C}$ in *Lomentaria australis* (current study), whereas growth rates and $\delta^{13}\text{C}$ simultaneously decreased in *Desmarestia aculeata* (Iñiguez et al., 2016a). This shows that it is important to measure physiological processes and parameters (e.g., net photosynthetic rates, C:N ratios, and $\delta^{13}\text{C}$) in addition to measuring the response (i.e., growth), if we are to understand how the response to CO_2 enrichment is regulated.

4.2 | The response of non-CCM species

The lack of response of *Craspedocarpus ramentaceus* to CO_2 enrichment suggests that growth and photosynthesis were saturated at

current CO_2 levels, as there was no evidence of nitrogen or light limitation in this experiment. C:N ratios were typical of N-sufficient Rhodophyta (Atkinson & Smith, 1983; Supporting information Table S4), the growth rates of the specimens in our experiment are within the healthy range of fleshy Rhodophyta (Kübler et al., 1999; Nishihara, Terada, & Noro, 2004), and the light levels were sufficient for saturating photosynthesis (Supporting information Figure S1, Table S2). Only three non-CCM species (including our work) have been studied in relation to CO_2 enrichment, with contrasting results (Table 2). This indicates that, as with CCM species, non-CCM species could have different affinities for CO_2 . For non-CCM species, the affinity for DIC will mainly depend on Rubisco activity and kinetics (as this is the enzyme involved in carbon fixation). The amount of Rubisco, the V_{max} (maximal Rubisco activity) and K_m (affinity of Rubisco for CO_2) vary greatly between species (Israel & Hophy, 2002). Red algae, for example, have a higher affinity for CO_2 relative to O_2 compared to green and brown algae, and, in general, studies show that Rubisco enzymes in CO_2 -only users have evolved a higher affinity for CO_2 than species with a CCM (Badger et al., 1998). In addition, environmental conditions play an important role in Rubisco kinetics. High temperatures, for example, decrease the affinity for CO_2 in algae (Beardall, Stojkovic, & Larsen, 2009) and the activation state of Rubisco changes with light and CO_2 concentrations, as is also known from terrestrial plants (Salvucci & Crafts-Brandner, 2004). The capacity of non-CCM species to fix carbon is often thought to be adapted to low rates of light absorption, in order to avoid photo-damage (Maberly et al., 1992), and it is possible that non-CCM species are therefore unable to take advantage of elevated CO_2 , rather than being limited in carbon. Studying how Rubisco kinetics change with CO_2 enrichment could give more insight into the response of non-CCM species. Given that 35% of red macroalgae are non-CCM (Kübler & Dudgeon, 2015), we emphasize the need to incorporate more non-CCM species in future laboratory studies if we are to determine how macroalgae will respond to future global change.

4.3 | Biochemical responses

Our study provides the first findings on the impacts of CO₂ enrichment on the lipid content, lipid class, and fatty acid composition in a CCM species compared to a non-CCM species. Both species invested in growth rather than storing lipids, which supported our hypothesis for *C. ramentaceus* (the non-CCM species), but did not support the hypothesis that lipid content and fatty acid composition would change in *L. australis* (the CCM species). There are several possible explanations for the lack of biochemical response of *L. australis*: (a) Previous work, mainly on microalgae, has shown that it is the combination of excess carbon and limiting nitrogen (i.e., insufficient nitrogen for protein synthesis) that provides optimal conditions for an increase in carbon flow toward acetyl coenzyme A (acetyl Co-A) and NADPH for FA synthesis (Sun et al., 2016). However, the nitrogen concentration used in this experiment was not likely to be limited, as the concentrations were that of ambient seawater and C:N ratios in the algal tissue were in the range consistent with nitrogen sufficiency (Atkinson & Smith, 1983). (b) Results from the lipid class composition showed that most of the lipid occurred as structural polar lipids. This indicates that both species accumulate storage lipids only in very low amounts. The ability to accumulate storage lipids might be exhibited only by certain macroalgal taxa, while others, including some red algae, generally exhibit low storage lipid concentrations and predominantly change the composition and concentration of structural polar lipids and therefore no changes were observed (Schmid, Guihéneuf, & Stengel, 2017).

4.4 | DIC limitation as a predictor of macroalgal responses to future high CO₂ oceans

The results from this study and prior studies (Table 2) highlight that—even though carbon use strategy will play a role—the responses of macroalgae to CO₂ enrichment cannot be inferred solely from the carbon uptake strategy as has been implicated in field studies (Cornwall et al., 2017, 2015; Diaz-Pulido et al., 2016; Hepburn et al., 2011). We suggest that the key to predicting the response(s) of macroalgae to a future high CO₂ ocean is to understand which species have growth rates that are limited by DIC availability under current CO₂ levels, and which species are saturated. In addition, the ability of CCM species to downregulate their CCM will be a decisive factor in their response to future CO₂ enrichment conditions. This view is summarized in Figure 5, which synthesizes the predicted responses of fleshy macroalgae based on their putative limitation by DIC (with high-affinity species being saturated and low-affinity species being limited) and carbon uptake strategy, expanding on the figure of Cornwall et al. (2017). We additionally suggest that some species may be sensitive to higher [H⁺] also predicted for the future, and a mechanistic understanding of the interactions between DIC and [H⁺] is needed (Bach et al., 2013; Roleda et al., 2012). To be able to predict the responses of macroalgae to CO₂ enrichment, future studies should focus on providing a better understanding of the physiological mechanisms that underlie DIC acquisition: Details

of physiological and biochemical regulatory mechanisms are known for only a few of the studied species listed in Table 2 (indicated with *), and these are mostly species with high commercial value or fouling species that inhibit growth of species with commercial value. This emphasizes an urgent need for targeted physiological experiments, as well as molecular studies, on a wide range of macroalgal species from the tropics to the poles, if we are to gain a mechanistic understanding of macroalgal responses to a future higher CO₂ world, and move beyond predictions based on field observational studies.

DATA ACCESSIBILITY STATEMENT

All data created during this research are available at the IMAS data management system and at FigShare (<https://doi.org/10.6084/m9.figshare.7189382> and <https://doi.org/10.6084/m9.figshare.7189292>).

ACKNOWLEDGMENTS

L.M.L. was funded by Holland Scholarship and the Groningen University Fund. M.S. was funded by the Deutsche Forschungsgemeinschaft (DFG, grant ID: SCHM 3335/1-1). We thank Dr Toby Bolton for his expert support. The authors thank the editor and two anonymous reviewers for their helpful comments on the manuscript.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTION

L.M.L. carried out laboratory experiment and analyzed the data with assistance by M.S, P.P.L., and D.B. L.M.L., M.S, P.P.L., and C.L.H. designed the study. L.M.L., M.S., and C.L.H. wrote the original draft, with subsequent contributions by all other authors. A.T.R., P.V., P.D.N., and C.M.M. collected laboratory data, contributed materials, and assisted with carbonate chemistry calculations.

ORCID

Luna M. van der Loos  <https://orcid.org/0000-0003-3686-2844>

Matthias Schmid  <https://orcid.org/0000-0002-7482-7356>

Pablo P. Leal  <https://orcid.org/0000-0002-7616-1850>

Christina M. McGraw  <https://orcid.org/0000-0001-5841-0748>

Damon Britton  <https://orcid.org/0000-0002-9029-7527>

Andrew T. Revill  <https://orcid.org/0000-0003-2486-5976>

Patti Virtue  <https://orcid.org/0000-0002-9870-1256>

Peter D. Nichols  <https://orcid.org/0000-0003-4658-4665>

Catriona L. Hurd  <https://orcid.org/0000-0001-9965-4917>

REFERENCES

- Al-Hasan, R. H., Hantash, F. M., & Radwan, S. S. (1991). Enriching marine macroalgae with eicosatetraenoic (arachidonic) and eicosapentaenoic acids by chilling. *Applied Microbiology and Biotechnology*, *35*, 530–535. <https://doi.org/10.1007/BF00169763>
- Atkinson, M. J., & Smith, S. V. (1983). C:N: P ratios of benthic marine plants. *Limnology and Oceanography*, *28*, 568–574. <https://doi.org/10.4319/lo.1983.28.3.0568>.
- Bach, L. T., Mackinder, L. C. M., Schulz, K. G., Wheeler, G., Schroeder, D. C., Brownlee, C., & Riebesell, U. (2013). Dissecting the impact of CO₂ and pH on the mechanisms of photosynthesis and calcification in the coccolithophore *Emiliania huxleyi*. *New Phytologist*, *199*, 121–134. <https://doi.org/10.1111/nph.12225>.
- Badger, M. R., Andrews, T. J., Whitney, S. M., Ludwig, M., Yellowlees, D. C., Leggat, W., & Price, G. D. (1998). The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae. *Canadian Journal of Botany*, *76*, <https://doi.org/10.1139/b98-074>.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*, 48. <https://doi.org/10.18637/jss.v067.i01>.
- Beardall, J., Stojkovic, S., & Larsen, S. (2009). Plant ecology & diversity living in a high CO₂ world: Impacts of global climate change on marine phytoplankton. *Plant Ecology & Diversity* *2*(2), 191–205. <https://doi.org/10.1080/17550870903271363>
- Bender-Champ, D., Diaz-Pulido, G., & Dove, S. (2017). Effects of elevated nutrients and CO₂ emission scenarios on three coral reef macroalgae. *Harmful Algae*, *65*, 40–51. <https://doi.org/10.1016/j.hal.2017.04.004>
- Bennett, S., Wernberg, T., Connell, S. D., Hobday, A. J., Johnson, C. R., & Poloczanska, E. S. (2015). The “Great Southern Reef”: Social, ecological and economic value of Australia’s neglected kelp forests. *Marine and Freshwater Research*, *67*, 47–56. <https://doi.org/10.1071/MF15232>
- Bermúdez, R., Feng, Y., Roleda, M. Y., Tatters, A. O., Hutchins, D. A., Larsen, T., ... Winder, M. (2015). Long-term conditioning to elevated pCO₂ and warming influences the fatty and amino acid composition of the diatom *Cylindrotheca fusiformis*. *PLoS ONE*, *10*, 1–15. <https://doi.org/10.1371/journal.pone.0123945>.
- Björk, M., Haglund, K., Ramazanov, Z., & Pedersén, M. (1993). Inducible mechanisms for HCO₃⁻ utilization and repression of photorespiration in protoplast and thalli of three species of *Ulva* (Chlorophyta). *Journal of Phycology*, *29*, 166–173.
- Bligh, E., & Dyer, W. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry*, *37*, 911–917. <https://doi.org/10.1139/y59-099>
- Bockmon, E. E., Frieder, C. A., Navarro, M. O., White-Kershek, L. A., & Dickson, A. G. (2013). Technical Note: Controlled experimental aquarium system for multi-stressor investigation of carbonate chemistry, oxygen saturation, and temperature. *Biogeosciences*, *10*, 5967–5975. <https://doi.org/10.5194/bg-10-5967-2013>.
- Britton, D., Cornwall, C. E., Revill, A. T., Hurd, C. L., & Johnson, C. R. (2016). Ocean acidification reverses the positive effects of seawater pH fluctuations on growth and photosynthesis of the habitat-forming kelp. *Ecklonia radiata*. *Scientific Reports*, *6*, 26036. <https://doi.org/10.1038/srep26036>.
- Chen, B., Zou, D., Zhu, M., & Yang, Y. (2017). Effects of CO₂ levels and light intensities on growth and amino acid contents in red seaweed *Gracilaria lemaneiformis*. *Aquaculture Research*, *48*, 2683–2690. <https://doi.org/10.1111/are.13100>.
- Chung, I. K., Beardall, J., Mehta, S., Sahoo, D., & Stojkovic, S. (2011). Using marine macroalgae for carbon sequestration: A critical appraisal. *Journal of Applied Phycology*, *23*, 877–886. <https://doi.org/10.1007/s10811-010-9604-9>.
- Cornwall, C. E., Revill, A. T., Hall-Spencer, J. M., Milazzo, M., Raven, J. A., & Hurd, C. L. (2017). Inorganic carbon physiology underpins macroalgal responses to elevated CO₂. *Scientific Reports*, *7*, <https://doi.org/10.1038/srep46297>.
- Cornwall, C. E., Revill, A. T., & Hurd, C. L. (2015). High prevalence of diffusive uptake of CO₂ by macroalgae in a temperate subtidal ecosystem. *Photosynthesis Research*, *124*, 181–190. <https://doi.org/10.1007/s11120-015-0114-0>.
- de Paula Silva, P. H., Paul, N. A., de Nys, R., & Mata, L. (2013). Enhanced production of green tide algal biomass through additional carbon supply. *PLoS ONE*, *8*, e81164. <https://doi.org/10.1371/journal.pone.0081164>.
- Diaz-Pulido, G., Cornwall, C., Gartrell, P., Hurd, C., & Tran, D. V. (2016). Strategies of dissolved inorganic carbon use in macroalgae across a gradient of terrestrial influence: Implications for the Great Barrier Reef in the context of ocean acidification. *Coral Reefs*, *35*, 1327–1341. <https://doi.org/10.1007/s00338-016-1481-5>.
- Dickson, A. G., & Millero, F. J. (1987). A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research Part A, Oceanographic Research Papers*, *34*, 1733–1743. [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5).
- Dickson, A. G., Sabine, C. L., & Christian, J. R. 2007. Guide to best practices for ocean CO₂ measurements. PICES Special Publication 3.
- Fernández, P. A., Roleda, M. Y., & Hurd, C. L. (2015). Effects of ocean acidification on the photosynthetic performance, carbonic anhydrase activity and growth of the giant kelp *Macrocystis pyrifera*. *Photosynthesis Research*, *124*, 293–304. <https://doi.org/10.1007/s11120-015-0138-5>.
- Gao, K., Aruga, Y., Asada, K., Ishihara, T., Akano, T., & Kiyohara, M. (1991). Enhanced growth of the red alga *Porphyra yezoensis* Ueda in high CO₂ concentrations. *Journal of Applied Phycology*, *3*, 355–362. <https://doi.org/10.1007/BF00026098>.
- Gao, K., Aruga, Y., Asada, K., & Kiyohara, M. (1993). Influence of enhanced CO₂ on growth and photosynthesis of the red algae *Gracilaria* sp. and *G. chilensis*. *Journal of Applied Phycology*, *5*, 563–571. <https://doi.org/10.1007/BF02184635>
- Gao, G., Beardall, J., Bao, M., Wang, C., Ren, W., & Xu, J. (2018). Ocean acidification and nutrient limitation synergistically reduce growth and photosynthetic performances of a green tide alga *Ulva linza*. *Biogeosciences*, *15*, 3409–3420. <https://doi.org/10.5194/bg-15-3409-2018>.
- García-Sánchez, M. J., Fernández, J. A., & Niell, X. (1994). Effect of inorganic carbon supply on the photosynthetic physiology of *Gracilaria tenuistipitata*. *Planta*, *194*, 55–61. <https://doi.org/10.1007/BF00201034>.
- Gordillo, F. J. L., Aguilera, J., Wiencke, C., & Jiménez, C. (2015). Ocean acidification modulates the response of two Arctic kelps to ultraviolet radiation. *Journal of Plant Physiology*, *173*, 41–50. <https://doi.org/10.1016/j.jplph.2014.09.008>.
- Gordillo, F. J. L., Carmona, R., Viñegla, B., Wiencke, C., & Jiménez, C. (2016). Effects of simultaneous increase in temperature and ocean acidification on biochemical composition and photosynthetic performance of common macroalgae from Kongsfjorden (Svalbard). *Polar Biology*, *39*, 1993–2007. <https://doi.org/10.1007/s00300-016-1897-y>.
- Gordillo, F. J. L., Figueroa, F. L., & Niell, F. X. (2003). Photon- and carbon-use efficiency in *Ulva rigida* at different CO₂ and N levels. *Planta*, *218*, 315–322. <https://doi.org/10.1007/s00425-003-1087-3>.
- Gordillo, F. J. L., Niell, F. X., & Figueroa, F. L. (2001). Non-photosynthetic enhancement of growth by high CO₂ level in the nitrophilic seaweed *Ulva rigida* C. Agardh (Chlorophyta). *Biomedical and Life Sciences*, *213*, 64–70. <https://doi.org/10.1007/s004250000468>.
- Gosch, B. J., Lawton, R. J., Paul, N. A., Nys, R. D., & Magnusson, M. (2015). Environmental effects on growth and fatty acids in three isolates of *Derbesia tenuissima* (Bryopsidales, Chlorophyta). *Algal Research*, *9*, 82–93. <https://doi.org/10.1016/j.algal.2015.02.022>

- Gutow, L., Rahman, M. M., Bartl, K., Saborowski, R., Bartsch, I., & Wiencke, C. (2014). Ocean acidification affects growth but not nutritional quality of the seaweed *Fucus vesiculosus* (Phaeophyceae, Fucales). *PANGAEA*. <https://doi.org/10.1594/PANGAEA.835334>
- Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., ... Buia, M.-C. (2008). Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*, *454*, 96–99. <https://doi.org/10.1038/nature07051>.
- Hepburn, C. D., Pritchard, D. W., Cornwall, C. E., Mcleod, R. J., Beardall, J., Raven, J. A., & Hurd, C. L. (2011). Diversity of carbon use strategies in a kelp forest community: Implications for a high CO₂ ocean. *Global Change Biology*, *17*, 2488–2497. <https://doi.org/10.1111/j.1365-2486.2011.02411.x>.
- Hibberd, J. M., & Covshoff, S. (2010). The regulation of gene expression required for C₄ photosynthesis. *Annual Review of Plant Biology*, *61*, 181–207. <https://doi.org/10.1146/annurev-arplant-042809-112238>.
- Ho, M., & Carpenter, R. C. (2017). Differential growth responses to water flow and reduced pH in tropical marine macroalgae. *Journal of Experimental Marine Biology and Ecology*, *491*, 58–65. <https://doi.org/10.1016/j.jembe.2017.03.009>.
- Hurd, C. L., Hepburn, C. D., Currie, K. I., Raven, J. A., & Hunter, K. A. (2009). Testing the effects of ocean acidification on algal metabolism: Considerations for experimental designs. *Journal of Phycology*, *45*, 1236–1251. <https://doi.org/10.1111/j.1529-8817.2009.00768.x>.
- Hurd, C. L., Lenton, A., Tilbrook, B., & Boyd, P. W. (2018). Current understanding and challenges for oceans in a higher-CO₂ world. *Nature Climate Change*, *8*, 686–694. <https://doi.org/10.1038/s41558-018-0211-0>.
- Iñiguez, C., Carmona, R., Lorenzo, M. R., Niell, F. X., Wiencke, C., & Gordillo, F. J. L. (2016a). Increased CO₂ modifies the carbon balance and the photosynthetic yield of two common Arctic brown seaweeds: *Desmarestia aculeata* and *Alaria esculenta*. *Polar Biology*, *39*, 1979–1991. <https://doi.org/10.1007/s00300-015-1724-x>.
- Iñiguez, C., Carmona, R., Lorenzo, M. R., Niell, F. X., Wiencke, C., & Gordillo, F. J. L. (2016b). Increased temperature, rather than elevated CO₂, modulates the carbon assimilation of the Arctic kelps *Saccharina latissima* and *Laminaria solidungula*. *Marine Biology*, *163*. <https://doi.org/10.1007/s00227-016-3024-6>.
- Iñiguez, C., Heinrich, S., Harms, L., & Gordillo, F. J. L. (2017). Increased temperature and CO₂ alleviate photoinhibition in *Desmarestia anceps*: From transcriptomics to carbon utilization. *Journal of Experimental Botany*, *68*, 3971–3984. <https://doi.org/10.1093/jxb/erx164>.
- IPCC. (2013). Climate Change 2013: the physical science basis. In: *Contribution of Working Group I to the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change* (pp. 1029–1136).
- Israel, A., & Hophy, M. (2002). Growth, photosynthetic properties and Rubisco activities and amounts of marine macroalgae grown under current and elevated seawater CO₂ concentrations. *Global Change Biology*, *8*, 831–840. <https://doi.org/10.1046/j.1365-2486.2002.00518.x>.
- Israel, A., Katz, S., Dubinsky, Z., Merrill, J. E., & Friedlander, M. (1999). Photosynthetic inorganic carbon utilization and growth of *Porphyra linearis* (Rhodophyta). *Journal of Applied Phycology*, *11*, 447–453. <https://doi.org/10.1023/A:1008122306268>.
- Jiang, H., Zou, D., Lou, W., & Gong, J. (2018). Effects of CO₂ supply on growth and photosynthetic ability of young sporophytes of the economic seaweed *Sargassum fusiforme* (Sargassaceae, Phaeophyta). *Journal of Applied Phycology*. <https://doi.org/10.1007/s10811-018-1569-0>.
- Johnson, M. D., Comeau, S., Lantz, C. A., & Smith, J. E. (2017). Complex and interactive effects of ocean acidification and temperature on epilithic and endolithic coral-reef turf algal assemblages. *Coral Reefs*, *36*, 1059–1070. <https://doi.org/10.1007/s00338-017-1597-2>.
- Jokiel, P. L., Rodgers, K. S., Kuffner, I. B., Andersson, A. J., Cox, E. F., & Mackenzie, F. T. (2008). Ocean acidification and calcifying reef organisms: A mesocosm investigation. *Coral Reefs*, *27*, 473–483. <https://doi.org/10.1007/s00338-008-0380-9>.
- Jungnick, N., Ma, Y., Mukherjee, B., Cronan, J. C., Speed, D. J., Laborde, S. M., ... Moroney, J. V. (2014). The carbon concentrating mechanism in *Chlamydomonas reinhardtii*: Finding the missing pieces. *Photosynthesis Research*, *121*, 159–173. <https://doi.org/10.1007/s11120-014-0004-x>.
- Kang, J. W., & Chung, I. K. (2017). The effects of eutrophication and acidification on the ecophysiology of *Ulva pertusa* Kjellman. *Journal of Applied Phycology*, *29*, 2675–2683. <https://doi.org/10.1007/s10811-017-1087-5>.
- Kang, J. W., & Chung, I. K. (2018). The interactive effects of elevated CO₂ and ammonium enrichment on the physiological performances of *Saccharina japonica* (Laminariales, Phaeophyta). *Ocean Science Journal*, *53*(3), 487–497. <https://doi.org/10.1007/s12601-018-0014-2>.
- Kang, J. W., Kambey, C., Shen, Z., Yang, Y., & Chung, I. K. (2017). The short-term effects of elevated CO₂ and ammonium concentrations on physiological responses in *Gracilariopsis lemaneiformis* (Rhodophyta). *Fisheries and Aquatic Sciences*, *20*. <https://doi.org/10.1186/s41240-017-0063-y>.
- Kang, E. J., & Kim, K. Y. (2016). Effects of future climate conditions on photosynthesis and biochemical component of *Ulva pertusa* (Chlorophyta). *Algae*, *31*, 49–59. <https://doi.org/10.4490/algae.2016.31.3.9>.
- Kawamitsu, Y., & Boyer, J. S. (1999). Photosynthesis and carbon storage between tides in a brown alga, *Fucus vesiculosus*. *Marine Biology*, *133*, 361–369. <https://doi.org/10.1007/s002270050475>.
- Kim, J. H., Kang, E. J., Edwards, M. S., Lee, K., Jeong, H. J., & Kim, K. Y. (2016). Species-specific responses of temperate macroalgae with different photosynthetic strategies to ocean acidification: A mesocosm study. *Algae*, *31*, 243–256. <https://doi.org/10.4490/algae.2016.31.8.20>.
- Koch, M., Bowes, G., Ross, C., & Zhang, X. H. (2013). Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Global Change Biology*, *19*, 103–132. <https://doi.org/10.1111/j.1365-2486.2012.02791.x>.
- Kram, S. L., Price, N. N., Donham, E. M., Johnson, M. D., Kelly, E. L. A., Hamilton, S. L., & Smith, J. E. (2016). Variable responses of temperate calcified and fleshy macroalgae to elevated pCO₂ and warming. *ICES Journal of Marine Science: Journal Du Conseil*, *73*, 693–703. <https://doi.org/10.1093/icesjms/fsv168>.
- Kübler, J. E., & Dudgeon, S. R. (2015). Predicting effects of ocean acidification and warming on algae lacking carbon concentrating mechanisms. *PLoS ONE*, *10*, 1–19. <https://doi.org/10.1371/journal.pone.0132806>.
- Kübler, J. E., Johnston, A. M., & Raven, J. A. (1999). The effects of reduced and elevated CO₂ and O₂ on the seaweed *Lomentaria articulata*. *Plant, Cell and Environment*, *22*, 1303–1310. <https://doi.org/10.1046/j.1365-3040.1999.00492.x>.
- Li, X., Xu, J., & He, P. (2016). Comparative research on inorganic carbon acquisition by the macroalgae *Ulva prolifera* (Chlorophyta) and *Pyropia yezoensis* (Rhodophyta). *Journal of Applied Phycology*, *28*, 491–497. <https://doi.org/10.1007/s10811-015-0603-8>.
- Li, Y., Zhong, J., Zheng, M., Zhuo, P., & Xu, N. (2018). Photoperiod mediates the effects of elevated CO₂ on the growth and physiological performance in the green tide alga *Ulva prolifera*. *Marine Environmental Research*. <https://doi.org/10.1016/j.marenvres.2018.07.015>.
- Lignell, Å., & Pedersén, M. (1989). Effects of pH and inorganic carbon concentration on growth of *Gracilaria secundata*. *British Phycological Journal*, *24*, 83–89. <https://doi.org/10.1080/00071618900650071>.
- Liu, C., & Zou, D. (2015a). Do increased temperature and CO₂ levels affect the growth, photosynthesis, and respiration of the marine macroalga *Pyropia haitanensis* (Rhodophyta)? An experimental study. *Hydrobiologia*, *745*, 285–296. <https://doi.org/10.1007/s10750-014-2113-0>.

- Liu, C., & Zou, D. (2015b). Responses of elevated CO₂ on photosynthesis and nitrogen metabolism in *Ulva lactuca* (Chlorophyta) at different temperature levels. *Marine Biology Research*, *11*, 1043–1052. <https://doi.org/10.1080/17451000.2015.1062520>
- Liu, C., Zou, D., & Yang, Y. (2018). Comparative physiological behaviors of *Ulva lactuca* and *Gracilariopsis lemaneiformis* in responses to elevated atmospheric CO₂ and temperature. *Environmental Science and Pollution Research*, *25*, 27493–27502. <https://doi.org/10.1007/s11356-018-2792-6>
- Maberly, S. C., Raven, J. A., & Johnston, A. M. (1992). Discrimination between ¹²C and ¹³C by marine plants. *Oecologia*, *91*, 481–492. <https://doi.org/10.1007/BF00650320>
- McGraw, C. M., Cornwall, C. E., Reid, M. R., Currie, K. I., Hepburn, C. D., Boyd, P. W., ... Hunter, K. A. (2010). An automated pH-controlled culture system for laboratory-based ocean acidification experiments. *Limnology and Oceanography: Methods*, *8*, 686–694. <https://doi.org/10.4319/lom.2010.8.686>
- Mehrbach, C., Culbertson, C. H., Hawley, J. E., & Pytkowicz, R. M. (1973). Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*, *18*, 897–907.
- Mensch, B., Neulinger, S. C., Graiff, A., Pansch, A., Künzel, S., Fischer, M. A., & Schmitz, R. A. (2016). Restructuring of epibacterial communities on *Fucus vesiculosus* forma *mytili* in response to elevated pCO and increased temperature levels. *Frontiers in Microbiology*, *7*, 1–15. <https://doi.org/10.3389/fmicb.2016.00434>
- Mercado, J. M., Javier, F., Gordillo, L., Xavier Niell, F., & Figueroa, F. L. (1999). Effects of different levels of CO₂ on photosynthesis and cell components of the red alga *Porphyra leucosticta*. *Journal of Applied Phycology*, *11*, 455–461.
- Nishihara, G. N., Terada, R., & Noro, T. (2004). Photosynthesis and growth rates of *Laurencia brongniartii* J. Agardh (Rhodophyta, Ceramiales) in preparation for cultivation. *Journal of Applied Phycology*, *16*, 303–308. <https://doi.org/10.1023/B>
- Nunes, J., McCoy, S. J., Findlay, H. S., Hopkins, F. E., Kitidis, V., Queiros, A. M., ... Widdicombe, S. (2016). Two intertidal, non-calcifying macroalgae (*Palmaria palmata* and *Saccharina latissima*) show complex and variable responses to short-term CO₂ acidification. *ICES Journal of Marine Science*, *73*, 887–896. <https://doi.org/10.1093/icesjms/fsv081>
- Ober, G. T., & Thornber, C. S. (2017). Divergent responses in growth and nutritional quality of coastal macroalgae to the combination of increased pCO₂ and nutrients. *Marine Environmental Research*, *131*, 69–79. <https://doi.org/10.1016/j.marenvres.2017.09.003>
- Olischläger, M., Iñiguez, C., Koch, K., Wiencke, C., & Gordillo, F. J. L. (2017). Increased pCO₂ and temperature reveal ecotypic differences in growth and photosynthetic performance of temperate and Arctic populations of *Saccharina latissima*. *Planta*, *245*, 119–136. <https://doi.org/10.1007/s00425-016-2594-3>
- Olischläger, M., & Wiencke, C. (2013). Ocean acidification alleviates low-temperature effects on growth and photosynthesis of the red alga *Neosiphonia harveyi* (Rhodophyta). *Journal of Experimental Botany*, *64*, 5587–5597. <https://doi.org/10.1093/jxb/ert329>
- Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ... Wagner, H. (2016). Vegan: Community ecology package. R package version 2.3-4. Retrieved from <http://CRAN.R-project.org/package=vegan>
- Rautenberger, R., Fernández, P. A., Strittmatter, M., Heesch, S., Cornwall, C. E., Hurd, C. L., & Roleda, M. Y. (2015). Saturating light and not increased carbon dioxide under ocean acidification drives photosynthesis and growth in *Ulva rigida* (Chlorophyta). *Ecology and Evolution*, *5*, 874–888. <https://doi.org/10.1002/ece3.1382>
- Raven, J. A., Ball, L. A., Beardall, J., Giordano, M., & Maberly, S. C. (2005). Algae lacking carbon-concentrating mechanisms. *Canadian Journal of Botany*, *83*, 879–890. <https://doi.org/10.1139/b05-074>
- Raven, J. A., Johnston, A. M., Kübler, J. E., Korb, R., McInroy, S. G., Handley, L. L., ... Walker, D. I. (2002). Mechanistic interpretation of carbon isotope discrimination by marine macroalgae and seagrasses. *Functional Plant Biol.*, *29*, 335–378. <https://doi.org/10.1071/PP01201>
- Reidenbach, L. B., Fernandez, P. A., Leal, P. P., Noiset, F., McGraw, C. M., Revill, A. T., ... Kübler, J. E. (2017). Growth, ammonium metabolism, and photosynthetic properties of *Ulva australis* (Chlorophyta) under decreasing pH and ammonium enrichment. *PLoS ONE*, *12*, 1–20. <https://doi.org/10.1371/journal.pone.0188389>
- Rivers, J. S., & Peckol, P. (1995). Interactive effects of nitrogen and dissolved inorganic carbon on photosynthesis, growth, and ammonium uptake of the macroalgae *Cladophora vagabunda* and *Gracilaria tikvahiae*. *Marine Biology*, *121*, 747–753. <https://doi.org/10.1007/BF00349311>
- Roleda, M. Y., Morris, J. N., McGraw, C. M., & Hurd, C. L. (2012). Ocean acidification and seaweed reproduction: Increased CO₂ ameliorates the negative effect of lowered pH on meiospore germination in the giant kelp *Macrocystis pyrifera* (Laminariales, Phaeophyceae). *Global Change Biology*, *18*, 854–864. <https://doi.org/10.1111/j.1365-2486.2011.02594.x>
- Salvucci, M. E., & Crafts-Brandner, S. J. (2004). Inhibition of photosynthesis by heat stress: The activation state of Rubisco as a limiting factor in photosynthesis. *Physiologia Plantarum*, *120*, 179–186. <https://doi.org/10.1111/j.0031-9317.2004.0173.x>
- Sampath-Wiley, P., & Neefus, C. D. (2007). An improved method for estimating R-phycoerythrin and R-phycoerythrin contents from crude aqueous extracts of *Porphyra* (Bangiales, Rhodophyta). *Journal of Applied Phycology*, *19*, 123–129. <https://doi.org/10.1007/s10811-006-9118-7>
- Sandrini, G., Matthijs, H. C. P., Verspagen, J. M. H., Muijzer, G., & Huisman, J. (2014). Genetic diversity of inorganic carbon uptake systems causes variation in CO₂ response of the cyanobacterium *Microcystis*. *The ISME Journal*, *8*, 589–600. <https://doi.org/10.1038/ismej.2013.179>
- Sarker, M. Y., Bartsch, I., Olischläger, M., Gutow, L., & Wiencke, C. (2013). Combined effects of CO₂, temperature, irradiance and time on the physiological performance of *Chondrus crispus* (Rhodophyta). *Botanica Marina*, *56*, 63–74. <https://doi.org/10.1515/bot-2012-0143>
- Schmid, M., Guihéneuf, F., & Stengel, D. B. (2017). Plasticity and remodelling of lipids support acclimation potential in two species of low-intertidal macroalgae, *Fucus serratus* (Phaeophyceae) and *Palmaria palmata* (Rhodophyta). *Algal Research*, *26*, 104–114. <https://doi.org/10.1016/j.algal.2017.07.004>
- Schmidt, E. C., Maraschin, M., & Bouzon, Z. L. (2010). Effects of UVB radiation on the carragenophyte *Kappaphycus alvarezii* (Rhodophyta, Gigartinales): Changes in ultrastructure, growth, and photosynthetic pigments. *Hydrobiologia*, *649*, 171–182. <https://doi.org/10.1007/s10750-010-0243-6>
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*, 671–675. <https://doi.org/10.1038/nmeth.2089>
- Sebök, S., Herppich, W. B., & Hanelt, D. (2017). Development of an innovative ring-shaped cultivation system for a land-based cultivation of marine macroalgae. *Aquacultural Engineering*, *77*, 33–41. <https://doi.org/10.1016/j.aquaeng.2017.01.005>
- Seely, G. R., Duncan, M. J., & Vidaver, W. E. (1972). Preparative and analytical extraction of pigments from brown algae with dimethyl sulfoxide. *Marine Biology*, *12*, 184–188. <https://doi.org/10.1007/BF00350754>
- Suárez-Álvarez, S., Gómez-Pinchetti, J. L., & García-Reina, G. (2012). Effects of increased CO₂ levels on growth, photosynthesis, ammonium uptake and cell composition in the macroalga *Hypnea spinella*

- (Gigartinales, Rhodophyta). *Journal of Applied Phycology*, 24, 815–823. <https://doi.org/10.1007/s10811-011-9700-5>.
- Sun, Z., Chen, Y. F., & Du, J. (2016). Elevated CO₂ improves lipid accumulation by increasing carbon metabolism in *Chlorella sorokiniana*. *Plant Biotechnology Journal*, 14, 557–566. <https://doi.org/10.1111/pbi.12398>.
- Sunday, J. M., Fabricius, K. E., Kroeker, K. J., Anderson, K. M., Brown, N. E., Barry, J. P., ... Harley, C. D. G. (2016). Ocean acidification can mediate biodiversity shifts by changing biogenic habitat. *Nature Climate Change*, 1, 1–6. <https://doi.org/10.1038/nclimate3161>.
- Swanson, A. K., & Fox, C. H. (2007). Altered kelp (Laminariales) phlorotannins and growth under elevated carbon dioxide and ultraviolet-B treatments can influence associated intertidal food webs. *Global Change Biology*, 13, 1696–1709. <https://doi.org/10.1111/j.1365-2486.2007.01384.x>
- The Royal Society (2005). Ocean acidification due to increasing atmospheric carbon dioxide. The Royal Society Policy document 12/05.
- Thom, R. M. (1996). CO₂-Enrichment effects on eelgrass (*Zostera marina* L.) and bull kelp (*Nereocystis luetkeana* (mert.) P & R.). *Water, Air, and Soil Pollution*, 88, 383–391. <https://doi.org/10.1007/BF00294113>.
- Tuya, F., Wernberg, T., & Thomsen, M. S. (2008). The spatial arrangement of reefs alters the ecological patterns of fauna between interspersed algal habitats. *Estuarine, Coastal and Shelf Science*, 78, 774–782. <https://doi.org/10.1016/j.ecss.2008.02.017>.
- Wickham, H. (2007). Reshaping data with the reshape package. *Journal of Statistical Software*, 21, 1–20.
- Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis* (pp. 1–213). New York, NY: Springer-Verlag.
- Wickham, H. (2011). The split-apply-combine strategy for data analysis. *Journal of Statistical Software*, 40, 1–29. <https://doi.org/10.1039/np9971400083>
- Xu, K., Chen, H., Wang, W., Xu, Y., Ji, D., Chen, C., & Xie, C. (2017). Responses of photosynthesis and CO₂ concentrating mechanisms of marine crop *Pyropia haitanensis thalli* to large pH variations at different time scales. *Algal Research*, 28, 200–210. <https://doi.org/10.1016/j.algal.2017.10.023>.
- Xu, J., & Gao, K. (2012). Future CO₂-induced ocean acidification mediates the physiological performance of a green tide alga. *Plant Physiology*, 160, 1762–1769. <https://doi.org/10.1104/pp.112.206961>
- Xu, Z., Gao, G., Xu, J., & Wu, H. (2017). Physiological response of a golden tide alga (*Sargassum muticum*) to the interaction of ocean acidification and phosphorus enrichment. *Biogeosciences*, 14, 671–681. <https://doi.org/10.5194/bg-14-671-2017>.
- Xu, Z., Zou, D., & Gao, K. (2010). Effects of elevated CO₂ and phosphorus supply on growth, photosynthesis and nutrient uptake in the marine macroalga *Gracilaria lemaneiformis* (Rhodophyta). *Botanica Marina*, 53, 123–129. <https://doi.org/10.1515/BOT.2010.012>
- Yong, Y. S., Yong, W. T. L., & Anton, A. (2013). Analysis of formulae for determination of seaweed growth rate. *Journal of Applied Phycology*, 25, 1831–1834. <https://doi.org/10.1007/s10811-013-0022-7>
- Zhu, Z., Piao, S., Myneni, R. B., Huang, M., Zeng, Z., Canadell, J. G., ... Zeng, N. (2016). Greening of the Earth and its drivers. *Nature Climate Change*, 6, 791–795. <https://doi.org/10.1038/nclimate3004>
- Zou, D. (2005). Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). *Aquaculture*, 250, 726–735. <https://doi.org/10.1016/j.aquaculture.2005.05.014>
- Zou, D., & Gao, K. (2009). Effects of elevated CO₂ on the red seaweed *Gracilaria lemaneiformis* (Gigartinales, Rhodophyta) grown at different irradiance levels. *Phycologia*, 48, 510–517. <https://doi.org/10.2216/08-99.1>
- Zou, D., Gao, K., & Luo, H. (2011). Short- and long-term effects of elevated CO₂ on photosynthesis and respiration in the marine macroalga *Hizikia fusiformis* (Sargassaceae, Phaeophyta) grown at low and high N supplies. *Journal of Phycology*, 47, 87–97. <https://doi.org/10.1111/j.1529-8817.2010.00929.x>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: van der Loos LM, Schmid M, Leal PP, et al. Responses of macroalgae to CO₂ enrichment cannot be inferred solely from their inorganic carbon uptake strategy. *Ecol Evol*. 2019;9:125–140. <https://doi.org/10.1002/ece3.4679>