

RESEARCH ARTICLE

RNaseq Transcriptional Profiling following Whip Development in Sugarcane Smut Disease

Patricia D. C. Schaker¹✉, Alessandra C. Palhares¹✉, Lucas M. Taniguti¹, Leila P. Peters¹, Silvana Creste², Karen S. Aitken³, Marie-Anne Van Sluys⁴, João P. Kitajima⁵, Maria L. C. Vieira¹, Claudia B. Monteiro-Vitorello¹*

1 Departamento de Genética, Universidade de São Paulo, Escola Superior de Agricultura “Luiz de Queiroz”, Piracicaba, São Paulo, Brazil, **2** Instituto Agronômico de Campinas, Centro de Cana, Ribeirão Preto, São Paulo, Brazil, **3** CSIRO Agriculture, Queensland Bioscience Precinct, St Lucia, Queensland, Australia, **4** Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil, **5** Mendelics Análise Genômica, São Paulo, São Paulo, Brazil

✉ These authors contributed equally to this work.

* cbrmontei@usp.br



OPEN ACCESS

Citation: Schaker PDC, Palhares AC, Taniguti LM, Peters LP, Creste S, Aitken KS, et al. (2016) RNaseq Transcriptional Profiling following Whip Development in Sugarcane Smut Disease. PLoS ONE 11(9): e0162237. doi:10.1371/journal.pone.0162237

Editor: Jason E Stajich, University of California Riverside, UNITED STATES

Received: April 20, 2016

Accepted: August 21, 2016

Published: September 1, 2016

Copyright: © 2016 Schaker et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The sequencing data has been deposited at DDBJ/EMBL/GenBank under the BioProject ID PRJNA291816. All other relevant data are within the paper and its Supporting Information files.

Funding: The authors acknowledge the support of the Brazilian institution FAPESP (grant numbers 2010/05591-9; 2015/07112-4 [C.B.M.-V.] and fellowships to A.C.P. (2012/09524-0); P.D.C.S (2013/25599-2); L.M.T (2014/17034-8); L.P.P (2013/15014-7). We thank the sequencing facility Center of Functional Genomics (USP, Campus “Luiz de Queiroz”); NAP/MEPA (Núcleo de Apoio à Pesquisa

Abstract

Sugarcane smut disease is caused by the biotrophic fungus *Sporisorium scitamineum*. The disease is characterized by the development of a whip-like structure from the primary meristems, where billions of teliospores are produced. Sugarcane smut also causes tillering and low sucrose and high fiber contents, reducing cane productivity. We investigated the biological events contributing to disease symptoms in a smut intermediate-resistant sugarcane genotype by examining the transcriptional profiles (RNaseq) shortly after inoculating the plants and immediately after whip emission. The overall picture of disease progression suggests that premature transcriptional reprogramming of the shoot meristem functions continues until the emergence of the whip. The guidance of this altered pattern is potentially primarily related to auxin mobilization in addition to the involvement of other hormonal imbalances. The consequences associated with whip emission are the modulation of typical meristematic functions toward reproductive organ differentiation, requiring strong changes in carbon partitioning and energy production. These changes include the overexpression of genes coding for invertases and trehalose-6P synthase, as well as other enzymes from key metabolic pathways, such as from lignin biosynthesis. This is the first report describing changes in the transcriptional profiles following whip development, providing a hypothetical model and candidate genes to further study sugarcane smut disease progression.

Introduction

Sugarcane (*Saccharum* spp.) is the fifth most important crop in the world [1]. In addition to being a source of sugar for food, the crop has the potential to generate clean and renewable

em Microscopia Eletrônica Aplicada a Agricultura) and Laboratórios de Genética Molecular, Departamento de Fitopatologia e Nematologia, and Plantas Hortícolas, Departamento de Produção Vegetal, ESALQ. Mendelics Análise Genômica provided support in the form of a salary for author JPK, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of JPK are articulated in the 'author contributions' section.

Competing Interests: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. Mendelics Análise Genômica provided support only in the form of a salary for author JPK. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

products, such as biofuels, bioplastics, bio-hydrocarbons, and bioelectricity. Due to its agronomic attributes, such as its high yield and ability to survive under adverse conditions [2], sugarcane is found in most of the 90 tropical and subtropical countries [1]. Nonetheless, the crop hosts several pathogens, including the fungus *Sporisorium scitamineum*, the causal agent of sugarcane smut disease [3] (Fig 1). Sugarcane smut is mainly characterized by the development of a long whip-like structure consisting of plant and fungal tissues in which billions of teliospores are produced. The name 'smut' derives from the black powdery mass of teliospores released by these structures that resemble soot. The whips originate in the primary meristems of the apex and lateral buds of infected stalks, and they are initially covered with a thin silvery membranous sheath [3], which detaches after the teliospores mature and are ready to disperse. In more susceptible varieties, whips can be detected as early as 2 to 4 months of age, with peak whip growth occurring in the 6th or 7th month [4]. Smut is mainly transmitted by wind-borne teliospores infecting standing canes but also by teliospores in the soil that infect planted setts. The germination of the teliospores leads to meiosis, which produces haploid sporidia. Mating-compatible sporidial cells produce infective hyphae through hyphal anastomosis, which initiates plant colonization [5].

The disease limits the crop yield and properties of sugarcane products, causing losses in cane tonnage and juice quality. Other disease symptoms include tillering and low sucrose and increased fiber contents [3]. Like most agronomic traits, smut resistance is a quantitative characteristic [6] that is difficult to genetically and functionally characterize. Moreover, modern varieties of sugarcane ($2n = 100-130$) have a complex genomic structure that derives from a highly polyploid and aneuploid interspecific hybridization [7, 8], hindering the understanding of the quantitative traits and mapping of their loci [9, 10]. Efforts to elucidate the molecular basis of sugarcane smut resistance have been made since James [11] proposed the existence of a chemical resistance mechanism. Lloyd and Pillay [12] identified some flavonoids, which are teliospore-germination inhibitors, and subsequently, a correlation between the resistance rating and concentration of glycosidic substances was established [13]. Later studies reported changes in the patterns of free polyamines and their conjugation in both susceptible and resistant sugarcane varieties infected by *S. scitamineum* [4, 14]. Changes in the sugarcane gene expression profile induced by the fungus have been identified by several authors using techniques such as suppression-subtractive hybridization-based sequencing and differential display of complementary DNA-amplified fragment-length polymorphisms [15–18].

Despite these attainments, more detailed studies are needed to precisely define the changes in the entire sugarcane gene repertoire when challenged with the pathogen, both at different




Fig 1. Sugarcane smut. (A) Scan electron microscopy of *S. scitamineum* hyphal growth in the sugarcane bud at 5 DAI (1, 2). Fungal sporogenesis and teliospore maturation in the base of the sugarcane whip at 200 DAI (3; 4). Bar = 10 μ m. (B) (1) Sugarcane bud at 5 DAI. (2) Base of the whip at 200 DAI. Bar = 1 cm.

doi:10.1371/journal.pone.0162237.g001

stages of fungal development and in different host tissues. Messenger RNA sequencing (RNA-seq) technology has the potential to explore the complete set of gene expression programs to a high level of accuracy and depth, providing further insights into plant-pathogen interactions [19]. This method has been applied to several mixed-model systems of plant-fungus interactions [18,20–24], and more recently, to elucidate the early stages of the sugarcane-smut pathosystem [25–27]. Continuing the study of this pathosystem, we used RNAseq technology to perform a comparative analysis of infected sugarcane tissues of a smut intermediate-resistant genotype at two time points: shortly after inoculation and later, when the whips appeared and disease symptoms were evident (Fig 1). In addition to confirming the existing data [25], this work addresses the molecular events following whip emission. The most relevant conclusions are: 1) the association with transcriptional reprogramming of shoot apical functions probably by restraining floral development; 2) the transcriptional changes in carbon partitioning, mostly pronounced in hexoses and lignin; and 3) the relatedness of auxin to whip emission as well as the response associated with oxidative stress.

Material and Methods

Ethics Statement

S. scitamineum SSC39 teliospores were collected from experimental network areas of IAC sugarcane breeding program (Instituto Agronômico, Centro de Cana, Ribeirão Preto, São Paulo, Brazil), as described by Taniguti et al. [26]. The healthy buds used to conduct the experiments were obtained from IAC sugarcane nursery. No special permits were necessary for teliospores and genotype used, because this project was developed in collaboration with IAC researchers. This work does not involve endangered or protected species.

Experimental Design

S. scitamineum SSC39teliospores were checked for viability and were inoculated as previously described using artificial wounding method [26]. The initial sugarcane response was analyzed based on pools of 10 breaking buds collected at 5 DAI (days after inoculation). The late response was evaluated using culms after whips emerged at 200 DAI. Sampling was at the base of the whips, up to 2 cm below the culm. This is a region of intensive sugarcane cell division and fungal sporogenesis. Infected plants were compared with control (mock-inoculated) plants of the same age. Three biological replicates were included for each inoculated and control plant using a completely randomized design maintained on greenhouse benches (S1A File). A PCR amplicon containing the rDNA internal transcribed spacer region (ITS1, 5.8S and ITS2) of *S. scitamineum* generated with the primers Hs (5' -AACACGGTTGCATCGGTTGGGTC- 3') and Ha (5' -GCTTCTTGCTCATCCTCACCACCAA- 3') according to Bueno [28] was used to confirm infection at 5 DAI.

RNA Extraction, Libraries, and Sequencing

Total RNA was extracted from the samples using distinct methods for each plant developmental stage as described by Taniguti et al. [26](S1A File). The quality of the total RNA was verified using an Agilent 2100 Bioanalyzer (Agilent Technologies, USA), and the libraries were constructed using a TruSeq RNA Sample Prep v2 Low Throughput (LT) kit as described in the manufacturer's instructions (Illumina, San Diego, CA). The libraries were paired-end sequenced using the Illumina system (HiScanSQ).

Pre-Processing and Mapping the Illumina Reads

The Illumina reads were treated as previously described [26] (S1A File). Two reference sequences were used to map the RNAseq data: the complete genome sequence of *S. scitamineum* [26] and a set of unigenes produced by the assemblage of RNAseq data from six sugarcane cultivars [29]. The software packages used for mapping were Bowtie2 V2.1.0 [30] and BWA [31]. Bowtie2 was used with the default parameters in the sensitive mode (-D 15; -R 2; -L 22; -i S, 1, 1.15), while BWA alignments were obtained using the default parameters (-n 0.04; -k 2; -O 11). The RNAseq reads that showed no similarities to the sugarcane unigenes using the above parameters were assembled using Trinity [32]. Clusters identified by the prefix “gg” were then selected by comparison to the Viridiplantae sequences of UniProtKB [33].

Sugarcane Gene Expression Analysis

The differentially expressed genes (DEGs) were identified using the DESeq2 package [34]. For the 5-DAI data, DEGs were considered to be statistically significant if they had a p-value less than 0.05 when compared with control buds. The multiple-test correction proposed by Benjamini and Hochberg [35] was used for the 200-DAI data by applying a FDR (False discovery rate) to generate a set of DEGs with the same significance level (<0.05). The DrawVenn webtool (<http://bioinformatics.psb.ugent.be/webtools/Venn/>) was used to produce Venn diagrams from the sets of DEGs obtained from the BWA/DESeq2 or Bowtie2/DESeq2 analyses as well as different reference sets.

Annotation and Gene Ontology Analysis

The BLAST2GO tool V2.7.2 [36] was used with the default parameters to assign GO (Gene Ontology) terms to the DEGs. Metabolic pathways analysis was performed based on the KEGG database [37]. GO enrichment analysis was conducted with the BLAST2GO tool using the two-sided Fisher’s exact test with the p-value set at ≤ 0.05 . The GenBank [38] and UniProt [33] databases and InterProScan [39], SignalP [40], TMHMM [41], ScanProsite, and MyDomains [42] tools were used to predict the function and features of the protein sequences.

Quantitative PCR (qPCR) Expression Analysis

Quantitative PCR analysis was used to confirm the gene expression profiling data obtained from RNAseq. Transcripts encoding: invertase, auxin transporter, trehalose-6P synthase, pyruvate decarboxylase, aldolase, S-adenosylmethionine synthetase (SAM), peroxidase and longifolia-like protein (LGN) were selected for reverse transcription-qPCR (RT-qPCR) reactions (S2 File). The primers were manually designed and the quality verified using Gene Runner (<http://www.generunner.net/>) and NetPrimer (<http://www.premierbiosoft.com/netprimer/>). All RT-qPCRs were performed in a 7500 Fast Real-Time PCR System (Applied Biosystems, Waltham, MA) using a GoTaq[®] One-Step RT-qPCR System Kit (Promega, Madison, WI). A reaction mixture containing 50 ng of RNA, 6.5 μ L of GoTaq[®] qPCR Master Mix, 0.2 μ M of each primer, 0.25 μ L of GoScript[™] RT Mix, and nuclease-free water to a final volume of 12.5 μ L was used for the three biological replicates and two technical replicates. The cycling conditions were as follows: 37°C for 15 min; 95°C for 10 min; 40 cycles of 95°C for 10 s, 60°C for 30 s, and 72°C for 30 s. Primer specificity was confirmed by obtaining the dissociation curve for each reaction. Sugarcane housekeeping genes encoding polyubiquitin [43] and GAPDH (d-glyceraldehyde-3-phosphate dehydrogenase [44] were used to normalize the expression signals. The PCR efficiencies and Cq values were obtained using the LinReg PCR program [45]. Relative changes in the gene expression ratios were calculated using REST software [46]. Control samples (mock-

inoculated plants) were used as calibrators. Student's *t*-test was used to estimate significant changes in the relative expression levels ($p < 0.05$).

Results and Discussion

General Analysis

In this study, the smut-intermediate resistant variety 'RB925345' developed whips and other disease symptoms beginning 127 days after inoculation (DAI). However, the plants were sampled at 200 DAI because this was the time when whips were detected in all three replicate plants used in the experimental design (S1A File). Of the total number of inoculated plants, 48 (53%) developed whip over the timeline of the experiment (334 days) (S1B File).

Samples of 5 DAI were used to amplify the 509-bp sequence that corresponds to the 5.8S ribosomal RNA gene and flanks internal transcribed spacers 1 and 2 in *S. scitamineum* [28], confirming the fungal infection (S1C File). It was particularly necessary to confirm infection 5 DAI, because the buds were collected with no smut disease symptoms.

A total of 225.2 million paired-end sequences (PEs) of ~100 bp (~22.5 Gbp) was obtained for the 12 RNAseq libraries (~18 million reads per library). Including the corresponding control libraries, 111,926,958 (49.7%) PEs were from the 5-DAI collection and 113,269,226 (50.3%) PEs were from the 200-DAI collection (S3A File). Fungal sequencing reads were screened after mapping them to the whole *S. scitamineum* SSC39B genome [26], leading to the removal of approximately 20% (2% 5 DAI; 18% 200 DAI) of the PEs (S3B File).

Count-Based Differential Expression Analysis of the RNAseq Data

A set of previously obtained sugarcane transcripts was used to describe the biological events underlying the interaction with *S. scitamineum*. The reference set of sugarcane unigenes consisted of 72,268 sequences obtained from a de novo RNAseq assembly and a transcriptome annotation for six cultivars collected in various sugarcane crop fields, including the 'RB925345' variety [29]. This set of unigenes was used to allow cross-comparisons among the sugarcane sequencing data. To define the best alignment of the RNAseq to the unigenes set we performed two analyses. Using both Burrows-Wheeler Aligner (BWA) [31] and Bowtie2 [30] softwares, approximately 73% of the high-quality sequence reads were aligned to 67% of the unigenes (S3 File). The remaining subset of reads (15,000,000 PEs), those that showed no similarity to the sugarcane unigenes, was clustered using Trinity v2.0.6 [32]. A total of 25,794 contigs of more than 500 bp was assembled, and 16,219 were defined as 'RB925345' transcripts based on the presence of orthologs in the Viridiplantae section of the UniProt database (UniProt release 2015_03) (S4 File). These transcripts were identified by the prefix "gg" and probably include those most related to the pathogen infection. This new set of transcripts (gg) was combined with the sugarcane unigenes [29] (88,487 transcripts) to define the final set of DEGs (S5 and S6 Files). Our goal was to assess differential gene expression that started shortly after inoculation (5-DAI) and continued through fungal sporogenesis and, subsequently whip emission (200-DAI). DEGs were defined for both 5-DAI and 200-DAI samples, and the intersection between the two transcript groups was used for the annotation processes (Fig 2).

It was not possible to detect DEGs using FDR in 5-DAI samples. However, we pursued the analysis cautiously using *p*-values set at less than 0.05 and 0.01. The subtle sugarcane response shortly after inoculation may be due to the low percentage of the fungus in buds and/or delayed plant response due to the susceptibility of the variety used in the experiment [25]. Only 2% of the total reads were detected as genes expressed by the fungus, which represents approximately 67% of the *S. scitamineum* complete set of genes [26].




Fig 2. Differential expression gene analysis of 5-DAI and 200-DAI samples of the smut infected plants. Venn diagrams show the number of genes detected as differentially expressed (DEGs) by the DESeq2 package [33] using two mapping software systems: 1) Bowtie2 and 2) BWA, as well as two sets of references: 1) sugarcane unigenes [29] and 2) the novel sugarcane transcripts assembled from sugarcane unigenes unmapped sequences using Trinity [32]. Intersections of the identified DEGs from the mapping results were considered for the following analysis.

doi:10.1371/journal.pone.0162237.g002

Enrichment Analysis of GO Terms

The molecular events underlying sugarcane response during infection were initially suggested based on GO terms assignment and GO enrichment analysis of DEGs (S7 File). The enrichment analysis of 5-DAI DEGs showed as expected two contrasting molecular responses as previously described [25]. The genes involved in general plant immunity were down-regulated, while those for epigenetic mechanisms were up-regulated (Fig 3). In addition, this same enrichment test included terms related to shoot apical activities with the identification of three GO terms: Regionalization, Organ boundary specification, and Specification of floral organ identity. These terms suggest that the plant meristem functions are prematurely modulated by the presence of the pathogen. Some genes related to this same functional group were also enriched after whip emission. The gene regulatory network for shoot apical functions known in plant models is responsible for the differentiation of cells and organs (leaves and inflorescences). In corn, smut fungi are known to prevent or modify floral organ differentiation inducing tumor-like galls [47,48]. We suggest that a similar modulation occurs in susceptible genotypes of sugarcane infected with *S. scitamineum* very shortly after colonization. However, although the GO term enrichment test was in agreement with this hypothesis, it needs to be further investigated because 5-DAI DEGs were not supported by FDR. A time-course experiment using sugarcane varieties showing various levels of smut-related responses, for instance, should be conducted to determinate the expression profile of genes associated with the meristematic functions identified here.

The meristem-related functions were remarkably affected considering the enrichment test of 200-DAI sample DEGs (S7 File). Several identified genes allowed us to propose a putative model (Fig 4) for the transition of the normal meristematic functions controlled by the interaction of auxin and cytokinin hormones [49] to the development of the whip. For instance, a *longifolia-like* gene (LNG, *comp200950_c0_seq1*) is up regulated at this time point. Mutants overexpressing this same gene in *A. thaliana* have long petioles, narrow but extremely long leaf




Fig 3. Enrichment analysis of GO terms. DEGs were submitted to enrichment analysis in BLAST2GO software, and a p-value ≤ 0.05 was used as the cut-off parameter. The gray bars represent the percentage of genes related to each selected GO term in the total set of sugarcane unigenes. The red bars represent the percentage of genes related to each selected GO term in the set of DEGs. The complete list of enriched GO terms in each set of DEGs can be found in [S6 File](#).

doi:10.1371/journal.pone.0162237.g003

blades with serrated margins, elongated floral organs, and elongated siliques as a result of polar-cell elongation [50]. This description resembles the whip development in sugarcane. The LNG-like gene was assayed by qPCR, confirming the RNAseq data (Fig 5). Transcripts encoding the homologs of VIN3 (*vernalization insensitive 3 protein*) were also up-regulated after whip emission. VIN3-like proteins are involved in both the vernalization and photoperiod pathways by regulating the expression of the floral repressors FLOWERING LOCUS C (FLC) and FLOWERING LOCUS M (FLM). In *A. thaliana*, the VIN3-LIKE protein epigenetically represses a member of the FLC family, enabling flowering in the non-inductive photoperiods [51]. Additional transcripts members of three gene classes responsible for floral development (A, B and C) were identified. The ABCDE model proposes that a certain combination of MADS proteins activates different groups of genes related to flower [52,53]. A MADS-box TF homologous to AP1 (APETALA1; class A) was detected to be highly expressed (*comp207551_c1_seq1*; \log_2 fold-change = 9.4). This gene is essential in *A. thaliana* for the transition from an inflorescence meristem to a floral meristem [54]. This same MADS-box TF,




Fig 4. Sugarcane DEGs related to meristem functions. (A) Expression profile of genes related to meristem functions represented as values of a log2 fold change (inoculated/control). The heatmap was constructed using the R software package. Blue squares represent down-regulated genes, and red squares represent up-regulated ones. The statistical significance of expression regulation is presented in S5 and S6 Files. (B) Model of probable events related to whip development in sugarcane. Increase in VIN3 expression early in infected plants may release FT expression, which in turn positively regulates Apetala-1 (AP1) expression, turning the vegetative growth program to reproductive, via the autonomous/vernalization pathway. The black arrow represents up-regulation at 5 DAI, and the gray arrows represent up-regulation at 200 DAI.

doi:10.1371/journal.pone.0162237.g004




Fig 5. RT-qPCR validation. Sugarcane unigenes selected for RT-qPCR analysis of 5-DAI and 200-DAI samples: longifolia-like protein (comp200950_c0_seq1); auxin transporter (comp205699_c0_seq1); SAM (comp194455_c0_seq1); invertase (comp201528_c0_seq1); trehalose 6P synthase (comp204716_c0_seq1); aldolase (comp196354_c1_seq1); pyruvate decarboxylase (comp200606_c0_seq1) and peroxidase (comp187834_c0_seq1). The reactions were performed using a one-step GoTaq[®] One-Step RT-qPCR System Kit (Promega) using a 7500 Fast Real-Time PCR System (Applied Biosystems). Statistical analysis was performed using REST[®] software. “*” indicates genes differentially expressed in the RT-qPCR reactions (p-value < 0.05).

doi:10.1371/journal.pone.0162237.g005

along with the product of FLOWERING LOCUS T (FT; *gg_11173*), which is also highly expressed in infected sugarcane plants (\log_2 fold-change = 2.74), promotes the transition from vegetative to reproductive growth. In *A. thaliana*, FT encodes a small peptide that is recognized as the major component of florigen, which induces the expression of other floral genes, such as AP1 [55]. Three other MADS-box TFs were up-regulated after whip emission, encoding the homologs of APETALA3 (AP3, *gg_00300*), the class B gene AGAMOUS (AG, *comp204141_c1_seq1*), the class C gene APETALA1 (AP1, *gg_05696*) and COL6 (C2C2-CO-like transcription factor; *comp194394_c0_seq1*). COL6 belongs to the CONSTANS family and encodes a putative zinc finger TF that promotes the induction of flowering in *A. thaliana* during long photoperiods [56] through activation of floral meristem-identity genes, such as LEAFY [57]. Regulatory switches coordinating these developmental changes have been extensively studied in *A. thaliana* [54]; they are very precise and may vary in sugarcane, but the enrichment of genes related to the transition in meristem functions led us to associate these events with the plant-pathogen interaction mode [54]. It seems reasonable to assume that a combination of MADS-box TFs that are up-regulated in smut-infected plants may coordinate the gene expression related to whip development as an alternative route instead of the normal flowering program.

Given the interaction of *S. reilianum* with maize [48] and the data presented here, both *Sporisorium* species may share a common trend of modifying the meristem identity. In maize, phyllody and tumor formation result from alterations in the floral developmental program at both the apex and axillary meristems [48]. In sugarcane, the whip may result from releasing the transition from vegetative to reproductive/flowering, potentially via the autonomous/vernalization pathway [51] (Fig 4B).

Hormonal Imbalance Plays a Role in Sugarcane Smut Disease

Changes in the expression profiles of genes related to the regulation, synthesis, and transport of hormones identified in the enrichment analysis were investigated. We detected that JA (jasmonic acid)-mediated as well as SA (salicylic acid) signaling are potentially restrained in the experiments at 5-DAI (Fig 3). Indeed Que et al. [25], using a smut-resistant sugarcane genotype, identified the overexpression of JA-associated genes.

In contrast to SA and JA, auxin-activated signaling pathways are up-regulated. In addition to acting as a negative regulator of the plant immune system [58, 59], we suspect that auxin-related DEGs are associated with the meristem transcriptional reprogramming during whip emission because an increased auxin transporter gene expression was confirmed by RT-qPCR in both 5- and 200-DAI samples (Fig 5). The balance of auxin-cytokinin is reported to be essential for typical meristem function [49]. Additionally, auxin is the hormone with the highest number of responsive genes up-regulated after whip emission (Fig 6 and S8 File), including those involved in auxin influx/efflux, auxin-amino acid hydrolase, and auxin-responsive proteins, such as Aux/IAA, SAUR, and auxin-induced β -glucosidase. In the *S. reilianum*-maize pathosystem, the floral reversion process is partially attributed to an increase in auxin concentration, contributing to the loss of apical dominance and a greater number of ears per branch [48]. Auxin-dependent signaling is likely necessary for whip emission, which also involves the loss of apical dominance and growth of secondary buds [3]. An increase in auxin-like substances was reported by Hector et al. [60] using smutted sugarcane extracts. The authors suggested that the balance between auxin and cytokinin is disrupted in infected sugarcane plants.

Although cytokinin is a hormone that is often related to tillering [61], a symptom often related to smut, genes related to its synthesis were not detected among those that were

differentially expressed (Fig 6). However, several type-A response regulators (RRs) were up-regulated at 200 DAI. Type-A RRs negatively regulate cytokinin signaling by repressing type-B RRs, and they are transcriptionally up-regulated in response to cytokinin [62]. Cytokinins are central regulators in maize smut caused by *U. maydis*. The fungus can synthesize cytokinins,




Fig 6. DEGs related to hormone biosynthesis and signaling. (A) Heatmap constructed using the R software package and log2 fold change values (inoculated/control). The left column represents the regulation of each gene at 5 DAI, and the right column represents the regulation of each gene at 200 DAI. **(B)** Model concerning the contribution of the main plant hormones with up-regulated DEGs in sugarcane after whip emission.

doi:10.1371/journal.pone.0162237.g006

an important virulence factor that is associated with uncoordinated cell division and tumor formation [63,64]. The *S. scitamineum* genome SSC39B [26] does not have cytokinin biosynthetic gene homologs. However, previously data have shown that *S. scitamineum* secretes some cytokinin-like substances that can potentially activate cytokinin-responsive genes [65].

Regarding ethylene (ET), DEGs related to biosynthesis, perception, and signal transduction were detected after whip emission (Fig 6). For instance, SAM (S-adenosyl-L-methionine) was identified as being up-regulated, which was confirmed by RT-qPCR analysis (Fig 5). ET is often related to the lignification of plant tissues by increasing the expression of genes involved in the phenylpropanoid pathway [66,67].

The genes related to brassinosteroids were also up-regulated. They included several BAK1 LRRs [68] and the transcription repressor BZR1, which binds directly to the promoters of feedback-regulated brassinosteroid biosynthetic genes [69]. The identification of GA2ox and DELLA proteins among the DEGs suggests a blockage of GA signaling because the products of these genes act by reducing the availability of active GAs and repressing GA-responsive genes, respectively [70,71] (Fig 6). For instance, rice plants overexpressing C20-GA2ox exhibit early tillering, adventitious root growth, and changes in plant architecture that generate semi-dwarfs [70]. Diseased sugarcane plants have reduced node distance and poor cane formation ([3], our own observations), similar to the symptoms described in rice with low GA levels.

Carbon Distribution Is Affected by *S. scitamineum* Colonization

In biotrophic interactions pathogen growth relies on host nutrients derived from active metabolism. Sucrose and its derivatives are central molecules involved in carbohydrate translocation, metabolism, and sensing in higher plants [72]. Invertases were among the DEGs up-regulated after whip emission and included a neutral alkaline invertase (*comp189016_c0_seq1*) and soluble acid invertase (*comp201528_c0_seq1*), both of which were confirmed by RT-qPCR analysis. Invertases catalyze the irreversible hydrolysis of sucrose (EC 3.2.1.26), in some cases leading to a shift of the apoplastic sucrose/hexose ratio in favor of hexoses [73], which regulate many aspects of plant metabolism such as carbohydrate partitioning, developmental processes, and hormonal responses to biotic stress [73,74]. Plant invertases are classified into three groups: alkaline/neutral invertases localized in the cytosol, mitochondria or plastids and two types of acid invertases, one insoluble and bound to the cell wall (cell wall invertase CWI) and the other soluble in the vacuole space (vacuolar invertase VI) [73,74]. The acid invertases CWI play a role in sucrose partitioning, plant development and cell differentiation, whereas the VIs determine the sucrose level stored in the vacuole and its remobilization for metabolic processes. The up-regulation of vacuolar and neutral invertases represents a shift in the plant's metabolism that targets carbon to pathways unrelated to sucrose storage, which can aggravate sugarcane smut symptoms. Additionally, it has relevant implications for the hexose-based sugar signaling system involved in plant immunity [75]. Increased hexose levels can also be related to the nutrients supplied to the pathogen during teliospore differentiation and whip formation (Fig 7). The importance of the sugar content in signaling for axillary bud growth was recently demonstrated and indicates that, in addition to auxin, an increased sugar supply is necessary and sufficient for suppressed buds to be released from apical dominance [76]. Redistribution of the host carbon in response to *S. scitamineum* sporogenesis is suggested by the transcriptional profiles of the genes related to glycolysis, the citric acid cycle, sucrose, starch, xylan, trehalose 6P, and cellulose biosynthesis (Fig 7), findings that were also confirmed by RT-qPCR analysis (Fig 5). It has been suggested that, rather than playing a metabolic role, the low concentration of trehalose-6P (T6P) in infected plants functions as a regulatory component. Trehalose-6P synthase can sense sucrose availability to generate T6P as a signal to promote growth [72,77,78]. There




Fig 7. Smutted sugarcane metabolism in the late moments of interaction. Schematic representation of smutted sugarcane metabolism in the whip base (200 DAI). Red and blue arrows represent up- and down-regulated expression, respectively, and black arrows are unchanged expression.

doi:10.1371/journal.pone.0162237.g007

is also evidence in *A. thaliana* that T6P acts as an endogenous signal to control the transition from vegetative growth to flowering by increasing trehalose-6P synthase transcript levels [72,79]. These findings should encourage new experiments to better understand the sugarcane metabolic response to smut as the disease progresses, the turning point at which the plant changes its metabolism to allow fungal sporogenesis, and the significance of this shift to teliospore/whip differentiation.

Several transcripts related to lignin biosynthesis (Fig 5) were detected as being up-regulated after whip emission such as shikimate hydroxycinnamoyl transferase (EC 2.3.1.133), cinnamoyl-CoA reductase (EC 1.2.1.44), and peroxidase (EC 1.11.1.7). In addition, up-regulated laccases were also detected, such as DEGs (*gg_01080*, *gg_14238*, *gg_10439*, *gg_15488*). Plant laccases (EC 1.10.3.2) are glycoproteins involved in lignin biosynthesis through the oxidation of lignin precursors [80]. An increase in the lignification of smut-resistant plants has been detected by measuring cinnamyl alcohol dehydrogenase levels and by the overexpression of genes in RNAseq experiments of resistant varieties in the early moments of interaction [25,81]. The increase in lignin content after whip emission is probably not related to a protective host response; instead, it is likely a stage in the formation of the whip, which is composed of lignified plant tissue [82]. Recently, a proteomic approach developed after whip emission revealed 53 proteins related to lignin accumulation and oxidative stress at this stage of disease symptoms [83]. Responses regarding ROS (*Reactive Oxygen Species*) in 200-DAI samples were also detected at the RNA level. Nineteen DEs were identified as being related to ROS-scavenging

enzymes (S9 File), including 16 up-regulated DEGs (one catalase, eight peroxidases, two thioredoxins, and five glutathione S-transferases) and three down-regulated DEGs (two peroxidases and one thioredoxin). These results suggest that the ROS level is high in *S. scitamineum*-colonized cells during sporogenesis.

Sequence Features of Resistance Gene Analogs (RGAs) Differentially Expressed

Resistance gene analogs (RGAs) were analyzed for both time points. The predicted domains and other sequence features that are potentially important for RGAs function were identified (Fig 8). Although this study was conducted with an intermediate-resistant genotype, we detected promising candidates associated with this particular biotrophic interaction and their potential role in the disease progression mechanism proposed here. Several RGAs containing leucine-rich repeat (LRR) domains have already been identified in sugarcane [84]. Three of them were also found here: the two BAM-related proteins *comp_188744* and *gg_06875* (RGA482) and a protein encoded by *comp_187876* (RGA367) (Fig 8). All of the other proteins predicted in this work are new discoveries in sugarcane.

BAM orthologs are leucine-rich repeat receptor-like serine/threonine-RD kinases (LRR-RLKs) [85] and receptors for signals to switch meristem identity. BAM proteins play roles equivalent to those of CLV (CLAVATA) proteins in *A. thaliana*, representing a functional redundancy within the program related to meristem functionality [86]. Other LRR-RLKs, such as ERECTA, were first described in relation to plant development, and only later were they positively associated with disease resistance [85].

Other kinases were identified sharing amino acid identities with the RLKs of different families (Fig 8). For instance, *comp205382_c0_seq1* encodes a transmembrane protein (non-RD kinase) harboring an LRR domain [85]. The translated amino acid sequence is most similar to the protein Xa21 in *O. sativa*, sharing all conserved residues [87]. The *xa21* gene is known to confer resistance to *Xanthomonas oryzae* pv. *oryzae* race 6 in rice [87]. LRR-RLK-harboring domains shared by plant S-locus glycoproteins and S-receptor kinases (SRK) (IPR003609) were also detected (*comp182409_c0_seq1*). SRKs were first described as being associated with *Brassica* self-incompatibility, but were later related to the perception of pathogen infection, probably by binding to a glycoprotein inducer, such as cellulose or chitin [88]. A transcript (EST) similar to *comp182409_c0_seq1* was detected by BLAST (GenBank) in *Oryza longistaminata*. *Comp182409_c0_seq1* shares an 87% identity to the 3' end of the *xa21* nucleotide sequence (FF359116), which is the portion coding for the kinase domain. No sugarcane ESTs similar to *comp182409_c0_seq1* were found in the NCBI-expressed sequence tags database.

S. scitamineum also induced the presence of cytoplasmic LRR proteins known as plant intracellular Ras group-related LRR proteins (PIRLs) [89]. The protein encoded by the *comp196006_c0_seq1* DEG at 200 DAI is most similar to other PIRLs and contains the conserved GxxxVxxYxxxxW ('GVYW') motif immediately following the LRR domain.

The expression of an F-box/LRR-related gene was also detected in 200-DAI samples. F-box proteins are part of the SCF (SKP1/Cullin/F-box) ubiquitin ligase complex involved in protein degradation (proteasome) [90]. In *O. sativa*, the F-box/LRR-repeat MAX2-homolog controls tillering by suppressing axillary bud activity, potentially by degrading specific proteins that activate axillary growth [91]. The *comp204415_c0_seq1* DEG encoded a protein that has an 83% amino acid identity with the rice MAX2-homolog (Q5VMP0). Tillering is one of the earlier disease symptoms related to smut [3].

Cytoplasmic proteins attached to the nucleotide binding-ARC (NB-ARC) domains and containing an ATPase and a nucleotide-binding site [92,93] compose the last class of LRR




Fig 8. Resistance gene analogs (RGAs) detected as DEGs in smut-infected sugarcane in early and late interaction. The guide tree was obtained based on translated amino acid sequence similarity using CLUSTALW2. The heatmap represents the respective expression log2 fold change values. Protein structural features: signal peptide sequences predicted by SignalP 4.1 (<http://www.cbs.dtu.dk/services/SignalP/>); transmembrane domains predicted by TMHMM Server v. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>). Other domains, motifs and active sites were predicted by InterProScan. CC, coiled coil; kinase (IPR011009); NB, nucleotide-binding (IPR002182); B, Bulb-type lectin domain (IPR001480); EGF, epidermal growth factor domain (IPR000742); PAN domain (IPR03609); LRR, leucine-rich-repeat (IPR001611; IPR003591); red pins represent kinase active sites, and gray arrows are P-loops of N-terminal NB-ARC proteins.

doi:10.1371/journal.pone.0162237.g008

proteins identified among the translated transcripts that were differentially accumulated. The translated protein sequence of *comp202536_c0_seq1* contains the conserved GLPLA and MHD motifs that are essential to the function of other resistance proteins [93]; although lacking an obvious CC-domain at the N-terminus, this gene probably belongs to one of the CC-domain-containing subfamilies. Its sequence is most similar to an *O. sativa* gene assigned to chromosome 4 (CAE03396). All of these RGA-like encoding proteins are

potential targets for functional characterization as receptors of signals due to the presence of *S. scitamineum*.

Conclusion

This work reveals transcriptional changes associated with the most characteristic symptom of sugarcane smut disease. We speculate that whip emission is a consequence of premature transcriptional changes in meristem function (5-DAI) that results in the restraint of floral development via the vernalization pathway by increasing VIN3, COL6, FT, and AP1 gene expression and other flowering-related transcriptional factors (200-DAI). Fungal sporogenesis and whip emission are most related to auxin mobilization followed by a strong response of ROS-scavenging enzymes. In addition, the role of other plant hormones is also suggested. Because the processes associated with fungal development and whip emission require energy, the carbon partitioning of sugarcane is the most affected. The gene expression profile indicates that smutted sugarcane metabolism shifts towards energy production, increasing the expression of genes involved in glycolysis and TCA. The synthesis of signaling molecules, such as trehalose 6P, is also among the results described. Increased expression of genes involved in lignin biosynthesis and sucrose breakdown are potential markers of whip development. We also described the RGA expression patterns involved in this particular interaction, leading to effective fungal colonization and disease establishment. *S. scitamineum* is known to colonize not only susceptible plants but also smut-resistant genotypes that, in response to unknown signals, allow unexpected fungal sporogenesis and whip emission. This detailed work is an attempt to expose molecular mechanisms and candidate genes that can possibly reveal ways to control sugarcane smut disease.

Database Accession Number

The sequencing data has been deposited at DDBJ/EMBL/GenBank under the BioProject ID PRJNA291816.

Supporting Information

S1 File. A) Experimental design used to produce biological insights of the interaction sugarcane-*S. scitamineum*. B) Number of whips developed each month after smut inoculation in the intermediate resistant genotype RB925345. C) Amplicons of primers Hs and Ha using total DNA of buds collected 5 DAI.

(PDF)

S2 File. Primers used for qPCR validation of RNAseq analysis.

(XLS)

S3 File. A) Total of RNAseq data obtained and mapping results of BWA and Bowtie2 softwares against sugarcane unigenes. B) Percentage of reads mapped in *S. scitamineum* SSC39B genome.

(XLSX)

S4 File. Fasta file of the new sugarcane transcripts assembled (gg).

(TXT)

S5 File. A) Mapping results and differential expression analysis using as reference sugarcane unigenes. B) IDS of sugarcane unigenes differentially expressed during interaction with *S. scitamineum* in early and late moments.

(XLSX)

S6 File. A) Mapping results and differential expression analysis considering the new sugarcane transcripts (gg) obtained from non-mapped reads in sugarcane unigenes. B) IDS of new sugarcane transcripts differentially expressed during interaction with *S. scitamineum* in early and late moments. C) Blast2GO annotation of new sugarcane transcripts differentially expressed during interaction with *S. scitamineum* in early and late moments.
(XLS)

S7 File. GO terms enrichment analysis of sugarcane. A) down-regulated DEGs at 5 DAI. B) GO terms enrichment analysis sugarcane up-regulated DEGs 5 DAI. C) GO terms enrichment analysis of sugarcane down-regulated DEGs 200 DAI. D) GO terms enrichment analysis of sugarcane up-regulated DEGs 200 DAI. E) GO terms enrichment analysis sugarcane down-regulated DEGs (gg) 5 DAI. F) GO terms enrichment analysis of sugarcane up-regulated DEGs (gg) 5 DAI. G) GO terms enrichment analysis of sugarcane down-regulated DEGs (gg) 200 DAI. H) GO terms enrichment analysis of sugarcane up-regulated DEGs (gg) 200 DAI.
(XLS)

S8 File. DEGs hormone-related at 200 DAI.
(XLS)

S9 File. DEGs ROS-related at 5 DAI and 200 DAI.
(XLS)

Acknowledgments

We thank the sequencing facility Center of Functional Genomics (USP, Campus “Luiz de Queiroz”); NAP/MEPA (Núcleo de Apoio à Pesquisa em Microscopia Eletrônica Aplicada a Agricultura) and Laboratórios de Genética Molecular, Departamento de Fitopatologia e Nematologia, and Plantas Hortícolas, Departamento de Produção Vegetal, ESALQ.

Author Contributions

Conceived and designed the experiments: CBMV MLCV JPK.

Performed the experiments: PDCS ACP LPP.

Analyzed the data: LMT PDCS LPP JPK.

Contributed reagents/materials/analysis tools: SC KSA.

Wrote the paper: PDCS ACP LMT LPP CBMV.

Provided expertise and editing: JPK MLCV MAVS.

References

1. FAOSTAT, 2013. Food and Agriculture Organization Statistics Database. Available: <http://faostat3.fao.org/faostat-getaway/go/to/home/E>. Accessed 10 November 2015.
2. Waclawovsky AJ, Sato PM, Lembke CG, Moore PH, Souza GM. Sugarcane for bioenergy production: an assessment of yield and regulation of sucrose content. *Plant Biotechnol J*.2010; 8:3. Available: <http://www.ncbi.nlm.nih.gov/pubmed/20388126>.
3. Sundar AR, Barnabas EL, Matathi P, Viswanathan R. A Mini-Review on Smut Disease of Sugarcane Caused by *Sporisorium scitamineum*. In: Mworio JK, editor. *Botany*. InTech; 2012. p. 107–28.
4. Legaz ME, Armas R d., Pinon D, Vicente C. Relationships between phenolics-conjugated polyamines and sensitivity of sugarcane to smut (*Ustilago scitaminea*). *J Exp Bot*. 1998, 49:327. Available: <http://jxb.oxfordjournals.org/cgi/content/long/49/327/1723>.

5. Bakkeren G, Kämper J, Schirawski J. Sex in smut fungi: Structure, function and evolution of mating-type complexes. *Fungal Genet Biol.* 2008; 45. Available: <http://www.sciencedirect.com/science/article/pii/S1087184508000650>.
6. Chao CP, Hoy JW, Saxton AM, Martin F.A. Heritability of resistance and repeatability of clone reactions to sugarcane smut in Louisiana. *Phytopathology.* 1990; 80: 622–626.
7. D'Hont A, Grivet L, Feldmann P, Rao S, Berding N, Glaszmann JC. Characterisation of the double genome structure of modern sugarcane cultivars (*Saccharum* spp.) by molecular cytogenetics. *Mol Gen Genet.* 1996; 250: 4. Available: <http://dx.doi.org/10.1007/BF02174028>.
8. de Setta N, Monteiro-Vitorello CB, Metcalfe CJ, Cruz GMQ, Del Bem LE, Vicentini R, et al. Building the sugarcane genome for biotechnology and identifying evolutionary trends. *BMC Genomics.* 2014; 15: 1. Available: <http://www.biomedcentral.com/1471-2164/15/540>.
9. Garcia AAF, Mollinari M, Marconi TG, Serang OR, Silva RR, Vieira MLC, et al. SNP genotyping allows an in-depth characterisation of the genome of sugarcane and other complex autopolyploids. *Sci Rep* 2013; 3: 3399. Available: <http://dx.doi.org/10.1038/srep03399>. PMID: 24292365
10. Palhares AC, Rodrigues-Morais TB, Van Sluys M-A, Domingues DS, Maccheroni W, Jordão H, et al. A novel linkage map of sugarcane with evidence for clustering of retrotransposon-based markers. *BMC Genet.* 2012; 13: 51. Available: <http://www.biomedcentral.com/1471-2156/13/51>. doi: 10.1186/1471-2156-13-51 PMID: 22742069
11. James GL. Smut spore germination on sugarcane internode surfaces. *Proc South African Sugar Technol Assoc.* 1973; 47: 179–80.
12. Pillay HLLM. The development of an improved method for evaluating sugarcane for resistance to smut. *Proc South African Sugar Technol Assoc.* 1980; 54: 168–72.
13. Lloyd HL, Naidoo G. Chemical-assay potentially suitable for determination of smut resistance of sugarcane cultivars. *Plant Dis.* 1983; 67:1103–5.
14. Piñon D, de Armas R, Vicente C, Legaz ME. Role of polyamines in the infection of sugarcane buds by *Ustilago scitaminea* spores. *Plant Physiol Biochem.* 1999; 37: 1 Available: <http://www.sciencedirect.com/science/article/pii/S0981942899800679>.
15. Borrás-Hidalgo O, Thomma BPHJ, Carmona E, Borroto CJ, Pujol M, Arencibia A, et al. Identification of sugarcane genes induced in disease-resistant somaclones upon inoculation with *Ustilago scitaminea* or *Bipolaris sacchari*. *Plant Physiol Biochem.* 2005; 43: 12. Available: <http://www.sciencedirect.com/science/article/pii/S0981942805002020>.
16. LaO M, Arencibia A, Carmona E, Acevedo R, Rodríguez E, León O, et al. Differential expression analysis by cDNA-AFLP of *Saccharum* spp. after inoculation with the host pathogen *Sporisorium scitamineum*. *Plant Cell Rep.* 2008; 27: 6. Available: <http://dx.doi.org/10.1007/s00299-008-0524-y>.
17. RUTHERFORD LTAR. cDNA-AFLP differential display of sugarcane (*Saccharum* spp. hybrids) genes induced by challenge with the fungal pathogen *Ustilago scitaminea* (sugarcane smut). *Proc South African Sugar Technol Assoc.* 2001; 75: 104–7.
18. You-Xiong Q, Jian-Wei L, Xian-Xian S, Li-Ping X, Ru-Kai C. Differential Gene Expression in Sugarcane in Response to Challenge by Fungal Pathogen *Ustilago scitaminea* Revealed by cDNA-AFLP. *J Biomed Biotechnol.* 2011; 14: 2011. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3142712/>.
19. Westermann AJ, Gorski SA, Vogel J. Dual RNA-seq of pathogen and host. *Nat Rev Micro.* 2012; 10: 9. Available: <http://dx.doi.org/10.1038/nrmicro2852>.
20. Kawahara Y, Oono Y, Kanamori H, Matsumoto T, Itoh T, Minami E. Simultaneous RNA-seq analysis of a mixed transcriptome of rice and blast fungus interaction. *PLoS One.* 2012; 7: 11. Available: <http://dx.doi.org/10.1371/journal.pone.0049423>.
21. Lowe RGT, Cassin A, Grandaubert J, Clark BL, Wouw Van de AP, Rouxel T, et al. Genomes and Transcriptomes of Partners in Plant-Fungal- Interactions between Canola (*Brassica napus*) and Two *Leptosphaeria* Species. *PLoS One.* 2014; 9: 7. Available: <http://dx.doi.org/10.1371/journal.pone.0103098>.
22. Tremblay A, Hosseini P, Alkharouf NW, Li S, Matthews BF. Gene Expression in Leaves of Susceptible Glycine max during Infection with *Phakopsora pachyrhizi* Using Next Generation Sequencing. *Sequencing.* 2011; 2011: 1. Available: <http://www.hindawi.com/archive/2011/827250/>.
23. Yazawa T, Kawahigashi H, Matsumoto T, Mizuno H. Simultaneous transcriptome analysis of *Sorghum* and *Bipolaris sorghicola* by using RNA-seq in combination with de novo transcriptome assembly. *PLoS One.* 2013; 8: 4. Available: <http://dx.doi.org/10.1371/journal.pone.0062460>.
24. Zhu Q-H, Stephen S, Kazan K, Jin G, Fan L, Taylor J, et al. Characterization of the defense transcriptome responsive to *Fusarium oxysporum*-infection in Arabidopsis using RNA-seq. *Gene.* 2013; 512: 2. Available: <http://www.sciencedirect.com/science/article/pii/S0378111912013145>.

25. Que Y, Su Y, Guo J, Wu Q, Xu L. A Global View of Transcriptome Dynamics during *Sporisorium scitamineum* Challenge in Sugarcane by RNA-seq. PLoS One.2014; 9: 8. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4149577/>.
26. Taniguti LM, Schaker PDC, Benevenuto J, Peters LP, Carvalho G, Palhares A, et al. Complete Genome Sequence of *Sporisorium scitamineum* and Biotrophic Interaction Transcriptome with Sugarcane. PLoS One.2015; 10: 6. Available: <http://dx.doi.org/10.1371/journal.pone.0129318>.
27. Wu Q, Xu L, Guo J, Su Y, Que Y. Transcriptome Profile Analysis of Sugarcane Responses to *Sporisorium scitamineum* Infection Using Solexa Sequencing Technology. Biomed Res Int. 2013; 2013: 1. Available: <http://www.hindawi.com/journals/bmri/2013/298920/>.
28. Bueno CRNC. Infection by *Sporisorium scitamineum* on sugarcane: influence of environmental variables and development of a method for early diagnosis. São Paulo: University of São Paulo, Ph.D. thesis. 2011. Available: <http://www.teses.usp.br/teses/disponiveis/1/1/11135/tde-17092010-164620/>.
29. Cardoso-Silva CB, Costa EA, Mancini MC, Balsalobre TWA, Canesin LEC, Pinto LR, et al. De novo assembly and transcriptome analysis of contrasting sugarcane varieties. PLoS One.2014; 9: 2. Available: <http://dx.doi.org/10.1371/journal.pone.0088462>.
30. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012; 9: 4. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3322381/>.
31. Li H, Durbin R. Fast and accurate short read alignment with Burrows—Wheeler transform. Bioinformatics.2009; 25: 14. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2705234/>.
32. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotech.2011; 29: 7. Available: <http://dx.doi.org/10.1038/nbt.1883>.
33. The UniProtConsortium: a hub for protein information. Nucleic Acids Res.2015; 28: 43. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4384041/>.
34. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol.2014; 15: 12. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4302049/>.
35. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J R Stat Soc Ser B. 1995; 57: 1. Available: <http://www.jstor.org/stable/2346101>.
36. Conesa A, Götz S. Blast2GO: A Comprehensive Suite for Functional Analysis in Plant Genomics. Int J Plant Genomics. 2008; 30: 2008. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2375974/>.
37. Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res. 2000; 28: 1. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC102409/>.
38. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Rapp BA, Wheeler DL. GenBank. Nucleic Acids Res.2000; 28: 1. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC102453/>. PMID: 10592167
39. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale protein function classification. Bioinformatics. 2014; 30: 9. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3998142/>.
40. Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from transmembrane regions. Nat Meth.2011; 8: 10. Available: <http://dx.doi.org/10.1038/nmeth.1701>.
41. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. J Mol Biol.2001; 305: 3. Available: <http://www.sciencedirect.com/science/article/pii/S0022283600943158>.
42. de Castro E, Sigrist CJA, Gattiker A, Bulliard V, Langendijk-Genevaux PS, Gasteiger E, et al. ScanProsite: detection of PROSITE signature matches and ProRule-associated functional and structural residues in proteins. Nucleic Acids Res.2006; 1: 34. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1538847/>.
43. Papini-Terzi FS. Transcription Profiling of Signal Transduction-Related Genes in Sugarcane Tissues. DNA Res. 2005; 12:1. Available: <http://dnaresearch.oxfordjournals.org/content/12/1/27.abstract>.
44. Iskandar H, Simpson R, Casu R, Bonnett G, Maclean D, Mannes J. Comparison of reference genes for quantitative real-time polymerase chain reaction analysis of gene expression in sugarcane. Plant Mol Biol Report. 2004; 22: 4. Available: <http://dx.doi.org/10.1007/BF02772676>.
45. Ramakers C, Ruijter JM, Deprez RHL, Moorman AF. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. Neurosci Lett.2003; 339: 1. Available: <http://www.sciencedirect.com/science/article/pii/S0304394002014234>.
46. Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. Nucleic Acids Res.2002; 30: 9. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC113859/>.

47. Gao L, Kelliher T, Nguyen L, Walbot V. *Ustilago maydis* reprograms cell proliferation in maize anthers. *Plant J.* 2013; 75: 6. Available: <http://onlinelibrary.wiley.com/doi/10.1111/tpl.12270/full>.
48. Ghareeb H, Becker A, Iven T, Feussner I, Schirawski J. *Sporisorium reilianum* Infection Changes Inflorescence and Branching Architectures of Maize. *Plant Physiol.* 2011; 156: 4. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3149921/>.
49. Su Y-H, Liu Y-B, Zhang X-S. Auxin—Cytokinin Interaction Regulates Meristem Development. *Mol Plant.* 2011; 4: 4. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3146736/>.
50. Lee YK, Kim G-T, Kim I-J, Park J, Kwak S-S, Choi G, et al. LONGIFOLIA1 and LONGIFOLIA2, two homologous genes, regulate longitudinal cell elongation in Arabidopsis. *Development.* 2006; 133: 21. Available: <http://dev.biologists.org/content/133/21/4305.abstract>.
51. Kim D-H, Sung S. Genetic and Epigenetic Mechanisms Underlying Vernalization. *Arabidopsis Book.* 2014; 12: e0171. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3952384/>. doi: [10.1199/tab.0171](https://doi.org/10.1199/tab.0171) PMID: [24653667](https://pubmed.ncbi.nlm.nih.gov/24653667/)
52. Honma T, Goto K. Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature.* 2001; 409: 6819. Available: <http://dx.doi.org/10.1038/35054083>.
53. Murai K. Homeotic Genes and the ABCDE Model for Floral Organ Formation in Wheat. *Plants.* 2013; 2: 3. Available: <http://www.mdpi.com/2223-7747/2/3/379/htm>.
54. Kaufmann K, Wellmer F, Muiño JM, Ferrier T, Wuest SE, Kumar V, et al. Orchestration of Floral Initiation by APETALA1. *Sci.* 2010; 328: 5974. Available: <http://www.sciencemag.org/content/328/5974/85.abstract>.
55. Eckardt NA. Dissecting cis-Regulation of FLOWERING LOCUS T. *Plant Cell.* 2010; 22: 5. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2899881/>.
56. Lagercrantz U, Axelsson T. Rapid Evolution of the Family of CONSTANS LIKE Genes in Plants. *Mol Biol Evol.* 2000; 17: 10. Available: <http://mbe.oxfordjournals.org/cgi/content/long/17/10/1499>.
57. Simon R, Igeno MI, Coupland G. Activation of floral meristem identity genes in Arabidopsis. *Nature.* 1996; 384: 6604. Available: <http://dx.doi.org/10.1038/384059a0>.
58. Robert-Seilantantz A, Navarro L, Bari R, Jones JDG. Pathological hormone imbalances. *Curr Opin Plant Biol.* 2007; 10: 4. Available: <http://www.sciencedirect.com/science/article/pii/S136952660700074X>.
59. Wang D, Pajeroska-Mukhtar K, Culler AH, Dong X. Salicylic acid inhibits pathogen growth in plants through repression of the auxin signaling pathway. *Curr Biol.* 2007; 17: 20. Available: <http://www.sciencedirect.com/science/article/pii/S0960982207019847>.
60. Hector E, Rodriguez R, de Prada F, Delmonte A G R. Experimental evidence for the presence of different smut resistance mechanisms in sugarcane. In: Napompeth B, editor. XXI Congress of the International Society of Sugarcane Technologists. Bangkok: International Society of Sugarcane Technologists; 1992. p. 565–73.
61. Liu Y, Ding Y, Gu D, Li G, Wang Q, Wang S. The Positional Effects of Auxin and Cytokinin on the Regulation of Rice Tiller Bud Growth. *Crop Sci.* 2011; 51:2749–58. Available: <http://dx.doi.org/10.2135/cropsci2011.02.0083>.
62. To JPC, Deruère J, Maxwell BB, Morris VF, Hutchison CE, Ferreira FJ, et al. Cytokinin Regulates Type-A Arabidopsis Response Regulator Activity and Protein Stability via Two-Component Phosphorelay. *Plant Cell.* 2007; 19: 12. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2217641/>.
63. Brefort T, Tanaka S, Neidig N, Doehlemann G, Vincon V, Kahmann R. Characterization of the Largest Effector Gene Cluster of *Ustilago maydis*. Xu J-R, editor. *PLoS Pathog.* 2014; 10: 7. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4081774/>.
64. Bruce S, Saville B, Neil Emery RJ. *Ustilago maydis* Produces Cytokinins and Abscisic Acid for Potential Regulation of Tumor Formation in Maize. *J Plant Growth Regul.* 2011; 30: 1. Available: <http://dx.doi.org/10.1007/s00344-010-9166-8>.
65. Peros J.P. and Chagvardieff P. Mise en evidence in vitro d'effets toxiques entre *Ustilago scitaminea* Syd. et des cats de canne a sucre; influence de deux facteurs de croissance sur les phenomenes observes. *Agronomie.* 1983; 3: 629–34.
66. Ecker JR, Davis RW. Plant defense genes are regulated by ethylene. *Proc Natl Acad Sci USA.* 1987; 84: 15. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC298822/>.
67. Guo H, Ecker JR. The ethylene signaling pathway: new insights. *Curr Opin Plant Biol.* 2004; 7: 1. Available: <http://www.sciencedirect.com/science/article/pii/S1369526603001420>.
68. Li J, Wen J, Lease KA, Doke JT, Tax FE, Walker JC. BAK1, an Arabidopsis LRR Receptor-like Protein Kinase, Interacts with BRI1 and Modulates Brassinosteroid Signaling. *Cell.* 2002; 110: 2. Available: <http://www.sciencedirect.com/science/article/pii/S0092867402008127>.

69. He J-X, Gendron JM, Sun Y, Gampala SSL, Gendron N, Sun CQ, et al. BZR1 Is a Transcriptional Repressor with Dual Roles in Brassinosteroid Homeostasis and Growth Responses. *Sci*.2005; 307: 5715. Available: <http://www.sciencemag.org/content/307/5715/1634.abstract>.
70. Lo S-F, Yang S-Y, Chen K-T, Hsing Y-I, Zeevaart JAD, Chen L-J, et al. A Novel Class of Gibberellin 2-Oxidases Control Semidwarfism, Tillering, and Root Development in Rice. *Plant Cell*.2008; 20: 10. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2590730/>.
71. Thomas SG, Phillips AL, Hedden P. Molecular cloning and functional expression of gibberellin 2-oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proc Natl Acad Sci*.1999; 96: 8. Available: <http://www.pnas.org/content/96/8/4698.abstract>.
72. Ruan Y-L. Sucrose Metabolism: Gateway to Diverse Carbon Use and Sugar Signaling. *Annu Rev Plant Biol*.2014; 65: 1. Available: <http://dx.doi.org/10.1146/annurev-arplant-050213-040251>.
73. Tausin AS, Giardina T. Sucrose and invertases, a part of the plant defense response to the biotic stresses. *Front Plant Sci*.2014; 5:293. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4066202/>. doi: 10.3389/fpls.2014.00293 PMID: 25002866
74. Roitsch T, González M-C. Function and regulation of plant invertases: sweet sensations. *Trends Plant Sci*.2004; 9: 12. Available: <http://www.sciencedirect.com/science/article/pii/S136013850400250X>.
75. Morkunas I, Ratajczak L. The role of sugar signaling in plant defense responses against fungal pathogens. *Acta Physiol Plant*.2014; 36: 7. Available: <http://dx.doi.org/10.1007/s11738-014-1559-z>.
76. Mason MG, Ross JJ, Babst BA, Wienclaw BN, Beveridge CA. Sugar demand, not auxin, is the initial regulator of apical dominance. *Proc Natl Acad Sci*.2014; 111: 16. Available: <http://www.pnas.org/content/111/16/6092.abstract>.
77. Gómez LD, Baud S, Gilday A, Li Y, Graham IA. Delayed embryo development in the ARABIDOPSIS TREHALOSE-6-PHOSPHATE SYNTHASE 1 mutant is associated with altered cell wall structure, decreased cell division and starch accumulation. *Plant J*. 2006; 46: 1. Available: <http://dx.doi.org/10.1111/j.1365-313X.2006.02662.x>.
78. Kolbe A, Tiessen A, Schluempmann H, Paul M, Ulrich S, Geigenberger P. Trehalose 6-phosphate regulates starch synthesis via posttranslational redox activation of ADP-glucose pyrophosphorylase. *Proc Natl Acad Sci United States Am*.2005; 102: 31. Available: <http://www.pnas.org/content/102/31/11118.abstract>.
79. Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, et al. Regulation of Flowering by Trehalose-6-Phosphate Signaling in Arabidopsis thaliana. *Sci*. 2013; 339: 6120. Available: <http://www.sciencemag.org/content/339/6120/704.abstract>.
80. Dean JFD, LaFayette PR, Rugh C, Tristram AH, Hoopes JT, Eriksson K-EL, et al. Laccases Associated with Lignifying Vascular Tissues. In: *Lignin and Lignan Biosynthesis*. American Chemical Society; 1998. p. 8–96. Available: <http://dx.doi.org/10.1021/bk-1998-0697.ch008>.
81. Santiago R, Alarcón B, de Armas R, Vicente C, Legaz ME. Changes in cinnamyl alcohol dehydrogenase activities from sugarcane cultivars inoculated with *Sporisorium scitamineum* sporidia. *Physiol Plant*. 2012; 145: 2. Available: <http://dx.doi.org/10.1111/j.1399-3054.2012.01577.x>.
82. Legaz ME, Santiago R, de Armas R, Alarcón B, Díaz EM, Sánchez-Elordi E, Sacristán M, Fontaniella B, Millanes AB B M. Molecular defence responses of sugarcane (*Saccharum officinarum* L.) to smut (*Sporisorium scitamineum* (Syd.)). In *Science against microbial pathogens: communicating current research and technological advances*. (M.-V A., ed), Piepenbr & Oberw.; 2011.
83. Barnabas L, Ashwin NMR, Kaverinathan K, Trentin AR, Pivato M, Sundar AR, et al. Proteomic analysis of a compatible interaction between sugarcane and *Sporisorium scitamineum*. *Proteomics*. 2016; 16: 7. Available: <http://onlinelibrary.wiley.com/doi/10.1002/pmic.201500245/full>.
84. Rossi M, Araujo PG, Paulet F, Garsmeur O, Dias VM, Chen H, et al. Genomic distribution and characterization of EST-derived resistance gene analogs (RGAs) in sugarcane. *Mol Genet Genomics*. 2003; 269: 3. Available: <http://dx.doi.org/10.1007/s00438-003-0849-8>.
85. Afzal AJ, Wood AJ, Lightfoot DA. Plant Receptor-Like Serine Threonine Kinases: Roles in Signaling and Plant Defense. *Mol Plant-Microbe Interact*.2008; 21: 5. Available from: <http://dx.doi.org/10.1094/MPMI-21-5-0507>.
86. DeYoung BJ, Bickle KL, Schrage KJ, Muskett P, Patel K, Clark SE. The CLAVATA1-related BAM1, BAM2 and BAM3 receptor kinase-like proteins are required for meristem function in Arabidopsis. *Plant J*. 2006; 45: 1. Available from: <http://dx.doi.org/10.1111/j.1365-313X.2005.02592.x>. PMID: 16367950
87. Song W-Y, Wang G-L, Chen L-L, Kim H-S, Pi L-Y, Holsten T, et al. A Receptor Kinase-Like Protein Encoded by the Rice Disease Resistance Gene, Xa21. *Sci*. 1995; 270: 5243. Available: <http://www.sciencemag.org/content/270/5243/1804.abstract>.

88. Pastuglia M, Ruffio-Châble V, Delorme V, Gaude T, Dumas C, Cock JM. A functional S locus anther gene is not required for the self-incompatibility response in *Brassica oleracea*. *Plant Cell*. 1997; 9: 11. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC157058/>.
89. Forsthoefel NR, Cutler K, Port MD, Yamamoto T, Vernon DM. PIRLs: a novel class of plant intracellular leucine-rich repeat proteins. *Plant Cell Physiol*. 2005; 46: 6. Available: <http://pcp.oxfordjournals.org/cgi/content/long/46/6/913>.
90. Jain M, Nijhawan A, Arora R, Agarwal P, Ray S, Sharma P, et al. F-Box Proteins in Rice. Genome-Wide Analysis, Classification, Temporal and Spatial Gene Expression during Panicle and Seed Development, and Regulation by Light and Abiotic Stress. *Plant Physiol*. 2007; 143: 4. Available: <http://www.plantphysiol.org/content/143/4/1467.abstract>.
91. Ishikawa S, Maekawa M, Arite T, Onishi K, Takamura I, Kyojuka J. Suppression of tiller bud activity in tillering dwarf mutants of rice. *Plant Cell Physiol*. 2005; 46: 1. Available: <http://pcp.oxfordjournals.org/cgi/content/long/46/1/79>.
92. McHale L, Tan X, Koehl P, Michelmore RW. Plant NBS-LRR proteins: adaptable guards. *Genome Biol*. 2006; 7: 212. Available: <http://genomebiology.com/content/7/4/212>. PMID: 16677430
93. van Ooijen G, Mayr G, Albrecht M, Cornelissen BJC, Takken FLW. Transcomplementation, but not physical association of the CC-NB-ARC and LRR domains of tomato R protein Mi-1.2 is altered by mutations in the ARC2 subdomain. *Mol Plant*. 2008; 1: 3. Available: <http://www.sciencedirect.com/science/article/pii/S1674205214603373>.