

CORRECTION

Correction: Are two internal thoracic grafts better than one in patients with chronic obstructive lung disease? Analysis of 387 cases between 1996-2011

Dmitry Pevni, Zahi Aizer, Rephael Mohr, Nahum Neshner, Amir Kremer, Yosef Paz, Nadav Taih, Yanai Ben-Gal

There are multiple errors in the author byline. Please view the correct author order, affiliations, and citation here:


Dmitry Pevni^{1,2}, Zahi Aizer², Rephael Mohr^{1,2}, Nahum Neshner^{1,2}, Amir Kremer^{1,2}, Yosef Paz^{1,2}, Nadav Taih^{1,2}, Yanai Ben-Gal^{1,2}

1 Department of Cardiothoracic Surgery Tel Aviv Sourasky Medical Center, Tel Aviv University, Tel Aviv, Israel, 2 Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

Pevni D, Aizer Z, Mohr R, Neshner N, Kremer A, Paz Y, et al. (2018) Are two internal thoracic grafts better than one in patients with chronic obstructive lung disease? Analysis of 387 cases between 1996–2011. PLoS ONE 13(8): e0201227. <https://doi.org/10.1371/journal.pone.0201227>.

Reference

1. Dmitry P, Zahi A, Rephael M, Nahum N, Amir K, Yosef P, et al. (2018) Are two internal thoracic grafts better than one in patients with chronic obstructive lung disease? Analysis of 387 cases between 1996–2011. PLoS ONE 13(8): e0201227. <https://doi.org/10.1371/journal.pone.0201227> PMID: 30102699



OPEN ACCESS

Citation: Pevni D, Aizer Z, Mohr R, Neshner N, Kremer A, Paz Y, et al. (2018) Correction: Are two internal thoracic grafts better than one in patients with chronic obstructive lung disease? Analysis of 387 cases between 1996-2011. PLoS ONE 13(11): e0208046. <https://doi.org/10.1371/journal.pone.0208046>

Published: November 19, 2018

Copyright: © 2018 Pevni et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.