



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Dataset on Catal's reagent: Sensitive detection of iron (II) sulfate using spectrophotometry

Funda Ozkok^{a,†}, Yesim Muge Sahin^{b,c,†}, Vildan Enisoglu Atalay^d, Kamala Asgarova^a, Nihal Onul^a, Tunc Catal^{d,e,*}^a Department of Chemistry, Istanbul University-Cerrahpasa, Avcilar, Istanbul, Turkey^b Department of Biomedical Engineering, Istanbul Arel University Turkey^c Polymer Technologies and Composite Application and Research Center (ArelPOTKAM), Istanbul Arel University Buyukcekmece, Istanbul, Turkey^d Istanbul Protein Research Application and Innovation Center (PROMER)^e Department of Molecular Biology and Genetics, Uskudar University 34662 Uskudar, Istanbul, Turkey

ARTICLE INFO

Article history:

Received 30 June 2020

Revised 25 July 2020

Accepted 4 August 2020

Available online 8 August 2020

Keywords:

Catal's reagent

Iron (II) sulfate

Spectrophotometer

1-(Dodecylthio)anthracene-9,10-dione

Thiols

ABSTRACT

Catal's reagent is characterized by spectroscopic methods such as fourier-transform infrared spectroscopy (FT-IR), nuclear magnetic resonance (NMR) spectroscopy, mass spectrometry (MS), ultraviolet (UV)-visible spectrophotometry. Effects of different solvents such as methanol and ethanol on absorption spectrum of 1-(Dodecylthio)anthracene-9,10-dione (**3**) were present. Detection range of iron (II) sulfate using Catal's reagent was analyzed. Synthesis of 1-(Dodecylthio)anthracene-9,10-dione (**3**) was explained, and absorbances of various concentrations of iron (II) sulfate (0-10 mg mL⁻¹) were measured. The possible detection mechanism was also explained. The dataset is useful to improve the detection of iron (II) sulfate in various application fields such as environmental, agricultural, sensor, food, textile and cement industries.

The study refers to: F. Ozkok, Y.M. Sahin, V. Enisoglu-Atalay, K. Asgarova, N. Onul, T. Catal, Sensitive Detection of Iron (II) Sulfate with a Novel Reagent using Spectrophotometry,

DOI of original article: [10.1016/j.saa.2020.118631](https://doi.org/10.1016/j.saa.2020.118631)

* Corresponding author.

E-mail addresses: ozkok@istanbul.edu.tr (F. Ozkok), ymugesahin@arel.edu.tr (Y.M. Sahin), tunc.catal@uskudar.edu.tr (T. Catal).

† Authors equally contributed to the study.

<https://doi.org/10.1016/j.dib.2020.106149>2352-3409/© 2020 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license. (<http://creativecommons.org/licenses/by/4.0/>)

Spectchim. Acta. A, 240 (2020), 118631. <https://doi.org/10.1016/j.saa.2020.118631>.

© 2020 The Author(s). Published by Elsevier Inc.
This is an open access article under the CC BY license.
(<http://creativecommons.org/licenses/by/4.0/>)

Specifications table

Subject	Chemistry
Specific subject area	Analytical chemistry
Type of data	Table and figure
How data were acquired	The data were acquired: FT-IR, NMR, mass spectrometry, UV-vis spectrophotometry
Data format	Raw and Analyzed
Parameters for data collection	1-(Dodecylthio)anthracene-9,10-dione was synthesized in the laboratory.
Description of data collection	Catal's reagent was prepared and examined using traditional methods. The data were collected after confirmation of the structure of 1-(Dodecylthio)anthracene-9,10-dione. UV-vis spectrophotometer, FT-IR, 1H-NMR, 13C-NMR, mass spectrometer were used in the data collection.
Data accessibility	With the article
Related research article	F. Ozkok, Y.M. Sahin, V. Enisoglu-Atalay, K. Asgarova, N. Onul, T. Catal, Sensitive Detection of Iron (II) Sulfate with a Novel Reagent using Spectrophotometry, Spectrochim. Acta. A, 240 (2020), 118631. https://doi.org/10.1016/j.saa.2020.118631 .

Value of the data

- A database of Catal's reagent is essential for characterization of 1-(Dodecylthio)anthracene-9,10-dione (3)
- The data are key for examining iron (II) sulfate in various samples.
- These data are an important reference source for research on developing novel studies to use Catal's reagent.

1. Data description

This research reports on a Catal's reagent data set for detection of iron (II) sulfate. Fig. 1 shows absorption spectra of Catal's reagent in methanol (A) and ethanol (B) solution. Fig. 2 shows absorption spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3) in acetonitrile solution with lower scan rate. Fig. 3 shows FT-IR spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3). Fig. 4 shows 1H-NMR spectra of 1-(Dodecylthio)anthracene-9,10-dione (3). Fig. 5 shows 13C-NMR spectra of 1-(Dodecylthio)anthracene-9,10-dione (3). Fig. 6 shows MS spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3). Fig. 7 shows synthesis of 1-(Dodecylthio)anthracene-9,10-dione (3). Fig. 8 shows oxidation of iron in presence of hydrogen peroxide (Fenton Reaction). Fig. 9 shows electrochemical redox reaction of anthraquinones. Fig. 10 shows oxidation of anthraquinone derivative in the presence of hydrogen peroxide. Fig. 11 shows absorbances of various concentrations of iron (II) sulfate. Table 1 shows absorbances of several compounds at the concentration of 10 mg mL⁻¹ in distilled water.

2. Experimental design, materials and methods

Novel thio anthraquinone derivative (1-(Dodecylthio)anthracene-9,10-dione) was synthesized by this novel method for scientific applications [1]. Chemical structure of novel thio anthraquinone compound (3) was characterized by spectroscopic methods such as FT-IR, NMR, MS,

Table 1

Absorbances of several compounds at the concentration of 10 mg mL⁻¹ in distilled water. The examined molecules were not reacted with Catal's reagent under the examined conditions.

Compound	Absorbance (304 nm)
Ammonium persulphate	-0.032
Ammonium sulphate	0.031
Aluminum potassium sulfate dodecahydrate	-0.003
Copper (III) sulphate	-0.008
Copper (III) sulphate pentahydrate	-0.056
Sodium dodecyl sulphate	-0.018
Sodium sulphate anhydrous	-0.022
Sodium thiosulphate pentahydrate	-0.002
Manganese (II) sulphate monohydrate	0.039
Magnesium sulphate heptahydrate	0.027
Sodium 2-bromoethanesulfonate hydrate	-0.029
Sulfanilic acid	0.033
Sodium sulfite anhydrous	0.001
Zinc sulphate heptahydrate	0.006
Sodium 2-chloroethane sulfonate hydrate	0.028
Potassium sulphate	0.037
Ammonium iron (III) citrate	0.012
Iron (III) citrate hydrate	0.038
Iron (II) sulfate heptahydrate	0.985

(UV)-visible spectrophotometry, and the structure of the compound was confirmed. The thio anthraquinone derivative, 1-(Dodecylthio)anthracene-9,10-dione (**3**), was dissolved in the following organic solvents to prepare the reactant named as Catal's reagent: Ethanol and methanol (Fig. 1). Tetra JASCO 6600 spectrometer used for fourier transform infrared (FT-IR) spectra, and Tetra JASCO V 750 spectrometer recorded Ultraviolet-visible spectra. A Varian UNITY INOVA at 500 MHz was used for ¹H NMR and ¹³C NMR spectra. Mass spectra was recorded on (Shimadzu, Kyoto-Japan) LCMS-8030 triple quadrupole spectrometer in ESI (+) polarity. The absorbances were measured at 304 nm of wavelength using a UV-visible spectrophotometer in air (Shimadzu UV-2600, Cat. No. 206-27600-45, Kyoto, Japan). The reaction mixture was prepared as follow; 1-(Dodecylthio)anthracene-9,10-dione (**3**) (20 mg) was added to either ethanol, methanol or acetonitrile (60 mL) in order to prepare Catal's reagent. Catal's reagent (50 μL) was mixed with iron (II) sulfate solution (100 μL). Finally, H₂O₂ solution (17.5 percent in distilled water, v:v) (50 μL) was added to the mixture. 30 mM of trisodium citrate dihydrate (9 mL) was then used to stabilize pH changes. In advance, different rations of the compounds could be used to enhance the sensitivity of the reaction with Catal's reagent. Catal's reagent can be used for spectrophotometric and colorimetric detection of iron (II) sulfate [4].




Fig. 1. Absorption spectra of Catal's reagent in methanol (A) and ethanol (B) solution. Three similar peaks were identified.




Fig. 2. Absorption spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3), in acetonitrile solution in lower scan rate.






Fig. 3. FT-IR spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3).



INDEX	FREQUENCY PPM	HEIGHT	INDEX	FREQUENCY PPM	HEIGHT
1	8234.437	12.220	2.1	3805.228	7.614
2	8225.709	12.514	88.8	3805.244	7.605
3	8180.455	11.287	8.1	3798.250	7.599
4	8124.891	8.254	7.3	3792.537	7.589
5	8123.251	8.252	89.5	3785.211	7.583
6	8114.498	8.240	114.0	3780.193	7.576
7	8114.226	8.239	89.5	3777.896	7.560
8	8114.442	8.234	114.0	3770.098	7.544
9	8107.809	8.220	246.2	3671.332	7.239
10	8104.346	8.217	232.0	3616.356	7.237
11	8104.881	8.214	20.2	3611.944	7.238
12	8104.441	8.209	27.6	3609.035	7.232
13	8098.537	8.201	4.0	3609.059	7.230
14	8097.561	8.199	2.2	3605.607	7.212
15	8094.354	8.189	32.6	3602.203	7.208
16	8090.929	8.180	9.1	3593.456	7.187
17	8094.120	8.113	11.4	3143.999	3.289
18	8093.149	8.112	13.2	3143.102	3.282
19	8079.468	7.743	24.3	289.061	1.179
20	8077.944	7.740	23.9	608.487	0.817
21	8071.440	7.737	43.0	602.143	0.805
22	8071.020	7.733	32.6	384.284	0.793
23	8071.989	7.747	43.0	389.454	0.779
24	8071.623	7.743	50.6	3.414	0.007
25	8069.150	7.742	89.8	0.000	0.000
26	8064.231	7.737	44.2	-1.004	-0.008
27	8064.279	7.733	111.0		
28	8064.815	7.730	44.2		
29	8060.843	7.724	80.3		
30	8059.864	7.724	80.3		
31	8058.422	7.721	80.3		
32	8058.484	7.714	80.4		
33	8053.084	7.710	13.9		
34	8052.078	7.708	18.6		
35	8050.614	7.705	7.3		
36	8049.643	7.703	3.3		

Fig. 4. $^1\text{H-NMR}$ spectra of 1-(Dodecylthio)anthracene-9,10-dione (3).




Fig. 5. ¹³C-NMR spectra of 1-(Dodecylthio)anthracene-9,10-dione (3).




Fig. 6. MS spectrum of 1-(Dodecylthio)anthracene-9,10-dione (3).




Fig. 7. Synthesis of 1-(Dodecylthio)anthracene-9,10-dione (3).




Fig. 8. Oxidation of iron in presence of hydrogen peroxide (Fenton Reaction) [2].




Fig. 11. Absorbances of various concentrations of iron (II) sulfate. R^2 values were over 0.9.

Declaration of Competing Interest

The authors (F. Ozkok, Y.M. Sahin, V. Enisoglu-Atalay, K. Asgarova, N. Onul, T. Catal) declare patent application (Turkish Patent and Trademark Office, PY2019-00552; PCT International Application, No: PCT/TR2020/050061– submitted).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.dib.2020.106149](https://doi.org/10.1016/j.dib.2020.106149).

References

- [1] F. Ozkok, Y. M. Şahin, Biyoaktif Antrakinon Anologlarının Sentezine Yönelik Özgün Metot Geliştirilmesi, TÜRKİYE, Patent, TR 2016/19610.
- [2] H.J.H. Fenton, Oxidation of tartaric acid in presence of iron, *J. Chem. Soc. Trans.* 65 (1894).
- [3] B. Yang, A. Murali, A. Nirmalchandar, B. Jayathilake, G.K.S. Prakash, S.R. Narayanan, A Durable, inexpensive and scalable redox flow battery based on iron sulfate and anthraquinone disulfonic acid, *J. Electrochem. Soc.* 167 (2020) 060520.
- [4] F. Ozkok, Y.M. Sahin, V. Enisoglu-Atalay, K. Asgarova, N. Onul, T. Catal, Sensitive detection of iron (ii) sulfate with a novel reagent using spectrophotometry, *Spectrochim. Acta. A* 240 (2020) 118631 <https://doi.org/10.1016/j.saa.2020.118631>.