



# Genome Sequence of Historical *Bacillus anthracis* Strain Tyrol 4675 Isolated from a Bovine Anthrax Case in Austria

Markus Antwerpen, Roman Wölfel, Gregor Grass

Bundeswehr Institute of Microbiology, Munich, Germany

**ABSTRACT** In 1988, an outbreak of anthrax occurred among cattle in the Austrian state of Tyrol. Since then, Austria has been declared anthrax-free. Here, we report the draft genome sequence of one of these last outbreak strains, *Bacillus anthracis* Tyrol 4675, isolated from a diseased cow.

The last outbreak of the zoonotic disease anthrax in Austria occurred in the state of Tyrol in 1988. *Bacillus anthracis* strain Tyrol 4675 was isolated from one of the diseased cows (1, 2). Strain Tyrol 4675 harbors both *B. anthracis* virulence plasmids pXO1 and pXO2, as confirmed by real-time PCR assays (1–4). Initial genotyping using canonical single-nucleotide polymorphisms (canSNP) (5) grouped strain Tyrol 4675 to branch B.Br. CNEVA (B2) (5). More specifically, it positions in subbranch B.Br.005/004 (6). This is the same subbranch as strain BF-1 (6, 7), isolated from the Bavarian northern foothills of the Alps neighboring Tyrol. Strains from this clade have been reported before from other central and western European countries, such as Germany or France, and this genotype appears to be autochthonous to this geographic region (5, 7, 8). Strain Tyrol 4675 represents the first representative genome sequence from the country of Austria.

Whole-genome shotgun (WGS) sequencing of *B. anthracis* Tyrol 4675 was performed by Illumina MiSeq sequencing technology (Illumina, Inc.) using Nextera version 3 with 2 × 300-bp chemistry. For the WGS library, 2 × 813,272 reads accumulating to a total of 464 Mb were generated. Burrows-Wheeler Aligner's Smith-Waterman Alignment (BWA-SW) (9) was used for mapping to the *B. anthracis* Ames Ancestor chromosome and plasmids pXO1 and pXO2 (accession numbers NC\_007530.2, NC\_007322.2, and AE017335.3), respectively. The G+C content was calculated using an in-house python script.

The total length of the genome draft sequence of *B. anthracis* Tyrol 4675 was 5,227,565 bp, with 62.3-fold coverage for the chromosome, 145-fold for pXO1, and 102-fold for pXO2, respectively. The mean G+C content was 35.4%. For initial annotation, assembled contigs were submitted to the RAST Annotation Pipeline RASTk (10). The *B. anthracis* Tyrol 4675 draft genome contains 5,693 putative coding sequences (CDS). Within the genome annotation, there are 11 copies of genes for the 16S rRNA, the 5S rRNAs, and the 23S rRNA; 95 tRNA loci were identified.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the GenBank accession numbers [CP018904](#) (pXO1), [CP018905](#) (pXO2), and [CP018903](#) (chromosome). The versions described in this paper are the first versions.

## ACKNOWLEDGMENTS

We thank Franz Allerberger, Austrian Agency for Health and Food Safety (Vienna, Austria), for the generous gift of strain *B. anthracis* Tyrol 4675. We also thank Linda Dobrzykowski and Claudia Kahlhofer for skillful technical assistance.

Received 3 January 2017 Accepted 11 January 2017 Published 9 March 2017

**Citation** Antwerpen M, Wölfel R, Grass G. 2017. Genome sequence of historical *Bacillus anthracis* strain Tyrol 4675 isolated from a bovine anthrax case in Austria. *Genome Announc* 5:e00002-17. <https://doi.org/10.1128/genomeA.00002-17>.

**Copyright** © 2017 Antwerpen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Gregor Grass, [gregorgrass@bundeswehr.org](mailto:gregorgrass@bundeswehr.org).

## REFERENCES

1. Grif K, Dierich MP, Much P, Hofer E, Allerberger F. 2003. Identifying and subtyping species of dangerous pathogens by automated ribotyping. *Diagn Microbiol Infect Dis* 47:313–320. [https://doi.org/10.1016/S0732-8893\(03\)00095-6](https://doi.org/10.1016/S0732-8893(03)00095-6).
2. Allerberger F, Grif K, Dierich MP, Wimmer A, Plicka H. 2002. Anthrax inhalation and lethal human infection. *Lancet* 359:711. [https://doi.org/10.1016/S0140-6736\(02\)07793-0](https://doi.org/10.1016/S0140-6736(02)07793-0).
3. Ellerbrok H, Nattermann H, Ozel M, Beutin L, Appel B, Pauli G. 2002. Rapid and sensitive identification of pathogenic and apathogenic *Bacillus anthracis* by real-time PCR. *FEMS Microbiol Lett* 214:51–59. <https://doi.org/10.1111/j.1574-6968.2002.tb11324.x>.
4. Antwerpen MH, Zimmermann P, Bewley K, Frangoulidis D, Meyer H. 2008. Real-time PCR system targeting a chromosomal marker specific for *Bacillus anthracis*. *Mol Cell Probes* 22:313–315. <https://doi.org/10.1016/j.mcp.2008.06.001>.
5. Van Ert MN, Easterday WR, Huynh LY, Okinaka RT, Hugh-Jones ME, Ravel J, Zanecki SR, Pearson T, Simonson TS, U'Ren JM, Kachur SM, Leadem-Dougherty RR, Rhoton SD, Zinser G, Farlow J, Coker PR, Smith KL, Wang B, Kenefic LJ, Fraser-Liggett CM, Wagner DM, Keim P. 2007. Global genetic population structure of *Bacillus anthracis*. *PLoS One* 2:e461. <https://doi.org/10.1371/journal.pone.0000461>.
6. Sahl JW, Pearson T, Okinaka R, Schupp JM, Gillece JD, Heaton H, Birdsell D, Hepp C, Fofanov V, Nosedá R, Fasanella A, Hoffmaster A, Wagner DM, Keim P. 2016. A *Bacillus anthracis* genome sequence from the Sverdlovsk 1979 autopsy specimens. *mBio* 7:e01501-16. <https://doi.org/10.1128/mBio.01501-16>.
7. Antwerpen M, Proença DN, Rückert C, Licht K, Kalinowski J, Hanczaruk M, Tiemann C, Grass G. 2012. Draft genome sequence of *Bacillus anthracis* BF-1, Isolated from Bavarian cattle. *J Bacteriol* 194:6360–6361. <https://doi.org/10.1128/JB.01676-12>.
8. Derzelle S, Laroche S, Le Flèche P, Hauck Y, Thierry S, Vergnaud G, Madani N. 2011. Characterization of genetic diversity of *Bacillus anthracis* in France by using high-resolution melting assays and multilocus variable-number tandem-repeat analysis. *J Clin Microbiol* 49:4286–4292. <https://doi.org/10.1128/JCM.05439-11>.
9. Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
10. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.