



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijidINTERNATIONAL
SOCIETY
FOR INFECTIOUS
DISEASES

Short Communication

Silent hypoxia in patients with SARS CoV-2 infection before hospital discharge



Natascha Josephine Ulstrand Fuglebjerg, Tomas Oestergaard Jensen, Nils Hoyer, Camilla Koch Ryrso, Birgitte Lindegaard, Zitta Barrella Harboe*

Department of Infectious Diseases and Pulmonary Medicine, Nordsjællands University Hospital, Denmark

ARTICLE INFO

Article history:

Received 4 June 2020

Received in revised form 6 July 2020

Accepted 9 July 2020

Keywords:

COVID-19
Hypoxia
Dyspnea
Exercise test
Discharge

ABSTRACT

Objective: To assess the degree of hypoxia and subjective dyspnea elicited by a 6-minute walking test (6MWT) in COVID-19 patients prior to discharge.**Methods:** A 6MWT was performed in 26 discharge-ready COVID-19 patients without chronic pulmonary disease or cardiac failure. Heart rate, oxyhemoglobin saturation (SpO₂), respiratory rate, and subjective dyspnea measured on the Borg CR-10 scale were measured before and immediately after the 6MWT, with continuous monitoring of SpO₂ and heart rate during the 6MWT. The 6MWT was terminated if SpO₂ dropped below 90%. A historical cohort of 204 patients with idiopathic pulmonary fibrosis (IPF) was used for comparison.**Results:** 13 (50%) of the COVID-19 patients developed exercise-induced hypoxia (SpO₂ < 90%) during the 6MWT, of which one third had pulmonary embolism. COVID-19 patients experienced less hypoxia-related dyspnea during the 6MWT compared with patients with IPF.**Conclusion:** The 6MWT is a potential tool in the diagnosis of asymptomatic exercise-induced hypoxia in hospitalized COVID-19 patients prior to discharge. Due to important methodological limitations, further studies are needed to confirm our findings and to investigate their clinical consequences.© 2020 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Manuscript

Anecdotal evidence is emerging that patients with COVID-19 may present with 'silent hypoxia', which refers to a discrepancy between the level of hypoxia and subjective dyspnea experienced by the patient (Couzín-Frankel, 2020). This phenomenon is of potential concern among patients with chronically compromised end-organ perfusion, for example in ischemic heart disease, where the absence of dyspnea may permit an acute ischemic event to progress to a critical level without the usual early warning symptoms. We aimed to investigate whether silent hypoxia in adult COVID-19 patients ready for discharge can be elicited by a 6-minute walking test (6MWT).

Patients hospitalized with COVID-19 confirmed by polymerase chain reaction (PCR) testing at the University Hospital of

Copenhagen, North Zealand, Denmark were eligible for inclusion at the time of planned discharge, which required resolution of fever and an oxyhemoglobin saturation (SpO₂) of 94% or above at rest, without supplementary oxygen. Patients with chronic lung diseases and New York Heart Association (NYHA) class II and above were excluded. The 6MWT was carried out according to the American Thoracic Society (ATS) statement from 2002 by trained pulmonary physiotherapists (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002). Patients were instructed to walk the greatest distance possible in 6 min, at a self-determined pace, pausing to rest if needed. Heart rate (HR), SpO₂, respiratory rate (RR), and dyspnoea rated between 1 and 10 on the Borg CR-10 dyspnea scale (Borg and Kaijser, 2006; Borg, 1982) were recorded before and immediately after the 6MWT. A PaO₂ below 80 mmHg (10.6 kPa) is generally considered pathological, while a PaO₂ of at least 60 mmHg (8 kPa), which correlates to an SpO₂ of 90%, is required to maintain sufficient tissue oxygenation (Hansen et al., 1984). SpO₂ and HR were continuously monitored during the 6MWT. We assumed that patients with no history of chronic lung disease should be able to maintain an SpO₂ of 96% or above, even during physical activity

* Corresponding author: Department of Infectious Diseases and Pulmonary Medicine, University Hospital of Copenhagen, North Zealand, Denmark, Durehavevej 29, 3400 Denmark.

E-mail address: zitta.barrella.harboe@regionh.dk (Z.B. Harboe).

Table 1

Baseline characteristics of 26 patients with COVID-19 undergoing the 6-minute walking test (6MWT) at hospital discharge and a historical cohort of 204 patients with idiopathic pulmonary fibrosis who underwent the same test.

	Total number of patients (n = 26)	COVID-19 patients		IPF patients (n = 204)
		6-MWT completed (n = 13)	6-MWT terminated early due to SpO ₂ < 90% (n = 13)	
Age in years, median (range)	63 (29–85)	62 (29–85)	64 (38–76)	74 (42–92)
Male sex, n (%)	16	8 (62)	8 (62)	158 (77)
Number of comorbidities per patient, median (range)	1 (1–5)	1 (1–5)	1 (1–4)	Unknown
Need for ICU during admission, n (%)	8 (31)	2 (15)	6 (46)	Unknown
Need for mechanical ventilation during admission, n (%)	2 (15)	0 (0)	2 (15)	Unknown
Infiltrate on chest X-ray during admission, n (%)	21 (81)	8 (62)	13 (100)	Unknown




Figure 1. Decline in oxygen saturation (SpO₂) and Borg CR-10 dyspnea scale rating after 6MWT in patients with COVID-19 (blue) and idiopathic pulmonary fibrosis (red). Each dot represents one patient. The lines are based on univariate linear regression.

(Casanova et al., 2011). A previous prospective cohort of 204 patients with idiopathic pulmonary fibrosis (IPF), who also underwent the 6MWT according to the ATS guidelines, was used for comparison (Hoyer et al., 2019).

Forty-seven consecutive patients were screened between April 16 and April 28, 2020, and all 26 eligible patients without exclusion criteria were included. The 6MWT was terminated early in 13 patients (50%) due to an SpO₂ below 90%. Baseline characteristics in patients who terminated the 6MWT early compared with those who completed the 6MWT, as well as baseline characteristics for the comparison group, are presented in Table 1. COVID-19 patients with an SpO₂ below 90% during the 6MWT had only discrete increase in perceived dyspnea (Figure 1). Six patients who terminated the 6MWT early due to hypoxia were investigated with CT angiography and/or ventilation-perfusion scintigraphy in order to exclude pulmonary embolism (PE), which was confirmed in four patients. As a comparison group, IPF patients had a similar decline in SpO₂, but they developed more pronounced concomitant worsening in the level of subjective dyspnea measured by the Borg scale (Figure 1). COVID-19 patients had a mean peak score of 2.5 on Borg CR-10 (range 0–5) during the 6MWT, compared with 4.1 (range 0–10) in IPF patients.

In summary, nearly half of hospitalized COVID-19 patients without chronic pulmonary disease developed marked exercise-induced hypoxia without accompanying subjective dyspnea at the time of discharge, and one third of patients who terminated the 6MWT early had PE. We interpret this as evidence of silent hypoxia.

Our study had several limitations. No previous studies have examined the role of the 6MWT in otherwise healthy adults with COVID-19 or other acute respiratory tract infections, although exercise-induced hypoxia has been used in the diagnosis and severity assessment of *Pneumocystis jirovecii* pneumonia in

HIV-positive adults (Sauleda et al., 1994). Clinical implications of this hypoxia are thus not well described, and we therefore decided to routinely terminate the 6MWT early if SpO₂ dropped below 90% for safety reasons. This could have led to underestimation of the relative degree of dyspnea in COVID-19 patients, since they were compared with a control group of IPF patients, who were all allowed to complete the 6MWT unless they developed chest pain, intolerable dyspnea, leg cramps, staggering, diaphoresis, or pale or ashen appearance, in accordance with the ATS guidelines. Furthermore, patients in our control group all had IPF, a condition resulting in hypoxic respiratory failure, and therefore increased tolerance to lower SpO₂ without subjective dyspnea was a possibility in this group.

We propose further, larger studies to confirm our findings using a more accurately matched prospective group of controls, and to investigate the clinical consequences of silent hypoxia in particular among patients at elevated risk of serious end-organ ischemic events after discharge to the community.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval

The study was approved by the Danish Health and Medicines Authority (ID: 31–1521–264) and by the Danish Data Protection Agency (P-2020-375).

References

- ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: Guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;166(1):111–7. doi:http://dx.doi.org/10.1164/ajrccm.166.1.at1102.
- Borg E, Kaijser L. A comparison between three rating scales for perceived exertion and two different work tests. *Scand J Med Sci Sport* 2006;16(1):57–69. doi: http://dx.doi.org/10.1111/j.1600-0838.2005.00448.x.
- Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982;14(5):377–81.
- Casanova C, Celli BR, Barria P, Casas A, Cote C, de Torres JP, et al. Six Minute Walk Distance Project (ALAT). The 6-min walk distance in healthy subjects: Reference standards from seven countries. *Eur Respir J* 2011;37:150–6. doi:http://dx.doi.org/10.1183/09031936.00194909.
- Couzin-Frankel J. The mystery of the pandemic's 'happy hypoxia'. *Science* 2020;368(6490):455–6. doi:http://dx.doi.org/10.1126/science.368.6490.455.
- Hansen JE, Sue DY, Wasserman K. Predicted Values for Clinical Exercise Testing 1–3. *Am Rev Respir Dis* 1984;129(2P2):S49–55. doi:http://dx.doi.org/10.1164/arrd.1984.129.2P2.S49.
- Hoyer N, Prior TS, Bendstrup E, Wilcke T, Shaker SB. Risk factors for diagnostic delay in idiopathic pulmonary fibrosis. *Respir Res* 2019;20(1):103. doi:http://dx.doi.org/10.1186/s12931-019-1076-0.
- Sauleda J, Gea J, Aran X, Aguar MC, Orozco-Levi M, Broquetas JM. Simplified exercise test for the initial differential diagnosis of *Pneumocystis carinii* pneumonia in HIV antibody positive patients. *Thorax* 1994;49(2):112–4. doi:http://dx.doi.org/10.1136/thx.49.2.112.