




Utilization of Integrated Community Case Management of Childhood Illnesses at Health Posts in Southern Ethiopia

This article was published in the following Dove Press journal:
Pediatric Health, Medicine and Therapeutics

Asefa Berhanu¹
Mihiretu Alemayehu ²
Kassa Daka²
Wakgari Binu ²
Mohammed Suleiman ³

¹Wolaita Zone Health Department, Wolaita Sodo, Ethiopia; ²Wolaita Sodo University, College of Health Sciences and Medicine, School of Public Health, Wolaita Sodo, Ethiopia; ³Wolaita Sodo University, College of Health Sciences and Medicine, Department of Anesthesia, Wolaita Sodo, Ethiopia

Purpose: Even if strategic actions were undertaken to tackle common childhood illnesses, Ethiopia still stood as one of the six countries accounting for half of the global under-five deaths. So this study aimed to assess the utilization of integrated community case management during childhood illness at health posts in the study area.

Methods: A community-based cross-sectional study was conducted using a multistage sampling technique that includes 633 sick under-five children with common childhood illnesses. The data was entered into EpiData 3.1 and exported to SPSS 20 for further analysis. Descriptive statistics, bivariate and multivariate logistic regression analyses were computed and adjusted odds ratio within 95% confidence interval was used to measure the statistical association between variables.

Results: Out of the planned 633 participant, 624 caregivers underwent all the study components making the response rate 98.6%. The study indicated that only 10.6% of the sick children sought care from health posts. Income (AOR = 2.99, 95% CI: 1.37–6.53), previous service utilization (AOR = 6.66, 95% CI: 1.81–24.04), awareness of service availability (AOR = 4.74, 95% CI: 1.39–12.10), ownership of health insurance (AOR = 2.63, 95% CI: 1.45–4.76), distance (AOR = 5.23, 95% CI: 1.69–10.19) and type of illness (AOR = 2.97, 95% CI: 1.41–6.25) were the associated factors.

Conclusion: The low utilization of integrated community case management insights to focus on availing additional well-equipped health posts, creating community awareness on services at health posts and at what time the services are given to address the problem of the majority.

Keywords: ICCM, health post utilization, fever, diarrhea, cough, Kindo Didaye

Plain Language Summary

Evidence showed that Ethiopia still stood as one of the six countries accounting for half of the global under-five deaths. Even if Ethiopia had initiated the integrated community case management (ICCM) program a long time ago, there is a scarcity of data dealing with ICCM utilization. Available studies ignored remote areas at which the community is suffering from poor infrastructures due to difficult topography like our study area.

We conducted the study in one of the harsh areas in southern Ethiopia, Kindo Didaye district, which is highly mountainous and hard to reach, with poor infrastructures. We used a multistage sampling strategy to recruit the required sample size. We conducted a survey using identification panel questions in six randomly selected kebeles (the smallest administrative unit in Ethiopia) to identify households (HHs) with under-five sick children. Then we recruited proportionally allocated HHs fulfilling criteria until we obtained the sample size using random sampling technique.

Correspondence: Wakgari Binu
Wolaita Sodo University, College of
Health Sciences and Medicine, School of
Public Health, P.o. Box: 138, Wolaita
Sodo, Ethiopia
Tel +251 910107704
Email nubonsa@gmail.com

EpiData v3.1 and SPSS v20 were used for data management. Bivariate and multivariate logistic regression analysis were computed to control confounders and to identify predictors of ICCM utilization. The overall ICCM utilization at health posts (HPs) in this study was very low while childhood illness was prevalent in the area. This study identified factors influencing the service utilization among participants. The community need to be continually aware of the time when health posts are open for health services. Also providing additional equipped HPs to address the problem of the majority is the cutting issue. Even if limited number of studies reported suboptimal use of HPs for ICCM in other parts of Ethiopia, this study will provide recent evidence on the issue for health managers and stakeholders.

Introduction

Globally under-five deaths dropped to 5.3 million in 2018 from 12.6 million in 1990. Despite exhaustive progress made, Sub-Saharan Africa continues to be the region with the highest child mortality rate in the world, 78 deaths per 1000 live births mostly from preventable causes such as pneumonia, diarrhea, and malaria.¹

Even though Ethiopia has significantly reduced under-five child deaths from 204 per 1000 live births in 1990 to 59 per 1000 live births in 2017, 518 children are dying every day, mostly of preventable or treatable diseases such as diarrhea, malaria, and pneumonia. Also, Ethiopia is one of the six countries (India, Nigeria, Pakistan, the Democratic Republic of the Congo, Ethiopia, and China) accounting for half of the global under-five deaths.^{1,2}

Countries identified integrated community case management (ICCM) as an effective evidence-based strategy to increase coverage of lifesaving interventions and to reduce preventable child deaths where access to health services is limited. The aim of ICCM strategy was to improve access to treatment of diarrhea, fever and pneumonia by providing evidence-based free treatments to the largely rural population, where there is a lack of access to private clinics, public health centers, and higher-level facilities.³⁻⁵

In response to the high child mortality, the government of Ethiopia had introduced the implementation of ICCM in 2010 after the health policy update.³ The government has been implementing ICCM strategy through the platform of the Health Extension Program (HEP). The government initiated HEP to scale up the essential health services by reaching the people at the grassroots level in particular to the underserved rural population. It was designed based on the concept and principles of comprehensive primary

health care.^{4,5} In this program, health extension workers (HEWs) were trained to assess, classify and treat common childhood illnesses like diarrhea, malaria, pneumonia and severe acute malnutrition among under-five children. In this regard, HEWs were providing the service in rural communities at HPs since 2010 to reduce under-five mortality to 29 per 1000 live births by 2020.⁶⁻⁸

The government of Ethiopia has also made a series of efforts like an integration of ICCM into health policy, scale-up of ICCM implementation, and removing the ICCM utilization barriers being with different local and global development partners (Non-Government Organizations) to reduce under-five mortality. Despite these efforts made, health facility-based evidence showed a low level of ICCM utilization among under-five sick children in the country.⁹⁻¹²

Even if the ICCM program was initiated a long time ago in the country, there is a scarcity of data in the study area that dealt with ICCM utilization. Available evidence ignored remote areas at which the community is suffering from poor infrastructures due to difficult topography like our study area. Therefore, this study has assessed the ICCM utilization and influencing factors. The findings will provide evidence-based information to regional and local health managers, and health extension workers. The findings will also provide recent evidence for concerned bodies to strengthen community-based interventions and minimize the barriers of ICCM utilization at HPs. So the utilization of the finding helps to reduce child morbidity and mortality in the community.

Materials and Methods

Study Setting and Design

A community-based cross-sectional study was conducted from May 1st to 21st, 2019 in Kindo Didaye district, Wolaita zone, southern Ethiopia. The district is one of the 16 districts and 6 urban administrative towns in the Wolaita zone, located at 475 kilometers from Addis, capital of Ethiopia. The district is highly mountainous and the hardest to reach, with poor infrastructure areas in the zone. There are 22 Kebeles in the district, of which 19 are rural kebeles and 3 are semi-urban kebeles with an estimated population of 129,246 in 2017 as projected from the 2007 census. The number of under-five aged children is 20,162 accounting for 15.6% of the total population.

Regarding a health facility, there are 25 HPs each with two HEWs implementing the 17 Ethiopian health extension

packages including ICCM and CBNC, 3 health centers, 1 primary hospital and 4 medium private clinics serving the community with a range of health services in the district (Kindo Didaye administration annual report, 2018).

Population and Sampling

We had included caregivers of children aged 0–59 months having an episode of coughing or fever or diarrhea during the two weeks preceding the survey. We had excluded under-five sick children whose caregivers were unable to hear and speak from the study.

The sample size was estimated using a single population formula based on the survey done by L10K in South Nations, Nationalities and People's Regional State of Ethiopia. We have considered diarrhea, fever and pneumonia to determine the sample size. We used the proportion of children with diarrhea-sought care from an appropriate provider (48.28%) as it has given a maximum sample size.¹³ Considering 5% marginal error at 95% confidence level, design effect of 1.5 (accounting for correlation of observations among kebeles) and 10% compensation for non-response giving 633 caregivers were planned to be interviewed.

Then a multistage sampling strategy used to recruit the representative sample of respondents. At the first stage, out of the 22 kebeles (smallest administrative unit), 6 kebeles were selected randomly. In the second stage, we conducted a survey using HH identification panel questions [Table S1] before actual data collection to identify the total number of HHs in each selected kebele. The aim of the survey was to identify total HHs with under-five sick children. We had given specific codes (i.e., 0001sc, 0002sc, etc.)(sc: specific code given to indicate under-five sick child fulfilling inclusion criteria is present in that HH.) for HHs fulfilling inclusion criteria and used as a sampling frame. We allocated total sample size proportionally among the selected six kebeles. Then we recruited HHs fulfilling criteria in each kebele until we obtained the representative sample size in the respective kebele. In a case when there was more than one under-five sick children in a HH, we selected one of the children using lottery method (Figure S1). A revisit was scheduled for those who were not at home during the first visit. Respondents were considered as non-respondent if missed after two revisits on different days.

Tools and Data Collection Process

The data was collected using structured interviewer-administered questionnaires from caregivers. The questionnaire was adapted from Andersen's Health Service

Utilization Model,¹⁴ and experts from the health extension program of the Wolaita zone were consulted. After preparing in English, it was translated to the local language (Wolaytato doona) and back-translated to English by different language experts to ensure consistency. Six diploma nurses as data collectors and two masters in public health as supervisors, with prior experience from other districts, participated in data collection. They were given two days of training on the objective of the study, the process of data collection, and how to handle data a day before the actual data collection.

Key Terms and Measurement

ICCM

ICCM refers to the concurrent management of more than one common childhood illness. In this study, we focused on the common contributors of childhood illness and deaths in Ethiopia: cough, fever and diarrhea.¹⁵

ICCM Utilization

The outcome variable which was measured as a percentage of caregivers, among the study participants, visited HPs for their sick children in the last two weeks for either cough or fever or diarrhea (n = 624). A caregiver had utilized the ICCM if she/he reported that she/he visited HPs after recognizing one of the three childhood illnesses on her/his child otherwise they were not utilized.

Data Quality Control and Data Analysis

To ensure the quality of the data, the questionnaire was pre-tested in a neighboring district a week before the actual data collection to ensure clarity, wordings, logical sequence, skipping patterns, and modifications were done accordingly. Moreover, we made adequate follow ups throughout the data collection period. Supervisors had checked the filled questionnaires on a daily basis for completeness and necessary corrections were made in a timely fashion while the overall activities were monitored by the principal investigator.

We designed the data entry format using the statistical software EpiData 3.1 and exported to the statistical package for social science (SPSS) version 20 for further analysis. We computed a principal component analysis (PCA) to generate the wealth index of HHs. We computed descriptive summary measures to describe the socio-demographic characteristics of the study participants. We had computed bivariate and multivariable logistic regression analysis to assess the association between variables.

The significance of the association between dependent and independent variables was declared using adjusted odds ratio along with their 95% CI. Hosmer–Lemeshow goodness-of-fit was done to test the model fitness ($P = 0.57$).¹⁶ Moreover, multi-collinearity among independent variables was checked using the variance inflation factor (VIF) in which all the variables in the final model were below the recommended cutoff value of 10 (VIF = 1.86).

Result

Socio-Economic and Demographic Characteristics

Out of the total, 624 caregivers of under-five sick children had participated in this study making the response rate 98.6%. The mean age of respondents was 28.75 (SD = 5.4) years and 62% were greater than 30 years old. Significant proportions of caregivers, 283 (45.4%) were living 1–2 hours away from the HP in their kebele. The mean age of the under-five sick children was 20.5 (SD = 4.8) months while the majority 180 (29%), were in the 24–35 months age category and 225 (36%) were sick with diarrhea (Table 1).

ICCM Service Utilization and Health Care-Seeking Behaviors

In this study, only half 312 (50%) of mothers sought health care services from different sources, 66 (10.6%) from HP, 78 (12.5%) from the health center, 58 (9.3%) from hospitals, 54 (8.7%) from traditional healers, 39 (6.3%) home treatment, and 17 (2.7%) from private clinics. The majority of caregivers, 525 (84%), had no awareness of ICCM availability at HP. Besides, only 174 (28%) and 140 (22.5%) of caregivers owned community health insurance and were knowledgeable about child danger signs, respectively ([Table 2).

The magnitude of ICCM utilization at HP among under-five sick children in the study area was found to be 66 (10.6%) (95% CI: 8–13%), while the majority, 558 (89.4%) were non-users of ICCM at HP ([Figure 1). Half of the participants had sought health cares from different sources while 246 (78.85%) of them had not sought health care from HPs for different reasons. The main reported reasons not to use ICCM from HPs were uncertainty of the time when HP is open 86 (35%; 95% CI: 29.4–38.2), HP located too far from their home 81 (33%; 95% CI: 27.0–37.8), HP has no necessary

Table 1 Socioeconomic and Demographic Characteristics of Respondents in Kindo Didaye District, Southern Ethiopia, 2019 (n = 624)

Variable Categories		Frequency	Percentage
Maternal age in years (SD = 5.4)	<30	238	38
	≥30	386	62
Marital status of a mother	Married	585	93.7
	Single	23	3.7
	Others ^a	16	2.6
Occupation of mother	Employ	10	1.6
	Merchant	102	16.3
	Housewife	507	81.3
	Daily laborer	5	0.8
The educational level of a mother	Unable to read and write	495	79.3
	Grade 1–8	103	16.5
	Grade ≥9	26	4.2
Educational status of the father	Unable to write and read	390	62.5
	Grade 1–8	209	33.5
	Grade ≥9	25	4
Occupation of father	Employ	20	3.2
	Merchant	56	9
	Farmer	522	83.7
	Daily laborer	26	4.1
Region	Protestant	418	66.9
	Orthodox	188	30.1
	Others ^b	18	3
HHs Wealth Status	Poor (25%)	195	31.8
	Medium (50%)	238	38
	Rich (25%)	189	30.2
Family size in HH	≤5 members	303	48.6
	>5 members	321	51.4
Age of child in months (SD±4.8)	0–11	57	9
	12–23	86	14
	24–35	180	29
	36–47	147	23.5
	48–59	154	24.5
Child gender	Male	299	48

(Continued)

Table 1 (Continued).

Variable Categories		Frequency	Percentage
	Female	325	52
Distance from home to HP (travel time in minutes)	0–30	117	18.9
	31–60	167	27
	61–120	283	45.4
	>120	57	8.7

Notes: ^aWidowed, Divorced, ^bCatholic, Adventist, Traditional.

medication 74 (30%), and alternative care-seeking 5 (2%; 95% CI: 1.6–3.0), [Figure 2].

Factors Associated with ICCM Utilization at HP

During multivariate logistic regression analysis, the income of the household, previous use of HP for child illness, ownership of community health insurance, awareness of ICCM availability at HP, walking distance from home to HP in their kebele and type of child illness were found to be significantly associated with ICCM utilization at HP (see Table 3).

Discussion

This study has attempted to measure the level of ICCM utilization and identified factors that are influencing ICCM utilization among caregivers with under-five sick children.

The findings of this study revealed that only 10.6% (95% CI: 8–13%) of under-five children sick with diarrhea, fever, and cough/pneumonia took treatment from HP. This result is in line with other studies conducted in the Oromia region of the Jimma zone (10.8%), Hararge zone (10.8%), and Agarfa woreda (10.5%).^{12,17} However, it was lower than the result of the survey conducted by L10K in Amhara, SNNP, and Tigray regions (16.1%).¹³ The possible reason for the difference might be due to socio-economic and demographic differences. This also might result from differences in the study settings because the previous study was conducted in districts supported by the L10K project on the ICCM program.

Households in the poor wealth index category were about three times more likely to utilize ICCM at HPs for their sick children than those in the rich wealth index category. This finding is similar to other studies done in Kenya and Uganda.^{18,19} This might be justified as health services at HPs are free of charge while different costs are

Table 2 Health Care-Seeking Behaviors of Caregivers for Under-Five Sick Children with Common Childhood Illnesses in Kindo Didaye District, Southern Ethiopia, 2019

Sources of Healthcare Seeking		Frequencies	Percentage (%)
Reported type of child illness	Cough	219	35
	Fever	180	29
	Diarrhea	225	36
Health care services sought	Yes	312	50
	No	312	50
Sources of health care seeking	Health post	66	10.6
	Health center	78	12.5
	Hospitals	58	9.3
	Traditional healers	54	8.7
	Home treatment	39	6.3
	Privet clinic	17	2.7
Awareness of ICCM services at HP	Yes	99	16
	No	525	84
Previous use of services at HP	Yes	104	16.8
	No	520	83.2
Owner of community health insurance	Yes	174	28
	No	450	72
Knowledge of child danger signs	Yes	140	22.5
	No	484	77.5
History of index child death	Yes	31	5
	No	593	95

needed to utilize health services from higher health facilities.

The current study also revealed that caregivers who had previously utilized health services at HPs for child illness were about seven times more likely to utilize ICCM at HPs for their sick children than those who never utilized HPs for child illness. This finding is supported by findings from other studies done in Ethiopia and Kenya.^{18,20} The




Figure 1 ICCM utilization at HPs in Kindo Didaye district, Ethiopia, 2019.

possible explanation for this might be exposure to HP for health services encourages caregivers to utilize ICCM.

Caregivers who were not a member of the community health insurance were about three times more likely to utilize ICCM at HP for their sick children. This is in contrast with the fact that members of the community health insurance were more likely to utilize health services than their counterparts.²¹ In Ethiopia, all health care services at HPs are free so that those families who cannot afford to pay for community health insurance every year might prefer to utilize ICCM at HP. This possible justification is also supported by the evidence in our finding that caregivers from poor HHs were more likely to utilize health services at HPs.

Mothers who have awareness of ICCM availability at HP for sick children were about five times more likely to utilize the service than those who were not aware of the ICCM service availability at HP for sick children. This

finding is congruent with findings from other studies done in Ethiopia and Kenya.^{18,20} This might indicate that more awareness of ICCM service availability at HP will scale-up the healthcare-seeking behavior of caregivers.

Caregivers residing within 30 minutes on foot traveling distance from home to HP in their kebele were about five times more likely to utilize ICCM at a HP as compared to those whose distance is more than 120 minutes from home to HP. This finding is in line with the results of other studies done in Ethiopia and Uganda.^{22–24} Although ICCM provision aims to scale-up the health care access in the community by minimizing the distance to travel, this still showed that distance is a challenge for caregivers to utilize ICCM at HPs in the study area.

Children who had symptoms of fever were about three times more likely to utilize the service than children who had diarrhea. This finding is different from the study finding in Jima zone in which children who had pneumonia




Figure 2 Reasons for not utilizing ICCM at HP in Kindo Didaye district, Ethiopia, 2019.

Table 3 Factors Associated with ICCM Service Utilization at HP Among Sick Under-Five Children in Kindo Didaye District, Southern Ethiopia, 2019

Variables		ICCM Utilization		OR (95% CI)	
		Yes	No	COR (95% CI)	AOR (95% CI)
Maternal age in years	<30	49	337	1	1
	≥30	17	221	0.53(0.18–0.95)	0.79(0.47–1.18)
The educational level of a mother	Unable to read and write	50	445	1	1
	Grade 1–8	10	93	1.04(0.51–2.14)	0.97(0.47–2.29)
	Grade ≥9	6	20	0.37(0.14–1.67)	0.33(0.10–1.07)
Wealth index	Poor	30	159	3.15(1.53–6.50)	2.99(1.37–6.53)*
	Medium	25	213	1.96(0.91–2.84)	1.68(0.88–3.20)
	Rich	11	184	1	1
Age of child in months	0–11	2	55	0.22(0.14–1.16)	0.63(0.48–1.69)
	12–23	12	74	0.97(0.88–2.19)	1.03(0.94–2.91)
	24–35	17	163	0.64(0.51–1.13)	0.97(0.74–2.30)
	36–47	13	134	0.59(0.43–1.55)	0.95(0.88–1.32)
	48–49	22	132	1	1
Previous use of services at HP	Yes	63	457	4.64(1.42–15.07)	6.66(1.81–24.04)*
	No	3	101	1	1
Owner of health insurance	Yes	37	413	1	1
	No	29	145	2.23(1.32,3.76)	2.63(1.45,4.76)*
Awareness of ICCM services at HP	Yes	62	458	3.36(1.34,14.18)	4.74(1.39,12.10)*
	No	4	100	1	1
Distance from home to HP (in minutes)	0–30	24	45	4.93(1.74,13.94)	5.23(1.69,10.19)*
	31–60	10	150	0.62(0.44,1.29)	1.02(0.98,2.99)
	61–120	20	252	0.73(0.63,1.53)	1.20(0.96,2.06)
	>120	12	111	1	1
Type of child illness	Fever	32	193	2.13(1.08,4.19)	2.97(1.41,6.25)*
	Cough	21	198	1.36(0.87,2.80)	1.67(0.87,3.20)
	Diarrhea	13	167	1	1

Note: *Statistically significant in multivariate logistic regression (AOR) at P-value <0.05.

were more likely to utilize service than children who had diarrhea.²⁰ Even though it is a plausible hypothesis, this might be due to seasonal and other socio-demographic differences linked with common childhood illnesses. This study was done during the high malaria transmission season in Ethiopia, May, so that fever might be more prevalent and needed the attention of caregivers.

This study also revealed that caregivers are not seeking ICCM from their respective HPs due to the HP being located too far from their home, uncertainty as to the time when the HP is open and whether the HP has no necessary medication, all mentioned as main reasons. This finding is supported by another study conducted in other part of the country.¹⁰ This indicates that still there are

ample challenges for caregivers, such as the accessibility of HP, lack of awareness on working hours, and lack of commodity supplies at HPs.

In this study, the authors tried to minimize recall bias by asking respondents for events that happened to their child within the last two weeks preceding the survey. Social desirability bias in which people might be afraid to admit that they are not visiting HPs in their respective kebele was fixed by recruiting health professional data collectors from other districts. However, future researches need to incorporate qualitative assessments to explore other factors that may support the quantitative assessment so that strong recommendations will be forwarded.

Conclusion

This study revealed that the utilization of ICCM at HPs for under-five sick children with cough, diarrhea, and fever in the district was very low, and half of the sick children in the study area were not seeking out ICCM. Household income, previous utilization of health services at HP for child illness, ownership of community health insurance, the caregiver's awareness of ICCM availability at HP for sick children, walking distance from home to HP in their kebele and type of child illness were found to be significantly associated with ICCM utilization at HP.

Therefore, it is crucial to improve community awareness on ICCM at HPs through community-based platforms (e.g. forums, home visits, meetings) and structures (e.g. women development army, 1-to-5 networks, and volunteer community health workers). It is also important to avail additional HPs as near to many and equip them with the necessary medical supplies to address the problems of the majority.

Abbreviations

HEWs, health extension workers; ICCM, integrated community case management of common childhood illnesses; HP, health post; HH, household; L10K, the last 10 kilometers of JSI project; COR, crude odds ratio; AOR, adjusted odds ratio; CI, confidence interval.

Data Sharing Statement

All the datasets used and/or analyzed during the current study are included in the manuscript and additional information is available from the corresponding author on reasonable request.

Ethics Approval and Informed Consent

We obtained ethical approval from Ethical Review Committee of the College of Health Science at Wolaita Sodo University with Ref. No. CARD:335/334/11. We had obtained verbal informed consent from participants after informing them of the purpose, benefit, risk, confidentiality of the information and voluntary nature of participation in the study. The aforementioned ethical review committee had approved the process of the consent and we had conducted the study in accordance with the Declaration of Helsinki. We linked under-five sick children, who were sick and not on treatment during the data collection time, to the nearby HP to initiate appropriate treatment.

Author Contributions

Asefa Berhanu had led the field data collection and analysis. All authors made substantial contributions to conception and design, analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

Authors received no financial support for the research, authorship, and/or publication of this article.

Disclosure

The authors declare no conflicts of interest and no potential conflicts of interest with respect to the research, authorship, or publication of this article.

References

1. United Nations Inter-agency Group for Child Mortality Estimation (UN IGME). *Levels & Trends in Child Mortality: Report 2019, Estimates Developed by the United Nations Inter-Agency Group for Child Mortality Estimation*. New York: United Nations Children's Fund; 2019. Available from: www.childmortality.org. Accessed November 13, 2020.
2. United Nations Inter-agency Group for Child Mortality Estimation (UN IGME). *Levels & Trends in Child Mortality: Report 2017, Estimates developed by the United Nations Inter-agency Group for Child Mortality Estimation*. New York: United Nations Children's Fund; 2017. Available from: www.childmortality.org. Accessed November 13, 2020.
3. Ethiopian Federal Ministry of Health. *National Implementation Plan for Community Management of Common Childhood Diseases*. Addis Ababa: Federal Ministry of Health; 2010.
4. Ethiopian Federal Ministry of Health. *Health Extension Program in Ethiopia Profile*. Addis Ababa: Health Extension and Education center. Ministry of Health; 2007.
5. Bernadette D, Awa S, Humphreys N, Shelby W, Mark Y. Integrated community case management of childhood illness. *Am J Trop Med Hyg*. 2016;94(3).
6. Bryce J, Friberg IK, Kraushaar D. LiST as a catalyst in program planning: experiences from Burkina Faso, Ghana and Malawi. *Int J Epidemiol*. 2010;39(Supplement 1):i40-i47. doi:10.1093/ije/dyq020
7. Schellenberg JA, Victora CG, Mushi A. Inequities among the very poor: health care for children in rural southern Tanzania. *Lancet*. 2003;361(9357):561-566. doi:10.1016/S0140-6736(03)12515-9
8. Ameha A, Legesse H, Sylla M, et al. The effect of community-based newborn care intervention on service utilization for sick newborn and children. *Ethiop Med J*. 2019;3.
9. Federal Ministry of Health. *National Strategy for Child Survival in Ethiopia (2015/16 – 2019/20)*. Addis Ababa: FMOH; 2015.
10. UNICEF/PATH (2016). Literature review on barriers to health extension services. Draft Report; 2016
11. Legesse H, Degefe T, Hiluf M, et al. National scale-up of integrated community case management in rural Ethiopia: implementation and early lessons learned. *Ethiop Med J*. 2014;52(Supp.3):15-26.

12. Shaw B, Amouzou A, Miller NP, Tafesse M, Bryce J, Surkan PJ. Access to integrated community case management of childhood illnesses services in rural Ethiopia: a qualitative study of the perspectives and experiences of caregivers. *Health Policy Plan.* 2015;31(5):656–666. doi:10.1093/heapol/czv115
13. A Survey of The Last Ten Kilometers Project. Integrated community case management survey in Amhara, SNNP, and Tigray Regions. 2013. Available from: www.110k.jsi.com. Accessed November 13, 2020.
14. Andersen RM, Newman JF. Andersen and newman's framework of health services utilization. *Health Serv Res.* 1995;8(3):184–199.
15. Miller NP, Agbessi A, Mengistu T, et al. Integrated community case management of childhood illness in Ethiopia: implementation strength and quality of care. *Am J Trop Med Hyg.* 2014;91(2):424–434. doi:10.4269/ajtmh.13-0751
16. Hosmer DW, Lemeshow S. *Applied Logistic Regression*. 2nd ed. ohn Wiley & Sons, Inc., New York. doi:10.1002/0471722146
17. Marsha B. *Factors Affecting Utilization of Integrated Community Case Management of Common Childhood Illness in Agarfa Woreda, Oromia Region, Ethiopia* [Unpublished Thesis]. 2014.
18. Kisia J, Nelima F, Otieno DO, et al. Factors associated with utilization of community health workers in improving access to malaria treatment among children in Kenya. *Malar J.* 2012;11(1):248. doi:10.1186/1475-2875-11-248
19. Mukanga D, Tibenderana JK, Peterson S, et al. Access, acceptability and utilization of community health workers using diagnostics for case management of fever in Ugandan children: a cross-sectional study. *Malar J.* 2012;11(1):121. doi:10.1186/1475-2875-11-121
20. Shaw B, Amouzou A, Miller NP, et al. Determinants of utilization of health extension workers in the context of scale-up of integrated community case management of childhood illnesses in Ethiopia. *Am J Trop Med Hyg.* 2015;93(3):636–647.
21. Atnafu DD, Tilahun H, Alemu YM. Community-based health insurance and healthcare service utilisation, North-West, Ethiopia: a comparative, cross-sectional study. *BMJ Open.* 2018;8(8):e019613. doi:10.1136/bmjopen-2017-019613
22. Addis Ashenafi AM, Ameha A, Erbo A, Getachew N, Betemariam W. Effect of the health extension program and other accessibility factors on care-seeking behaviors fore common childhood illnesses in rural Ethiopia. *Ethiop Med J.* 2014;52(Supp 3):57.
23. Mebratie AD, Van de Poel E, Yilma Z, et al. Healthcare-seeking behaviour in rural Ethiopia: evidence from clinical vignettes. *BMJ Open.* 2014;4(2):e004020. doi:10.1136/bmjopen-2013-004020
24. Rutebemberwa E, Kallander K, Tomson G, Peterson S, Pariyo G. Determinants of delay in care-seeking for febrile children in eastern Uganda. *Trop Med Int Health.* 2009;14(4):472–479. doi:10.1111/j.1365-3156.2009.02237.x

Pediatric Health, Medicine and Therapeutics

Dovepress

Publish your work in this journal

Pediatric Health, Medicine and Therapeutics is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries. All aspects of health maintenance, preventative measures and disease treatment interventions are addressed within the journal. Practitioners from all disciplines are

invited to submit their work as well as healthcare researchers and patient support groups. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/pediatric-health-medicine-and-therapeutics-journal>