



OPEN ACCESS

Measures of exposure to the *Well London* Phase-1 intervention and their association with health well-being and social outcomes

Gemma Phillips,¹ Christian Bottomley,² Elena Schmidt,¹ Patrick Tobi,¹ Shahana Lais,¹ Ge Yu,¹ Rebecca Lynch,³ Karen Lock,⁴ Alizon Draper,⁵ Derek Moore,⁶ Angela Clow,³ Mark Petticrew,⁴ Richard Hayes,² Adrian Renton¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jech-2013-202507>).

For numbered affiliations see end of article.

Correspondence to

Professor Adrian Renton, Institute for Health and Human Development, University of East London, Suite 250, University House, The Green, Water Lane, London E15 4LZ, UK; a.renton@uel.ac.uk

Received 15 February 2013
Revised 18 December 2013
Accepted 19 December 2013
Published Online First
10 February 2014

INTRODUCTION

Measuring variation in implementation of public health interventions and levels of population exposure is an important part of evaluation.^{1–4} In the context of effectiveness trials, exposure data can be used to interpret the impact of the intervention on the study outcomes.⁴ In interventions where participation is voluntary or self-selected, measures of exposure can highlight inequity of access to the intervention among population subgroups of interest in public health.⁵

A substantial body of work exists on measuring exposure to media and other blanket health promotion campaigns.⁴ Less attention has been paid to measuring ‘exposure to’, or perhaps more appropriately ‘participation in’, complex public health interventions delivered through community engagement approaches. During the 1980s and 1990s, a handful of large-scale, multicomponent, community-delivered public health interventions were conducted in the USA, and evaluations of these programmes included extensive process measures to estimate the levels of exposure to and/or participation in different components of the intervention, ranging from mass media to small-group health education classes.^{6–8} These studies used the amount of potential exposure events or participation time offered and the actual number of exposure events or duration of participation as measures of implementation. They also attempted to weigh exposure to different intervention elements according to theoretically derived estimates of their relative effectiveness. More recently, Oakley and colleagues⁹ have suggested that quantitative and qualitative process evaluation of exposure to interventions should be considered an essential component of randomised controlled trials of complex public health interventions. This is because they can help, inter alia, to monitor the dose and reach of the intervention, to investigate contextual factors that influence effect, to explore the relationship between trial outcomes and variation in the extent and quality of implementation and to identify processes that might mediate the relationship between intervention and outcomes. However, high-quality process evaluation and measurement of exposure to intervention in complex public health interventions that use community engagement approaches is often difficult to achieve and difficult to resource.

In this paper, we describe the measures of intervention exposure used in the cluster randomised trial of the *Well London* programme, a public

health intervention using community engagement and community-based projects to increase physical activity, healthy eating and mental health and well-being in 20 of the most deprived neighbourhoods in London.¹⁰ No earmarked resources to support the development of these measures and associated data collection were provided to either the research team or to those delivering the interventions on the ground. Instead, these were derived from contractually specified performance management information reported quarterly by partners and by inclusion of questions seeking information about participation in the follow-up questionnaires used to measure the main trial outcomes. The exposure measures are consequently considerably less sophisticated than those used in the US studies, where earmarked funding was available.

The main trial analysis did not provide evidence of neighbourhood-level effects on primary outcomes arising from delivery of the *Well London* programme,¹¹ while the nested qualitative study indicated that active participation in the programme was key to improvements in subjective well-being.¹²

This paper aims (a) to describe the measures of exposure to *Well London* available for the trial analysis, (b) report the levels (and where possible sociodemographic correlates of) exposure and (c) examine the relationship between the trial outcomes and exposure measures. These analyses were prespecified in our trial analysis plan. In addition, we consider the limitations of our exposure measures and the importance and methodological challenges of measuring exposure in the evaluation of a community-level intervention.

METHODS

Well London cluster randomised trial

Methods for the CRT design and analysis are described in detail elsewhere,^{10 11} but information relevant to this paper is summarised here. Twenty matched pairs of deprived neighbourhoods in London were randomised to receive the *Well London* programme or be a control area. Trial outcomes were measured before (baseline) and after (follow-up) intervention delivery by an adult household survey and a school-based adolescent survey in intervention and control neighbourhoods. In these analyses, we used data from the adult household survey and programme monitoring activities.



Open Access
Scan to access more
free content



CrossMark

To cite: Phillips G, Bottomley C, Schmidt E, et al. *J Epidemiol Community Health* 2014;**68**:597–605.

The intervention

Well London Phase-1 was supported by the Big Lottery Well-being Fund and delivered between October 2007 and March 2011. The aim of the *Well London* programme is to support the development of social networks and resilience in communities, with additional changes to the food supply and the neighbourhood environment, to address the local structural and social determinants of health^{13 14} and facilitate individual health behaviour change (nutrition and physical activity) and improve mental well-being. Delivery of the *Well London* programme was coordinated by the London Health Commission (at the Greater London Authority) and conducted in partnership by London Sustainability Exchange (LSx), Groundwork London, Central YMCA, Arts Council England, South London and Maudsley NHS Mental Health Trust (SLaM) and the University of East London.

Each of these partners led a key theme of the programme, delivering their specific projects in all of the 20 intervention neighbourhoods. Each organisation also took an oversight role in three to four neighbourhoods, coordinating delivery of projects by the other partners. This local delivery was organised in collaboration with a local cohosting statutory or voluntary sector organisation with knowledge about, and experience of working with, the community in that specific neighbourhood.

Well London comprised 14 interlinked projects (detailed in table 1), some of which specifically focused on the main health outcomes (physical activity, healthy eating, mental health and well-being), while others sought to improve the local environment (eg, green spaces), provide arts and cultural activities, or employment and training opportunities to residents. At the beginning of the programme, there was an extensive community consultation and engagement process in each neighbourhood (using World Cafe Appreciative Inquiry methodologies^{15 16}) to identify community needs and tailor the suite of projects. The funding conditions of *Well London* required that specified projects be delivered in every neighbourhood addressing the health and well-being goals of the Big Lottery Well-being Fund (projects involving healthy eating/cooking, arts and culture, physical activity, healthy spaces, positive mental well-being), although the number and content of sessions of the individual projects delivered was tailored to local population needs and preferences, particularly the arts and culture, physical activity and healthy spaces projects. A central part of the intervention was the recruitment of a core group of volunteers in each neighbourhood who were trained to support other residents to participate in the *Well London* projects and to access health and social services and to improve health behaviours.

The number of sessions and spaces available varied depending on the format of the activities. Table 1 provides an indication of the volume of participation opportunities in each project, although each was locally tailored. Data on sessions and spaces were not reliably provided by all delivery organisations, and hence were not considered in the analyses presented here. The timing and frequency of activity on each project theme in each neighbourhood is shown in the online supplementary file 1.

Programme monitoring data

During intervention delivery, monitoring data were collected from participants in project activities and from the delivery organisations. Three types of data were collected: (a) *what was delivered*—delivery organisations provided summaries of the activities they had delivered; (b) *how many people participated*—delivery organisations counted and reported the total number

of participants for each project activity or session; and (c) *who participated*—at the end of each project session, participants were invited to complete a brief evaluation form requesting age, postcode, gender, number of activity sessions attended within the project and the number of other *Well London* projects in which they had participated. These data were collated for each 3-month period of programme delivery.

Adult household survey data

Two separate cross-sectional surveys were conducted with no follow-up of individuals between baseline and the end of the intervention. Households were selected at random from the Post Office Address File for each neighbourhood¹¹ and interviews requested from all adults (≥ 16 years) during fieldwork visits. Written informed consent was obtained from all participants. The domains covered in the questionnaire are presented in the online supplementary file 2, and the questions used to capture participation in *Well London* in the intervention and control neighbourhoods are shown in the online supplementary file 3. A copy of the questionnaire is available from the authors upon request.

Trial outcomes

The primary outcomes were eating at least five portions of fruit/vegetables per day, doing at least five sessions of 30 min of moderate intensity physical activity per week (or equivalent), a normal score on the GHQ-12 and the Warwick Edinburgh Mental Well-being Scale score. A range of secondary health and social outcomes were also examined (see online supplementary files 4 and 5).

Measures of exposure to *Well London*

Personal recall of participation

Individual-level participation of respondents in the household adult survey was represented as a binary indicator (0=no participation at any time during the 3.5-year programme; 1=individual had reported participating in any activities.) This was used in the individual-level regression analyses examining the association between self-reported participation and the trial outcomes.

The prevalence of self-reporting any participation in *Well London* in the household adult survey was used as a neighbourhood-level measure of participation.

Intensity of intervention delivery

Two measures of intervention delivery intensity were created:

Project contacts: The number of project contacts per 1000 neighbourhood residents, estimated from the monitoring data as described in the online supplementary file 6. The measure constructed from the monitoring data is not a population rate of participation because residents could take part in more than one project during a reporting period and across the lifetime of the programme and no unique identifiers were available for de-duplication. These estimates were also not adjusted for the length and capacity of specific projects because these data were not reliably reported by delivery organisations.

Project quarters: The number of calendar quarters (3-month periods) that projects were offered (from the programme monitoring data). The *Well London* programme comprised 13 separate projects and delivered over a 42-month period (14 quarters), so the total possible number of project quarters delivered was 182, although each project was not delivered in every month of the programme. The delivery of each project

Table 1 Summary of project content and delivery

Project title	Project description	Delivery lead	Example capacity/duration
Heart of the community projects			
CADBE	Consultation, assessment, design, brokerage, enterprise—includes community cafe needs assessments and appreciative inquiry workshops for design of suite of intervention projects that comprised the initial community engagement activities	University of East London	Open to whole community. Ongoing from the project initiation. Community cafes (start of year 1 and year 2)
Training Communities	Training on a variety of topics to support delivery of the other <i>Well London</i> projects by residents in the LSOAs, for example, facilitation, community engagement	South London and Maudsley NHS Mental Health Trust	Ongoing, individual opportunities on application
<i>Well London</i> Delivery Teams	Training for local volunteers in each LSOA to act in a similar role to NHS Health Trainers—to support people to develop healthier lifestyles through signposting to increase uptake of local services and peer support; the delivery team also act as advocates in interactions with local service providers	London Sustainability Exchange & Central YMCA	Ongoing, individual opportunities on application
Youth.com Unity	Engaging young people to be actively involved in decision making in their local community and in transforming the community to improve health and well-being—youth ambassadors were recruited and trained in each LSOA	Central YMCA	Ongoing from project initiation, a few individuals in each neighbourhood selected
Wellnet	<i>Well London</i> learning network for communities and professionals in London to share practice ideas and experience of delivering community-led interventions for improving health and well-being—it is not limited to delivery partners or areas involved in <i>Well London</i>	London Sustainability Exchange	Ongoing from project initiation
Active Living Maps	Maps of facilities and opportunities for healthy activities/lifestyle, for example, maps show sports facilities, parks, allotments—made for each LSOA and delivered in paper format to all residents	Groundwork London	Delivered to all homes
Themed projects			
Eatwell	Healthy cooking classes (cook and eat) and community feasts to provide engaging education about healthy eating and good nutrition	London Sustainability Exchange	Small groups (~10 people), repeated delivery in each 3-month period
Buywell	Working with local retail outlets and with local community members to improve access to affordable healthy food that is sustainably produced	London Sustainability Exchange	Ongoing from project initiation
Activate London	Range of activities for both young people and adults to engage in physical activity; this involves one or more of signposting to existing local facilities and activities, capacity building by providing training to residents to run physical activity sessions in the LSOA, or direct delivery of, for example, taster sessions and courses and joint initiatives with residents and other providers	Central YMCA	Variable format (small group exercises, football tournaments, existing provision) and according to need. Some provision in every 3-month period
Be Creative, Be Well	Arts activities are used to engage residents in the LSOAs in a process of change to improve, health, well-being, community cohesion and the environment; uses intercultural and intergenerational approaches	Arts Council England	Variable format (community sculptures, parades, small group activities). Activity throughout the programme
Changing Minds	Recruits and trains local residents who have direct experience of mental ill health to deliver awareness training in the LSOAs to reduce stigma and discrimination	South London and Maudsley NHS Mental Health Trust	Small group, high-intensity training (~5 people per area), delivered once
DIY Happiness	Uses humour, creativity and positive psychology approaches to increase psychological resilience; workshops of eight participants, targeted at women	South London and Maudsley NHS Mental Health Trust	Small groups (~10 people)
Healthy Spaces	Improve physical environments through development of community gardens and allotments and redevelopment of greenspaces and greenery	Groundwork London	Projects in all areas, variable format (community allotments, new gardens and changes to urban space)
Mental Well-being Impact Assessment	Local residents are trained to understand, assess and demonstrate the impact of projects, activities and organisations in the LSOA on mental well-being	South London and Maudsley NHS Mental Health Trust	Small groups (~10 people) trained and undertook assessment, two per neighbourhood

CADBE, Community Engagement, Assessment, Design, Brokerage, Enterprise; LSOA, Lower super output area.

across the intervention neighbourhoods is summarised in the online supplementary file 1.

The majority of projects required active, identifiable involvement from participants, for example, exercise or cooking classes. Residents were passively exposed to a few intervention components, such as changes to the green space and local environment, or changes to the commercial provision of fruit and vegetables or other aspects of the local food environment.

These environmental aspects of the intervention programme, and the small number of larger scale community events, are not captured in the project contact or survey-based estimates of participation.

Data analysis

Prevalence of self-reported participation in *Well London* across all intervention neighbourhoods was calculated overall and for

Table 2 Prevalence of participation in any *Well London* activities reported in the postintervention adult household survey in intervention neighbourhoods

	Total sample size	Men (n=802) (95% CI)	Women (n=1084) (95% CI)	Total (n=1886) (95% CI)
Total	1886	2.4 (1.2 to 3.6)	3.7 (1.8 to 5.6)	3.1 (1.6 to 4.6)
Age group (years)				
16–24	458	2.0 (0.2 to 3.7)	2.3 (0.5 to 4.2)	2.2 (0.7 to 3.7)
25–34	482	2.0 (0.0 to 3.9)	2.1 (0.0 to 4.4)	2.1 (0.3 to 3.9)
35–44	411	1.8 (0.0 to 3.8)	4.5 (2.0 to 7.0)	3.4 (1.4 to 5.4)
45–54	271	0.8 (0.0 to 2.4)	4.7 (1.1 to 8.3)	3.0 (0.7 to 5.2)
55–64	126	9.8 (3.0 to 16.6)	8.0 (0.0 to 18.8)	8.7 (2.5 to 15.0)
65+	138	3.4 (0.0 to 8.5)	5.1 (0.0 to 11.0)	4.3 (0.4 to 8.3)
Ethnicity				
White British	484	3.4 (0.7 to 6.2)	3.6 (1.3 to 5.9)	3.5 (1.4 to 5.6)
White other	330	1.4 (0.5 to 3.2)	2.1 (0.0 to 4.6)	1.8 (0.0 to 3.7)
Black Caribbean	196	4.2 (0.0 to 8.9)	8.0 (2.0 to 14.0)	6.6 (2.0 to 11.3)
Black African	399	0.6 (0.0 to 1.9)	2.9 (0.0 to 6.0)	2.0 (0.0 to 4.2)
Indian/Pakistani/Bangladeshi	221	0.8 (0.0 to 2.4)	5.9 (0.0 to 12.6)	3.2 (0.0 to 6.5)
Other Asian	104	5.9 (0.0 to 12.8)	5.7 (0.0 to 11.8)	5.8 (1.3 to 10.3)
Mixed	69	4.8 (0.0 to 13.4)	0.0	1.4 (0.0 to 4.0)
Other	83	2.6 (0.0 to 6.7)	0.0	1.2 (0.0 to 3.3)
English as first language	1013	2.9 (0.9 to 4.89)	4.4 (1.8 to 7.0)	3.8 (1.6 to 5.9)
English as second language	873	1.8 (0.5 to 3.2)	2.9 (1.4 to 4.4)	2.4 (1.2 to 3.6)
Level of educational attainment				
No formal qualifications	196	0.0	0.8 (0.0 to 2.4)	0.5 (0.0 to 1.6)
GCSE or equivalent	643	2.2 (0.2 to 4.2)	3.2 (1.3 to 5.2)	2.8 (1.2 to 4.4)
A-level or equivalent	417	3.5 (1.0 to 6.0)	4.5 (1.6 to 7.5)	4.1 (2.3 to 5.9)
University degree	591	2.6 (0.5 to 4.7)	4.4 (0.1 to 7.8)	3.6 (0.1 to 6.1)
Other	39	0.0	10.0 (0.0 to 22.0)	5.1 (0.0 to 11.0)
Employment status				
In paid employment	797	0.98 (0.0 to 1.9)	4.7 (1.0 to 8.3)	2.8 (1.0 to 4.5)
ILO unemployed (seeking work)	225	3.7 (0.0 to 8.0)	0.9 (0.0 to 2.6)	2.2 (0.0 to 4.8)
Full time education	309	2.2 (0.0 to 4.4)	2.3 (0.2 to 4.4)	2.3 (0.4 to 4.1)
Unable to work (disability/illness)	115	7.0 (0.0 to 17)	3.5 (1.6 to 8.5)	5.2 (0.1 to 10.3)
Not employed not seeking/retired/	440	4.4 (0.2 to 8.4)	4.3 (2.0 to 6.7)	4.3 (2.1 to 6.5)
Carer/other				
Mean number of projects attended (participants)	59	1.4 (0.9 to 1.9)	1.4 (1.1 to 1.6)	1.4 (1.1 to 1.6)
Mean number of sessions attended across all projects (participants)	59	4.9 (2.1 to 7.7)	9.3 (3.3 to 15.3)	7.8 (3.5 to 12.2)

GCSE, General certificate of school education; ILO, International Labour organisation.

different sociodemographic subgroups. Regression models were fitted to each of the individual-level trial outcomes, separately examining their association with the individual-level self-reported participation and each of the neighbourhood-level measures of exposure. All hypothesis tests and CIs were based on robust SEs to account for clustering. All analyses were conducted using complete cases, defined as those survey respondents with data for all sociodemographic variables used for adjustment and for the outcome of interest. Hypothesis tests were considered significant at $p < 0.01$ to account for multiple tests.

RESULTS

Survey response rate

The response rate for the adult household survey calculated at the household level was 26% and the household refusal rate was 51%.¹¹

Exposure to the *Well London* intervention

The prevalence of participation (in any *Well London* activity) reported by household survey respondents was 3.1% (SE 0.7%)

and was higher among women than men (table 2). The mean number of sessions attended across all projects was also higher among women. Self-reported participation increased with age and was highest among men and women aged 55–64 years. The greatest gender differential in participation was at age 35–54 years, with men participating substantially less than women. ‘Other Asians’ were the highest participating male group, while Black Caribbean women participated most often. Those with higher levels of educational qualifications were more likely to report participation, although the prevalence of participation was substantially higher among women with university degrees than men with the same qualifications. Self-reported participation was also higher among those who were unable to work/not seeking employment compared with those who were working or seeking work, and was higher among individuals with English as their first language. Employed women were much more likely than employed men to participate, but the pattern was reversed for men and women who were unemployed and seeking work.

There was variation between intervention neighbourhoods in project contact events per 1000 residents (data not shown),

which ranged from 143 to 1204 per 1000 (mean 436, SE 58) and in the number of project quarters of delivery, ranging from 21 to 47 (mean 32, maximum possible 182). Spearman's rank correlation showed that project contact events and project quarters of delivery were correlated ($\rho=0.5$, $p=0.03$), but that neither was correlated with the survey participation rate (data not shown).

Associations between measures of exposure and trial outcomes

A matrix of associations of individual and neighbourhood-level measures of exposure with trial health and well-being outcomes is shown in table 3 and with trial social outcomes in table 4. Full results are provided in the online supplementary file 7.

Individual-level measures of exposure

After adjustment for sociodemographic characteristics and baseline measures, there were no associations between individual-level participation and the primary trial health outcomes (table 3, column 1), but self-reported *Well London* participants were more likely to report volunteering and taking action to improve their local neighbourhood in the last 12 months than non-participants (table 4, column 1).

Neighbourhood-level measures of exposure

After adjustment for sociodemographic characteristics and baseline measures, there were significant (at $p < 0.01$) associations between exposure measures and some outcomes, although these were not consistent across the different exposure measures. Survey respondents living in areas where there was a higher prevalence of self-report participation from the survey (table 3, column 2) reported eating more portions of fruit and vegetables and were more likely to report meeting government guidelines on physical activity for obesity prevention (at least seven sessions of 60 min of moderate intensity physical activity per week or equivalent). They also reported (table 4, column 2) higher levels of social support and were more likely to report that they had taken action to solve a problem in their area in the past 12 months. After the same adjustments, survey respondents in areas with higher project headcounts reported (table 3, column 3) higher levels of positive mental well-being (Warwick Edinburgh scale). They reported (table 4, column 3) greater social connectedness (social network score) and were also more likely to report that people in their neighbourhood 'pull together to improve it', that individuals in their neighbourhoods 'do things to help each other' and that residents from different backgrounds in their neighbourhood 'get on'. Finally, after adjustment, survey respondents living in areas with more project quarters of intervention delivery were more likely to report (table 4, column 4) that racial harassment was a problem in their area and to report higher levels of incivilities and social disorder.

DISCUSSION

In this paper, we have described one individual-level measure of 'exposure' to the *Well London* programme: self-report participation at the time of trial outcome assessment survey; and three neighbourhood-level measures of exposure: neighbourhood prevalence of self-report participation (from survey); and project-contact event rate and the amount of project time delivered during the 3.5-year programme, calculated from programme monitoring data. The results from the adult household survey conducted at follow-up show a modest level of self-reported exposure and that individuals with lower levels of education, men who were in work or aged 35–54 years and women who were unemployed and seeking work, were less

likely to report having taken part in the *Well London* programme. This indicates potential inequity of access for those who might benefit most from the intervention. At the neighbourhood level, there was correlation between the two programme monitoring-derived measures, but neither was correlated with the survey-derived neighbourhood prevalence of self-report participation.

While it is commonly reported that individuals in lower socioeconomic groups are less likely to engage with and/or benefit from health improvement interventions,^{17–20} it is not possible to distinguish between likelihood of participation and likelihood of reporting participation. Problems with recall may also mean that the self-report prevalence of participation from the survey under-represents the levels of participation in the programme. There were 14 different projects in which residents could participate, which were run at different times during the 3.5-year programme (see online supplementary file 2); at the time of interview it may have been several years since a survey respondent took part in a *Well London* healthy cooking class or went to a community festival or football tournament. Brand recognition may also have been poor in some areas, despite efforts in the survey questionnaire to provide examples of *Well London* activities and branding, so that respondents may have taken part in an activity but not associated it with the *Well London* programme.

Turning to the association between exposure measures and outcomes, at the individual level, self-report participation was only associated with volunteering and taking local action, both of which could be subject to reverse causality, because individuals who already do these things might be more likely to participate in the intervention and/or to recall doing so. While there were significant and strong associations between the neighbourhood-level/exposure measures and some secondary health and social outcomes, there was no consistent trend of association across the three participation measures or across related outcomes. Although the finding of such associations is consistent with the findings of the qualitative study¹² that active participation in the programme was key to improvements in well-being, these results should be viewed with caution as they could clearly result from ecological bias, reverse causality or both, especially given that there were no effects on any of these outcomes in the main trial analysis.

There are significant limitations to all of the exposure measures that we have presented here. These mirror the challenges experienced in previous efforts to monitor and evaluate community health initiatives that used a similar philosophy and approach to *Well London*.^{6–8} When working to engage citizens in such deprived and disenfranchised communities, it might be counterproductive to ask participants repeatedly to complete forms and provide personal information to uniquely identify them. However, this information is needed to track their contact with the programme and their pathway through it to estimate prevalence of participation. Residents in these communities are often unwilling to give this information because of poor relationships with state or authoritative institutions, and to ask for it would risk their disengagement from the programme where the major aim is to reduce such alienation. In relation to the results presented here, we are also not able to say how comprehensively and consistently the anonymous programme monitoring questionnaires were administered by the project delivery staff, across different projects and across different intervention sites. It is therefore possible that the large variability in project time delivered and process evaluation participation prevalence could be due to variability in data collection, not the underlying

Table 3 Associations between individual and neighbourhood-level measures of exposure and trial health and well-being outcomes

	Individual-level self-report participation		Neighbourhood prevalence of self-report participation (per 10% percentage pt increase)		Project contact events per 1000 population (per 10% percentage pt increase)		Project quarters delivered	
	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value
<i>Primary health outcomes</i>								
Healthy eating—meeting five-a-day (fruit and vegetable portions) %	0.8 (0.5 to 1.4)	0.5	1.2 (0.7 to 1.8)	0.5	1.0 (1.0 to 1.1)	0.067	0.99 (0.97 to 1.01)	0.2
Physical activity—meeting 5×30 min moderate intensity activity per week %	1.1 (0.6 to 2.0)	0.7	1.3 (0.9 to 2.1)	0.2	1.0 (1.0 to 1.1)	0.3	0.99 (0.96 to 1.01)	0.3
Abnormal/borderline GHQ12 score %	1.4 (0.5 to 3.7)	0.5	1.3 (0.8 to 2.3)	0.3	1.0 (0.9 to 1.1)	1.0	1.02 (0.99 to 1.05)	0.2
Warwick Edinburgh Mental Well-being Scale mean score†	−1.5 (−3.7 to 0.8)	0.2	−4.7 (−8.8 to −0.5)	0.03	0.7 (0.3 to 1.1)	0.002	0.07 (−0.19 to 0.34)	0.6
<i>Secondary health outcomes</i>								
Unhealthy eating—mean score‡	0.1 (−0.1 to 0.3)	0.4	0.1 (−0.1 to 0.3)	0.4	0.0 (0.0 to 0.0)	0.6	0.00 (−0.01 to 0.01)	0.5
Healthy eating—number of portions of fruit and vegetables per day—mean	0.2 (−0.6 to 1.0)	0.7	1.0 (0.3 to 1.7)	0.009	0.0 (−0.1 to 0.1)	0.6	−0.02 (−0.05 to 0.01)	0.1
Meeting 7×60 min moderate intensity activity per week %	1.2 (0.6 to 2.4)	0.6	2.8 (1.6 to 5.1)	<0.001	10. (0.9 to 1.1)	0.8	0.98 (0.96 to 1.01)	0.2
Doing 150 min of moderate intensity activity per week %	1.2 (0.7 to 1.2)	0.5	1.1 (0.7 to 1.7)	0.8	1.1 (1 to 1.1)	0.096	0.98 (0.96 to 1.01)	0.3
Mean MET minutes per week—mean	654 (−85 to 139)	0.08	1615.3 (413.2 to 2817.4)	0.011	−46.7 (−204.6 to 111.2)	0.5	−22.1 (−61.1 to 16.9)	0.3
Mental Health—GHQ 12—mean score§	0.1 (−0.4 to 0.45)	0.8	0.1 (−0.2 to 0.5)	0.3	0.0 (0.0 to 0.0)	0.7	0.01 (−0.01 to 0.02)	0.3

All analyses adjusted for clustering at LSOA level.

*ORs (binary outcomes) and mean differences (continuous outcomes) adjusted for sociodemographic characteristics: age, gender, ethnicity, employment status, educational attainment and area summary health outcome measures (collected cross-sectionally at baseline.).

†Higher score indicates better mental health.

‡Higher score indicates more unhealthy food consumption.

§Higher score indicates poorer mental health.

GHQ, general health questionnaire; MET, metabolic equivalent.

Table 4 Individual and neighbourhood-level measures of exposure and trial social outcomes

	Individual-level self-report participation		Neighbourhood prevalence of self-report participation (per 10% percentage pt increase)		Project contact events per 1000 population (per 10% percentage pt increase)		Project quarters delivered	
	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value	Adjusted OR or mean difference* (95% CI)	p Value
Social networks score‡	−0.1 (−7.25 to 7.0)	1.0	−26.7 (−50.2 to −3.2)	0.028	3.2 (1.8 to 4.6)	<0.001	0.41 (−0.47 to 1.29)	0.3
Social support score§	1.1 (−0.1 to 2.4)	0.07	3.5 (1.7 to 5.2)	0.001	−0.2 (−0.4 to 0)	0.03	−0.03 (−0.10 to 0.04)	0.4
Social integration								
Some/most people in neighbourhood can be trusted	0.8 (0.4 to 1.7)	0.5	1.1 (0.6 to 2.0)	0.9	1.1 (1 to 1.2)	0.037	1.01 (0.96 to 1.06)	0.8
People from different backgrounds in the neighbourhood get on	0.6 (0.2 to 1.9)	0.4	1.3 (0.4 to 4.0)	0.7	1.1 (1 to 1.2)	0.003	1.01 (0.97 to 1.05)	0.7
Racial harassment is a problem	1.3 (0.5 to 3.2)	0.6	0.9 (0.2 to 4.5)	0.9	1.0 (0.8 to 1.1)	0.6	1.10 (1.04 to 1.18)	0.002
Collective efficacy								
People in the neighbourhood pull together to improve it	0.6 (0.3 to 1.3)	0.2	0.3 (0.1 to 0.9)	0.038	1.3 (1.1 to 1.4)	<0.001	1.04 (0.98 to 1.10)	0.2
People in the neighbourhood help each other and do things together	1.8 (0.8 to 4.2)	0.2	0.9 (0.4 to 2.4)	0.9	1.1 (1 to 1.2)	0.005	1.02 (0.99 to 1.06)	0.3
Taken any action to solve problems in the local area in past 12 months	3.6 (2.0 to 6.5)	<0.001	4.1 (1.5 to 11.3)	0.007	1.0 (0.9 to 1.1)	0.5	1.01 (0.97 to 1.06)	0.6
Volunteering—any activity in last 12 months	3.9 (2.0 to 7.7)	<0.001	2.1 (0.7 to 5.9)	0.2	0.9 (0.8 to 1)	0.061	1.01 (0.97 to 1.04)	0.8
Antisocial behaviour—resident perceptions score¶	0.2 (−0.2 to 0.7)	0.3	−0.3 (−1.2 to 0.6)	0.5	0.1 (−0.1 to 0.2)	0.2	0.06 (0.03 to 0.10)	<0.001
Fear of crime								
Feel safe in the neighbourhood (day)	0.4 (0.1 to 0.9)	0.03	0.6 (0.3 to 1.3)	0.2	1.0 (0.9 to 1.1)	0.9	0.96 (0.93 to 1.00)	0.05
Feel safe in the neighbourhood (night)	1.5 (0.9 to 2.6)	0.1	0.9 (0.6 to 1.5)	0.8	1.0 (0.9 to 1.0)	0.4	0.97 (0.94 to 1.01)	0.1

All analyses adjusted for clustering at lower super output area level.

*ORs (binary outcomes) and mean differences (continuous outcomes) adjusted for sociodemographic characteristics: age, gender, ethnicity, employment status, educational attainment.

‡Higher score indicates greater social connectedness.

§Higher score indicates greater social support.

¶Higher score indicates higher levels of perceived incivilities (survey respondents).

phenomena of delivery and engagement we assume these data represent.

In multilevel interventions that combine components conceptualised as traditional behaviour change activities (where ‘exposure’ is clearly observable and quantifiable) with activities that attempt to address the more indirect but perhaps more fundamentally important influences on health and well-being such as social norms, social support and the health architecture of the local environment (food options, urban layout), there is a bias towards measuring exposure to the individual-level activities. Even where previous studies have attempted to weigh the relative impact of intervention components acting at these different levels to provide a universal measure of programme exposure,⁶ these weights are somewhat arbitrary and may not be grounded in any substantial health-related theory. Other studies have used more qualitative measures to represent the degree of ‘community activation’, but these do not lend themselves to the types of exposure analyses we have attempted here.⁷

The qualitative study embedded in this CRT¹² has shown that interviewees were sometimes unsure whether they had taken part in *Well London*, identifying themselves initially as participants or non-participants but then revealing information during in-depth interviews that suggested the opposite. This suggests that ‘participation’ in a multicomponent, community-based intervention is exceptionally difficult to define, and therefore to measure, because it is differently interpreted by the research team, the intervention delivery staff and members of the community.

Some might suggest that attempting to measure the ‘dose’ of community engagement that individuals or communities receive completely misses the point of this approach,²¹ or at least that the way that individuals engage with such programmes is far less mechanistic than the way we have conceptualised participation in the measures presented here. Simple volume of contact, even when weighed for different levels of educational intensity,⁶ does not capture the complex and multifaceted processes by which the behaviour and self-concept of individuals may be affected and changed by such socially oriented health interventions.

Despite these criticisms, it is important to investigate levels of participation in some way, to highlight inequities of access, and whether the data from the self-report survey are taken at face value, they certainly warn us not to see deprived communities as homogeneous and point to the need to be vigilant so that the most disadvantaged and socially excluded individuals within them are not overlooked. In future studies it may be beneficial to explore the pathways that people take through complex multicomponent interventions such as *Well London* to help examine the ‘active ingredients’ of the programme for different individuals and different outcomes, rather than trying to capture all contact events. While there will still be limitations to these approaches (eg, Hawthorne effect, recruitment bias), recruiting a cohort of residents in intervention areas to be followed through the intervention, whether they participate or not, and/or recruiting a cohort of engaged participants after programme initiation are two possibilities. There is a need to experiment with using new mixed methods and participatory approaches to understand in more detail the ways in which individuals engage with community programmes and the way that this engagement may, or may not, impact on health and social outcomes that are the focus of public health and public policy. However, funders of interventions and evaluations of these must become more realistic about the resources that will be needed to achieve this.

What is already known on this subject

It has been suggested that process measures, including measures of exposure, should be considered as essential components of trials of complex public health interventions. This is because they can help to monitor dose and reach of the intervention, investigate contextual factors that influence effect, explore the relationship between trial outcomes and variation in the extent and quality of the implementation of the intervention and identify processes that might mediate the relationship between intervention and outcomes. A substantial body of work exists on measuring exposure to social marketing and other blanket media health promotion campaigns, which focus on survey recall. Less attention has been paid to measuring ‘exposure’ to complex public health interventions delivered through community engagement approaches. Previous studies of community interventions have looked at the level of provision and actual contact with the programme using management systems during delivery. However, these interventions have focused on educational and screening activities in community healthcare settings.

What this study adds

This study describes survey and programme management data-based approaches to measuring exposure to the *Well London* health improvement programme. We found sociodemographic variation in survey-based exposure measures. We found correlation between our two management data-derived neighbourhood-level measures of exposure but no correlation between these and survey-derived measures. We found some significant associations between the neighbourhood-level exposure measures and some outcomes that might suggest intervention effects not detected in the main trial but were not consistent across measures or related outcomes that might suggest reverse-causation or ecological bias. We discuss some of the implications of these findings for collecting, using and interpreting both types of data, with particular reference to community engagement in deprived urban areas.

Author affiliations

¹Institute for Health and Human Development, University of East London, London, England

²Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, England

³Department of Psychology, University of Westminster, London, England

⁴London School of Hygiene and Tropical Medicine, London, England

⁵Department of Human and Health Sciences, University of Westminster, London, England

⁶Institute for Research on Child Development, University of East London, London, England

Contributors The study was designed by AR, RH, AD, RL, MP, DM, GP and CB. RH designed the analysis strategy. GP and GY carried out statistical analysis. GP drafted the manuscript. AR, MP and RH made critical edits and revisions to the manuscript. All authors read and approved the final version of the manuscript. PT, ES, SL and KL carried out the fieldwork for the surveys.

Funding This work was supported by a grant from the Wellcome Trust (Grant Number 083679), which paid the salary GY and GP. The Greater London Authority funded the fieldwork for the surveys in the intervention sites. The University of East London provided core funding for the IHHD. The views expressed in this paper are

those of the authors and not necessarily those of any funding body or others whose support is acknowledged. The funders had no role in study design, data analysis, decision to publish or preparation of the manuscript.

Competing interests IHHD was funded by Big Lottery to deliver the Community Engagement component of the Well London Intervention.

Ethics approval Research Ethics Committees of University of East London, London School of Hygiene and Tropical Medicine and University of Westminster.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All datasets will be available through the UK Data Archive.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 3.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/3.0/>

REFERENCES

- Craig P, Cooper C, Gunnell D, *et al.* *Using natural experiments to evaluate population health interventions: guidance for producers and users of evidence.* Medical Research Council, 2011.
- Craig P, Dieppe P, Macintyre S, *et al.* Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;337:a1655.
- Hawe P, Shiell A, Riley T, *et al.* Methods for exploring implementation variation and local context within a cluster randomised community intervention trial. *J Epidemiol Community Health* 2004;58:788–93.
- Morris DS, Rooney MP, Wray RJ, *et al.* Measuring exposure to health messages in community-based intervention studies: a systematic review of current practices. *Health Educ Behav* 2009;36:979–98.
- Tugwell P, Petticrew M, Kristjansson E, *et al.* Assessing equity in systematic reviews: realising the recommendations of the Commission on Social Determinants of Health. *BMJ* 2010;341:c4739.
- Finnegan JR Jr., Murray DM, Kurth C, *et al.* Measuring and tracking education program implementation: the Minnesota Heart Health Program experience. *Health Educ Q* 1989;16:77–90.
- Lefebvre RC, Lasater TM, Carleton RA, *et al.* Theory and delivery of health programming in the community: the Pawtucket Heart Health Program. *J Prev Med* 1987;16:80–95.
- Wickizer TM, Wagner E, Cheadle A, *et al.* Implementation of the Henry J. Kaiser Family Foundation's Community Health Promotion Grant Program: a process evaluation. *Milbank Q* 1998;76:121–47.
- Oakley A, Strange V, Bonell C, *et al.* Process evaluation in randomised controlled trials of complex interventions. *BMJ* 2006;332:413–16.
- Wall M, Hayes R, Moore D, *et al.* Evaluation of community level interventions to address social and structural determinants of health: a cluster randomised controlled trial. *BMC Public Health* 2009;9:207.
- Phillips G, Bottomley C, Schmidt E, *et al.* Well London Phase-1: Results among adults of a cluster-randomised trial of a community engagement approach to improving health behaviours and mental well-being in deprived inner-city neighbourhoods. *J Epidemiol Community Health* 2014;68:606–14.
- Derges J, Clow A, Lynch R, *et al.* Well London and the benefits of participation: results of a qualitative study nested in a cluster randomised controlled trial. *BMJ Open* (in press).
- Marmot M. *The Marmot Review: Fair Society, Healthy Lives—Strategic Review of Health Inequalities in England post-2010.* <http://www.instituteofhealthequity.org/projects/fair-society-healthy-lives-the-marmot-review/fair-society-healthy-lives-full-report>.
- Marmot M, Wilkinson RG. *Social determinants of health.* USA: Oxford University Press, 2005.
- Brown J, Isaacs D. The world cafe: Llicing knowledge through conversations that matter. *Syst Thinker* 2001;12:1–5.
- Cooperrider DL, Whitney D, Stavros L. *Appreciative Inquiry Handbook.* San Francisco: Berrett-Koehler, 2003.
- Maheswaran R, Pearson T, Jordan H, *et al.* Socioeconomic deprivation, travel distance, location of service, and uptake of breast cancer screening in North Derbyshire, UK. *J Epidemiol Community Health* 2006;60:208–12.
- Stimpson JP, Nash AC, Ju H, *et al.* Neighborhood Deprivation is associated with lower levels of serum carotenoids among adults participating in the Third National Health and Nutrition Examination Survey. *J Am Diet Assoc* 2007;107:1895–902.
- Giskes K, van Lenthe FJ, Turrell G, *et al.* Smokers living in deprived areas are less likely to quit: a longitudinal follow-up. *Tob Control* 2006;15:485–8.
- Chivu CM, Reidpath DD. Social deprivation and exposure to health promotion. A study of the distribution of health promotion resources to schools in England. *BMC Public Health* 2010;10:473.
- Draper AK, Hewitt G, Rifkin S. Chasing the dragon: developing indicators for the assessment of community participation in health programmes. *Soc Sci Med* 2010;71:1102–9.