

Il-Kang Na^{1,2} and Philipp le Coutre¹

¹Department of Hematology, Oncology and Tumor Immunology, Charité University Medicine, Berlin, Germany. ²Experimental and Clinical Research Center (ECRC), Berlin, Germany.

Supplementary Issue: Tyrosine Kinases in Cancer

ABSTRACT: Tyrosine kinases (TKs) play a significant role in cancerogenesis and cancer cell function. Initial developments in this field go back to the early 80s, but the success story really started with the selective BCR-ABL inhibitor, imatinib. Owing to the cancer-driving role of BCR-ABL in chronic myeloid leukemia (CML), excellent response rates lead to fast FDA approval in both the first and second treatments of CML patients. Since then, numerous TKs were identified. TK inhibitors have been developed accordingly, and technology to test for ideal drug–target interactions has profoundly improved. By now, medical oncologists and hematologists struggle to have a pool of potential TK inhibitors, where the most efficient one could be picked out to treat a specific cancer patient, which might also help overcome the occurring resistance mechanisms against TK inhibitors. Whether disease eradication can be achieved via single or sequential TK inhibitor treatment(s) needs to be tested in the present and in the future.

KEYWORDS: tyrosine kinase inhibitors, targeted therapy, cancer

SUPPLEMENT: Tyrosine Kinases in Cancer

CITATION: Na and le Coutre. Emerging Role of Tyrosine Kinases as Drugable Targets in Cancer. *Biomarker Insights* 2015;10(S3) 29–31 doi: 10.4137/BMI.S22432.

TYPE: Commentary

RECEIVED: April 14, 2015. **RESUBMITTED:** June 21, 2015. **ACCEPTED FOR PUBLICATION:** June 22, 2015.

ACADEMIC EDITOR: Karen Pulford, Editor in Chief

PEER REVIEW: Three peer reviewers contributed to the peer review report. Reviewers' reports totaled 448 words, excluding any confidential comments to the academic editor.

FUNDING: Authors disclose no funding sources.

COMPETING INTERESTS: Authors disclose no potential conflicts of interest.

CORRESPONDENCE: il-kang.na@charite.de

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

Early Steps

Identification of tyrosine kinases with a critical role in cancer. As cloning technologies improved over the past few decades, cloning of large gene transcripts became available and allowed for the identification of cDNA, encoding for biologically relevant physiological peptide kinases and growth factors, including the group of tyrosine kinases (TKs). TKs are involved in multiple aspects of a “normal cell life.” They take part in essential processes, including cell signaling, differentiation, proliferation, and survival. Most receptor TKs are transmembrane proteins that bind extracellular ligands and hand over the signal to cytoplasmic effector and adaptor proteins to regulate biological processes. Nonreceptor TKs, typically, are located in the cytoplasm and include many well-characterized proteins, such as the Src family kinases, c-Abl, and Jak kinases.

Their potential participation in cancer development and cancer cell function was suggested when a close relationship of epidermal growth factor receptor (EGFR) and *the viral oncogene v-erbB*^{1,2} was revealed, and cloning of chromosomal translocations associated with malignancies enabled the characterization of oncogenic gene alterations.

Enhanced TK activity or constitutively active kinase activity with quantitatively and qualitatively altered downstream signaling has been regularly found in human malignancies.

These dysregulations are caused by gene amplification or overexpression of protein tyrosine kinases (PTKs) as in the case of HER2 and breast cancer and genomic rearrangements or chromosomal translocations resulting in fusion proteins as in the case of BCR-ABL and chronic myeloid leukemia (CML). Another example in this direction is the anaplastic lymphoma kinase in anaplastic large cell lymphoma.³ Furthermore, gain of function mutations or deletions in PTKs within the kinase domain or extracellular domain play a role as in the case of *EGFR* mutations and certain types of lung cancer. Mutations causing aberrant B-cell receptor activation are also relevant as in the case of Bruton's tyrosine kinase (BTK) and B-cell lymphomas.⁴

Past Era

Development of specific TK inhibitors for known TK targets. The first targeted antikinase therapeutic agent was the monoclonal antibody trastuzumab used in the treatment of breast cancer. The discovery of *HER2* gene amplification in breast and ovarian cancers was the basic finding that triggered the development of this antibody.

With advanced technologies, analysis of TK transforming properties became possible, and structure–function studies facilitated the development of small molecule inhibitors, specifically targeting constitutively activated protein kinases.



Specific TK inhibitors were developed to interfere with TK enzymes that are aberrantly activated in tumor cells and are critical to tumor growth. Antibodies against TK receptors or their ligands interrupt TK signaling through interference of the ligand-receptor binding. Many other TK inhibitors are designed to directly block the catalytic activity of the kinase by interfering with the binding of ATP or substrates. In 2001, imatinib was the first selective TK inhibitor that was approved by the FDA for use in patients with CML.^{5,6} This small molecule revolutionized the treatment and prognosis of Philadelphia chromosome-positive CML.⁷ Since imatinib potently inhibited the TKs such as Platelet-derived growth factor receptor (PDGFR) and KIT, the FDA approved its use also for patients with advanced GIST. Up to now, numerous anti-TK small molecules or monoclonal antibodies were integrated into therapeutic guidelines in different cancer entities, solid tumors (eg, bevacizumab, sunitinib, and gefitinib), and hematological malignancies (eg, ruxolitinib,⁸ ibrutinib, and idelalisib⁹).

TK inhibitors, typically, are orally administered and generally exhibit good tolerability that distinguishes these agents from conventional cytotoxic chemotherapies. However, as also seen with other drugs, efficacy of TK inhibitors is limited by the development of resistance. Mechanisms leading to drug resistance of tumor cells are manifold, often described are mutations, mostly point mutations that impede TK inhibitor binding,^{10,11} altered gene copy numbers, and protein expression level, which lead to restoration of oncogenic signaling in the presence of a given drug concentration (eg, *bcr-abl* amplification)¹² or increased drug efflux via increased MDR-1 expression.¹³ In order to overcome mutational escape routes, TK inhibitor designs were optimized and second/third-generation TK inhibitors became available.¹⁴ Since second-generation TK inhibitors induced better responses compared to imatinib in patients with newly diagnosed chronic-phase CML, nilotinib and dasatinib were also approved for first-line treatment. However, there is less clinical experience and long-term effects need to be compared to imatinib. Ponatinib, as a third-generation TK inhibitor, was the effective TK inhibitor in patients with T315I so far.

Present and Future Steps

Application of available TK inhibitors across different tumor stages and cancer entities. Some TKs like BCR-ABL are restricted to specific types of cancer, but others are commonly expressed and exhibit tumor-promoting activity in many solid tumors, such as EGFR. Targeting EGFR, therefore, affects various tumor cell types. However, efficiency is not as striking compared to BCR-ABL, because EGFR is not the only molecule driving the tumor growth.

As TKs are essential in many physiological functions and are involved, if dysregulated, in many general tumorigenic processes like invasion, metastasis, and prolonged survival, targeting TK activity is successful across disease stages and also across different cancer entities (eg, anti-EGFR in NSCLC, colon cancer, etc, as well as anti-BTK in diffuse large

cell lymphoma, mantle cell lymphoma, etc). Therefore, current studies address the search for new TK targets to broaden TK inhibitor application by analyzing tyrosine kinomes in different cancers using high-throughput sequencing technologies or microarray-based technologies combined with bioinformatics.

Another option could be “drug repositioning” on the search for the ideal TK inhibitor out of the already existing pool of TK inhibitors. An unbiased drug sensitivity and resistance testing of patient-derived cancer cells could reveal new drug–target interactions and, thereby, provide a basis for different and/or broader clinical applicability of TK inhibitors. One example is axitinib, an antiangiogenic drug already approved for renal cancer, that has recently been shown to potently inhibit the highly resistant T315I mutation of the *bcr-abl* gene.¹⁵

Beside the “effectiveness” criterion, safety/clinical applicability is also of great importance. Different TK inhibitors exhibit unique toxicity profiles and – given a comparable efficiency – selection of the ideal TK inhibitor can be adapted to individual patients with certain comorbidities, for example, in the case of a CML with T315I mutation: ponatinib with side effects like hypertension, rash, and abdominal pain versus axitinib with side effects like cardiac insufficiency, embolic/thrombotic events, and bleedings.

Perspective

TK inhibitors proved to be very efficient and established a new treatment option in various cancer entities. However, to which extent TK inhibitors can facilitate disease eradication is still unclear. Results from both the STOP-Imatinib study and the TWISTER study showed that molecular remissions sustained in 40%–50% of the patients for at least 12–24 months after imatinib cessation.^{16,17} To estimate the long-term efficacy of TK inhibitor cessation, more discontinuation studies and especially longer follow-ups are necessary. More studies are necessary to determine the best time point to restart TK inhibitor treatment for optimal outcome.¹⁸ Moreover, trials addressing this issue for solid tumor entities are missing so far. Apart from their imminent clinical relevance, these strategies are also of biological interest as they address the question, whether TKI may ultimately result in chronic disease control or may even have the potential to trigger cure.

Author Contributions

Wrote the first draft of the manuscript: IKN. Contributed to the writing of the manuscript: PIC. Jointly developed the structure and arguments for the paper: IKN, PIC. Made critical revisions and approved final version: IKN, PIC. Both authors reviewed and approved of the final manuscript.

REFERENCES

1. Downward J, Yarden Y, Mayes E, et al. Close similarity of epidermal growth factor receptor and v-erb-B oncogene protein sequences. *Nature*. 1984;307(5951):521–7.
2. Ullrich A, Coussens L, Hayflick JS, et al. Human epidermal growth factor receptor cDNA sequence and aberrant expression of the amplified gene in A431 epidermoid carcinoma cells. *Nature*. 1984;309(5967):418–25.



3. Gambacorti Passerini C, Farina F, Stasia A, et al. Crizotinib in advanced, chemoresistant anaplastic lymphoma kinase-positive lymphoma patients. *J Natl Cancer Inst.* 2014;106(2):djt378.
4. Maddocks K, Blum KA. Ibrutinib in B-cell lymphomas. *Curr Treat Options Oncol.* 2014;15(2):226–37.
5. Druker BJ, Tamura S, Buchdunger E, et al. Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. *Nat Med.* 1996;2(5):561–6.
6. Gambacorti-Passerini C, le Coutre P, Mologni L, et al. Inhibition of the ABL kinase activity blocks the proliferation of BCR/ABL+ leukemic cells and induces apoptosis. *Blood Cells Mol Dis.* 1997;23(3):380–94.
7. Woessner DW, Lim CS, Deininger MW. Development of an effective therapy for chronic myelogenous leukemia. *Cancer J.* 2011;17(6):477–86.
8. Vannucchi AM, Kantarjian HM, Kiladjian JJ, et al. A pooled analysis of overall survival in COMFORT-I and COMFORT-II, 2 randomized phase 3 trials of ruxolitinib for the treatment of myelofibrosis. *Haematologica.* 2015. [Ahead of print.]
9. Furman RR, Sharman JP, Coutre SE, et al. Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N Engl J Med.* 2014;370(11):997–1007.
10. Gorre ME, Mohammed M, Ellwood K, et al. Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. *Science.* 2001;293(5531):876–80.
11. Daver N, Cortes J, Ravandi F, et al. Secondary mutations as mediators of resistance to targeted therapy in leukemia. *Blood.* 2015;125(21):3236–45.
12. Volpe G, Panuzzo C, Ulisciani S, Cilloni D. Imatinib resistance in CML. *Cancer Lett.* 2009;274(1):1–9.
13. Leslie EM, Deeley RG, Cole SP. Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol Appl Pharmacol.* 2005;204(3):216–37.
14. Jabbour E, Kantarjian H, Cortes J. Use of second- and third-generation tyrosine kinase inhibitors in the treatment of chronic myeloid leukemia: an evolving treatment paradigm. *Clin Lymphoma Myeloma Leuk.* 2015;15(6):323–34.
15. Pemovska T, Johnson E, Kontro M, et al. Axitinib effectively inhibits BCR-ABL1(T315I) with a distinct binding conformation. *Nature.* 2015;519(7541):102–5.
16. Mahon FX, Réa D, Guilhot J, et al; Intergroupe Français des Leucémies Myéloïdes Chroniques. Discontinuation of imatinib in patients with chronic myeloid leukemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. *Lancet Oncol.* 2010;11:1029–35.
17. Ross DM, Branford S, Seymour JF, et al. Safety and efficacy of imatinib cessation for CML patients with stable undetectable minimal residual disease: results from the TWISTER Study. *Blood.* 2013;25:515–22.
18. Rousselot P, Charbonnier A, Cony-Makhoul P, et al. Loss of major molecular response as a trigger for restarting tyrosine kinase inhibitor therapy in patients with chronic-phase chronic myelogenous leukemia who have stopped imatinib after durable undetectable disease. *J Clin Oncol.* 2014;32(5):424–30.