

ORAL PRESENTATION

Open Access

Disentangling malaria anaemia

Kathrin Schuldt^{1*}, Jennifer Evans¹, Julia Lenzen¹, Christa Ehmen¹, Birgit Muntau¹, Jurgen Sievertsen¹, Yeetey Enuameh², Kingsley Osei-Kwakye², Ernest Cudjoe Opoku^{2,3}, Edmund NL Browne⁴, Rolf Horstmann¹, Christian Timmann¹

From Challanges in malaria research: Core science and innovation
Oxford, UK. 22-24 September 2014

Background

Anaemia is a major health burden in sub-Saharan Africa. With an estimated two-thirds of children being affected, childhood anaemia causes poor cognitive development, reduced growth and impaired immune functions associated with increased mortality. In malaria-endemic regions, particularly in those with perennial *Plasmodium falciparum* transmission, most of the childhood anaemia disease burden is being attributed to malaria. The pathogenesis of malaria anaemia, which may well be the single most frequent cause for childhood mortality worldwide, has been debated for many decades.

Materials and methods

In order to identify underlying mechanisms for malaria anaemia two studies were evaluated: a cohort study including 465 children that were followed weekly over eight months and a severe malaria case-control study including >4,500 children, both carried out in a hyper-endemic malaria area of Ghana, West Africa. Human genetic polymorphisms of candidate genes were genotyped and tested for association with malaria anaemia in both studies.

Results

We found several genotype-phenotype associations, which had differing effects after stratifying for anaemia with and without complications. For instance, uncomplicated anaemia was found associated with the A allele of rs196854, which was associated previously with an increased expression of interleukin-6, a key mediator of the systemic inflammatory response.

Conclusions

Our results suggest that there are at least two different underlying mechanisms of malaria anaemia, uncomplicated anaemia characterized by a slow and continuous development towards a deficit of red blood cells, and anaemia with complications characterized by a rapid and substantial loss of red blood cells. The uncomplicated form appears to be caused by chronic systemic inflammatory processes rather than malaria episodes or parasite densities, similar to what is found in other forms of anaemia of chronic disease (ACD).

Authors' details

¹Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany. ²Kumasi Centre for Collaborative Research in Tropical Medicine, Kumasi, Ghana.
³University of Copenhagen, Copenhagen, Denmark. ⁴Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

Published: 22 September 2014

doi:10.1186/1475-2875-13-S1-O21

Cite this article as: Schuldt et al.: Disentangling malaria anaemia. *Malaria Journal* 2014 13(Suppl 1):O21.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany
Full list of author information is available at the end of the article