

RESEARCH ARTICLE

Antagonistic Antiviral Activity between IFN-Lambda and IFN-Alpha against Lethal Crimean-Congo Hemorrhagic Fever Virus *In Vitro*

Licia Bordi^{1‡}, Eleonora Lalle^{1‡}, Claudia Caglioti¹, Damiano Travaglini¹, Daniele Lapa¹, Patrizia Marsella¹, Serena Quartu¹, Zoltan Kis², Kevin K. Arien³, Hartwig P. Huemer⁴, Silvia Meschi¹, Giuseppe Ippolito⁵, Antonino Di Caro⁶, Maria R. Capobianchi^{1*}, Concetta Castilletti¹

1 Laboratory of Virology, National Institute for Infectious Diseases “L. Spallanzani”, Rome, Italy, **2** Division of Virology, National Center for Epidemiology, Budapest, Hungary, **3** Unit of Virology, Institute of Tropical Medicine, Antwerp, Belgium, **4** Department of Hygiene, Microbiology & Social Medicine, Medical University Innsbruck, Innsbruck, Austria, **5** Scientific Direction, National Institute for Infectious Diseases “L. Spallanzani”, Rome, Italy, **6** Laboratory of Microbiology and Biorepository, National Institute for Infectious Diseases “L. Spallanzani”, Rome, Italy

‡ These authors contributed equally to this work.

* maria.capobianchi@inmi.it



OPEN ACCESS

Citation: Bordi L, Lalle E, Caglioti C, Travaglini D, Lapa D, Marsella P, et al. (2015) Antagonistic Antiviral Activity between IFN-Lambda and IFN-Alpha against Lethal Crimean-Congo Hemorrhagic Fever Virus *In Vitro*. PLoS ONE 10(2): e0116816. doi:10.1371/journal.pone.0116816

Academic Editor: Jens H. Kuhn, Division of Clinical Research, UNITED STATES

Received: May 21, 2014

Accepted: November 29, 2014

Published: February 18, 2015

Copyright: © 2015 Bordi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This study has been partly supported by grants from Italian Ministry of Health, Ricerca Corrente and Ricerca Finalizzata

Competing Interests: The authors have declared that no competing interests exist

Abstract

Background and Aims

Crimean Congo Hemorrhagic fever virus (CCHFV) is the causative agent of Crimean-Congo hemorrhagic fever, a severe disease with a mortality rate of around 30% in humans. Previous studies demonstrate that pre-treatment with type I IFNs have an antiviral effect against CCHFV, while established CCHFV infection is almost insensitive to subsequent IFN- α treatment. No data concerning type III IFNs antiviral activity against CCHFV are available so far. The aim of the present study was to explore the capability of IFN- λ 1 to inhibit the replication of CCHFV and the possible synergism/antagonism between IFN- α and IFN- λ 1 both in the inhibition of CCHFV replication and in the activation of intracellular pathways of IFN response.

Methods

Human A549 and HuH7 cells were treated with increasing amounts of IFN- λ 1, or IFN- α or a combination of them, infected with CCHFV; the extent of virus yield inhibition and the induction of MxA and 2'-5'OAS mRNA was measured.

Results and Conclusions

Our study pointed out that type III IFN possess an antiviral activity against CCHFV, even if lower than type I IFN. Moreover, a clear antagonism between IFN- λ and IFN- α was observed in both cell lines (A549 and HuH7 cells), in terms of antiviral effect and activation of pivotal ISGs, i.e. MxA and 2'-5'OAS. Elucidating the interplay between type I and III IFNs

will help to better understand innate defence mechanisms against viral infections and may provide novel scientific evidence for a more rational planning of available and future treatments, particularly against human diseases caused by high concern viruses.

Introduction

Crimean-Congo Hemorrhagic fever virus (CCHFV) is an emerging virus belonging to the Nairovirus genus of the *Bunyaviridae* family. It is the causative agent of Crimean-Congo hemorrhagic fever, a severe disease with a mortality rate of around 30% in humans, most of the deaths occurring 5 to 14 days after the onset of illness [1,2]. CCHFV is maintained in vertical and horizontal transmission cycles involving ixodid ticks and a variety of wild and domestic vertebrates, which do not show signs of illness. The virus circulates in a number of tick genera, but *Hyalomma* ticks are the principal source of human infection. Human-to-human transmission can occur, resulting from close contact with blood, secretions, or other body fluids from infected persons or by direct contact with virus-contaminated blood; nosocomial transmission and organ transplant-related transmission have been reported as well [3–6]. The virus shows a wide geographic distribution and is regarded as a public health threat in many regions of the world, including Asia, Eastern Europe, the Middle East, Africa and Russia [1,6,7]. Despite a wide distribution, the pathogenesis of CCHF remains poorly understood, because of limited human pathology and the need for high-containment facilities to handle CCHFV-infected specimens. There is currently no FDA-approved vaccine or specific antiviral therapy for CCHF. The classification of CCHFV as a WHO Risk Group IV pathogen and the lack of suitable animal models has caused progress in developing new prophylactic and therapeutic measures to be slow [8]. The innate immune response represents the first line of defence against viral infections in mammalian cells [9,10]. Upon sensing molecules produced during viral replication, signalling pathways are activated leading to induction and secretion of type I interferons (IFNs), mainly IFN- α and IFN- β , and subsequent upregulation of interferon stimulated genes (ISGs) [11]. Previous studies demonstrate that pre-treatment with type I IFNs have an antiviral effect against CCHFV [12], most likely due to the activities of IFN-induced antiviral proteins such as MxA [13,14], while established CCHFV infection is almost insensitive to subsequent IFN- α treatment [15]. Recently, a novel group of IFNs has been discovered (IFN- λ 1/interleukin-29 [IL-29], IFN- λ 2/IL-28A, and IFN- λ 3/IL-28B) [16], and assigned to a new type (type III) of IFN. The biological activity of type III IFN overlaps to some extent with that of type I IFN, and similar subsets of ISGs are induced, although type III IFN exerts its action through a receptor complex distinct from that of type I IFN [17,18]. However, no data concerning type III IFNs antiviral activity against CCHFV are available so far.

Moreover, little is known about the effect of type I and type III IFNs combination, as available studies report contrasting information [19–21]. In a previous study from our group an antagonistic activity of IFN- λ 1 and IFN- λ 2 towards both the IFN- α -driven inhibition of replication of several viruses (EMCV, WNV lineage 1 and 2, HSV-1 and CHIKV) and the induction of ISG mRNA was described [22]. Furthermore, in a recently published study IFN- λ pre-treatment of human fibroblasts resulted in lower IFN- α signalling and pro-inflammatory ISGs induction in response to CMV Infection [23].

The aim of the present study was to explore the capability of IFN- λ 1 to inhibit the replication of CCHFV in cell cultures and to investigate the possible synergism/antagonism between

IFN- α and IFN- λ 1 both in the inhibition of CCHFV replication and in the activation of intracellular pathways of IFN response.

Materials and Methods

A549 (ICLC Cell Factory IST, Genova, Italy) and HuH7 (kindly provided by Marco Tripodi, Rome, Italy and purchased from the American Type Culture Collection) were maintained in Dulbecco's Modified Eagle Medium (D-MEM), both containing 10% Foetal Calf Serum (FCS) and penicillin/streptomycin at 37°C in a humidified atmosphere of 5% CO₂.

For virus stock preparation, A549 and HuH7 cells were infected with CCHFV strain 10200 (kindly provided by Anna Papa, Thessaloniki, Greece). Cell lysates were clarified, aliquoted, and stored at -80°C until use. Virus titration was performed on the respective host cell line by limiting dilution assay, by determining tissue culture infectious dose (TCID₅₀) with the method of Reed & Muench. [24]. Virus stock titers were: 10^{8.25}TCID₅₀/ml for CCHFV on A549 and 10^{6.75}TCID₅₀/ml CCHF on HuH7. All procedures involving infectious CCHFV were performed in a biosafety level 4 (BSL4) facility, according to standard operating procedures approved by the institutional biosafety committee. This study was carried out as part of an international training programme, involving Biosafety Level 4 Laboratories from different countries, aimed at sharing and improving Group 4 viruses safety management.

The following IFN preparations were used: human recombinant IFN- α 2b (Intron; Schering Corp., Kenilworth, NJ, USA; specific activity: 400MIU/mg, 1IU corresponding to 2.5pg), human recombinant IL-29/IFN- λ 1 (R&D Systems; Inc., Minneapolis, USA; ED₅₀ 1–5ng/mL, 1IU corresponding to 1–5 ng).

A549 and HuH7 cells were seeded at about 4/5x10⁴ cells per well in 96-well plates and after 1 day were incubated with increasing amounts of either recombinant IFN- α (1, 10, 10² and 10³IU/mL) or IFN- λ 1 (0.01, 0.1, 1 and 10ng/mL) for 18–20h. In parallel cultures, the cells were incubated with mixtures of IFN- α +IFN- λ 1 (1IU/mL+0.01ng/mL, 10IU/mL+0.1ng/mL, 10²IU/mL+1ng/mL and 10³IU/mL+10ng/mL, respectively). Cell monolayers were then infected with CCHFV at a multiplicity of infection (MOI) of 0.01 TCID₅₀/cell and after 24h culture supernatants were collected and titrated to assess virus yield. These experimental conditions (low MOI and overnight incubation, implying multiple replication cycles) were selected in order to maximize the inhibitory effect of IFNs, and to obtain a higher resolution in the virus yield inhibition curves.

Combination index (CI) for constant ratio combinations was calculated through the CompuSyn software, which uses the CI isobologram method of Chou and Talalay [25]. The interpretation of CI value in quantifying two drug antiviral interactions is as follows: CI = 1, additive; CI > 1, antagonism; CI < 1, synergism.

The quantification of MxA and 2'-5'OAS mRNA was performed on total cellular RNA, extracted using Trizol (Gibco BRL, Grand Island, NY, USA), by a Taqman real-time RT-PCR method previously established in our laboratory. In particular, for each gene (including β -actin as housekeeping gene) a standard curve was prepared with serial dilutions of a recombinant plasmid containing the region of interest [26]. For each experimental condition, the ratio of mRNA for MxA and 2'-5'OAS to β -actin was calculated, and the results were expressed as fold induction over untreated cells

Results and Discussion

The ability of human recombinant IFN- α and IFN- λ 1 to inhibit the replication of CCHFV in different cell lines was initially investigated. Although virus yield varied between the 3 experiments performed, the overall trend of the results was uniform, showing a substantially different




Fig 1. Dose-dependent inhibition of CCHFV replication by recombinant IFN- α , IFN- λ 1 and IFN- α +IFN- λ 1. A549 (a) and HuH7 cells (b) were treated for 1 day with increasing amounts of either IFN type, alone or in combination, then infected with CCHFV at MOI 0.01; infectious virus yield was measured after overnight incubation. One out of three experiments is shown. Dotted lines: IFN- α (●) or IFN- λ 1 (▲) used alone; continuous line: IFN- α and IFN- λ 1 (■) used in combination.

doi:10.1371/journal.pone.0116816.g001

ability of the two IFN types to inhibit CCHFV replication, both in A549 (Fig. 1a) and HuH7 cells (Fig. 1b). In Fig. 1a and b, one representative experiment is reported, showing that both IFN- α and IFN- λ 1, used alone, reduced CCHFV yield in a dose-dependent manner, being IFN- α more potent than IFN- λ 1.

When considering the antiviral effects of combinations of IFN- α with IFN- λ 1, for each combination the inhibition of CCHF yield was invariably smaller than that obtained for IFN- α alone (Fig. 1a and b) indicating antagonistic effect between the two IFN types; notably, for the highest dose (10³ IU/mL IFN- α +10 ng/mL IFN- λ 1) the inhibition was even smaller than that obtained with IFN- λ 1 alone in both A549 and HuH7 cells (Fig. 1a and b).

The antagonistic effect was statistically confirmed by calculating the Combination index for each combination. As shown in Table 1, CI was well above 1 for all IFN- α + IFN- λ 1 combinations, both in A549 and HuH7 cells, adding strong statistical support to the graphical results shown in Fig. 1.

In order to characterize possible mechanisms underlying the observed antagonism between IFN- α and IFN- λ 1, we addressed the expression of two well-know IFN-inducible proteins, MxA and 2'-5'OAS. To this aim, HuH7 and A549 cells were incubated for 3, 6 an 24h, a with the same IFN doses (either alone or in combination) used for the virus inhibition experiments. At the indicated time points the quantification of MxA and 2'-5'OAS mRNA was performed.

A dose- and time-dependent induction of mRNA for both ISGs was observed in both cell lines treated with the single IFN types, being again IFN- α much more effective than IFN- λ 1.

Table 1. Combination index for IFN- α and IFN- λ 1 against CCHF replication respectively in A549 and HuH-7 cells.

IFN- α (IU/ml)	IFN- λ 1 (ng/ml)	Combination Index*	
		A549	HuH-7
1.0	0.01	582.8	1.08E14
10.0	0.1	103,469	3.04E13
100.0	1.0	9.25E7	3.55E20
1000.0	10.0	5,008,013	1.19E14

*The Combination Index (CI) was calculated using the CompuSyn software (Chou, T.-C. and Martin, N. CompuSyn software for drug combinations and for general dose effect analysis, and user's guide. ComboSyn, Inc. Paramus, NJ 2007. [www.combosyn.com]) which uses the method of Chou & Talalay. CI values <1, 1 and >1 indicate synergism, additive effect and antagonism, respectively.

doi:10.1371/journal.pone.0116816.t001




Fig 2. Dose-dependent induction of MxA and 2'-5'OAS mRNA levels following treatment with recombinant IFN- α , IFN- λ 1 and IFN- α +IFN- λ 1. A549 (a), (c) and HuH7 cells (b), (d) were exposed 3h to increasing amounts of either IFN type, alone or in combination, then mRNA levels for MxA (a), (b) and 2'-5'OAS (c), (d) were measured, expressed as fold induction vs. untreated cells. One out of three experiments is shown. Dotted lines: IFN- α (●) or IFN- λ 1 (▲) used alone; continuous line: IFN- α and IFN- λ 1 (■) used in combination.

doi:10.1371/journal.pone.0116816.g002

One representative (out of 3) experiment is shown in Fig. 2. The results were in close agreement with those obtained in virus yield inhibition experiments, indicating a weaker effect of IFN- λ 1 as compared to IFN- α ; once again, all the IFN combinations were less effective than IFN- α alone, supporting antagonism between the two IFN types.

In conclusion our study pointed out, for the first time, that type III IFN (IFN- λ 1) possess an antiviral activity against CCHFV, even if lower than type I IFN (IFN- α 2b). Moreover, a clear antagonism between IFN- λ and IFN- α was observed in two different cell lines (A549 and HuH7 cells), in terms of both antiviral effect and activation of pivotal ISGs, i.e. MxA and 2'-5'OAS.

Although possible mechanisms underlying the antagonistic effects of IFN- λ remain to be elucidated, it is possible that differences in the kinetics of activation of intracellular pathways involved in antiviral activity may play an important role. In fact, the kinetics of induction of ISGs differs between the 2 cytokines, as IFN- λ acts slowly but leads to a steady increase in ISGs induction, whereas IFN- α not only induces ISGs rapidly, but also leads to their rapid decline [19,27,28]. In turn, the more rapid, but less intense, STAT activation and ISGs induction by IFN- λ may lead to a partial desensitization to IFN- α , consistent with a recent study where UPS18 was found to mediate the inhibition of IFN- α activities by IFN- λ [29].

Elucidating the interplay between type I and III IFNs will help to better understand innate defence mechanisms against viral infections and may provide novel scientific evidence for a more rationale planning of available and future treatments, particularly against human diseases caused by high concern viruses.

Acknowledgments

Most of the activities were performed in the context of a training exercise within the ERINHA project.

Author Contributions

Conceived and designed the experiments: LB EL C. Castilletti. Performed the experiments: LB EL PM SQ ZK KKA HPH SM. Analyzed the data: LB EL C. Castilletti MRC C. Caglioti. Contributed reagents/materials/analysis tools: DL SM. Wrote the paper: LB EL C. Castilletti MRC C. Caglioti. Provided scientific support: DT GI ADC MRC.

References

- Nichol ST (2001) Bunyaviridae. In *Fields Virology*, pp. 1603–1633. 4th edn. Edited by Lippincott, Williams and Wilkins, Philadelphia.
- Pavlovic J, Arzet HA, Hefti HP, Frese M, Rost D, et al. (1995) Enhanced virus resistance of transgenic mice expressing the human MxA protein. *J Virol* 69: 4506–4510. PMID: [7769712](#)
- Bente DA, Forrester NL, Watts DM, McAuley AJ, Whitehouse CA, et al. (2013) Crimean-Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and genetic diversity. *Antiviral Res* 100: 159–89. doi: [10.1016/j.antiviral.2013.07.006](#) PMID: [23906741](#)
- Burney MI, Ghafoor A, Saleen M, Webb PA, Casals J (1980) Nosocomial outbreak of viral hemorrhagic fever caused by Crimean Hemorrhagic fever-Congo virus in Pakistan, January 1976. *Am J Trop Med Hyg* 29: 941–947. PMID: [7435795](#)
- Suleiman MN, Muscat-Baron JM, Harries JR, Satti AG, Platt GS, et al. (1980) Congo/Crimean haemorrhagic fever in Dubai. An outbreak at the Rashid Hospital. *Lancet* 2: 939–941. PMID: [6107588](#)
- Whitehouse CA (2004) Crimean-Congo hemorrhagic fever. *Antiviral Res* 64: 145–60. PMID: [15550268](#)
- Chinikar S, Persson SM, Johansson M, Bladh L, Goya M, et al. (2004) Genetic analysis of Crimean-congo hemorrhagic fever virus in Iran. *J Med Virol* 73: 404–411. PMID: [15170636](#)
- Keshkar-Jahromi M, Kuhn JH, Christova I, Bradfute SB, Jahrling PB, et al. (2011) Crimean-Congo hemorrhagic fever: current and future prospects of vaccines and therapies. *Antiviral Res* 90: 85–92. doi: [10.1016/j.antiviral.2011.02.010](#) PMID: [21362441](#)
- Bonjardim CA, Ferreira PC, Kroon EG (2009) Interferons: signaling, antiviral and viral evasion. *Immunol Lett* 122: 1–11. doi: [10.1016/j.imlet.2008.11.002](#) PMID: [19059436](#)
- García-Sastre A, Biron CA (2006) Type 1 interferons and the virus-host relationship: a lesson in détente. *Science* 312: 879–82. PMID: [16690858](#)
- Randall RE, Goodbourn S (2008) Interferons and viruses: an interplay between induction, signalling, antiviral responses and virus countermeasures. *J Gen Virol* 89: 1–47. PMID: [18089727](#)
- Andersson I, Lundkvist A, Haller O, Mirazimi A (2006) Type I interferon inhibits Crimean-Congo hemorrhagic fever virus in human target cells. *J Med Virol* 78: 216–22. PMID: [16372299](#)
- Frese M, Kochs G, Feldmann H, Hertkorn C, Haller O (1996) Inhibition of bunyaviruses, phleboviruses, and hantaviruses by human MxA protein. *J Virol* 70: 915–23. PMID: [8551631](#)
- Andersson I, Bladh L, Mousavi-Jazi M, Magnusson KE, Lundkvist A, et al. (2004) Human MxA protein inhibits the replication of Crimean-Congo hemorrhagic fever virus. *J Virol* 78: 4323–4329. PMID: [15047845](#)
- Andersson I, Karlberg H, Mousavi-Jazi M, Martínez-Sobrido L, Weber F, et al. (2008) Crimean-Congo hemorrhagic fever virus delays activation of the innate immune response. *J Med Virol* 80: 1397–404. doi: [10.1002/jmv.21222](#) PMID: [18551619](#)
- Ank N, West H, Paludan SR (2006) IFN-lambda: novel antiviral cytokines. *J Interferon Cytokine Res* 26: 373–379. PMID: [16734557](#)
- Sheppard P, Kindsvogel W, Xu W, Henderson K, Schlutsmeyer S, et al. (2003) IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* 4: 63–68. PMID: [12469119](#)
- Kotenko SV, Gallagher G, Baurin VV, Lewis-Antes A, Shen M, et al. (2003) IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat Immunol* 4: 69–77. PMID: [12483210](#)
- Marcello T, Grakoui A, Barba-Spaeth G, Machlin ES, Kotenko SV, et al. (2006) Interferons alpha and lambda inhibit hepatitis C virus replication with distinct signal transduction and gene regulation kinetics. *Gastroenterology* 131: 1887–98. PMID: [17087946](#)
- Pagliaccetti NE, Eduardo R, Kleinstein SH, Mu XJ, Bandi P, et al. (2008) Interleukin-29 functions cooperatively with interferon to induce antiviral gene expression and inhibit hepatitis C virus replication. *J Biol Chem* 283: 30079–30089. doi: [10.1074/jbc.M804296200](#) PMID: [18757365](#)

21. Ank N, West H, Bartholdy C, Eriksson K, Thomsen AR, et al. (2006) Lambda interferon (IFN-lambda), a type III IFN, is induced by viruses and IFNs and displays potent antiviral activity against select virus infections in vivo. *J Virol* 80: 4501–4509. PMID: [16611910](#)
22. Bordi L, Lalle E, Lapa D, Caglioti C, Quartu S, et al. (2013) Type III interferon (IFN-lambda) antagonizes the antiviral activity of interferon-alpha in vitro. *J Biol Regul Homeost Agents* 27: 1001–9. PMID: [24382181](#)
23. Egli A, Levin A, Santer DM, Joyce M, O'Shea D, et al. (2014) Immunomodulatory Function of Interleukin 28B During Primary Infection With Cytomegalovirus. *J Infect Dis*. In press
24. Reed LJ, Muench HA (1938) A simple method of estimating fifty percent endpoints. *Am J Hyg* 27: 493–497.
25. Chou TC, Talalay YP (1984) Quantitative analysis of dose-effect relationships: the combined effects of multiple drugs or enzyme inhibitors. *Adv Enzyme Regul* 22: 27–55. PMID: [6382953](#)
26. Capobianchi MR, Lalle E, Martini F, Poccia F, D'Offizi G, et al. (2006) Influence of GBV-C infection on the endogenous activation of the IFN system in HIV-1 co-infected patients. *Cell Mol Biol* 52: 3–8. PMID: [17543192](#)
27. Kohli A, Zhang X, Yang J, Russell RS, Donnelly RP, et al. (2012) Distinct and overlapping genomic profiles and antiviral effects of Interferon- λ and α on HCV-infected and noninfected hepatoma cells. *J Viral Hepat* 19:843–53. doi: [10.1111/j.1365-2893.2012.01610.x](#) PMID: [23121362](#)
28. Shindo H, Maekawa S, Komase K, Miura M, Kadokura M, et al. (2013) IL-28B (IFN- λ 3) and IFN- α synergistically inhibit HCV replication. *J Viral Hepat* 20:281–9. doi: [10.1111/j.1365-2893.2012.01649.x](#) PMID: [23490373](#)
29. François-Newton V, Magno de Freitas Almeida G, Payelle-Brogard B, Monneron D, Pichard-Garcia L, et al. (2011) USP18-based negative feedback control is induced by type I and type III interferons and specifically inactivates interferon α response. *PLoS One* 6: e22200. doi: [10.1371/journal.pone.0022200](#) PMID: [21779393](#)