



The miR-183~96~182 cluster promotes tumorigenesis in a mouse model of medulloblastoma

Zengdi Zhang, Sanen Li, Steven Y Cheng[✉]

Department of Developmental Genetics, Nanjing Medical University, Nanjing, Jiangsu 210029, China.

Received 25 January 2013, Revised 17 February 2013, Accepted 08 June 2013, Epub 10 July 2013

Abstract

Medulloblastoma is the most common malignant pediatric brain tumor. Some are thought to originate from cerebellar granule neuron progenitors (CGNPs) that fail to undergo normal cell cycle exit and differentiation. The contribution of microRNAs to the initiation and progression of medulloblastoma remains poorly understood. Increased expression of the miR-183~96~182 cluster of microRNAs has been noted in several aggressive subgroups. We identified that expression of miR-183~96~182 was higher in medulloblastomas with *Pten* gene loss in the background of the activated sonic hedgehog (Shh) signaling pathway. Ectopic miR-183~96~182 expression in CGNPs synergized with exogenous Shh to increase proliferation and its role depended on hedgehog signaling activation. Our findings suggest a new microRNA cluster, the miR-183~96~182, functionally collaborates with the Shh signaling pathway in the development of medulloblastomas in mice.

Keywords: medulloblastoma, microRNAs, Shh pathway, granule neuron progenitors, proliferation

INTRODUCTION

Medulloblastoma is a tumor of the cerebellum and the most common malignant pediatric brain tumor. Medulloblastoma occurs most frequently in children between the ages of 5 and 10 years but may occur in adults as well^[1-3]. Although 5-year in overall survival rates have reached 60%-80%, the prognosis for many medulloblastoma patients remains bleak. Many patients who develop the disease die from it and survivors, after conventional treatment regimens (surgery, radiotherapy, and chemotherapy), often face a variety of long-term neurological, neuroendocrine, and social sequelae^[2-4]. It is therefore imperative for us to gain a deeper understanding of the molecular pathogenesis driving medulloblastomas, in order to improve ap-

proaches for treating medulloblastomas.

MicroRNAs (miR) are small endogenous noncoding RNAs that play important roles in many biological processes, including tumorigenesis^[5]. These 21-23 nucleotide (nt) RNAs are matured through a 2-step biogenesis mechanism from long, RNA polIII transcripts^[6]. Mature microRNAs are incorporated into the RNA-induced silencing complex, where they anneal to their recognition sequences in the 3'-UTR of mRNA genes to attenuate gene expression through translational repression or mRNA degradation^[7]. Several genomic studies have revealed the differential expression of microRNAs across various molecular subgroups of medulloblastoma. As previously reported, the miR-17/92 polycistron was identified as a putative medulloblastoma oncomir and associated with the

✉ Corresponding author: Steven Y Cheng, Ph.D., Department of Developmental Genetics, Nanjing Medical University, 140 Hanzhong Road, Nanjing, Jiangsu 210029, China. Tel: +86-025-86862034.

E-mail: xaslp@126.com.

The authors reported no conflict of interests.

sonic hedgehog-driven medulloblastoma compared with other subgroups^[8,9].

In different comprehensive analyses of the human and mouse medulloblastoma oncomir, the miR-183~96~182 cluster is upregulated in several subgroups, including particularly aggressive clinical course involving high rates of metastasis and overall poor survival^[10]. Based on the studies of cell lines, Weeraratne et al. found that the knockdown of the full miR-183~96~182 cluster results in the enrichment of genes associated with apoptosis and dysregulation of the PI3K/AKT/mTOR signaling axis. However, the specific role of miR-183~96~182 cluster in medulloblastoma and cerebellar development has not been described^[11,12].

Cerebellar granule neuron progenitors (CGNPs) are proposed cells of origin for a subset of medulloblastomas. CGNPs undergo rapid Shh-dependent expansion perinatally in mice and humans, and excessive Shh pathway activity promotes medulloblastoma^[13,14]. We used the *Ptch1*^{+/-}; *Pten*^{Floxp/+}; *GFAP-Cre* mouse model of medulloblastoma to study the effect of the PI3K/Akt signaling pathway activated on medulloblastoma tumorigenesis. We found that the miR-183~96~182 cluster was highly expressed in this *Pten* and *Ptch1* loss associated medulloblastoma and promote tumorigenesis by controlling CGNP proliferation.

MATERIALS AND METHODS

Animal

Ptch1^{+/-} mice were purchased from Jackson Laboratory^[15]. C57BL/6 and *Pten*^{Floxp/Floxp} mice were purchased from the Model Animal Research Center of Nanjing University^[16]. *GFAP-Cre* mice were gifts from Minsheng Zhu's laboratory (Nanjing University, Nanjing, Jiangsu, China). *Pten*^{Floxp/Floxp} mice were crossed with *Ptch1*^{+/-} mice. The, *n Ptch1*^{+/-}; *Pten*^{Floxp/Floxp} offspring mice were crossed with *GFAP-Cre* mice to obtain *Ptch1*^{+/-}; *Pten*^{Floxp/+}; *GFAP-Cre* mice. Mice were observed for symptoms of medulloblastoma at least twice-weekly for 12 months. Transgenic expression patterns of *GFAP-Cre* were examined with a R26R reporter line (129S4/SvJaeSortGt(ROSA)26Sortm1(FLP1)Dym, (Jackson Laboratory, Bar Harbor, Maine, USA) carrying a *lacZ* gene whose expression requires excision of loxP-flanked stop sequences. All mice were housed in a specific pathogen free animal room. The study protocol was approved by the local institutional review board at the authors' affiliated institutions. Animal welfare and the experimental procedures were carried out strictly in accordance with the Guide for Care and Use of Laboratory Animals (National Research Council of USA, 1996).

Histological study

For histological analysis, mice were perfused with PBS followed by 4% paraformaldehyde. The cerebella were removed, fixed in 4% paraformaldehyde overnight, and transferred to gradient ethanol for dehydration and embedded in paraffin. Sections (5 μm) were stained with hematoxylin and eosin (Sigma, St Louis, MO, USA). Frozen sections (20 μm) were stained for β-galactosidase according to a previously reported method^[17].

RT-PCR

Total RNA from mouse tissues was extracted using the RNAiso reagent (TaKaRa, Osaka, Japan). Stem-loop RT-PCR primers were designed as previously reported^[18]. The U6 RNA was used for normalization. Stem-loop RT primers and PCR primers were synthesized by Invitrogen (Carlsbad, CA, USA). Reverse transcription was performed using the PrimeScript™ RT reagent Kit with gDNA Eraser (TaKaRa). The cycling condition of PCR was 95°C for 5 minutes, followed by 30 amplification cycles of 95°C for 15 seconds and 60°C for 1 minute. Stem-loop RT primers sequences for these microRNA were listed as follows: mmu-mir-183-RT: 5'-GTCG-TATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACAGTGAA-3'; mmu-mir-96-RT: 5'-GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACAGCAA-3'; and mmu-mir-182-RT: 5'-GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACACCGGTGT-3'. Forward PCR primers sequences for these microRNAs are: mmu-mir-183-FR: 5'-AGCCGTATGGCACTGGTAGAA-3'; mmu-mir-96-FR: 5'-AGCCGTTTGGCACTAGCACATT-3'; mmu-mir-182-FR: 5'-AGCCGTTTGGCAATGGTAGAACTC-3'.

Culture of CGNPs

Granule neuron progenitors were purified from the cerebellum of 7-day-old C57BL/6 pups by using a modified protocol as previously reported^[19,20]. Briefly, postnatal cerebella were triturated into single-cell suspensions that were loaded onto a step gradient of 35% and 65% Percoll (Amersham-Bioscience, Uppsala, Sweden) and separated by high-speed centrifugation for 10 minutes at 4°C. Granule cells and precursors were harvested from the 35/65% interface, washed in PBS/DNase, and further purified by depleting adherent cells with 2-hour incubation on normal tissue culture dishes. Their purity (> 95%) was assessed by immunostaining with markers of neurons and glia. Cells (5 × 10⁵ per well) were plated in four-well Lab-Tek

chamber slides precoated with 100 µg/mL poly-D-lysine (Millipore, Billerica, MA, USA) and Matrigel (Beckton Dickinson, Bedford, MA, USA) in neural basal medium supplemented with 0.45% glucose, SPITE (Sigma), oleic acid albumin/linoleic acid (Sigma), B27 (Invitrogen), and N-acetyl cysteine (Sigma).

Retrovirus production and infection

To generate a mmu-miR-183~96~182 cluster expression clone, the microRNA cluster was PCR-amplified from genomic DNA of C57BL/6 mouse and cloned into pENTR/D-TOPO plasmid (Invitrogen), then transferred into MSCV-P1G-gateway vector (Clontech, Mountain View, CA, USA) by Gateway LR Clonase II enzyme mix (Invitrogen). When CGNPs became attached, 250 µL medium was removed and 250 µL medium containing virus was added. Infection was repeated 3 times with 2 hours each time. The next day, the medium was changed for fresh complete medium with or without 10% ShhN conditional medium (ShhN CM), which was generated by transfecting 293T cells with a ShhN expression construct^[21].

EdU staining

Identification of purified granule cells in the S phase was performed by incubation with Click-iT EdU Alexa Fluor 555 Imaging Kit (Invitrogen). EdU was added for 12 hours before fixation. EdU detection was performed according to the manufacturer's instructions (Invitrogen). Stained chambers covered with mounting medium containing DAPI (Sigma) were enumerated under a microscope. TIFF images of 4 random fields were taken for each experimental group by using the 20 × objective lens. The counting of EdU-positive and DAPI-positive CGNPs number was quantified by using ImageJ software. The kinetics of the proliferative response to mmu-miR-183~96~182 was analyzed by using GraphPad Prism 5.

Statistical analysis

CGNPs proliferation at each time point was measured as the mean ratio (EdU-labeling to DAPI-labeling cell number) from 4 random fields. Statistical significance of proliferation was determined by using Student's *t*-test.

RESULTS

Heterozygosity for *Pten* promotes medulloblastoma in cooperation with Hedgehog signaling pathway

The founders of GFAP-Cre and their offspring were identified by PCR genotyping. Transgenic expression

patterns of Cre were examined with a Rosa26 reporter line, 129S4/SvJaeSor-Gt(ROSA)26Sortm1(FLP1) Dym, carrying a *lacZ* gene whose expression requires excision of *Floxp*-flanked stop sequences. β-galactosidase staining revealed GFAP-Cre mediated recombination both in astrocytes and immature precursor cells of the external granular layer (EGL) in the developing cerebellum of 7-day old mice (**Fig. 1A**). To explore the role of *Pten* loss in the background of *Ptch1*^{+/-} mice, we administered the GFAP-Cre transgenic mice to delete *Pten* in CNS lineages by using conditional knockout mice *Pten*^{Floxp/Floxp}. Broad CNS deletion of *Pten* results in lethal hydrocephalus in early postnatal life; modeling efforts henceforth emphasized the *Pten*^{Floxp/+} GFAP-Cre genotype.

In *Pten*^{Floxp/+} mice, part of *Ptch1* exon 1 (including putative start codon) and all of exon 2 were replaced with *lacZ* and a neomycin resistance gene in one allele^[15]. Approximately 17.6% (3 out of 17 *Pten*^{Floxp/+}) of mice expressing one allele of *Ptch1* developed symptoms of medulloblastoma by 1 year of age. Most of these mice died 5 months after birth^[22,23]. However, *Pten*^{Floxp/+} *Pten*^{Floxp/+} GFAP-Cre mice died of medulloblastoma from 2 months after birth. We found that, between 2 to 8 months of ages, 18/23 (78.2%) of the *Pten*^{Floxp/+} *Pten*^{Floxp/+} GFAP-Cre mice presented with acute-onset neurological symptoms. Similarly to *Ptch1*^{+/-} mice, these mice began to display signs of illness, including domed head, hunched back, abnormal gait and/or decreased movement (data not shown). Histopathologically, these mice harbored malignant medulloblastoma. Tumor cells, regardless of whether the mouse was *Ptch1*^{+/-} or *Ptch1*^{+/-} *Pten*^{Floxp/+} or GFAP-Cre, arose from the surface of the cerebellum and invaded the cortex and deep white matter of the cerebellum (**Fig. 1B**, panel a and b). The tumors in the *Ptch1*^{+/-} mice showed small round or ellipsoid cells with a high nuclear-to-cytoplasmic ratio and round-to-oval or triangular hyperchromatic nuclei, corresponding to patterns of classic medulloblastomas in humans (**Fig. 1B**, panel c). In contrast, the tumors from *Ptch1*^{+/-} *Pten*^{Floxp/+} GFAP-Cre mice had a biphasic architecture that nodular areas consisted of differentiated cells with round nuclei and less condensed chromatin, and were intermingled with the undifferentiated hyperchromatic cells (**Fig. 1B**, panel d). The nodular areas were surrounded with desmoplasia, a pericellular deposition of collagen in this context, similarly to the appearance of desmoplastic medulloblastomas in humans. However, none of the *Pten*^{Floxp/+} GFAP-Cre mice developed medulloblastoma. Therefore, in cooperation with the hedgehog signaling pathway, heterozygosity for *Pten* promoted medulloblastoma. Medul-




Fig. 1 Heterozygosity for *Pten* promotes medulloblastoma in the background of *Ptch1*^{+/-}. A: β -galactosidase staining in GFAP-Cre; Rosa26-lacZ mice of P7. B: Higher magnification of (A). Positive for β -galactosidase staining (blue reaction product) was found both in astrocytes and in immature precursor cells of the external granular layer (arrow) in the developing cerebellum. C: H&E staining of histological sections of representative malignancies in *Ptch1*^{+/-} medulloblastomas. D: H&E staining of histological sections of representative malignancies in *Ptch1*^{+/-}; *Pten*^{Flox/+}; GFAP-Cre medulloblastomas. E: Higher magnification of (C). F: Higher magnification of (D).

loblastomas induced by mutations in the Shh pathway has been reported to originate from long-lasting proliferative CGNPs and are located within the cerebellar hemispheres^[13,24,25]. Therefore, we speculated that the *Ptch1*^{+/-} *Pten*^{Flox/+} GFAP-Cre medulloblastomas arise from CGNPs.

Mouse medulloblastoma with heterozygosity for *Pten* and *Ptch1* overexpresses microRNAs from the miR-183~96~182 cluster

The miR-183~96~182 cluster is a highly conserved cluster of microRNAs residing on an intergenic region

of human chromosome 7q, which is commonly gained in medulloblastoma. The miR-183~96~182 cluster has been largely reported as pro-tumorigenic/prometastatic effects in cancers. Increased expression of the miR-183~96~182 cluster of microRNAs has been reported in several subgroups of medulloblastoma, including the most clinically aggressive subgroup associated with genetic amplification of *MYC* and activation of phosphoreceptor transcriptional programs^[11,12]. Therefore, we investigated the expression of mmu-miR-183, mmu-miR-96 and mmu-miR-182 in these medulloblastomas. The genes for 3 microRNAs of the cluster, including miR-183, miR-96 and miR-182, were clustered within 4 kb on mouse chr6qA3, and transcribed in the same direction (telomere→centromere) (Fig. 2A). All the 3 microRNAs had a high degree of sequence homology and potentially overlapping mRNA targets (Fig. 2B). For the stem-loop RT-PCR assay, we found that the mmu-miR-183, mmu-miR-96 and mmu-miR-182

expressions were increased in *Ptch1*^{+/-};*Pten*^{Floxp/+} *GFAP-Cre* medulloblastoma compared with *Ptch1*^{+/-} medulloblastoma, especially the mmu-miR-183 and mmu-miR-96. However, as the same PCR cycles, the expression of these microRNAs had very low level in adult normal cerebellum of WT, *GFAP-Cre*, *Pten*^{Floxp/+};*GFAP-Cre*, *Ptch1*^{+/-} and *Ptch1*^{+/-} *Pten*^{Floxp/+} *GFAP-Cre* mice (Fig. 2C). Therefore, we speculated that high expression of the miR-183~96~182 cluster was an important factor in promoting tumorigenesis in *Ptch1*^{+/-} *Pten*^{Floxp/+} *GFAP-Cre* mice.

Enforced expression of miR-183~96~182 promotes CGNP proliferation in cooperation with the hedgehog signaling pathway

As CGNPs are the source of Shh-subtype medulloblastomas, we asked that whether the overexpression of miR-183~96~182 promoted CGNP proliferation, and dysregulation of miR-183~96~182 was one of the




Fig. 2 Overexpresses miRNAs from the miR-183~96~182 Cluster in *Ptch1*^{+/-}; *Pten*^{Floxp/+}; *GFAP-Cre* medulloblastomas. A: Genomic organization of the miR-183~96~182 cluster on mouse chr6qA3. The numbers show the spacing in each of the three species in genes. B: Sequence alignment for mouse miR-183, miR-96, and miR-182. C: RT-PCR detection of mature miR-183, miR-96, and miR-182 in adult normal cerebellum of WT, *GFAP-Cre*, *Pten*^{Floxp/+};*GFAP-Cre*, *Ptch1*^{+/-} and *Ptch1*^{+/-};*Pten*^{Floxp/+};*GFAP-Cre* mouse and medulloblastoma of *Ptch1*^{+/-} and *Ptch1*^{+/-};*Pten*^{Floxp/+};*GFAP-Cre* mouse. The U6 RNA was used for normalization. Lane 1: WT cerebellum; Lane 2: *GFAP-Cre* cerebellum; Lane 3: *Pten*^{fl/+};*GFAP-Cre* cerebellum; Lane 4: *Ptch1*^{+/-} cerebellum; Lane 5: *Ptch1*^{+/-} medulloblastoma; Lane 6: *Ptch1*^{+/-} *Pten*^{fl/+};*GFAP-Cre* cerebellum; Lane 7: *Ptch1*^{+/-} *Pten*^{fl/+};*GFAP-Cre* medulloblastoma.

most important factors in *Ptch1*^{+/-} *Pten*^{Flox/+} *GFAP-Cre* medulloblastomas. We isolated CGNPs from the cerebellum of 7-day-old C57BL/6 pups and infected them with retroviruses expressing the mmu-miR-183~96~182 cluster or control with Retro-GFP for 12 hours and then cultured CGNPs with or without ShhN CM. After infecting retrovirus for 36, 48, 60 and 72 hours, we measured each CGNP proliferation by quantifying EdU staining, respectively (**Fig. 3A**). Consistent with previous studies, we found that cells maintained proliferation in the presence of shhN. After infection for 36 hours, the overexpression of miR-183~96~182 had little effect on high concentration of ShhN. During 48 to 60 hours, the overexpression of miR-183~96~182 cluster increased proliferation in ShhN CM-treated cells (**Fig. 3B**). With the prolongation of the culture time, CGNPs grew with contact inhibition. The proliferation was reduced in the group of overexpressing miR-183~96~182 with ShhN CM at 72 hours. However, when cells were cultured in the absence of ShhN, they showed a decline in proliferation. In the beginning, the proliferation of miR-183~96~182-infected CGNPs was decreased slower than control at 36 hours, which then showed a continuous decline. There were no significant differences for proliferation between miR-183~96~182-infected CGNPs and control-infected CGNPs from 48 hours (**Fig. 3C**). Therefore, miR-183~96~182 was unable to maintain cell proliferation in the absence of Shh, suggesting that its role in promoting CGNPs proliferation is dependent on hedgehog signaling activation.

DISCUSSION

Chromosome 10q deletion is frequently identified in human medulloblastoma, and is shown to be highly associated with survival probability. Chromosome 10q loss was primarily limited to Shh and group C tumors^[26]. Adult Shh-medulloblastomas harboring 10q loss show particularly poor progression-free survival and overall survival probabilities^[27]. *PTEN*, the tumor suppressor gene of phosphatase and tensin homolog, is located on the chromosomal region 10q23.31. We established a mouse model to present the important role for cross talk between *Pten* loss and sonic hedgehog signaling pathways in the pathogenesis of medulloblastoma.

We inactivated *Pten* during brain development in neuronal and glial cell populations in a precisely defined area of the cerebellum. It was achieved by using *GFAP-Cre* transgenic mice, in which *Cre* is active in EGL, the Bergmann glia and white matter of the cerebellum. *Pten*^{Flox/Flox} *GFAP-Cre* mice developed seizures and ataxia early in life and died prematurely,

which is similar to other previous studies^[28,29]. Although *PTEN* is frequently inactivated in malignant human brain tumors, inactivation of *Pten* in the mouse brain does not lead to tumor development, which may be due to the fact that additional genetic alterations are necessary. We used the *Ptch1-lacZ* mouse model of medulloblastoma to study the effect of *Pten* loss on medulloblastoma tumorigenesis. We found that the heterozygosity for *Pten*, in the context of heterozygosity for *Ptch1*, altered tumor histology and accelerated medulloblastoma tumorigenesis in *Ptch1*^{+/-} *Pten*^{Flox/+} *GFAP-Cre* mice. Although spontaneous medulloblastoma harbor only monoallelic mutations of *Pten* possess at least one functional wildtype *Pten* allele, they further or completely lose *Pten* protein immunoreactivity^[30]. These results showed that epigenetic silencing by aberrant promoter methylation, deregulated transcription and/or increased degradation may disrupt function and promote tumorigenesis. As a result, loss of *Pten* expression leads to activation of PI3K signaling: increased expression of activated Akt, phosphorylated on serine 473, in *Ptch1*^{+/-} *Pten*^{Flox/+} *GFAP-Cre* medulloblastoma (data not shown).

Here we examined the role of miR-183~96~182 in the context of heterozygosity for *Pten* and *Ptch1* associated medulloblastoma. The miR-183~96~182 cluster that upregulated across *Ptch1*^{+/-} *Pten*^{Flox/+} *GFAP-Cre* medulloblastoma samples is relative to *Ptch1*^{+/-} medulloblastoma. MiR-183~96~182 has been implicated in retinal development and stem-cell maintenance^[31-33]. With the expression of *GFAP-Cre* and recombination occur in scattered EGL cells before neoplastic transformation, we concluded that in *Ptch1*^{+/-} *Pten*^{Flox/+} *GFAP-Cre* mouse model, medulloblastomas arise from cerebellar granule neuron precursors retained on the cerebellar surface. CGNPs taken from the P6~7 cerebellums have greater proliferative capacity than those harvested later, when neuronal progenitors have already begun to exit the cell division cycle and migrate into the internal granular layer. When explanted into culture in the absence of ShhN, CGNP rapidly exit the cell cycle and differentiate, but ShhN additionally extends their proliferative potential^[19]. CGNPs harvested from C57BL/6 mice and engineered to ectopically express miR-183~96~182 by retroviral transfer demonstrated a greater proliferative advantage than their Retro-GFP counterparts in the presence of ShhN. However, its role in promoting CGNPs proliferation is dependent on hedgehog signaling activation. When cultured in vitro for 48 hours, CGNPs without ShhN-CM lost the effort of endogenous ShhN and declined sharply in the proliferation, despite enforced expression of miR-183~96~182. Our




Fig. 3 Enforced expression of the miR-183~96~182 cluster promotes CGNPs proliferation. A: Immunofluorescence staining for EdU incorporation. Freshly isolated CGNPs were grown in poly-D-lysine and matrigel coated chamber slides in the presence of ShhN CM or 293T CM. The cells were infected with retroviral miR-183~96~182 or empty-control. B: Kinetics of proliferative response to miR-183~96~182 or empty-control in the presence ShhN CM or 293T CM. The numbers at each data point were counted from 4 random fields. C: Column diagram shows quantification of EdU-positive CGNPs after retroviral infection for 48 hours and 60 hours. The bars denote standard deviation.

data concluded that miR-183~96~182 was unable to maintain cell proliferation in the absence of ShhN. miR-183~96~182 could amplify the effects of Shh signaling, perhaps by inhibiting a transcriptional repressor of a subset of Shh signaling target genes.

In conclusion, our study provides evidence for an important role of miR-183~96~182 cluster in heterozygosity for *Pten* and *Ptch1* associated medulloblastomas. Antagomirs to the miR-183~96~182 cluster might provide a potential new therapeutic strategy for patients with medulloblastomas harboring a constitutively activated PI3K signaling pathway.

References

- [1] Ellison D. Classifying the medulloblastoma: insights from morphology and molecular genetics. *Neuropathol Appl Neurobiol* 2002; 28: 257-82.
- [2] Packer RJ, Cogen P, Vezina G, Rorke LB. Medulloblastoma: clinical and biologic aspects. *Neuro Oncol* 1999; 1: 232-50.
- [3] Heikens J, Michiels EM, Behrendt H, Endert E, Bakker PJ, Fliers E. Long-term neuro-endocrine sequelae after treatment for childhood medulloblastoma. *Eur J Cancer* 1998; 34: 1592-7.
- [4] Mulhern RK, Palmer SL, Merchant TE, Wallace D, Kocak M, Brouwers P, et al. Neurocognitive consequences of risk-adapted therapy for childhood medulloblastoma. *J Clin Oncol* 2005; 23: 5511-9.
- [5] Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006; 6: 857-66.
- [6] He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, et al. A microRNA polycistron as a potential human oncogene. *Nature* 2005; 435: 828-33.
- [7] Carmell MA, Hannon GJ. RNase III enzymes and the initiation of gene silencing. *Nat Struct Mol Biol* 2004; 11: 214-8.
- [8] Northcott PA, Fernandez LA, Hagan JP, Ellison DW, Grajkowska W, Gillespie Y, et al. The miR-17/92 polycistron is up-regulated in sonic hedgehog-driven medulloblastomas and induced by N-myc in sonic hedgehog-treated cerebellar neural precursors. *Cancer Res* 2009; 69: 3249-55.
- [9] Uziel T, Karginov FV, Xie S, Parker JS, Wang YD, Gajjar A, et al. The miR-17~92 cluster collaborates with the Sonic Hedgehog pathway in medulloblastoma. *Proc Natl Acad Sci U S A* 2009; 106: 2812-7.
- [10] Cho YJ, Tsherniak A, Tamayo P, Santagata S, Ligon A, Greulich H, et al. Integrative genomic analysis of medulloblastoma identifies a molecular subgroup that drives poor clinical outcome. *J Clin Oncol* 2011; 29: 1424-30.
- [11] Bai AH, Milde T, Remke M, Rolli CG, Hielscher T, Cho YJ, et al. MicroRNA-182 promotes leptomeningeal spread of non-sonic hedgehog-medulloblastoma. *Acta Neuropathol* 2012; 123: 529-38.
- [12] Weeraratne SD, Amani V, Teider N, Pierre-Francois J, Winter D, Kye MJ, et al. Pleiotropic effects of miR-183~96~182 converge to regulate cell survival, proliferation and migration in medulloblastoma. *Acta Neuropathol* 2012; 123: 539-52.
- [13] Yang ZJ, Ellis T, Markant SL, Read TA, Kessler JD, Bourboulas M, et al. Medulloblastoma can be initiated by deletion of Patched in lineage-restricted progenitors or stem cells. *Cancer Cell* 2008; 14: 135-45.
- [14] Rubin JB, Rowitch DH. Medulloblastoma: a problem of developmental biology. *Cancer Cell* 2002; 2: 7-8.
- [15] Goodrich LV, Milenkovic L, Higgins KM, Scott MP. Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science* 1997; 277: 1109-13.
- [16] Suzuki A, Yamaguchi MT, Ohteki T, Sasaki T, Kaisho T, Kimura Y, et al. T cell-specific loss of Pten leads to defects in central and peripheral tolerance. *Immunity* 2001; 14: 523-34.
- [17] Peng YJ, He WQ, Tang J, Tao T, Chen C, Gao YQ, et al. Trio is a key guanine nucleotide exchange factor coordinating regulation of the migration and morphogenesis of granule cells in the developing cerebellum. *J Biol Chem* 2010; 285: 24834-44.
- [18] Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, et al. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res* 2005; 33: e179.
- [19] Wechsler-Reya RJ, Scott MP. Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron* 1999; 22: 103-14.
- [20] Zindy F, Uziel T, Ayrault O, Calabrese C, Valentine M, Reh J, et al. Genetic alterations in mouse medulloblastomas and generation of tumors de novo from primary cerebellar granule neuron precursors. *Cancer Res* 2007; 67: 2676-84.
- [21] Yue S, Chen Y, Cheng SY. Hedgehog signaling promotes the degradation of tumor suppressor Sufu through the ubiquitin-proteasome pathway. *Oncogene* 2009; 28: 492-9.
- [22] Read TA, Fogarty MP, Markant SL, McLendon RE, Wei Z, Ellison DW, et al. Identification of CD15 as a marker for tumor-propagating cells in a mouse model of medulloblastoma. *Cancer Cell* 2009; 15: 135-47.
- [23] Uziel T, Zindy F, Xie S, Lee Y, Forget A, Magdaleno S, et al. The tumor suppressors Ink4c and p53 collaborate independently with Patched to suppress medulloblastoma formation. *Genes Dev* 2005; 19: 2656-67.
- [24] Schuller U, Heine VM, Mao J, Kho AT, Dillon AK, Han YG, et al. Acquisition of granule neuron precursor identity is a critical determinant of progenitor cell competence to form Shh-induced medulloblastoma. *Cancer Cell* 2008; 14: 123-34.
- [25] Gibson P, Tong Y, Robinson G, Thompson MC, Curre DS, Eden C, et al. Subtypes of medulloblastoma have distinct developmental origins. *Nature* 2010; 468: 1095-9.
- [26] Northcott PA, Korshunov A, Witt H, Hielscher T, Eber-

- hart CG, Mack S, et al. Medulloblastoma comprises four distinct molecular variants. *J Clin Oncol* 2011; 29: 1408-14.
- [27] Northcott PA, Hielscher T, Dubuc A, Mack S, Shih D, Remke M, et al. Pediatric and adult sonic hedgehog medulloblastomas are clinically and molecularly distinct. *Acta Neuropathol* 2011; 122: 231-40.
- [28] Backman SA, Stambolic V, Suzuki A, Haight J, Elia A, Pretorius J, et al. Deletion of Pten in mouse brain causes seizures, ataxia and defects in soma size resembling Lhermitte-Duclos disease. *Nat Genet* 2001; 29: 396-403.
- [29] Kwon CH, Zhu X, Zhang J, Knoop LL, Tharp R, Smeyne RJ, et al. Pten regulates neuronal soma size: a mouse model of Lhermitte-Duclos disease. *Nat Genet* 2001; 29: 404-11.
- [30] Castellino RC, Barwick BG, Schniederjan M, Buss MC, Becher O, Hambardzumyan D, et al. Heterozygosity for Pten promotes tumorigenesis in a mouse model of medulloblastoma. *PLoS One* 2010; 5: e10849.
- [31] Loscher CJ, Hokamp K, Kenna PF, Ivens AC, Humphries P, Palfi A, et al. Altered retinal microRNA expression profile in a mouse model of retinitis pigmentosa. *Genome Biol* 2007; 8: R248.
- [32] Loscher CJ, Hokamp K, Wilson JH, Li T, Humphries P, Farrar GJ, et al. A common microRNA signature in mouse models of retinal degeneration. *Exp Eye Res* 2008; 87: 529-34.
- [33] Viswanathan SR, Mermel CH, Lu J, Lu CW, Golub TR, Daley GQ. microRNA expression during trophectoderm specification. *PLoS One* 2009; 4: e6143.

CLINICAL TRIAL REGISTRATION

The *Journal* requires investigators to register their clinical trials in a public trials registry for publication of reports of clinical trials in the *Journal*. Information on requirements and acceptable registries is available at www.icmje.org/faq_clinical.html.