



Received 4 February 2016

Accepted 16 February 2016

Edited by R. F. Baggio, Comisión Nacional de  
Energía Atómica, Argentina**Keywords:** crystal structure; cisplatin; platinum(II); benzothiazole; anticancer.**CCDC references:** 1441324; 1441327;  
1441326; 1441325**Supporting information:** this article has  
supporting information at journals.iucr.org/e

## New platinum(II) complexes with benzothiazole ligands

José A. Carmona-Negrón,<sup>a</sup> Mayra E. Cádiz,<sup>a</sup> Curtis E. Moore,<sup>b</sup> Arnold L. Rheingold<sup>b</sup>  
and Enrique Meléndez<sup>a\*</sup>


<sup>a</sup>University of Puerto Rico, Department of Chemistry, PO Box 9019, Mayaguez, PR 00681, Puerto Rico, and <sup>b</sup>University of California-San Diego, Department of Chemistry, Urey Hall 5128, 9500 Gilman Drive, La Jolla, CA 92093-0358, USA.

\*Correspondence e-mail: enrique.melendez@upr.edu


Four new platinum(II) complexes, namely tetraethylammonium tribromido(2-methyl-1,3-benzothiazole- $\kappa N$ )platinate(II), [NEt<sub>4</sub>][PtBr<sub>3</sub>(C<sub>8</sub>H<sub>7</sub>NS)] (**1**), tetraethylammonium tribromido(6-methoxy-2-methyl-1,3-benzothiazole- $\kappa N$ )platinate(II), [NEt<sub>4</sub>][PtBr<sub>3</sub>(C<sub>9</sub>H<sub>9</sub>NOS)] (**2**), tetraethylammonium tribromido(2,5,6-trimethyl-1,3-benzothiazole- $\kappa N$ )platinate(II), [NEt<sub>4</sub>][PtBr<sub>3</sub>(C<sub>10</sub>H<sub>11</sub>NS)] (**3**), and tetraethylammonium tribromido(2-methyl-5-nitro-1,3-benzothiazole- $\kappa N$ )platinate(II), [NEt<sub>4</sub>][PtBr<sub>3</sub>(C<sub>8</sub>H<sub>6</sub>N<sub>2</sub>O<sub>2</sub>S)] (**4**), have been synthesized and structurally characterized by single-crystal X-ray diffraction techniques. These species are precursors of compounds with potential application in cancer chemotherapy. All four platinum(II) complexes adopt the expected square-planar coordination geometry, and the benzothiazole ligand is engaged in bonding to the metal atom through the imine N atom (Pt–N). The Pt–N bond lengths are normal: 2.035 (5), 2.025 (4), 2.027 (5) and 2.041 (4) Å for complexes **1**, **2**, **3** and **4**, respectively. The benzothiazole ligands are positioned out of the square plane, with dihedral angles ranging from 76.4 (4) to 88.1 (4)°. The NEt<sub>4</sub> cation in **3** is disordered with 0.57/0.43 occupancies.


### 1. Chemical context

The synthesis of new platinum complexes as potential drugs for cancer is still of interest for medicinal chemists. The structural details of these complexes provide the opportunity to predict, to a certain extent, the potential biological activity of these species. In this regard, four new platinum(II) complexes with benzothiazole ligands of general formula [PtBr<sub>3</sub>L]<sup>−</sup> have been synthesized according to the equation below and their structures characterized.



*L* = 2-methyl-1,3-benzothiazole (**1**), 6-methoxy-2-methyl-1,3-benzothiazole (**2**), 2,5,6-trimethyl-1,3-benzothiazole (**3**), and 2-methyl-5-nitro-1,3-benzothiazole (**4**). All complexes showed the benzothiazoles to coordinate the Pt<sup>II</sup> atom through the imino nitrogen atom. Also, the benzothiazole is positioned out of the square plane with dihedral angles between 76.4 (4) and 88.1 (4)°, as previously reported in other platinum–benzothiazole complexes. Given that benzothiazoles have anticancer properties, these platinum complexes may have enhanced properties as a result of potential synergism between the ligand and Pt<sup>II</sup>. This deserves further studies as suggested by Noolvi *et al.* (2012)






- 1**  $R_1 = R_2 = \text{H}$ ;    **2**  $R_1 = \text{H}$ ,     $R_2 = \text{OCH}_3$   
**3**  $R_1 = R_2 = \text{CH}_3$ ;    **4**  $R_1 = \text{NO}_2$ ,     $R_2 = \text{H}$

## 2. Structural commentary

To elucidate with certainty and accurately the platinum coordination patterns, the structural determination of the complexes was performed by single crystal X-ray diffraction technique. Table 1 contains selected bond lengths, dihedral angles and torsion angles. All of the title complexes adopt a square-planar coordination geometry about the  $\text{Pt}^{\text{II}}$  atom with a deviation of no more than  $4^\circ$  from ideal  $180^\circ$  and  $90^\circ$  angles. As reported previously, although not predicted using Pearson's hard-soft acid base theory, the benzothiazole is engaged in bonding to the metal through the imine nitrogen (Pt–N) instead of Pt–S coordination (Muir *et al.*, 1987, 1988*a,b*, 1990; Gomez *et al.*, 1988; Lozano *et al.*, 1994). Also the benzothia-



**Figure 1**  
 The molecular structure of  $[\text{NEt}_4][\text{PtBr}_3(2\text{-Me-benzothiazole})]$  (**1**), with displacement ellipsoids drawn at the 50% probability level.

zole ligand is positioned out of the square plane as discussed below.


Figs. 1–4 show the molecular structures of the four new complexes.  $[\text{NEt}_4][\text{PtBr}_3(2\text{-Me-benzothiazole})]$  (**1**) crystallizes in an orthorhombic unit cell with eight formula units. It is a square-planar complex with Pt–N and average Pt–Br bond lengths of 2.035 (5) and 2.433 (6) Å, respectively, which are within the expected range for  $\text{Pt}^{\text{II}}$  complexes. There is no *trans*-influence observed in the Pt–Br bond *trans* to the Pt–N bond. The benzothiazole ligand is planar and the methyl group resides in the ligand plane. The dihedral angle between the  $\text{PtBr}_3\text{N}$  unit and the benzothiazole ring is  $88.1 (4)^\circ$ , similar to those observed in other  $\text{Pt}^{\text{II}}$ –benzothiazole complexes, as a result of reducing the steric strain between  $\text{PtBr}_3$  and the benzothiazole ligand (Muir *et al.*, 1987, 1988*a,b*, 1990; Gomez *et al.*, 1988; Lozano *et al.*, 1994). Two types of N–C bonds are present, one long [N–C2 1.408 (7) Å] and one short [N–C1

**Table 1**


Selected bond distances and angles (Å, °).

The dihedral angle is between the Pt–Br<sub>3</sub>N unit and the benzothiazole ring. The torsion angle is between the benzothiazole ring and the R group.

	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Pt–Br <sub>average</sub>	2.433 (6)	2.430 (6)	2.425 (6)	2.431 (8)
Pt–N	2.035 (5)	2.025 (4)	2.027 (5)	2.041 (4)
N1–C2	1.408 (7)	1.396 (6)	1.401 (8)	1.383 (6)
N1–C1	1.309 (7)	1.309 (6)	1.303 (8)	1.315 (6)
Pt–Br1	2.4375 (8)	2.4352 (5)	2.4309 (7)	2.4335 (6)
Pt–Br2	2.4349 (8)	2.4241 (7)	2.4198 (7)	2.4216 (5)
Pt–Br3	2.4268 (7)	2.4309 (5)	2.4240 (7)	2.4367 (5)
S–C7	1.744 (6)	1.743 (5)	1.739 (7)	1.738 (5)
S–C1	1.735 (6)	1.730 (5)	1.727 (6)	1.724 (5)
C1–N1–C2	113.0 (5)	112.6 (4)	112.3 (5)	111.9 (4)
C1–S–C7	90.3 (3)	89.9 (2)	89.8 (3)	90.0 (2)
N1–Pt–Br1	90.6 (1)	87.0 (1)	89.2 (1)	88.6 (1)
N1–Pt–Br3	86.4 (1)	89.3 (1)	88.5 (1)	89.3 (1)
N1–Pt–Br2	177.7 (1)	177.4 (1)	178.8 (1)	178.4 (1)
Br1–Pt–Br3	176.85 (2)	176.30 (2)	177.45 (3)	176.23 (2)
Br2–Pt–Br3	91.69 (2)	92.51 (2)	91.23 (2)	91.18 (2)
Br1–Pt–Br2	91.31 (2)	91.17 (2)	91.10 (2)	90.99 (2)
Dihedral angle	88.1 (4)	86.7 (3)	78.6 (4)	76.4 (4)
Torsion angle	0.72 (1) (CH <sub>3</sub> )	11.9 (7) (OCH <sub>3</sub> )	1.5 (5) (C <sub>8</sub> H <sub>3</sub> ) 0.2 (6) (C <sub>9</sub> H <sub>3</sub> ) 0.3 (6) (C <sub>10</sub> H <sub>3</sub> )	1.1 (5) (CH <sub>3</sub> ) 7.5 (7) (NO <sub>2</sub> )




**Figure 2**  
The molecular structure of  $[\text{NEt}_4][\text{PtBr}_3(6\text{-OMe-2-Me-benzothiazole})]$  (**2**), with displacement ellipsoids drawn at the 50% probability level.



**Figure 3**  
The molecular structure of  $[\text{NEt}_4][\text{PtBr}_3(2,5,6\text{-Me-benzothiazole})]$  (**3**), with displacement ellipsoids drawn at the 50% probability level. The  $\text{NEt}_4$  cation in **3** presented disorder with 0.57/0.43 occupancies. Only the major fraction is shown for clarity.



1.309 (7) Å], indicating the presence of single- and double-bond character in the thiazole ring. The angle at the S atom in the thiazole ring is  $90.3(3)^\circ$  suggesting it is using unhybridized  $p$  orbitals for bonding.

$[\text{NEt}_4][\text{PtBr}_3(6\text{-OMe-2-Me-benzothiazole})]$  (**2**),  $[\text{NEt}_4][\text{PtBr}_3(2,5,6\text{-Me-benzothiazole})]$  (**3**) and  $[\text{NEt}_4][\text{PtBr}_3(5\text{-NO}_2\text{-2-Me-benzothiazole})]$  (**4**) crystallize in the same type of unit cell and space group, monoclinic  $P2_1/n$ , containing four



**Figure 4**  
The molecular structure of  $[\text{NEt}_4][\text{PtBr}_3(5\text{-NO}_2\text{-2-Me-benzothiazole})]$  (**4**), with displacement ellipsoids drawn at the 50% probability level.

formula units. The Pt–N and average Pt–Br bond lengths for **2**, **3**, and **4** are 2.025 (4)/2.430 (6) Å, 2.027 (5)/2.425 (6) Å and 2.041 (4)/2.431 (8) Å, respectively, which are within the expected range. The dihedral angle between  $\text{PtBr}_3\text{N}$  and the benzothiazole in **2** is  $86.7(3)^\circ$  and the torsion angle between the aromatic ring and the  $\text{OCH}_3$  group is  $11.9(7)^\circ$ . The C–O ( $\text{OCH}_3$ ) bond length is 1.427 (7) Å, and the C–O– $\text{CH}_3$  angle is  $116.3(5)^\circ$ . In contrast to **1** and **2**,  $[\text{NEt}_4][\text{PtBr}_3(2,5,6\text{-Me-benzothiazole})]$  and  $[\text{NEt}_4][\text{PtBr}_3(5\text{-NO}_2\text{-2-Me-benzothiazole})]$  have lower dihedral angles between the  $\text{PtBr}_3\text{N}$  unit and the benzothiazole ring,  $78.6(4)$  and  $76.4(4)^\circ$ , respectively. The methyl groups on **3** and **4** are almost co-planar with the benzothiazole plane with deviations  $\leq 1.60^\circ$  but in **4**, the  $\text{NO}_2$  group is out of the benzothiazole plane with a torsion angle of



**Figure 5**  
Details of the packing interactions in (a)  $[\text{NEt}_4][\text{PtBr}_3(2\text{-Me-benzothiazole})]$  and (b)  $[\text{NEt}_4][\text{PtBr}_3(6\text{-Ome-2-Me-benzothiazole})]$ .

**Table 2**  
Experimental details.

	(1)	(2)	(3)	(4)
<b>Crystal data</b>				
Chemical formula	(C <sub>8</sub> H <sub>20</sub> N)[PtBr <sub>3</sub> (C <sub>8</sub> H <sub>7</sub> NS)]	(C <sub>8</sub> H <sub>20</sub> N)[PtBr <sub>3</sub> (C <sub>9</sub> H <sub>9</sub> NOS)]	(C <sub>8</sub> H <sub>20</sub> N)[PtBr <sub>3</sub> (C <sub>10</sub> H <sub>11</sub> NS)]	(C <sub>8</sub> H <sub>20</sub> N)[PtBr <sub>3</sub> (C <sub>8</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> S)]
<i>M<sub>r</sub></i>	714.27	744.30	742.33	759.28
Crystal system, space group	Orthorhombic, <i>Pbca</i>	Monoclinic, <i>P2<sub>1</sub>/n</i>	Monoclinic, <i>P2<sub>1</sub>/n</i>	Monoclinic, <i>P2<sub>1</sub>/n</i>
Temperature (K)	100	100	100	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	12.114 (3), 10.656 (3), 34.043 (9)	7.7591 (2), 30.4214 (8), 9.6551 (3)	7.9742 (4), 30.2807 (14), 9.6427 (5)	8.1170 (3), 29.2717 (12), 9.5102 (4)
$\alpha$ , $\beta$ , $\gamma$ (°)	90, 90, 90	90, 94.539 (1), 90	90, 100.151 (3), 90	90, 100.720 (1), 90
<i>V</i> (Å <sup>3</sup> )	4394 (2)	2271.87 (11)	2291.9 (2)	2220.17 (15)
<i>Z</i>	8	4	4	4
Radiation type	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$
$\mu$ (mm <sup>-1</sup> )	11.94	11.55	11.45	11.83
Crystal size (mm)	0.18 × 0.16 × 0.12	0.32 × 0.30 × 0.24	0.50 × 0.36 × 0.25	0.32 × 0.30 × 0.25
<b>Data collection</b>				
Diffractometer	Bruker APEXII CCD	Bruker APEXII CCD	Bruker APEXII CCD	Bruker APEXII CCD
Absorption correction	Multi-scan ( <i>SADABS</i> ; Bruker, 2014)	Multi-scan ( <i>SADABS</i> ; Bruker, 2014)	Multi-scan ( <i>SADABS</i> ; Bruker, 2014)	Multi-scan ( <i>SADABS</i> ; Bruker, 2014)
<i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>	0.052, 0.093	0.056, 0.093	0.003, 0.028	0.020, 0.045
No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections	16951, 4418, 3675	12741, 4650, 4377	10729, 4692, 4120	15975, 4550, 4254
<i>R<sub>int</sub></i> ( $\sin \theta/\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )	0.047 0.623	0.017 0.626	0.048 0.627	0.028 0.627
<b>Refinement</b>				
<i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.031, 0.081, 1.03	0.027, 0.066, 1.08	0.039, 0.106, 1.05	0.029, 0.060, 1.18
No. of reflections	4418	4650	4692	4550
No. of parameters	213	232	266	240
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\max}$ , $\Delta\rho_{\min}$ (e Å <sup>-3</sup> )	2.38, -0.93	1.25, -1.36	1.88, -1.02	1.25, -1.37

Computer programs: *APEX2* and *SAINT* (Bruker, 2013), *SHELXT* (Sheldrick, 2015a), *SIR2004* (Burla *et al.*, 2007), *SHELXL2014* (Sheldrick, 2015b) and *OLEX2* (Dolomanov *et al.*, 2009).

7.5 (7)°. The C—NO<sub>2</sub> bond length is 1.476 (7) Å, and the O—N—O angle is 117.4 (5)°. The C—NO<sub>2</sub> bond length and O—N—O angle in **4** are smaller than those observed in nitrobenzene [C—NO<sub>2</sub> = 1.486 (2) Å and O—N—O = 123.9 (5)°], which suggests higher electron delocalization between the nitro group and the aromatic ring in **4** (Johnson, 2015). The angles at the S atom in **2**, **3** and **4** are also near 90°, suggesting the use of pure *p* orbitals for bonding.

### 3. Supramolecular features

Analysis of the packing diagrams of all of the complexes showed their packings consist of [NEt<sub>4</sub>]<sup>+</sup> cations and [PtBr<sub>3</sub>(*L*)]<sup>-</sup> anions. The [NEt<sub>4</sub>][PtBr<sub>3</sub>(2-Me-benzothiazole)] and [NEt<sub>4</sub>][PtBr<sub>3</sub>(6-OMe-2-Me-benzothiazole)] complexes showed partial  $\pi$ -stacking between the phenyl and the thiazole rings (Fig. 5).

### 4. Synthesis and crystallization

The parent complex [NEt<sub>4</sub>]<sub>2</sub>[Pt<sub>2</sub>Br<sub>6</sub>] was prepared as reported in the literature (Livingstone & Whitley, 1962). Ligands were purchased from Sigma–Aldrich and were used without further purification.

Acetone solutions of [NEt<sub>4</sub>]<sub>2</sub>[Pt<sub>2</sub>Br<sub>6</sub>] were prepared (0.075 g, 0.068 mmol) and the corresponding amount of ligand was added with stirring. For 2-methyl-1,3-benzothiazole (99%) 18  $\mu$ L (0.021 g, 0.14 mmol) were added; for 2-methyl-5-nitro-1,3-benzothiazole (98%) (0.027 g, 0.14 mmol) were added, and for 2-methyl-6-methoxy-1,3-benzothiazole (97%) (0.024 g, 0.14 mmol) were added. The reaction mixtures were stirred without heating until the volume reduced considerably; then the samples were placed in desiccators containing CaCl<sub>2</sub> at room temperature to evaporate slowly, leading to the formation of X-ray quality single crystals. For the synthesis with 2,5,6-trimethyl-1,3-benzothiazole (99%), the ligand (0.0227 g, 0.128 mmol) was added to 20 mL of an acetone solution (0.07515 g, 0.0677 mmol) of [NEt<sub>4</sub>]<sub>2</sub>[Pt<sub>2</sub>Br<sub>6</sub>] with stirring, and a portion of the reaction mixture was slowly evaporated at 277 K in a small beaker in a secondary container which also contained CaCl<sub>2</sub> to form X-ray quality single crystals.

### 5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms were positioned in idealized locations: *d*(C—H) = 0.95 Å, *U*<sub>iso</sub>(H) = 1.2*U*<sub>eq</sub>(C); *d*(C—H<sub>2</sub>) = 0.99 Å, *U*<sub>iso</sub>(H) = 1.2*U*<sub>eq</sub>(C); *d*(C—H<sub>3</sub>) = 0.98 Å,

$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ . The  $\text{NEt}_4$  cation in **3** presented disorder with 0.57/0.43 occupancies.

### Acknowledgements

We thank Ms Lorraine Hernández and Ms Nivia Ruiz-Alago for their help with the synthesis of the platinum compounds. We are grateful to Dr Jorge Rios-Steiner and Mr Daniel J. Vallés-Cádiz for their assistance in the crystallization process. EM thanks the NIH for financial support and JACN acknowledges the financial support of Sloan Program.

### References

- Bruker (2013). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2014). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G., Siliqi, D. & Spagna, R. (2007). *J. Appl. Cryst.* **40**, 609–613.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Gomez, G. M., Muir, M. M., Muir, J. A. & Cox, O. (1988). *Acta Cryst. C* **44**, 1554–1557.
- Johnson, R. D. III (2015). *NIST Computational Chemistry Comparison and Benchmark Database, NIST Standard Reference Database Number 101 Release 17b*, September 2015, edited by Russell D. Johnson III. <http://cccbdb.NIST.gov/exp2.asp?casno=98953> (accessed on November 24, 2015).
- Livingstone, S. E. & Whitley, A. (1962). *Aust. J. Chem.* **15**, 175–180.
- Lozano, C. M., Muir, M. M., Tang, X. & Li, Y. (1994). *J. Chem. Cryst.* **24**, 639–642.
- Muir, M. M., Cadiz, M. E. & Baez, A. (1988a). *Inorg. Chim. Acta*, **151**, 209–213.
- Muir, M. M., Gomez, G. M., Cadiz, M. E. & Muir, J. E. (1990). *Inorg. Chim. Acta*, **168**, 47–57.
- Muir, M. M., Gomez, G., Muir, J. A., Cadiz, M. E., Cox, O. & Barnes, C. L. (1988b). *Acta Cryst. C* **44**, 803–806.
- Muir, J. A., Gomez, G. M., Muir, M. M., Cox, O. & Cadiz, M. E. (1987). *Acta Cryst. C* **43**, 1258–1261.
- Noolvi, M. N., Patel, H. M. & Kaur, M. (2012). *Eur. J. Med. Chem.* **54**, 447–462.
- Sheldrick, G. M. (2015a). *Acta Cryst. A* **71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst. C* **71**, 3–8.

## supporting information

*Acta Cryst.* (2016). E72, 412-416 [doi:10.1107/S2056989016002826]

## New platinum(II) complexes with benzothiazole ligands

José A. Carmona-Negrón, Mayra E. Cádiz, Curtis E. Moore, Arnold L. Rheingold and Enrique Meléndez

## Computing details

For all compounds, data collection: *APEX2* (Bruker, 2013); cell refinement: *S SAINT* (Bruker, 2013); data reduction: *S SAINT* (Bruker, 2013). Program(s) used to solve structure: *SHELXT* (Sheldrick, 2015a) for (1), (2), (3); *SIR2004* (Burla *et al.*, 2007) for (4). For all compounds, program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

(1) Tetraethylammonium tribromido(2-methyl-1,3-benzothiazole- $\kappa$ N)platinate(II)*Crystal data*

(C<sub>8</sub>H<sub>20</sub>N)[PtBr<sub>3</sub>(C<sub>8</sub>H<sub>7</sub>NS)]

$M_r = 714.27$

Orthorhombic, *Pbca*

$a = 12.114$  (3) Å

$b = 10.656$  (3) Å

$c = 34.043$  (9) Å

$V = 4394$  (2) Å<sup>3</sup>

$Z = 8$

$F(000) = 2688$

$D_x = 2.159$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 5330 reflections

$\theta = 2.4$ – $26.3^\circ$

$\mu = 11.94$  mm<sup>-1</sup>

$T = 100$  K

Block, bronze

$0.18 \times 0.16 \times 0.12$  mm

*Data collection*

Bruker APEXII CCD  
diffractometer

Radiation source: Micro Focus Rotating Anode,  
Bruker TXS

Double Bounce Multilayer Mirrors  
monochromator

Detector resolution: 7.9 pixels mm<sup>-1</sup>

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2014)

$T_{\min} = 0.052$ ,  $T_{\max} = 0.093$

16951 measured reflections

4418 independent reflections

3675 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 26.3^\circ$ ,  $\theta_{\min} = 2.1^\circ$

$h = -14 \rightarrow 15$

$k = -13 \rightarrow 10$

$l = -32 \rightarrow 42$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.031$

$wR(F^2) = 0.081$

$S = 1.03$

4418 reflections

213 parameters

0 restraints

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0352P)^2 + 9.4131P]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.003$

$$\Delta\rho_{\max} = 2.38 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.93 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N2	1.0097 (4)	0.4493 (4)	0.31862 (12)	0.0213 (9)
C9	1.1117 (4)	0.3738 (5)	0.30557 (16)	0.0242 (12)
H9A	1.1049	0.2872	0.3158	0.029*
H9B	1.1781	0.4117	0.3177	0.029*
C10	1.1289 (4)	0.3677 (5)	0.26030 (15)	0.0251 (12)
H10A	1.0624	0.3336	0.2479	0.038*
H10B	1.1920	0.3134	0.2544	0.038*
H10C	1.1431	0.4523	0.2502	0.038*
C11	1.0130 (5)	0.5845 (5)	0.30315 (17)	0.0259 (12)
H11A	1.0120	0.5820	0.2741	0.031*
H11B	0.9452	0.6282	0.3119	0.031*
C12	1.1120 (5)	0.6608 (5)	0.31618 (18)	0.0339 (14)
H12A	1.1104	0.7430	0.3033	0.051*
H12B	1.1799	0.6166	0.3089	0.051*
H12C	1.1097	0.6720	0.3447	0.051*
C13	1.0124 (5)	0.4436 (6)	0.36271 (15)	0.0259 (12)
H13A	1.0827	0.4809	0.3719	0.031*
H13B	1.0121	0.3543	0.3708	0.031*
C14	0.9162 (5)	0.5110 (5)	0.38363 (18)	0.0305 (13)
H14A	0.9172	0.6003	0.3768	0.046*
H14B	0.9243	0.5017	0.4121	0.046*
H14C	0.8460	0.4739	0.3753	0.046*
C15	0.9038 (4)	0.3934 (5)	0.30181 (16)	0.0248 (12)
H15A	0.8405	0.4417	0.3122	0.030*
H15B	0.9048	0.4048	0.2730	0.030*
C16	0.8844 (4)	0.2555 (5)	0.31049 (17)	0.0303 (13)
H16A	0.9447	0.2056	0.2993	0.045*
H16B	0.8142	0.2291	0.2988	0.045*
H16C	0.8819	0.2426	0.3390	0.045*
Pt1	0.47417 (2)	0.49411 (2)	0.37148 (2)	0.01931 (8)
Br1	0.31934 (5)	0.60895 (7)	0.39974 (2)	0.04565 (19)
Br2	0.39703 (4)	0.51437 (5)	0.30574 (2)	0.02401 (13)
Br3	0.63067 (4)	0.37553 (6)	0.34693 (2)	0.03153 (15)
S1	0.67084 (13)	0.48262 (14)	0.48520 (4)	0.0321 (3)
N1	0.5409 (4)	0.4702 (4)	0.42583 (13)	0.0221 (10)
C1	0.6192 (4)	0.5394 (5)	0.44109 (15)	0.0247 (12)

C2	0.5158 (4)	0.3638 (5)	0.44867 (15)	0.0246 (12)
C3	0.4356 (5)	0.2744 (5)	0.44047 (16)	0.0290 (12)
H3	0.3925	0.2796	0.4172	0.035*
C4	0.4200 (6)	0.1769 (6)	0.46718 (16)	0.0351 (14)
H4	0.3642	0.1162	0.4624	0.042*
C5	0.4853 (5)	0.1671 (7)	0.50112 (16)	0.0398 (16)
H5	0.4740	0.0987	0.5186	0.048*
C6	0.5666 (6)	0.2558 (6)	0.50970 (17)	0.0379 (15)
H6	0.6106	0.2495	0.5327	0.045*
C7	0.5807 (5)	0.3552 (5)	0.48283 (16)	0.0315 (13)
C8	0.6643 (5)	0.6567 (5)	0.42301 (16)	0.0290 (13)
H8A	0.7204	0.6929	0.4404	0.044*
H8B	0.6043	0.7173	0.4193	0.044*
H8C	0.6976	0.6368	0.3975	0.044*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N2	0.022 (2)	0.022 (2)	0.020 (2)	0.0039 (19)	0.0002 (18)	0.0024 (19)
C9	0.016 (3)	0.024 (3)	0.032 (3)	0.001 (2)	0.000 (2)	0.000 (2)
C10	0.019 (3)	0.025 (3)	0.032 (3)	0.003 (2)	0.004 (2)	-0.003 (2)
C11	0.026 (3)	0.021 (3)	0.031 (3)	0.002 (2)	0.004 (2)	0.006 (2)
C12	0.031 (3)	0.027 (3)	0.044 (4)	0.000 (3)	-0.004 (3)	0.002 (3)
C13	0.030 (3)	0.028 (3)	0.020 (3)	0.000 (3)	-0.001 (2)	-0.001 (2)
C14	0.030 (3)	0.037 (3)	0.025 (3)	-0.001 (3)	0.000 (2)	-0.003 (2)
C15	0.018 (3)	0.032 (3)	0.025 (3)	0.003 (2)	-0.005 (2)	-0.002 (2)
C16	0.021 (3)	0.031 (3)	0.039 (3)	-0.003 (2)	-0.001 (2)	0.002 (3)
Pt1	0.01841 (13)	0.02210 (12)	0.01742 (13)	0.00011 (8)	-0.00120 (7)	-0.00100 (8)
Br1	0.0399 (4)	0.0675 (5)	0.0296 (3)	0.0248 (3)	-0.0059 (3)	-0.0135 (3)
Br2	0.0241 (3)	0.0255 (3)	0.0224 (3)	-0.0012 (2)	-0.0022 (2)	-0.0005 (2)
Br3	0.0235 (3)	0.0411 (3)	0.0300 (3)	0.0027 (3)	0.0004 (2)	-0.0050 (3)
S1	0.0337 (8)	0.0381 (8)	0.0245 (7)	0.0055 (7)	-0.0105 (6)	-0.0028 (6)
N1	0.022 (2)	0.025 (2)	0.019 (2)	0.0030 (19)	0.0017 (18)	0.0000 (19)
C1	0.022 (3)	0.031 (3)	0.021 (3)	0.006 (2)	-0.004 (2)	-0.009 (2)
C2	0.030 (3)	0.026 (3)	0.018 (3)	0.005 (2)	0.001 (2)	-0.005 (2)
C3	0.033 (3)	0.032 (3)	0.021 (3)	-0.003 (3)	0.004 (2)	-0.005 (2)
C4	0.051 (4)	0.032 (3)	0.022 (3)	-0.004 (3)	0.006 (3)	-0.002 (2)
C5	0.062 (4)	0.032 (4)	0.026 (3)	0.003 (3)	0.005 (3)	0.003 (2)
C6	0.054 (4)	0.035 (3)	0.025 (3)	0.011 (3)	-0.004 (3)	0.001 (3)
C7	0.036 (3)	0.033 (3)	0.025 (3)	0.006 (3)	0.002 (3)	-0.010 (2)
C8	0.034 (3)	0.032 (3)	0.021 (3)	-0.001 (3)	-0.010 (2)	-0.004 (2)

*Geometric parameters (Å, °)*

N2—C9	1.540 (6)	C16—H16A	0.9800
N2—C11	1.534 (7)	C16—H16B	0.9800
N2—C13	1.503 (7)	C16—H16C	0.9800
N2—C15	1.526 (7)	Pt1—Br1	2.4375 (8)



C9—H9A	0.9900	Pt1—Br2	2.4349 (8)
C9—H9B	0.9900	Pt1—Br3	2.4268 (7)
C9—C10	1.557 (7)	Pt1—N1	2.035 (5)
C10—H10A	0.9800	S1—C1	1.735 (6)
C10—H10B	0.9800	S1—C7	1.744 (6)
C10—H10C	0.9800	N1—C1	1.309 (7)
C11—H11A	0.9900	N1—C2	1.408 (7)
C11—H11B	0.9900	C1—C8	1.497 (8)
C11—C12	1.515 (8)	C2—C3	1.389 (8)
C12—H12A	0.9800	C2—C7	1.407 (8)
C12—H12B	0.9800	C3—H3	0.9500
C12—H12C	0.9800	C3—C4	1.393 (8)
C13—H13A	0.9900	C4—H4	0.9500
C13—H13B	0.9900	C4—C5	1.405 (8)
C13—C14	1.543 (8)	C5—H5	0.9500
C14—H14A	0.9800	C5—C6	1.396 (9)
C14—H14B	0.9800	C6—H6	0.9500
C14—H14C	0.9800	C6—C7	1.410 (8)
C15—H15A	0.9900	C8—H8A	0.9800
C15—H15B	0.9900	C8—H8B	0.9800
C15—C16	1.517 (8)	C8—H8C	0.9800
C11—N2—C9	111.8 (4)	C16—C15—H15A	108.3
C13—N2—C9	104.5 (4)	C16—C15—H15B	108.3
C13—N2—C11	112.4 (4)	C15—C16—H16A	109.5
C13—N2—C15	112.1 (4)	C15—C16—H16B	109.5
C15—N2—C9	111.2 (4)	C15—C16—H16C	109.5
C15—N2—C11	105.1 (4)	H16A—C16—H16B	109.5
N2—C9—H9A	108.6	H16A—C16—H16C	109.5
N2—C9—H9B	108.6	H16B—C16—H16C	109.5
N2—C9—C10	114.5 (4)	Br2—Pt1—Br1	91.31 (2)
H9A—C9—H9B	107.6	Br3—Pt1—Br1	176.85 (2)
C10—C9—H9A	108.6	Br3—Pt1—Br2	91.69 (2)
C10—C9—H9B	108.6	N1—Pt1—Br1	90.56 (12)
C9—C10—H10A	109.5	N1—Pt1—Br2	177.68 (13)
C9—C10—H10B	109.5	N1—Pt1—Br3	86.41 (12)
C9—C10—H10C	109.5	C1—S1—C7	90.3 (3)
H10A—C10—H10B	109.5	C1—N1—Pt1	125.3 (4)
H10A—C10—H10C	109.5	C1—N1—C2	113.0 (5)
H10B—C10—H10C	109.5	C2—N1—Pt1	121.2 (4)
N2—C11—H11A	108.5	N1—C1—S1	114.1 (4)
N2—C11—H11B	108.5	N1—C1—C8	124.9 (5)
H11A—C11—H11B	107.5	C8—C1—S1	121.0 (4)
C12—C11—N2	115.1 (5)	C3—C2—N1	126.3 (5)
C12—C11—H11A	108.5	C3—C2—C7	120.8 (5)
C12—C11—H11B	108.5	C7—C2—N1	112.8 (5)
C11—C12—H12A	109.5	C2—C3—H3	120.8
C11—C12—H12B	109.5	C2—C3—C4	118.4 (5)

C11—C12—H12C	109.5	C4—C3—H3	120.8
H12A—C12—H12B	109.5	C3—C4—H4	119.5
H12A—C12—H12C	109.5	C3—C4—C5	121.0 (6)
H12B—C12—H12C	109.5	C5—C4—H4	119.5
N2—C13—H13A	108.5	C4—C5—H5	119.4
N2—C13—H13B	108.5	C6—C5—C4	121.3 (6)
N2—C13—C14	115.2 (5)	C6—C5—H5	119.4
H13A—C13—H13B	107.5	C5—C6—H6	121.4
C14—C13—H13A	108.5	C5—C6—C7	117.3 (6)
C14—C13—H13B	108.5	C7—C6—H6	121.4
C13—C14—H14A	109.5	C2—C7—S1	109.7 (4)
C13—C14—H14B	109.5	C2—C7—C6	121.2 (6)
C13—C14—H14C	109.5	C6—C7—S1	129.1 (5)
H14A—C14—H14B	109.5	C1—C8—H8A	109.5
H14A—C14—H14C	109.5	C1—C8—H8B	109.5
H14B—C14—H14C	109.5	C1—C8—H8C	109.5
N2—C15—H15A	108.3	H8A—C8—H8B	109.5
N2—C15—H15B	108.3	H8A—C8—H8C	109.5
H15A—C15—H15B	107.4	H8B—C8—H8C	109.5
C16—C15—N2	115.8 (4)		

(2) Tetraethylammonium tribromido(6-methoxy-2-methyl-1,3-benzothiazole- $\kappa$ N)platinate(II)*Crystal data* $(\text{C}_8\text{H}_{20}\text{N})[\text{PtBr}_3(\text{C}_9\text{H}_9\text{NOS})]$  $M_r = 744.30$ Monoclinic,  $P2_1/n$  $a = 7.7591$  (2) Å $b = 30.4214$  (8) Å $c = 9.6551$  (3) Å $\beta = 94.539$  (1)° $V = 2271.87$  (11) Å<sup>3</sup> $Z = 4$  $F(000) = 1408$  $D_x = 2.176$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7838 reflections

 $\theta = 2.2$ – $26.4$ ° $\mu = 11.55$  mm<sup>-1</sup> $T = 100$  K

Block, bronze

 $0.32 \times 0.3 \times 0.24$  mm*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: Micro Focus Rotating Anode,

Bruker TXS

Double Bounce Multilayer Mirrors

monochromator

Detector resolution: 7.9 pixels mm<sup>-1</sup> $\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2014)

 $T_{\min} = 0.056$ ,  $T_{\max} = 0.093$ 

12741 measured reflections

4650 independent reflections

4377 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.017$  $\theta_{\max} = 26.4$ °,  $\theta_{\min} = 2.2$ ° $h = -9 \rightarrow 9$  $k = -32 \rightarrow 38$  $l = -12 \rightarrow 7$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.027$  $wR(F^2) = 0.066$  $S = 1.08$ 

4650 reflections

232 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0227P)^2 + 15.6321P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.25 \text{ e } \text{Å}^{-3}$   
 $\Delta\rho_{\min} = -1.36 \text{ e } \text{Å}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{Å}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N2	0.4744 (6)	0.68841 (14)	1.0072 (4)	0.0224 (9)
C10	0.4365 (10)	0.6636 (2)	0.8727 (7)	0.0442 (16)
H10A	0.3248	0.6744	0.8289	0.053*
H10B	0.5266	0.6715	0.8099	0.053*
C11	0.4277 (12)	0.6153 (2)	0.8795 (10)	0.065 (3)
H11A	0.5326	0.6039	0.9305	0.097*
H11B	0.4183	0.6032	0.7851	0.097*
H11C	0.3263	0.6065	0.9274	0.097*
C12	0.6550 (9)	0.6759 (3)	1.0680 (8)	0.0529 (19)
H12A	0.6514	0.6451	1.1004	0.064*
H12B	0.7336	0.6769	0.9921	0.064*
C13	0.7308 (9)	0.7028 (3)	1.1828 (7)	0.061 (2)
H13A	0.6425	0.7094	1.2470	0.092*
H13B	0.7743	0.7304	1.1462	0.092*
H13C	0.8264	0.6868	1.2322	0.092*
C14	0.3484 (8)	0.6760 (2)	1.1136 (7)	0.0403 (15)
H14A	0.3582	0.6440	1.1319	0.048*
H14B	0.3825	0.6915	1.2018	0.048*
C15	0.1616 (7)	0.6868 (2)	1.0702 (6)	0.0338 (13)
H15A	0.0883	0.6767	1.1421	0.051*
H15B	0.1267	0.6719	0.9824	0.051*
H15C	0.1488	0.7186	1.0581	0.051*
C16	0.4651 (9)	0.7366 (2)	0.9800 (8)	0.0426 (15)
H16A	0.4764	0.7522	1.0703	0.051*
H16B	0.3493	0.7435	0.9349	0.051*
C17	0.6004 (9)	0.7545 (2)	0.8898 (7)	0.0425 (15)
H17A	0.7160	0.7485	0.9340	0.064*
H17B	0.5849	0.7864	0.8787	0.064*
H17C	0.5876	0.7404	0.7984	0.064*
Pt1	0.73847 (2)	0.62959 (2)	0.53521 (2)	0.01668 (6)
Br1	0.90363 (7)	0.66953 (2)	0.71969 (5)	0.02744 (12)
Br2	0.52471 (7)	0.68778 (2)	0.51179 (5)	0.02766 (12)
Br3	0.58868 (7)	0.58627 (2)	0.35144 (6)	0.02867 (12)
S1	1.20643 (16)	0.54060 (4)	0.58464 (13)	0.0233 (3)
O1	0.8812 (6)	0.43871 (13)	0.9169 (4)	0.0337 (9)
N1	0.9166 (5)	0.58111 (13)	0.5642 (4)	0.0184 (8)

C1	1.0717 (6)	0.58239 (17)	0.5202 (5)	0.0213 (10)
C2	0.8935 (6)	0.54544 (15)	0.6519 (5)	0.0191 (10)
C3	0.7394 (7)	0.53338 (17)	0.7090 (5)	0.0222 (10)
H3	0.6361	0.5497	0.6889	0.027*
C4	0.7421 (7)	0.49725 (17)	0.7951 (5)	0.0255 (11)
H4	0.6390	0.4885	0.8344	0.031*
C5	0.8937 (8)	0.47324 (17)	0.8256 (5)	0.0262 (11)
C6	1.0457 (7)	0.48345 (17)	0.7664 (5)	0.0250 (11)
H6	1.1476	0.4664	0.7841	0.030*
C7	1.0411 (6)	0.52035 (16)	0.6788 (5)	0.0204 (10)
C8	1.1316 (7)	0.61649 (18)	0.4252 (6)	0.0277 (11)
H8A	1.1838	0.6409	0.4799	0.042*
H8B	1.0329	0.6273	0.3652	0.042*
H8C	1.2175	0.6038	0.3678	0.042*
C9	1.0403 (9)	0.4191 (2)	0.9692 (7)	0.0393 (15)
H9A	1.0908	0.4030	0.8943	0.059*
H9B	1.0187	0.3987	1.0445	0.059*
H9C	1.1206	0.4420	1.0046	0.059*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N2	0.024 (2)	0.024 (2)	0.019 (2)	0.0044 (18)	0.0044 (17)	0.0013 (17)
C10	0.056 (4)	0.044 (4)	0.035 (3)	-0.010 (3)	0.015 (3)	-0.011 (3)
C11	0.076 (6)	0.040 (4)	0.086 (6)	-0.017 (4)	0.048 (5)	-0.026 (4)
C12	0.033 (4)	0.079 (6)	0.047 (4)	0.012 (4)	0.008 (3)	0.020 (4)
C13	0.026 (3)	0.130 (8)	0.027 (3)	-0.007 (4)	-0.004 (3)	0.005 (4)
C14	0.037 (3)	0.051 (4)	0.033 (3)	0.003 (3)	0.009 (3)	0.003 (3)
C15	0.026 (3)	0.042 (4)	0.034 (3)	0.000 (2)	0.003 (2)	-0.008 (3)
C16	0.047 (4)	0.029 (3)	0.052 (4)	-0.001 (3)	0.003 (3)	-0.004 (3)
C17	0.045 (4)	0.034 (3)	0.049 (4)	-0.007 (3)	0.007 (3)	0.005 (3)
Pt1	0.01677 (10)	0.01604 (10)	0.01721 (10)	0.00084 (7)	0.00130 (7)	0.00005 (7)
Br1	0.0303 (3)	0.0270 (3)	0.0244 (2)	-0.0012 (2)	-0.0021 (2)	-0.0027 (2)
Br2	0.0295 (3)	0.0267 (3)	0.0266 (3)	0.0050 (2)	0.0015 (2)	-0.0004 (2)
Br3	0.0262 (3)	0.0286 (3)	0.0303 (3)	0.0029 (2)	-0.0035 (2)	-0.0062 (2)
S1	0.0177 (6)	0.0245 (6)	0.0275 (6)	0.0036 (5)	0.0004 (5)	-0.0025 (5)
O1	0.047 (2)	0.022 (2)	0.032 (2)	0.0026 (18)	0.0023 (18)	0.0078 (16)
N1	0.019 (2)	0.018 (2)	0.0171 (19)	-0.0008 (16)	-0.0012 (16)	-0.0010 (16)
C1	0.018 (2)	0.022 (3)	0.023 (2)	0.0007 (19)	-0.0009 (19)	-0.003 (2)
C2	0.024 (2)	0.013 (2)	0.020 (2)	0.0017 (19)	0.0000 (19)	-0.0010 (18)
C3	0.021 (2)	0.020 (3)	0.026 (2)	0.001 (2)	0.004 (2)	-0.002 (2)
C4	0.028 (3)	0.024 (3)	0.025 (3)	-0.001 (2)	0.007 (2)	0.001 (2)
C5	0.041 (3)	0.016 (2)	0.021 (2)	0.000 (2)	0.000 (2)	-0.0010 (19)
C6	0.033 (3)	0.017 (2)	0.025 (2)	0.005 (2)	-0.004 (2)	-0.002 (2)
C7	0.021 (2)	0.018 (2)	0.022 (2)	0.0006 (19)	-0.0022 (19)	-0.0056 (19)
C8	0.024 (3)	0.027 (3)	0.033 (3)	-0.001 (2)	0.008 (2)	0.000 (2)
C9	0.053 (4)	0.025 (3)	0.037 (3)	-0.001 (3)	-0.011 (3)	0.007 (2)

*Geometric parameters (Å, °)*

N2—C10	1.511 (7)	C17—H17C	0.9800
N2—C12	1.523 (8)	Pt1—Br1	2.4352 (5)
N2—C14	1.521 (7)	Pt1—Br2	2.4241 (6)
N2—C16	1.491 (8)	Pt1—Br3	2.4309 (5)
C10—H10A	0.9900	Pt1—N1	2.025 (4)
C10—H10B	0.9900	S1—C1	1.730 (5)
C10—C11	1.474 (10)	S1—C7	1.743 (5)
C11—H11A	0.9800	O1—C5	1.379 (6)
C11—H11B	0.9800	O1—C9	1.427 (7)
C11—H11C	0.9800	N1—C1	1.309 (6)
C12—H12A	0.9900	N1—C2	1.396 (6)
C12—H12B	0.9900	C1—C8	1.483 (7)
C12—C13	1.464 (11)	C2—C3	1.405 (7)
C13—H13A	0.9800	C2—C7	1.383 (7)
C13—H13B	0.9800	C3—H3	0.9500
C13—H13C	0.9800	C3—C4	1.377 (7)
C14—H14A	0.9900	C4—H4	0.9500
C14—H14B	0.9900	C4—C5	1.396 (8)
C14—C15	1.513 (8)	C5—C6	1.386 (8)
C15—H15A	0.9800	C6—H6	0.9500
C15—H15B	0.9800	C6—C7	1.404 (7)
C15—H15C	0.9800	C8—H8A	0.9800
C16—H16A	0.9900	C8—H8B	0.9800
C16—H16B	0.9900	C8—H8C	0.9800
C16—C17	1.517 (9)	C9—H9A	0.9800
C17—H17A	0.9800	C9—H9B	0.9800
C17—H17B	0.9800	C9—H9C	0.9800
C10—N2—C12	108.5 (5)	C16—C17—H17B	109.5
C10—N2—C14	111.4 (5)	C16—C17—H17C	109.5
C14—N2—C12	107.4 (4)	H17A—C17—H17B	109.5
C16—N2—C10	109.6 (5)	H17A—C17—H17C	109.5
C16—N2—C12	110.1 (5)	H17B—C17—H17C	109.5
C16—N2—C14	109.8 (5)	Br2—Pt1—Br1	91.171 (19)
N2—C10—H10A	107.9	Br2—Pt1—Br3	92.507 (19)
N2—C10—H10B	107.9	Br3—Pt1—Br1	176.30 (2)
H10A—C10—H10B	107.2	N1—Pt1—Br1	87.04 (11)
C11—C10—N2	117.8 (6)	N1—Pt1—Br2	177.41 (11)
C11—C10—H10A	107.9	N1—Pt1—Br3	89.29 (11)
C11—C10—H10B	107.9	C1—S1—C7	89.9 (2)
C10—C11—H11A	109.5	C5—O1—C9	116.3 (5)
C10—C11—H11B	109.5	C1—N1—Pt1	124.7 (3)
C10—C11—H11C	109.5	C1—N1—C2	112.6 (4)
H11A—C11—H11B	109.5	C2—N1—Pt1	122.1 (3)
H11A—C11—H11C	109.5	N1—C1—S1	114.0 (4)
H11B—C11—H11C	109.5	N1—C1—C8	124.2 (5)

N2—C12—H12A	108.0	C8—C1—S1	121.8 (4)
N2—C12—H12B	108.0	N1—C2—C3	126.5 (4)
H12A—C12—H12B	107.3	C7—C2—N1	113.5 (4)
C13—C12—N2	117.1 (6)	C7—C2—C3	120.0 (5)
C13—C12—H12A	108.0	C2—C3—H3	120.9
C13—C12—H12B	108.0	C4—C3—C2	118.2 (5)
C12—C13—H13A	109.5	C4—C3—H3	120.9
C12—C13—H13B	109.5	C3—C4—H4	119.4
C12—C13—H13C	109.5	C3—C4—C5	121.2 (5)
H13A—C13—H13B	109.5	C5—C4—H4	119.4
H13A—C13—H13C	109.5	O1—C5—C4	115.7 (5)
H13B—C13—H13C	109.5	O1—C5—C6	122.6 (5)
N2—C14—H14A	108.7	C6—C5—C4	121.7 (5)
N2—C14—H14B	108.7	C5—C6—H6	121.8
H14A—C14—H14B	107.6	C5—C6—C7	116.5 (5)
C15—C14—N2	114.3 (5)	C7—C6—H6	121.8
C15—C14—H14A	108.7	C2—C7—S1	109.9 (4)
C15—C14—H14B	108.7	C2—C7—C6	122.4 (5)
C14—C15—H15A	109.5	C6—C7—S1	127.7 (4)
C14—C15—H15B	109.5	C1—C8—H8A	109.5
C14—C15—H15C	109.5	C1—C8—H8B	109.5
H15A—C15—H15B	109.5	C1—C8—H8C	109.5
H15A—C15—H15C	109.5	H8A—C8—H8B	109.5
H15B—C15—H15C	109.5	H8A—C8—H8C	109.5
N2—C16—H16A	108.4	H8B—C8—H8C	109.5
N2—C16—H16B	108.4	O1—C9—H9A	109.5
N2—C16—C17	115.4 (5)	O1—C9—H9B	109.5
H16A—C16—H16B	107.5	O1—C9—H9C	109.5
C17—C16—H16A	108.4	H9A—C9—H9B	109.5
C17—C16—H16B	108.4	H9A—C9—H9C	109.5
C16—C17—H17A	109.5	H9B—C9—H9C	109.5

**(3) Tetraethylammonium tribromido(2,5,6-trimethyl-1,3-benzothiazole- $\kappa$ N)platinate(II)***Crystal data* $(\text{C}_8\text{H}_{20}\text{N})[\text{PtBr}_3(\text{C}_{10}\text{H}_{11}\text{NS})]$  $M_r = 742.33$ Monoclinic,  $P2_1/n$  $a = 7.9742$  (4) Å $b = 30.2807$  (14) Å $c = 9.6427$  (5) Å $\beta = 100.151$  (3)° $V = 2291.9$  (2) Å<sup>3</sup> $Z = 4$  $F(000) = 1408$  $D_x = 2.151$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 5770 reflections

 $\theta = 2.3$ – $26.4$ ° $\mu = 11.45$  mm<sup>-1</sup> $T = 100$  K

Block, red

 $0.5 \times 0.36 \times 0.25$  mm

*Data collection*

Bruker APEXII CCD diffractometer	$T_{\min} = 0.003$ , $T_{\max} = 0.028$
Radiation source: Micro Focus Rotating Anode, Bruker TXS	10729 measured reflections
Double Bounce Multilayer Mirrors monochromator	4692 independent reflections
Detector resolution: 7.9 pixels $\text{mm}^{-1}$	4120 reflections with $I > 2\sigma(I)$
$\varphi$ and $\omega$ scans	$R_{\text{int}} = 0.048$
Absorption correction: multi-scan (SADABS; Bruker, 2014)	$\theta_{\max} = 26.5^\circ$ , $\theta_{\min} = 2.3^\circ$
	$h = -9 \rightarrow 5$
	$k = -37 \rightarrow 31$
	$l = -10 \rightarrow 12$

*Refinement*

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.039$	$w = 1/[\sigma^2(F_o^2) + (0.056P)^2 + 1.6623P]$
$wR(F^2) = 0.106$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\max} = 0.002$
4692 reflections	$\Delta\rho_{\max} = 1.88 \text{ e } \text{\AA}^{-3}$
266 parameters	$\Delta\rho_{\min} = -1.02 \text{ e } \text{\AA}^{-3}$
0 restraints	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N2	0.5342 (7)	0.18676 (17)	0.5028 (5)	0.0281 (11)	
C12	0.8178 (11)	0.1847 (4)	0.6760 (9)	0.060 (2)	
H12A	0.8673	0.1911	0.7743	0.089*	0.566 (9)
H12B	0.8346	0.1535	0.6560	0.089*	0.566 (9)
H12C	0.8736	0.2030	0.6137	0.089*	0.566 (9)
H12D	0.9301	0.1706	0.6901	0.089*	0.434 (9)
H12E	0.8313	0.2168	0.6706	0.089*	0.434 (9)
H12F	0.7606	0.1775	0.7551	0.089*	0.434 (9)
C14	0.4260 (12)	0.1207 (3)	0.3440 (10)	0.057 (2)	
H14A	0.4285	0.0884	0.3448	0.086*	0.566 (9)
H14B	0.3076	0.1309	0.3310	0.086*	0.566 (9)
H14C	0.4792	0.1316	0.2665	0.086*	0.566 (9)
H14D	0.3786	0.1145	0.2451	0.086*	0.434 (9)
H14E	0.5321	0.1041	0.3718	0.086*	0.434 (9)
H14F	0.3439	0.1119	0.4033	0.086*	0.434 (9)
C16	0.2437 (10)	0.1856 (2)	0.5783 (8)	0.0404 (17)	
H16A	0.1334	0.2008	0.5672	0.061*	0.566 (9)
H16B	0.2262	0.1549	0.5468	0.061*	0.566 (9)
H16C	0.2981	0.1862	0.6777	0.061*	0.566 (9)
H16D	0.1836	0.1759	0.6532	0.061*	0.434 (9)

H16E	0.2345	0.2178	0.5684	0.061*	0.434 (9)
H16F	0.1925	0.1716	0.4893	0.061*	0.434 (9)
C18	0.6616 (12)	0.2550 (3)	0.4092 (10)	0.054 (2)	
H18A	0.7199	0.2649	0.3334	0.081*	0.566 (9)
H18B	0.5547	0.2713	0.4045	0.081*	0.566 (9)
H18C	0.7348	0.2603	0.5004	0.081*	0.566 (9)
H18D	0.6529	0.2872	0.4073	0.081*	0.434 (9)
H18E	0.7769	0.2463	0.4540	0.081*	0.434 (9)
H18F	0.6378	0.2435	0.3126	0.081*	0.434 (9)
C11	0.6256 (17)	0.1951 (4)	0.6507 (12)	0.033 (3)	0.566 (9)
H11A	0.5715	0.1771	0.7162	0.040*	0.566 (9)
H11B	0.6104	0.2266	0.6739	0.040*	0.566 (9)
C13	0.5186 (18)	0.1373 (4)	0.4765 (14)	0.037 (3)	0.566 (9)
H13A	0.6356	0.1252	0.4883	0.044*	0.566 (9)
H13B	0.4650	0.1245	0.5524	0.044*	0.566 (9)
C15	0.3598 (15)	0.2095 (4)	0.4884 (12)	0.032 (3)	0.566 (9)
H15A	0.3039	0.2093	0.3882	0.038*	0.566 (9)
H15B	0.3758	0.2406	0.5193	0.038*	0.566 (9)
C17A	0.6257 (15)	0.2086 (5)	0.3936 (13)	0.040 (3)	0.566 (9)
H17A	0.7349	0.1930	0.3950	0.048*	0.566 (9)
H17B	0.5555	0.2039	0.2993	0.048*	0.566 (9)
C11A	0.716 (2)	0.1689 (6)	0.546 (2)	0.044 (5)	0.434 (9)
H11C	0.7089	0.1363	0.5527	0.053*	0.434 (9)
H11D	0.7792	0.1755	0.4687	0.053*	0.434 (9)
C13A	0.460 (2)	0.1664 (5)	0.3608 (15)	0.033 (4)	0.434 (9)
H13C	0.3509	0.1819	0.3266	0.040*	0.434 (9)
H13D	0.5377	0.1742	0.2953	0.040*	0.434 (9)
C15A	0.435 (2)	0.1723 (6)	0.6165 (14)	0.031 (4)	0.434 (9)
H15C	0.4863	0.1860	0.7073	0.038*	0.434 (9)
H15D	0.4442	0.1398	0.6281	0.038*	0.434 (9)
C17	0.535 (2)	0.2363 (5)	0.4914 (17)	0.033 (4)	0.434 (9)
H17C	0.4195	0.2461	0.4466	0.040*	0.434 (9)
H17D	0.5582	0.2489	0.5876	0.040*	0.434 (9)
Pt1	0.28820 (3)	0.36639 (2)	0.54603 (2)	0.02321 (10)	
Br1	0.11056 (8)	0.41268 (2)	0.37347 (7)	0.03333 (17)	
Br2	0.08097 (9)	0.30729 (2)	0.50744 (7)	0.03324 (17)	
Br3	0.47121 (9)	0.32318 (2)	0.72302 (8)	0.04046 (19)	
S1	0.7243 (2)	0.46490 (5)	0.58781 (16)	0.0279 (3)	
N1	0.4604 (6)	0.41615 (16)	0.5825 (5)	0.0239 (10)	
C1	0.6036 (8)	0.4185 (2)	0.5354 (6)	0.0271 (13)	
C2	0.4361 (8)	0.4523 (2)	0.6672 (6)	0.0258 (13)	
C3	0.2953 (8)	0.4599 (2)	0.7319 (6)	0.0263 (13)	
H3	0.2031	0.4396	0.7202	0.032*	
C4	0.2917 (8)	0.4974 (2)	0.8136 (6)	0.0285 (13)	
C5	0.4298 (9)	0.5269 (2)	0.8313 (7)	0.0319 (14)	
C6	0.5667 (9)	0.5201 (2)	0.7669 (6)	0.0294 (14)	
H6	0.6578	0.5408	0.7782	0.035*	
C7	0.5713 (8)	0.4824 (2)	0.6840 (6)	0.0263 (13)	



C8	0.6633 (9)	0.3844 (2)	0.4455 (7)	0.0342 (15)
H8A	0.5737	0.3784	0.3642	0.051*
H8B	0.6900	0.3572	0.5001	0.051*
H8C	0.7657	0.3949	0.4126	0.051*
C9	0.1406 (9)	0.5053 (2)	0.8848 (7)	0.0346 (15)
H9A	0.0846	0.5330	0.8503	0.052*
H9B	0.1792	0.5073	0.9869	0.052*
H9C	0.0599	0.4808	0.8636	0.052*
C10	0.4240 (10)	0.5680 (2)	0.9223 (8)	0.0391 (16)
H10A	0.3289	0.5869	0.8798	0.059*
H10B	0.5311	0.5844	0.9286	0.059*
H10C	0.4084	0.5591	1.0170	0.059*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N2	0.028 (3)	0.026 (3)	0.031 (3)	0.001 (2)	0.007 (2)	-0.001 (2)
C12	0.036 (4)	0.093 (7)	0.047 (5)	0.003 (4)	-0.002 (4)	0.015 (5)
C14	0.053 (5)	0.052 (5)	0.072 (6)	-0.005 (4)	0.026 (5)	-0.035 (5)
C16	0.038 (4)	0.037 (4)	0.051 (4)	0.000 (3)	0.023 (3)	-0.005 (3)
C18	0.061 (5)	0.045 (5)	0.060 (5)	-0.009 (4)	0.021 (4)	0.014 (4)
C11	0.045 (7)	0.027 (6)	0.027 (5)	-0.004 (5)	0.004 (5)	-0.001 (5)
C13	0.038 (7)	0.028 (6)	0.043 (7)	0.009 (5)	0.004 (6)	-0.008 (5)
C15	0.037 (6)	0.023 (6)	0.034 (6)	0.003 (5)	0.007 (5)	-0.002 (5)
C17A	0.024 (6)	0.063 (9)	0.034 (6)	0.000 (6)	0.010 (5)	0.003 (6)
C11A	0.035 (9)	0.044 (10)	0.058 (11)	0.017 (8)	0.020 (8)	0.022 (8)
C13A	0.049 (9)	0.032 (8)	0.023 (7)	-0.013 (7)	0.014 (7)	-0.013 (6)
C15A	0.032 (8)	0.044 (9)	0.020 (7)	0.000 (7)	0.009 (6)	0.006 (6)
C17	0.037 (8)	0.031 (8)	0.037 (8)	-0.009 (7)	0.018 (7)	0.001 (6)
Pt1	0.02518 (15)	0.01967 (15)	0.02593 (15)	-0.00120 (8)	0.00770 (10)	0.00056 (8)
Br1	0.0312 (3)	0.0284 (3)	0.0401 (4)	-0.0015 (3)	0.0056 (3)	0.0073 (3)
Br2	0.0415 (4)	0.0266 (3)	0.0321 (3)	-0.0074 (3)	0.0077 (3)	0.0004 (2)
Br3	0.0389 (4)	0.0386 (4)	0.0420 (4)	0.0009 (3)	0.0019 (3)	0.0113 (3)
S1	0.0268 (7)	0.0263 (8)	0.0320 (8)	-0.0029 (6)	0.0091 (6)	0.0027 (6)
N1	0.026 (3)	0.025 (3)	0.021 (2)	0.001 (2)	0.006 (2)	0.002 (2)
C1	0.030 (3)	0.025 (3)	0.026 (3)	-0.004 (2)	0.006 (3)	0.000 (2)
C2	0.023 (3)	0.027 (3)	0.028 (3)	-0.001 (2)	0.007 (2)	0.006 (2)
C3	0.028 (3)	0.022 (3)	0.030 (3)	-0.003 (2)	0.007 (3)	0.002 (2)
C4	0.033 (3)	0.023 (3)	0.031 (3)	0.004 (3)	0.009 (3)	-0.001 (3)
C5	0.046 (4)	0.025 (3)	0.025 (3)	0.004 (3)	0.005 (3)	0.000 (2)
C6	0.037 (4)	0.021 (3)	0.030 (3)	-0.005 (3)	0.005 (3)	0.005 (2)
C7	0.032 (3)	0.022 (3)	0.026 (3)	0.002 (2)	0.007 (3)	0.007 (2)
C8	0.030 (3)	0.037 (4)	0.039 (4)	0.001 (3)	0.013 (3)	-0.005 (3)
C9	0.045 (4)	0.027 (3)	0.034 (3)	0.001 (3)	0.012 (3)	0.001 (3)
C10	0.050 (4)	0.020 (3)	0.048 (4)	-0.004 (3)	0.013 (3)	-0.004 (3)

*Geometric parameters (Å, °)*

N2—C11	1.504 (12)	C13—H13A	0.9900
N2—C13	1.520 (12)	C13—H13B	0.9900
N2—C15	1.535 (13)	C15—H15A	0.9900
N2—C17A	1.533 (13)	C15—H15B	0.9900
N2—C11A	1.537 (16)	C17A—H17A	0.9900
N2—C13A	1.523 (14)	C17A—H17B	0.9900
N2—C15A	1.523 (15)	C11A—H11C	0.9900
N2—C17	1.505 (16)	C11A—H11D	0.9900
C12—H12A	0.9800	C13A—H13C	0.9900
C12—H12B	0.9800	C13A—H13D	0.9900
C12—H12C	0.9800	C15A—H15C	0.9900
C12—H12D	0.9800	C15A—H15D	0.9900
C12—H12E	0.9800	C17—H17C	0.9900
C12—H12F	0.9800	C17—H17D	0.9900
C12—C11	1.542 (15)	Pt1—Br1	2.4309 (7)
C12—C11A	1.45 (2)	Pt1—Br2	2.4198 (7)
C14—H14A	0.9800	Pt1—Br3	2.4240 (7)
C14—H14B	0.9800	Pt1—N1	2.027 (5)
C14—H14C	0.9800	S1—C1	1.727 (6)
C14—H14D	0.9800	S1—C7	1.739 (7)
C14—H14E	0.9800	N1—C1	1.303 (8)
C14—H14F	0.9800	N1—C2	1.401 (8)
C14—C13	1.449 (15)	C1—C8	1.482 (9)
C14—C13A	1.415 (16)	C2—C3	1.397 (9)
C16—H16A	0.9800	C2—C7	1.398 (9)
C16—H16B	0.9800	C3—H3	0.9500
C16—H16C	0.9800	C3—C4	1.385 (9)
C16—H16D	0.9800	C4—C5	1.404 (9)
C16—H16E	0.9800	C4—C9	1.507 (9)
C16—H16F	0.9800	C5—C6	1.363 (10)
C16—C15	1.553 (14)	C5—C10	1.528 (9)
C16—C15A	1.561 (17)	C6—H6	0.9500
C18—H18A	0.9800	C6—C7	1.400 (9)
C18—H18B	0.9800	C8—H8A	0.9800
C18—H18C	0.9800	C8—H8B	0.9800
C18—H18D	0.9800	C8—H8C	0.9800
C18—H18E	0.9800	C9—H9A	0.9800
C18—H18F	0.9800	C9—H9B	0.9800
C18—C17A	1.435 (17)	C9—H9C	0.9800
C18—C17	1.500 (16)	C10—H10A	0.9800
C11—H11A	0.9900	C10—H10B	0.9800
C11—H11B	0.9900	C10—H10C	0.9800
C11—N2—C13	109.7 (7)	N2—C17A—H17B	107.9
C11—N2—C15	106.9 (8)	C18—C17A—N2	117.4 (10)
C11—N2—C17A	111.6 (8)	C18—C17A—H17A	107.9

C13—N2—C15	112.3 (8)	C18—C17A—H17B	107.9
C13—N2—C17A	110.2 (8)	H17A—C17A—H17B	107.2
C17A—N2—C15	106.1 (7)	N2—C11A—H11C	107.7
C13A—N2—C11A	107.5 (11)	N2—C11A—H11D	107.7
C13A—N2—C15A	111.2 (9)	C12—C11A—N2	118.3 (13)
C15A—N2—C11A	106.7 (9)	C12—C11A—H11C	107.7
C17—N2—C11A	110.9 (10)	C12—C11A—H11D	107.7
C17—N2—C13A	110.0 (9)	H11C—C11A—H11D	107.1
C17—N2—C15A	110.4 (9)	N2—C13A—H13C	106.7
H12A—C12—H12B	109.5	N2—C13A—H13D	106.7
H12A—C12—H12C	109.5	C14—C13A—N2	122.3 (12)
H12B—C12—H12C	109.5	C14—C13A—H13C	106.7
H12D—C12—H12E	109.5	C14—C13A—H13D	106.7
H12D—C12—H12F	109.5	H13C—C13A—H13D	106.6
H12E—C12—H12F	109.5	N2—C15A—C16	111.4 (9)
C11—C12—H12A	109.5	N2—C15A—H15C	109.4
C11—C12—H12B	109.5	N2—C15A—H15D	109.4
C11—C12—H12C	109.5	C16—C15A—H15C	109.4
C11A—C12—H12D	109.5	C16—C15A—H15D	109.4
C11A—C12—H12E	109.5	H15C—C15A—H15D	108.0
C11A—C12—H12F	109.5	N2—C17—H17C	108.5
H14A—C14—H14B	109.5	N2—C17—H17D	108.5
H14A—C14—H14C	109.5	C18—C17—N2	115.2 (11)
H14B—C14—H14C	109.5	C18—C17—H17C	108.5
H14D—C14—H14E	109.5	C18—C17—H17D	108.5
H14D—C14—H14F	109.5	H17C—C17—H17D	107.5
H14E—C14—H14F	109.5	Br2—Pt1—Br1	91.23 (2)
C13—C14—H14A	109.5	Br2—Pt1—Br3	91.10 (2)
C13—C14—H14B	109.5	Br3—Pt1—Br1	177.45 (3)
C13—C14—H14C	109.5	N1—Pt1—Br1	89.16 (14)
C13A—C14—H14D	109.5	N1—Pt1—Br2	178.76 (14)
C13A—C14—H14E	109.5	N1—Pt1—Br3	88.50 (14)
C13A—C14—H14F	109.5	C1—S1—C7	89.8 (3)
H16A—C16—H16B	109.5	C1—N1—Pt1	126.1 (4)
H16A—C16—H16C	109.5	C1—N1—C2	112.3 (5)
H16B—C16—H16C	109.5	C2—N1—Pt1	121.6 (4)
H16D—C16—H16E	109.5	N1—C1—S1	114.9 (5)
H16D—C16—H16F	109.5	N1—C1—C8	123.9 (6)
H16E—C16—H16F	109.5	C8—C1—S1	121.2 (5)
C15—C16—H16A	109.5	C3—C2—N1	126.5 (6)
C15—C16—H16B	109.5	C3—C2—C7	120.4 (6)
C15—C16—H16C	109.5	C7—C2—N1	113.1 (5)
C15A—C16—H16D	109.5	C2—C3—H3	120.4
C15A—C16—H16E	109.5	C4—C3—C2	119.2 (6)
C15A—C16—H16F	109.5	C4—C3—H3	120.4
H18A—C18—H18B	109.5	C3—C4—C5	119.8 (6)
H18A—C18—H18C	109.5	C3—C4—C9	119.1 (6)
H18B—C18—H18C	109.5	C5—C4—C9	121.1 (6)

H18D—C18—H18E	109.5	C4—C5—C10	119.0 (6)
H18D—C18—H18F	109.5	C6—C5—C4	121.5 (6)
H18E—C18—H18F	109.5	C6—C5—C10	119.5 (6)
C17A—C18—H18A	109.5	C5—C6—H6	120.4
C17A—C18—H18B	109.5	C5—C6—C7	119.1 (6)
C17A—C18—H18C	109.5	C7—C6—H6	120.4
C17—C18—H18D	109.5	C2—C7—S1	109.9 (5)
C17—C18—H18E	109.5	C2—C7—C6	120.0 (6)
C17—C18—H18F	109.5	C6—C7—S1	130.1 (5)
N2—C11—C12	114.7 (9)	C1—C8—H8A	109.5
N2—C11—H11A	108.6	C1—C8—H8B	109.5
N2—C11—H11B	108.6	C1—C8—H8C	109.5
C12—C11—H11A	108.6	H8A—C8—H8B	109.5
C12—C11—H11B	108.6	H8A—C8—H8C	109.5
H11A—C11—H11B	107.6	H8B—C8—H8C	109.5
N2—C13—H13A	107.3	C4—C9—H9A	109.5
N2—C13—H13B	107.3	C4—C9—H9B	109.5
C14—C13—N2	120.2 (10)	C4—C9—H9C	109.5
C14—C13—H13A	107.3	H9A—C9—H9B	109.5
C14—C13—H13B	107.3	H9A—C9—H9C	109.5
H13A—C13—H13B	106.9	H9B—C9—H9C	109.5
N2—C15—C16	111.2 (8)	C5—C10—H10A	109.5
N2—C15—H15A	109.4	C5—C10—H10B	109.5
N2—C15—H15B	109.4	C5—C10—H10C	109.5
C16—C15—H15A	109.4	H10A—C10—H10B	109.5
C16—C15—H15B	109.4	H10A—C10—H10C	109.5
H15A—C15—H15B	108.0	H10B—C10—H10C	109.5
N2—C17A—H17A	107.9		

**(4) Tetraethylammonium tribromido(2-methyl-5-nitro-1,3-benzothiazole- $\kappa$ N)platinate(II)***Crystal data* $(C_8H_{20}N)[PtBr_3(C_8H_6N_2O_2S)]$  $M_r = 759.28$ Monoclinic,  $P2_1/n$  $a = 8.1170$  (3) Å $b = 29.2717$  (12) Å $c = 9.5102$  (4) Å $\beta = 100.720$  (1)° $V = 2220.17$  (15) Å<sup>3</sup> $Z = 4$  $F(000) = 1432$  $D_x = 2.272$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 9483 reflections

 $\theta = 2.6$ – $26.4$ ° $\mu = 11.83$  mm<sup>-1</sup> $T = 100$  K

Block, bronze

 $0.32 \times 0.3 \times 0.25$  mm*Data collection*Bruker APEXII CCD  
diffractometerRadiation source: Micro Focus Rotating Anode,  
Bruker TXSDouble Bounce Multilayer Mirrors  
monochromatorDetector resolution: 7.9 pixels mm<sup>-1</sup> $\varphi$  and  $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Bruker, 2014) $T_{\min} = 0.020$ ,  $T_{\max} = 0.045$ 

15975 measured reflections

4550 independent reflections

4254 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.028$  $\theta_{\max} = 26.5$ °,  $\theta_{\min} = 2.3$ °

$h = -10 \rightarrow 10$   
 $k = -31 \rightarrow 36$

$l = -11 \rightarrow 7$

### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.060$   
 $S = 1.18$   
 4550 reflections  
 240 parameters  
 0 restraints

Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0044P)^2 + 13.0832P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 1.25 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.37 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N3	0.5548 (5)	0.31569 (15)	0.5135 (4)	0.0187 (9)
C9	0.3795 (6)	0.30025 (19)	0.5223 (6)	0.0247 (12)
H9A	0.3850	0.2685	0.5585	0.030*
H9B	0.3112	0.3000	0.4245	0.030*
C10	0.2911 (7)	0.3294 (2)	0.6170 (7)	0.0320 (13)
H10A	0.2786	0.3606	0.5790	0.048*
H10B	0.3575	0.3300	0.7143	0.048*
H10C	0.1802	0.3165	0.6191	0.048*
C11	0.5608 (7)	0.36508 (18)	0.4693 (6)	0.0257 (12)
H11A	0.6776	0.3727	0.4612	0.031*
H11B	0.5304	0.3844	0.5459	0.031*
C12	0.4466 (7)	0.3771 (2)	0.3292 (6)	0.0281 (12)
H12A	0.4815	0.3601	0.2510	0.042*
H12B	0.4534	0.4099	0.3115	0.042*
H12C	0.3308	0.3689	0.3347	0.042*
C13	0.6205 (7)	0.28611 (19)	0.4041 (6)	0.0262 (12)
H13A	0.5380	0.2869	0.3133	0.031*
H13B	0.7258	0.2998	0.3853	0.031*
C14	0.6543 (8)	0.2369 (2)	0.4470 (7)	0.0323 (13)
H14A	0.6871	0.2201	0.3675	0.048*
H14B	0.5527	0.2232	0.4707	0.048*
H14C	0.7452	0.2354	0.5307	0.048*
C15	0.6624 (7)	0.3097 (2)	0.6607 (6)	0.0266 (12)
H15A	0.6503	0.2780	0.6929	0.032*
H15B	0.6197	0.3303	0.7284	0.032*
C16	0.8519 (7)	0.3199 (3)	0.6674 (7)	0.0373 (15)
H16A	0.8974	0.2984	0.6054	0.056*
H16B	0.9122	0.3164	0.7661	0.056*

H16C	0.8655	0.3512	0.6348	0.056*
Pt1	0.19959 (2)	0.63310 (2)	0.93287 (2)	0.01542 (6)
Br1	0.01278 (6)	0.67668 (2)	0.75173 (6)	0.02534 (12)
Br2	0.39391 (7)	0.69652 (2)	0.97693 (6)	0.02338 (12)
Br3	0.38687 (6)	0.58556 (2)	1.10249 (6)	0.02324 (12)
S1	-0.23297 (16)	0.53228 (4)	0.89580 (14)	0.0211 (3)
O1	0.4596 (5)	0.50291 (14)	0.6504 (4)	0.0304 (9)
O2	0.3206 (5)	0.44662 (14)	0.5385 (5)	0.0343 (10)
N1	0.0320 (5)	0.58046 (14)	0.8995 (4)	0.0168 (8)
N2	0.3324 (6)	0.48001 (15)	0.6180 (5)	0.0247 (10)
C1	-0.1103 (6)	0.57965 (17)	0.9469 (6)	0.0187 (10)
C2	0.0503 (6)	0.54253 (17)	0.8172 (5)	0.0187 (10)
C3	0.1902 (6)	0.53230 (17)	0.7554 (5)	0.0192 (10)
H3	0.2846	0.5520	0.7661	0.023*
C4	0.1839 (7)	0.49242 (17)	0.6787 (6)	0.0213 (11)
C5	0.0465 (7)	0.46309 (19)	0.6567 (6)	0.0263 (12)
H5	0.0473	0.4365	0.5994	0.032*
C6	-0.0899 (7)	0.47253 (18)	0.7177 (6)	0.0245 (12)
H6	-0.1844	0.4528	0.7047	0.029*
C7	-0.0852 (6)	0.51233 (18)	0.7997 (6)	0.0212 (11)
C8	-0.1675 (7)	0.61580 (18)	1.0364 (6)	0.0239 (11)
H8A	-0.1978	0.6433	0.9784	0.036*
H8B	-0.0770	0.6231	1.1168	0.036*
H8C	-0.2655	0.6049	1.0732	0.036*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N3	0.018 (2)	0.024 (2)	0.014 (2)	0.0008 (17)	0.0023 (17)	0.0010 (17)
C9	0.020 (3)	0.027 (3)	0.027 (3)	-0.003 (2)	0.006 (2)	0.004 (2)
C10	0.032 (3)	0.036 (3)	0.031 (3)	0.002 (3)	0.013 (3)	0.005 (3)
C11	0.025 (3)	0.024 (3)	0.029 (3)	-0.004 (2)	0.007 (2)	0.002 (2)
C12	0.035 (3)	0.028 (3)	0.022 (3)	0.005 (2)	0.009 (2)	0.007 (2)
C13	0.030 (3)	0.032 (3)	0.020 (3)	-0.002 (2)	0.011 (2)	-0.004 (2)
C14	0.034 (3)	0.031 (3)	0.033 (3)	0.011 (2)	0.008 (3)	-0.004 (3)
C15	0.025 (3)	0.033 (3)	0.022 (3)	0.000 (2)	0.002 (2)	-0.003 (2)
C16	0.018 (3)	0.064 (4)	0.026 (3)	0.004 (3)	-0.006 (2)	-0.011 (3)
Pt1	0.01462 (10)	0.01661 (9)	0.01517 (10)	-0.00278 (7)	0.00313 (7)	-0.00054 (7)
Br1	0.0203 (3)	0.0302 (3)	0.0244 (3)	-0.0025 (2)	0.0012 (2)	0.0072 (2)
Br2	0.0250 (3)	0.0238 (3)	0.0213 (3)	-0.0058 (2)	0.0043 (2)	-0.0007 (2)
Br3	0.0194 (2)	0.0225 (3)	0.0263 (3)	-0.0023 (2)	-0.0001 (2)	0.0038 (2)
S1	0.0183 (6)	0.0210 (6)	0.0235 (7)	-0.0059 (5)	0.0029 (5)	0.0014 (5)
O1	0.026 (2)	0.034 (2)	0.032 (2)	-0.0023 (17)	0.0070 (18)	-0.0038 (18)
O2	0.046 (3)	0.024 (2)	0.037 (2)	-0.0025 (18)	0.018 (2)	-0.0089 (18)
N1	0.014 (2)	0.019 (2)	0.016 (2)	-0.0005 (16)	0.0018 (17)	0.0022 (17)
N2	0.035 (3)	0.019 (2)	0.021 (2)	0.002 (2)	0.008 (2)	0.0032 (19)
C1	0.017 (2)	0.018 (2)	0.022 (3)	-0.0015 (19)	0.004 (2)	0.005 (2)
C2	0.023 (3)	0.017 (2)	0.014 (2)	-0.003 (2)	-0.001 (2)	0.0027 (19)

C3	0.017 (2)	0.021 (2)	0.018 (3)	-0.003 (2)	0.000 (2)	0.004 (2)
C4	0.026 (3)	0.019 (2)	0.018 (3)	0.000 (2)	0.002 (2)	0.004 (2)
C5	0.036 (3)	0.020 (3)	0.023 (3)	-0.006 (2)	0.005 (2)	-0.001 (2)
C6	0.026 (3)	0.023 (3)	0.022 (3)	-0.010 (2)	-0.001 (2)	0.000 (2)
C7	0.022 (3)	0.021 (3)	0.020 (3)	0.000 (2)	0.002 (2)	0.006 (2)
C8	0.020 (3)	0.022 (3)	0.030 (3)	-0.002 (2)	0.006 (2)	0.000 (2)

*Geometric parameters (Å, °)*

N3—C9	1.510 (6)	C16—H16B	0.9800
N3—C11	1.509 (7)	C16—H16C	0.9800
N3—C13	1.524 (6)	Pt1—Br1	2.4335 (6)
N3—C15	1.516 (7)	Pt1—Br2	2.4216 (5)
C9—H9A	0.9900	Pt1—Br3	2.4367 (5)
C9—H9B	0.9900	Pt1—N1	2.041 (4)
C9—C10	1.515 (8)	S1—C1	1.724 (5)
C10—H10A	0.9800	S1—C7	1.738 (5)
C10—H10B	0.9800	O1—N2	1.221 (6)
C10—H10C	0.9800	O2—N2	1.228 (6)
C11—H11A	0.9900	N1—C1	1.315 (6)
C11—H11B	0.9900	N1—C2	1.383 (6)
C11—C12	1.516 (8)	N2—C4	1.476 (7)
C12—H12A	0.9800	C1—C8	1.486 (7)
C12—H12B	0.9800	C2—C3	1.405 (7)
C12—H12C	0.9800	C2—C7	1.397 (7)
C13—H13A	0.9900	C3—H3	0.9500
C13—H13B	0.9900	C3—C4	1.372 (7)
C13—C14	1.509 (8)	C4—C5	1.392 (7)
C14—H14A	0.9800	C5—H5	0.9500
C14—H14B	0.9800	C5—C6	1.370 (8)
C14—H14C	0.9800	C6—H6	0.9500
C15—H15A	0.9900	C6—C7	1.399 (7)
C15—H15B	0.9900	C8—H8A	0.9800
C15—C16	1.557 (8)	C8—H8B	0.9800
C16—H16A	0.9800	C8—H8C	0.9800
C9—N3—C13	108.7 (4)	C16—C15—H15B	108.7
C9—N3—C15	107.5 (4)	C15—C16—H16A	109.5
C11—N3—C9	112.4 (4)	C15—C16—H16B	109.5
C11—N3—C13	108.7 (4)	C15—C16—H16C	109.5
C11—N3—C15	108.9 (4)	H16A—C16—H16B	109.5
C15—N3—C13	110.5 (4)	H16A—C16—H16C	109.5
N3—C9—H9A	108.6	H16B—C16—H16C	109.5
N3—C9—H9B	108.6	Br1—Pt1—Br3	176.23 (2)
N3—C9—C10	114.8 (5)	Br2—Pt1—Br1	91.183 (19)
H9A—C9—H9B	107.5	Br2—Pt1—Br3	90.989 (19)
C10—C9—H9A	108.6	N1—Pt1—Br1	88.64 (11)
C10—C9—H9B	108.6	N1—Pt1—Br2	178.40 (12)

C9—C10—H10A	109.5	N1—Pt1—Br3	89.28 (11)
C9—C10—H10B	109.5	C1—S1—C7	90.0 (2)
C9—C10—H10C	109.5	C1—N1—Pt1	124.2 (3)
H10A—C10—H10B	109.5	C1—N1—C2	111.9 (4)
H10A—C10—H10C	109.5	C2—N1—Pt1	123.8 (3)
H10B—C10—H10C	109.5	O1—N2—O2	123.9 (5)
N3—C11—H11A	108.6	O1—N2—C4	118.7 (4)
N3—C11—H11B	108.6	O2—N2—C4	117.4 (5)
N3—C11—C12	114.9 (5)	N1—C1—S1	114.6 (4)
H11A—C11—H11B	107.5	N1—C1—C8	124.8 (5)
C12—C11—H11A	108.6	C8—C1—S1	120.6 (4)
C12—C11—H11B	108.6	N1—C2—C3	126.0 (5)
C11—C12—H12A	109.5	N1—C2—C7	114.2 (4)
C11—C12—H12B	109.5	C7—C2—C3	119.7 (5)
C11—C12—H12C	109.5	C2—C3—H3	121.6
H12A—C12—H12B	109.5	C4—C3—C2	116.8 (5)
H12A—C12—H12C	109.5	C4—C3—H3	121.6
H12B—C12—H12C	109.5	C3—C4—N2	117.7 (5)
N3—C13—H13A	108.5	C3—C4—C5	123.6 (5)
N3—C13—H13B	108.5	C5—C4—N2	118.6 (5)
H13A—C13—H13B	107.5	C4—C5—H5	120.0
C14—C13—N3	115.3 (4)	C6—C5—C4	120.0 (5)
C14—C13—H13A	108.5	C6—C5—H5	120.0
C14—C13—H13B	108.5	C5—C6—H6	121.2
C13—C14—H14A	109.5	C5—C6—C7	117.6 (5)
C13—C14—H14B	109.5	C7—C6—H6	121.2
C13—C14—H14C	109.5	C2—C7—S1	109.3 (4)
H14A—C14—H14B	109.5	C2—C7—C6	122.2 (5)
H14A—C14—H14C	109.5	C6—C7—S1	128.5 (4)
H14B—C14—H14C	109.5	C1—C8—H8A	109.5
N3—C15—H15A	108.7	C1—C8—H8B	109.5
N3—C15—H15B	108.7	C1—C8—H8C	109.5
N3—C15—C16	114.3 (5)	H8A—C8—H8B	109.5
H15A—C15—H15B	107.6	H8A—C8—H8C	109.5
C16—C15—H15A	108.7	H8B—C8—H8C	109.5
C9—N3—C11—C12	-55.8 (6)	N1—C2—C7—S1	-2.1 (6)
C9—N3—C13—C14	-68.6 (6)	N1—C2—C7—C6	178.8 (5)
C9—N3—C15—C16	174.6 (5)	N2—C4—C5—C6	176.8 (5)
C11—N3—C9—C10	-54.2 (6)	C1—S1—C7—C2	1.9 (4)
C11—N3—C13—C14	168.7 (5)	C1—S1—C7—C6	-179.1 (5)
C11—N3—C15—C16	-63.3 (6)	C1—N1—C2—C3	-177.5 (5)
C13—N3—C9—C10	-174.6 (5)	C1—N1—C2—C7	1.1 (6)
C13—N3—C11—C12	64.6 (6)	C2—N1—C1—S1	0.4 (6)
C13—N3—C15—C16	56.0 (6)	C2—N1—C1—C8	-179.1 (5)
C15—N3—C9—C10	65.7 (6)	C2—C3—C4—N2	-177.5 (4)
C15—N3—C11—C12	-174.9 (4)	C2—C3—C4—C5	1.9 (8)
C15—N3—C13—C14	49.2 (6)	C3—C2—C7—S1	176.6 (4)



---

Pt1—N1—C1—S1	176.9 (2)	C3—C2—C7—C6	-2.4 (8)
Pt1—N1—C1—C8	-2.6 (7)	C3—C4—C5—C6	-2.6 (8)
Pt1—N1—C2—C3	6.0 (7)	C4—C5—C6—C7	0.7 (8)
Pt1—N1—C2—C7	-175.4 (3)	C5—C6—C7—S1	-177.1 (4)
O1—N2—C4—C3	7.5 (7)	C5—C6—C7—C2	1.7 (8)
O1—N2—C4—C5	-171.9 (5)	C7—S1—C1—N1	-1.4 (4)
O2—N2—C4—C3	-173.4 (5)	C7—S1—C1—C8	178.2 (5)
O2—N2—C4—C5	7.2 (7)	C7—C2—C3—C4	0.6 (7)
N1—C2—C3—C4	179.2 (5)		

---