

RESEARCH ARTICLE

Open Access

Antibacterial activity of *Artemisia nilagirica* leaf extracts against clinical and phytopathogenic bacteria

Abdul R Ahameethunisa, Waheeta Hopper*

Abstract

Background: The six organic solvent extracts of *Artemisia nilagirica* were screened for the potential antimicrobial activity against phytopathogens and clinically important standard reference bacterial strains.

Methods: The agar disk diffusion method was used to study the antibacterial activity of *A. nilagirica* extracts against 15 bacterial strains. The Minimum Inhibitory Concentration (MIC) of the plant extracts were tested using two fold agar dilution method at concentrations ranging from 32 to 512 µg/ml. The phytochemical screening of extracts was carried out for major phytochemical derivatives in *A. nilagirica*.

Results: All the extracts showed inhibitory activity for gram-positive and gram-negative bacteria except for *Klebsiella pneumoniae*, *Enterococcus faecalis* and *Staphylococcus aureus*. The hexane extract was found to be effective against all phytopathogens with low MIC of 32 µg/ml and the methanol extract exhibited a higher inhibition activity against *Escherichia coli*, *Yersinia enterocolitica*, *Salmonella typhi*, *Enterobacter aerogenes*, *Proteus vulgaris*, *Pseudomonas aeruginosa* (32 µg/ml), *Bacillus subtilis* (64 µg/ml) and *Shigella flaxneri* (128 µg/ml). The phytochemical screening of extracts answered for the major derivative of alkaloids, amino acids, flavonoids, phenol, quinines, tannins and terpenoids.

Conclusion: All the extracts showed antibacterial activity against the tested strains. Of all, methanol and hexane extracts showed high inhibition against clinical and phytopathogens, respectively. The results also indicate the presence of major phytochemical derivatives in the *A. nilagirica* extracts. Hence, the isolation and purification of therapeutic potential compounds from *A. nilagirica* could be used as an effective source against bacterial diseases in human and plants.

Background

Artemisia is one of the diverse genera of *Asteraceae* family with many important medicinally valuable essential oils and secondary metabolites. Essential oils of *Artemisia* spp. have been widely used for a variety of medicinal purposes for many years. *Artemisia nilagirica* (Clarke) pamp commonly called Indian wormwood, is widely found in the hilly areas of India. *A. nilagirica* has been reported to exhibit insecticidal activities [1]. Around 59 compounds were identified from essential oil of *A. nilagirica* which showed an inhibitory activity on *Phytophthora capsici*, causing "foot rot" in pepper [2].

Various species of *Artemisia* have been characterized for their biological activities. It is considered to produce most medicinally important secondary metabolites [3,4]. Several interesting studies using *Artemisia* spp. showed a series of antimicrobial and antioxidant activities [5-9]. The qualitative determination of various secondary metabolites like flavonoids, terpenoids, saponins and polysaccharides of *Artemisia* spp. were detected by HPLC, GC-MS and NMR [10,11]. Few considerable secondary metabolites were successfully isolated and used in food industry as an alternative to synthetic antimicrobials [12,13]. Furthermore, extracts of *Artemisia* spp. were used as natural pesticide and also in the treatment of few human diseases [14-17]. The determination of potential antimicrobial activity of *Artemisia nilagirica* extracts could be more informative for the future use in

* Correspondence: bioinfresearch2009@gmail.com
Department of Bioinformatics, School of Bioengineering, SRM University, Kattankulathur, Tamil Nadu, India

controlling phytopathogens and also in clinical treatment as natural antimicrobial agents.

The organisms like *Escherichia*, *Enterobacter*, *Klebsiella*, *Proteus*, *Shigella* and *Staphylococcus* species are implicated to cause severe infections in human, as they are found in multiple environmental habitats [18,19]. *Erwinia* spp., *Clavibacter michiganense*, *Pseudomonas syringae* and *Xanthomonas campestris* were reported to be severe phytopathogens, causing damage in carrot, potato, tomato, leafy greens, onion, green pepper, squash and other cucurbits. Furthermore, these phytopathogens cause disease in any plant tissue it invades [20].

In the present study, the antimicrobial potency of chloroform, diethyl ether, ethanol, hexane, methanol and petroleum ether extracts of *Artemisia nilagirica* was investigated. The antibacterial activity was determined by disk diffusion method and minimum inhibitory concentration (MIC) test. Four plant pathogens and 11 clinically important CLSI [21] reference bacterial strains from American Type Culture Collection (ATCC), Microbial Type Culture Collection (MTCC) and a local isolate from SPIC Science Foundation Patholab (SSFP) were used as test cultures. The preliminary phytochemical screening was carried out to identify the derivatives in the extracts.

Methods

Plant material and extraction

Fresh leaves of *Artemisia nilagirica* were collected from Nilgiri district, Tamil Nadu, India. Plant leaves were cleaned with deionized water and dried at shade for a week. Blotted leaves were grounded and filtered using four layers of gauss cloth. The plant powder was stored in air tight container and maintained at 4°C until use.

Solvent systems used for the extractions were chloroform, diethyl ether, ethanol, hexane, methanol and petroleum ether. Soxhlet and flask extraction procedures were adapted for extraction. Ten grams of the powdered samples were packed in muslin cloth and used for extraction by soxhlet apparatus at a temperature below the boiling temperature of each solvent. A portion of the powdered plant samples was soaked in the conical flask containing solvent, wrapped with aluminum foil and placed in shaker for 48 hours at 120-130 rpm.

After 48 hours, the extracts were filtered using Whatman filter paper No: 1. The solvent was evaporated and the residue was dissolved in sterile dimethylsulfoxide (DMSO-9:1) in 50 mg/ml concentration. The extract was filtered using 0.22 micro filter (Type GV- Millipore) and stored at 4°C for further antimicrobial activity study.

Test microorganisms

The 15 bacterial cultures of both gram-positive and gram-negative bacterial strains used for screening are:

Erwinia sp. (MTCC 2760), *Xanthomonas campestris* (MTCC 2286), *Pseudomonas syringae* (ATCC 7386), *Clavibacter michiganense* (ATCC 27822), *Escherichia coli* (ATCC 25922), *Yersinia enterocolitica* (MTCC 840), *Klebsiella pneumoniae* (ATCC 15380), *Salmonella typhi* (SSFP 4S), *Enterobacter aerogenes* (MTCC 111), *Proteus vulgaris* (MTCC 1771), *Pseudomonas aeruginosa* (ATCC 27853), *Shigella flaxneri* (MTCC 1457), *Bacillus subtilis* (MTCC 441), *Enterococcus faecalis* (ATCC 29212) and *Staphylococcus aureus* (MTCC 29212). The stock cultures were maintained in nutrient agar (NA) slant at 4°C and sub-cultured monthly. Working cultures were prepared by inoculating a loopful of each test microorganism in 3 ml of nutrient broth (NB) from NA slants. Broths were incubated at 37°C for 12 hours. The suspension was diluted with sterile distilled water to obtain approximately 10⁶ CFU/ml.

Screening for antibacterial activity

Disk diffusion: 5 mm of sterile disks were incorporated in 100 µl of plant extracts (5 mg/disk). The disk was completely saturated with the extract and allowed to dry. Mueller Hinton (MH) agar plates were swabbed with test bacteria and six extract disks with one of the standard positive control disks (ampicillin, streptomycin or gentamycin) was placed on the MH agar plate. DMSO was taken as the negative control. Plates were incubated overnight at 37°C.

Agar dilution susceptibility test

The agar dilution susceptibility test was performed based on modified method of NCCLS and CLSI [22,23] to determine the MIC. Extracts dissolved in sterilized DMSO (5120 µg/ml concentration) were taken as standard stock. A series of two fold dilutions of each extract in the final concentration of 512, 256, 128, 64 and 32 µg/ml were prepared in MH agar. After solidification, the plates were spotted with 2 µl of overnight grown bacterial cultures approximately containing 1 × 10⁴ CFU/ml. The test was carried out in triplicates. The plates were incubated overnight at 37°C. After 18 - 24 hours, the MIC was determined and the percentage of growth inhibition was calculated by,

$$\text{Percentage of inhibition (Pi)} = \frac{T-SC}{PC} \times 100$$

T: Test; SC: Solvent control; PC: positive control

Phytochemical screening

To identify the phytochemical derivatives in the extracts, standard phytochemical screening was performed [24,25]. Alkaloids test was performed by Dragendorff's and Meyer's tests, amino acids by ninhydrin, carbohydrates by Barfoed's and Fehling tests, flavonoids by FeCl₃, Pew's and Shinoda's tests, glycosides by Killer-Killanis test, saponin by frothing test, tannins by FeCl₃

Table 1 Antibacterial activity screening of *A. nilagirica* by disk diffusion

Bacterial culture	Positive control	Solvent control	Zone of inhibition (mm diameter)					
			Ethanol	Methanol	Hexane	Petroleum Ether	Chloroform	Diethyl ether
<i>Erwinia sp.</i>	15	-	10 ± 0.0	12 ± 0.5	13 ± 1.0	10 ± 0.0	12 ± 0.0	10 ± 0.5
<i>X. campestris</i>	16	-	11 ± 0.5	12 ± 0.0	14 ± 0.0	11 ± 0.5	12 ± 0.1	11 ± 0.5
<i>P. syringae</i>	14	-	11.8 ± 0.5	11 ± 0.0	12 ± 0.0	10 ± 0.0	10 ± 0.0	12 ± 0.5
<i>C. michiganense</i>	15	-	12 ± 1.0	12 ± 0.0	13 ± 0.0	8 ± 0.0	11 ± 0.0	12.4 ± 0.5
<i>E. coli</i>	13	-	14 ± 1.0	12 ± 0.5	8 ± 0.0	8 ± 0.0	11 ± 0.5	11 ± 0.0
<i>Y. enterocolitica</i>	14	-	11 ± 1.0	12 ± 0.0	11 ± 0.0	9 ± 0.5	13 ± 0.0	14 ± 0.0
<i>B. subtilis</i>	13	-	8 ± 0.0	8 ± 0.0	-	8 ± 0.0	10 ± 0.0	10 ± 0.0
<i>E. faecalis</i>	14	-	-	-	-	-	7 ± 0.0	8 ± 0.0
<i>K. pneumonia</i>	12	-	-	8 ± 0.0	-	-	-	-
<i>S. typhi</i>	10	-	8 ± 0.0	10 ± 0.0	13 ± 1.0	12 ± 1.0	8 ± 0.0	8 ± 0.0
<i>S. aureus</i>	12	-	-	-	-	-	7 ± 0.0	-
<i>E. acrogens</i>	14	-	8 ± 0.0	11 ± 1.0	8 ± 0.0	8 ± 0.0	10 ± 0.0	8 ± 0.5
<i>P. vulgaris</i>	14	-	8 ± 0.0	12 ± 1.0	14 ± 0.0	14 ± 0.5	10 ± 0.0	8 ± 0.0
<i>P. aeruginosa</i>	13	-	10 ± 0.0	12 ± 0.0	13 ± 0.0	12 ± 0.0	10 ± 0.0	8 ± 0.0
<i>S. flexneri</i>	14	-	8 ± 0.0	10 ± 1.0	8 ± 0.0	8 ± 0.0	10 ± 0.0	8 ± 0.0

± - mean standard deviation of triplicates, Concentration of extract- 5 mg/disk, (-) - No zone of inhibition observed, Positive controls - ampicillin or streptomycin or gentamycin (10 µg/ml), Solvent control - 10% DMSO.

and lead acetate & terpenoids by Salkowski test. The test for hydrolysable tannins, phlobatannins, phenol, quinones and volatile oils were also carried out as in literature [26-28].

Statistical analysis

Analysis of variance (ANOVA) was performed using GeneSpring GX - 7.3 microarray software to determine the significance of MIC values between the extracts against bacterial culture. The probability value (p) ≤ 0.05 was considered as significance in this study.

Results & Discussion

The antibacterial activity of *A. nilagirica* leaf extracts were examined against 11 clinical and four phytopathogens causing illness in human and damage in major crops [20], respectively. The extractions were carried out using chloroform, diethyl ether, ethanol, hexane, methanol and petroleum ether solvents. The ethanol and methanol extracts gave the high yield of 2.3% (% concentration w/v) and hexane gave 1.6% w/v. While, other extracts provide much low yield of 0.6% w/v in soxhlet and flask extraction procedures. The antibacterial activity of the organic solvent extracts showed varying magnitudes of inhibition patterns with standard positive control depending on the susceptibility of the tested microorganism. Out of 15 bacterial strains tested, 12 showed inhibition activity to one or more extracts. The mean inhibitory zone of six solvent extracts against 15 bacterial species is summarized in Table 1.

The analysis of methanol and chloroform extracts against phytopathogens showed a significant level of inhibition against *Erwinia sp.* and *X. campestris*. On the

other hand, ethanol and diethyl ether extracts showed high activity against *C. michiganense* and *P. syringae* (p ≤ 0.05). Also, the petroleum ether extracts showed 11-12 mm zone of inhibition to *C. michiganense*. Interestingly, hexane extract of *A. nilagirica* exhibits maximum inhibitory activity against all the phytopathogens in comparison to other extracts [Table 1]. Further, hexane extracts showed the significant inhibitory effect against *Clavibacter michiganense* (13 mm), *Erwinia sp* (13 mm), *Pseudomonas syringae* (12 mm) and *Xanthomonas campestris* (14 mm). It is understandable that hexane extract is more potent showing a higher degree of antimicrobial activity to phytopathogens in comparison to other extracts. Also, the results of hexane extract against *X. campestris* was appeared to be two-fold better than the previous study of *Phyllanthus emblica*, *Acacia nilotica*, *Sapindus mukorossi* and *Terminalia chebula* which shows 6.00 mm zone of inhibition at 50 gm/l concentration reported as the most effective for *X. campestris* [29]. In addition, moderate effects were seen in chloroform, diethyl ether, ethanol and methanol extracts against all tested phytopathogens except petroleum ether which showed comparatively minimum area of inhibition. This possibly means that the compound responsible for the antibacterial activity was least in concentration.

Similar analysis of *A. nilagirica* leaf extracts were carried out on clinical bacterial pathogens. The hexane, methanol and petroleum ether extracts exhibited significant high inhibitory zones against *P. aeruginosa*, *P. vulgaris* and *S. typhi*. The chloroform and diethyl ether extracts showed maximum area zone of inhibition (10

Table 2 Minimum inhibitory concentration (MIC) of various solvent extracts of *A. nilagirica* (% v/v) against micro-organisms

Test organism		Plant extracts					
		Chloroform	Diethyl ether	Ethanol	Hexane	Methanol	Petroleum ether
		Concentration (µg/ml) showing Minimum inhibitor concentration					
<i>Erwinia sp.</i>	Phytopathogens	32	64	64	32	64	64
<i>Xanthomonas campestris</i>		32	64	32	32	32	128
<i>Pseudomonas syringae</i>		64	32	128	32	64	64
<i>Clavibacter michiganense</i>		128	32	32	32	32	32
<i>Escherichia coli</i>	Clinical pathogens	256	256	32	256	32	128
<i>Yersinia enterocolitica</i>		64	256	128	128	32	512
<i>Bacillus subtilis</i>		32	32	32	32	64	32
<i>Enterococcus faecalis</i>		*	*	*	*	512	*
<i>Klebsiella pneumonia</i>		*	*	*	*	*	*
<i>Salmonella typhi</i>		32	64	128	64	32	32
<i>Staphylococcus aureus</i>		*	*	*	512	*	*
<i>Enterobacter aerogenes</i>		512	128	32	128	32	32
<i>Proteus vulgaris</i>		32	256	128	32	32	64
<i>Pseudomonas aeruginosa</i>		512	128	128	128	32	512
<i>Shigella flexneri</i>		512	256	256	256	128	128

* No inhibition observed upto: 512 µg/ml.

mm) for *B. subtilis*, compared to other extracts (8 mm). The ethanol extract exhibited 14 mm zone for *E. coli*, which is the maximum with respect to the positive standard streptomycin. Subsequently, antibacterial activity of methanol (12 mm), chloroform (13 mm) and diethyl ether (14 mm) extracts were found as effective for *Y. enterocolitica*. Moderate activities were observed against *S. flexneri* and *E. aerogenes* for chloroform and methanol extracts. Among the 11 clinical bacterial strains, *Escherichia coli*, *Yersinia enterocolitica*, *Bacillus subtilis*, *Salmonella typhi*, *Enterobacter aerogenes*, *Proteus vulgaris*, *Pseudomonas aeruginosa* and *Shigella flexneri* were the most susceptible bacteria to all solvent extracts. Nearly, 40% growth inhibitions were observed in concentration of 5 mg/ml. Surprisingly, no activity were observed against *S. aureus*, *E. faecalis* and *K. pneumoniae* suggesting their resistance to *A. nilagirica* extracts.

The MIC tests of *A. nilagirica* organic solvent extracts against 15 bacterial species were carried out using the micro dilution technique. The MIC values of 6 extracts ranged from 32 to 512 µg/ml [Table 2]. While considering phytopathogens, chloroform extract showed maximum activity with MIC 32 µg/ml for *Erwinia sp.* and *X. campestris*. Diethyl ether extracts showed 32 µg/ml MIC for *P. syringae* and *C. michiganense*. In general, all extracts showed less than 128 µg/ml MIC for the tested phytopathogens. Interestingly, hexane extract had higher activity in all the phytopathogens even at a low concentration of 32 µg/ml. The MIC analysis of clinical pathogens showed that the methanol extract was highly active

in comparison to other extracts, which inhibited the series of study organisms at a low concentration (32 µg/ml) except *B. subtilis* (64 µg/ml) and *Shigella flexneri* (128 µg/ml) [Table 2]. In supportive to the susceptible test, *A. nilagirica* extracts had very low or no activity, even in the highest concentration (512 µg/ml) MIC against *S. aureus*, *E. faecalis* and *K. pneumoniae*. Hence, we conclude that these organisms are resistant to *A. nilagirica* extracts. According to the results from zone of inhibition and MIC studies [Table 1 & 2], hexane and methanol extracts are considerably good inhibitors for the phytopathogens and clinical pathogenic bacteria, respectively.

Phytochemical test were carried out on the *A. nilagirica* extracts to determine the natural bioactive compound. By studying the presence of phytochemical in *A. nilagirica*, the medicinal value of the plant can be explained scientifically. The phytochemical screening of extracts showed the presence of major derivatives and their results were summarized [Table 3]. The analysis showed the occurrence of alkaloids, flavonoids, phenol, quinines and terpenoids in all extracts. Tannins were present in ethanol, methanol and diethyl ether. Volatile oils were present in methanol, hexane and petroleum ether. Phlobatannins metabolites were found to be present in hexane and petroleum ether and absent in other extracts. Also, saponins and amino acid were present in ethanol and methanol extracts with carbohydrates particularly present in methanol extract. Surprisingly, glycosides and hydrolysable tannins were absent in all the extracts.

Table 3 Phytochemical screening of *A. nilagirica* extracts

S. no.	Test	Chloroform	Diethyl ether	Ethanol	Hexane	Methanol	Petroleum ether
1	Alkaloids	++	++	++	++	++	+
2	Amino acids	-	-	+	-	++	-
3	Carbohydrates	-	-	-	-	+	-
4	Flavonoids	++	+	++	++	++	+
5	Glycosides	-	-	-	-	-	-
6	Hydrolysable tannins	-	-	-	-	-	-
7	Phenol	++	+	++	+	++	+
8	Phlobatannins	-	-	-	+	-	+
9	Quinines	+	+	+	+	+	+
10	Saponin	-	-	+	-	+	-
11	Tannins	-	+	++	-	++	-
12	Terpenoids	++	++	++	++	++	++
13	Volatile oils	-	-	-	+	+	-

(++) abundant, (+) present, (-) absent

In susceptibility test, the hexane extracts showed the considerable levels of inhibition against phytopathogens. The phytopathogens test of petroleum ether extract showed low inhibition range (8 to 10 mm) in comparison to other extracts (10 to 14 mm). In conjugation with phytochemical screening of all the extracts with petroleum ether, showed the variations in abundance in alkaloids derivatives. Hence, it is suggested that reduction of alkaloid abundance in petroleum ether may be the cause of decreased activity in phytopathogens. Supportive to our finding, previous studies indicate the effective role of alkaloid against phytopathogens [30-32]. The MIC analyses of clinical pathogens showed an activity against Gram-positive and Gram-negative bacteria may be indicative of the presence of the broad spectrum antibiotic compounds. The methanol extracts showed high inhibition at the minimal concentration for most of the clinical pathogens in comparison to other extracts. The MIC value of methanol extract ranges from 32 to 64 µg/ml for *Escherichia coli*, *Bacillus subtilis*, *Yersinia enterocolitica*, *Salmonella typhi*, *Enterobacter aerogenes*, *Proteus vulgaris* and *Pseudomonas aeruginosa*. Also, the phytochemical screening of menthol extract showed the presence of most of the derivatives like flavonoids, terpenoids, phenol, amino acids, alkaloids and tannins. Furthermore, alkaloids [33,34], amino acids [35], flavonoids [36-38], phenols [39], tannins [40-42], terpenoids [43] of various plants extracts proven to be effective antimicrobials [44]. Our results are also in agreement with these studies suggesting the efficacy of methanol extract of *A. nilagirica* against clinical pathogens.

Conclusion

Extracts of *A. nilagirica* showed the broad spectrum of antibacterial activity on the tested microorganisms. Hexane extract exhibited high inhibitory potency against

phytopathogens and methanol extract showed maximum inhibition against clinical pathogens except *S. aureus*, *E. faecalis* and *K. pneumoniae*. The phytochemical analysis showed the presence of effective biological compounds like alkaloids, amino acids, flavonoids, phenols, tannins and terpenoids. These derivatives could be potential alternatives to the traditional chemical control of clinical pathogen and phytopathogenic bacteria. Furthermore, the development of natural antimicrobials will help to decrease the negative effects of synthetic drugs. Fractionation and characterization of these active compounds will be the future work to investigate.

Acknowledgements

We thank Dr. Murugesh KA, Assistant professor, Horticultural Research Station, Udthagamandalam, India, for the collection of the plant. Ahameethunisa thank, SRM University for financial support.

Authors' contributions

AA has carried out the experimental part such as extraction, inoculums preparation and antimicrobial evaluation. WH supervised the work, evaluated the results and corrected the manuscript for publication. Authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 23 July 2009

Accepted: 29 January 2010 Published: 29 January 2010

References

- Banerji A, Luthriya DL, Kokate SD: **Toxicity of capillin, the insecticidal principle of *Artemisia nilagirica* Clarke.** *Indian J Exp Biol* 1990, **28**:588-589.
- Shafi PM, Nambiar MKG, Clery RA, Sarma YR, Veena SS: **Composition and antifungal activity of the oil of *Artemisia nilagirica* (Clarke) Pamp.** *JEOR* 2004, **16**:377-379.
- Ambasta SP: **The useful plants of India.** Publications & Information Directorate. CSIR, New Delhi 1986, 55-56.
- Priscila IU, Mariama TNS, Luiz CDS, Luciano B, Fernandes AJ: **Antibacterial activity of medicinal plant extracts.** *Braz J Microbiol* 2007, **38**:717-719.
- Juteau F, Masotti V, Bessiere JM, Dherbomez M, Viano J: **Antibacterial and antioxidant activities of *Artemisia annua* essential oil.** *Fitoterapia* 2002, **73**:532-535.

6. Kordali S, Kotan R, Mavi A, Cakir A, Ala A, Yildirim A: **Determination of the Chemical Composition and Antioxidant Activity of the Essential Oil of Artemisia dracunculoides and the Antifungal and Antibacterial Activities of Turkish Artemisia absinthium, A. dracunculoides, Artemisia santonicum, and Artemisia spicigera Essential Oils.** *J Agric Food Chem* 2005, **53**:9452-9458.
7. Kordali S, Cakir A, Mavi A, Kilic H, Yildirim A: **Screening of chemical composition and antifungal and antioxidant activities of the essential oils from three Turkish Artemisia Species.** *J Agric Food Chem* 2005, **53**:1408-1416.
8. Curini M, Epifano F, Genovese S, Tammaro F, Menghini L: **Composition and antimicrobial activity of the essential oil of Artemisia dracunculoides "piemontese" from Italy.** *Chem Nat Comp* 2006, **42**:738-740.
9. Cha JD, Jung EK, Kil BS, Lee KY: **Chemical composition and antibacterial activity of essential oil from Artemisia feddei.** *J Microbiol Biotechnol* 2007, **17**:2061-2065.
10. Xie G, Schepetkin IA, Siemsen DW, Kirpotina LN, Wiley JA, Quinn MT: **Fractionation and Characterization of Biologically-active Polysaccharides from Artemisia tripartite.** *Phytochemistry* 2008, **69**:1359-1371.
11. Avula B, Wang YH, Smillie TJ, Mabusele W, Vincent L, Weitz F, Khan IA: **Quantitative determination of flavonoids by column high-performance liquid chromatography with mass spectrometry and ultraviolet absorption detection in Artemisia afar and comparative studies with various species of Artemisia plants.** *J AOAC Int* 2009, **92**:633-44.
12. Ng TB: **Antifungal proteins and peptides of leguminous and non-leguminous origins.** *Peptides* 2004, **25**:1215-22.
13. Pattison LT, Lindsay D, von Holy A: **Natural antimicrobial as potential replacements for calcium propionate in bread.** *S Afr J Sci* 2004, **100**:339-342.
14. Mueller MS, Karhagomba IB, Hirt HM, Wemakor E: **The potential of Artemisia annua L. as a locally produced remedy for malaria in the tropics: agricultural, chemical and clinical aspects.** *J Ethnopharmacol* 2000, **73**:487-493.
15. Rahman AF, Saleh MA: **Nematicidal activity of phytochemicals from some arid land plants.** Society of nematologists 45th annual meeting, Lihue, Hawaii. *J Nematol* 2006, **38**:258-303.
16. Saddi M, Sanna A, Cottiglia F, Chisu L, Casu L, Bonsignore L, De Logu A: **Antitherpevirus activity of Artemisia arborescens essential oil and inhibition of lateral diffusion in Vero cells.** *Ann Clin Microbiol Antimicrob* 2007, **6**:10.
17. Meneses R, Ocazonez RE, Martínez JR, Stashenko EE: **Inhibitory effect of essential oils obtained from plants grown in Colombia on yellow fever virus replication in vitro.** *Ann Clin Microbiol Antimicrob* 2009, **8**:8.
18. Emori TG, Gaynes RP: **An overview of nosocomial infections, including the role of the microbiology laboratory.** *Clin Microbiol Rev* 1993, **6**:428-442.
19. Maleki S, Seyyednejad SM, Damabi NM, Motamedi H: **Antibacterial activity of the fruits of Iranian torilis leptophylla against some clinical pathogens.** *Pak J Biol Sci* 2008, **11**:1286-1289.
20. Setubal JC, Moreira LM, da Silva AC: **Bacterial phytopathogens and genome science.** *Curr Opin Microbiol* 2005, **8**:595-600.
21. Clinical and Laboratory Standards Institute (CLSI): **Performance standards for antimicrobial susceptibility testing. Eighteenth informational supplement CLSI document M100-S18, Wayne, PA, USA** 2008.
22. National Committee for Clinical Laboratory Standards (NCCLS): **Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved Standard - NCCLS document M7-A6, Wayne, PA, USA, Sixth** 2003.
23. Clinical and Laboratory Standards Institute (CLSI): **Method for dilution antimicrobial susceptibility tests for bacterial that grow aerobically; approved standard - Eighth Edition. CLSI document M07-A8, Wayne, PA, USA** 2009.
24. Harborne JB: **Phytochemical methods: a guide to modern techniques of plant analysis.** London, Chapman and Hall, Ltd, 2 1973, 49-188.
25. Trease GE, Evans WC: **Orders and families of plant in pharmacognosy.** Oxford University Press 1983, **12**:343-383.
26. Sofowora A: **Recent trends in research into African medicinal plants.** *J Ethnopharmacol* 1993, **38**(2-3):209-214.
27. Sofowora A: **Research on Medicinal Plants and Traditional Medicine in Africa.** *J Altern Complement Med* 1996, **2**:365-372.
28. Krishnaiah D, Devi T, Bono A, Sarbatly R: **Studies on phytochemical constituents of six Malaysian medicinal plants.** *J Med Plants Res* 2009, **3**:067-072.
29. Akhtar MA, Rahber-Bhatti MH, Aslam M: **Antibacterial activity of plant diffusate against Xanthomonas campestris PV. Citri.** *Int J Pest Manag* 1997, **43**:149-153.
30. Torres YR, Berlinck RGS, Nascimento GGF, Fortier SC, Pessoa C, de Moraes MO: **Antibacterial activity against resistant bacteria and cytotoxicity of four alkaloid toxins isolated from the marine sponge Arenosciera brasiliensis.** *Toxicon* 2002, **40**:885-891.
31. Raghavendra MP, Sathis S, Raveesha Ka: **In vitro antibacterial potential of alkaloids of Samanea saman (Jacq.) merr. Against Xanthomonas and human pathogenic bacteria.** *World J Agric Sci* 2008, **4**:101-105.
32. Urgeova E, Polivka L: **Secondary metabolites with antibacterial effects from leaves of different hop cultivars during vegetal periods.** *Nova Biotechnologica* 2009, **9**:327-332.
33. Shaheen F, Khan RA, Soobia A, Khan SA, Saima T, Aqeel A: **New antimicrobial alkaloids from the roots of Polyalthia longifolia var. pendula.** *Planta medica* 2003, **69**:350-355.
34. Chakraborty A, Brantner AH: **Antibacterial steroid alkaloids from the stem bark of Holarrhena pubescens villosum leaves.** *J Ethnopharmacol* 1999, **68**:339-344.
35. Chowdhury N, Laskar S, Chandra G: **Mosquito larvicidal and antimicrobial activity of protein of Solanum.** *BMC Compl Alt Med* 2008, **8**:62-67.
36. Brandao MGL, Krettl AU, Soares LSR, Nery CGC, Marinuzzi HC: **Antimalarial activity of extracts and fractions from Bidens pilosa and other Bidens species (Asteraceae) correlated with the presence of acetylene and flavonoids compounds.** *J Ethnopharmacol* 1997, **57**:131-138.
37. Hernandez NE, Tereschuk ML, Abdala LR: **Antimicrobial activity of flavonoids in medicinal plants from Tafi Del Valle (Tucuman, Argentina).** *J Ethnopharmacol* 2000, **73**:317-322.
38. Mendoza L, Wilkens M, Urzua A: **Antimicrobial study of the resinous exudates and of diterpenoids and flavonoids isolated from some Chilean Pseudognaphalium (Asteraceae).** *J Ethnopharmacol* 1997, **58**:85-88.
39. Fernandez MA, Garcia MD, Saenz MT: **Antibacterial activity of the phenolic acids fractions of Scrophularia frutescens and Scrophularia sambucifolia.** *J Ethnopharmacol* 1996, **53**:11-14.
40. Akiyama H, Fujii K, Yamasaki O, Oono T, Iwatsuki K: **Antibacterial action of several tannins against Staphylococcus aureus.** *J Antimicrob Chemother* 2001, **48**:487-491.
41. Doss A, Mubarak MH, Dhanabalan R: **Antibacterial activity of tannins from the leaves of Solanum trilobatum Linn.** *Indian J Sci Technol* 2009, **2**:41-43.
42. Amarowicz R, Dykes GA, Pegg RB: **Antibacterial activity of tannin constituents from Phaseolus vulgaris, Fagopyrum esculentum, Corylus avellana and Juglans nigra.** *Fitoterapia* 2008, **79**:217-219.
43. Amaral JA, Ekins A, Richards SR, Knowles R: **Effect of selected monoterpenes on methane oxidation, denitrification, and aerobic metabolism by bacteria in pure culture.** *Appl Environ Microbiol* 1998, **64**:520-525.
44. Cowan MM: **Plant Products as Antimicrobial Agents.** *Clin Microbiol Rev* 1999, **12**:564-582.

Pre-publication history

The pre-publication history for this paper can be accessed here:<http://www.biomedcentral.com/1472-6882/10/6/prepub>

doi:10.1186/1472-6882-10-6

Cite this article as: Ahameethunisa and Hopper: Antibacterial activity of Artemisia nilagirica leaf extracts against clinical and phytopathogenic bacteria. *BMC Complementary and Alternative Medicine* 2010 **10**:6.