

COMMENTARY

Endogenous H₂S in hemorrhagic shock: innocent bystander or central player?

Enrico Calzia*¹, Peter Radermacher¹ and Kenneth R Olson²

See related research by Van de Louw and Haouzi, <http://ccforum.com/content/16/5/R178>

Abstract

The role of the gaseous mediator hydrogen sulfide (H₂S) in hemorrhagic shock is still a matter of debate. This debate is emphasized by the fact that available literature data on blood and tissue H₂S concentrations vary by three orders of magnitude, both under physiological conditions as well as during stress states. Therefore, in a rat model of unresuscitated, lethal hemorrhagic shock, Van de Louw and Haouzi tested the two hypotheses of whether blood and tissue H₂S levels would increase due to the shock-related tissue hypoxia, and whether vitamin B12 would attenuate organ injury and improve survival as a result of enhanced H₂S oxidation. Hemorrhage did not affect the blood and tissue H₂S content, and, despite the increased capacity to oxidize H₂S, vitamin B12 did not affect any parameter of shock severity. The authors concluded that H₂S concentrations cannot be used as a marker of shock, most probably as a result of tissue's capacity to oxidize H₂S even under conditions of severe oxygen debt. This research paper elegantly re-adjusts the currently available data on blood and tissue H₂S levels, and thereby adds an important piece to the puzzle of whether H₂S release should be enhanced or lowered during stress conditions associated with tissue hypoxia.

In a recent issue of *Critical Care*, Van de Louw and Haouzi report on the effects of lethal hemorrhage on blood and tissue levels of hydrogen sulfide (H₂S) [1]. The role of H₂S during hemorrhage is a matter of debate: while both inhaled H₂S and intravenous sodium sulfide

and sodium hydrosulfide improved survival [2-4], other authors reported that sodium sulfide did not exert any beneficial effects [5]. Moreover, blocking H₂S biosynthesis by inhibiting cystathione-γ-lyase attenuated circulatory failure and organ injury [6,7]. Since hypoxic conditions decrease [8,9] and supplemental vitamin B12 (hydroxocobalamin) increases (due to the rise in oxidative capacity) the rate of H₂S metabolism, the authors hypothesized that hemorrhage would increase plasma and tissue H₂S levels, and that vitamin B12 would improve survival. Rats were hemorrhaged by five times withdrawal of 5 ml/kg blood (that is, approximately 30% of the calculated blood volume). The total H₂S content was measured in the first and last blood samples, using the methylene blue assay [10]. Indirect calorimetry for oxygen uptake and carbon dioxide production before and at the end of the hemorrhage period allowed determination of the shock-induced oxygen deficit. The major finding was that, despite a severe cumulative oxygen debt (100 to 140 ml/kg), H₂S blood and tissue concentrations did not change, rendering them useless as markers of shock severity. In line with this finding, vitamin B12 failed to exert any therapeutic effects despite an increased capacity to oxidize H₂S.

What do we learn from this study? According to the authors' standard curve for the methylene blue assay, the plasma light absorbance peak at 670 nm would correspond to ~8 μM H₂S [1]. This absorbance, however, was due to turbidity rather than the presence of the blue dye. The true H₂S concentrations were most probably therefore much lower, possibly even below the detection limit of 1.5 μM.

There is considerable discrepancy in the literature on blood H₂S concentrations. In rats, baseline values of 25 to 50 μM have been reported, which increased up to 80 μM after hemorrhage, endotoxin exposure and injection of sulfide donors [6,11,12]. However, bolus (4 mg/kg) or continuous intravenous (20 mg/kg/hour) sodium sulfide only increased blood H₂S levels from 0.4 to 0.9 μM to 4.0 to 4.5 μM when the monobromobamine assay [10] was used to determine H₂S concentrations [13]. In mice, 10 mg/kg endotoxin either decreased (from ~2.3 to

*Correspondence: enrico.calzia@uni-ulm.de

¹Sektion Anästhesiologische Pathophysiologie und Verfahrensentwicklung, Klinik für Anästhesiologie, Universitätsklinikum Ulm, Helmholzstrasse 8-1, D-89081 Ulm, Germany

Full list of author information is available at the end of the article

~1.8 μM [14]) or increased (from ~34 to ~65 μM [15]) the blood sulfide content. Finally, inhaling up to 200 ppm gaseous H_2S in mice increased the sulfide content by <1.5 μM [13,16,17].

According to the available literature, the blood H_2S content may vary by three orders of magnitude – so which H_2S concentrations are real? At physiological pH, dissolved H_2S gas represents 20 to 50% of the total sulfide [9,10,17], which can of course escape into the headspace [9,10]. Van de Louw and Haouzi carefully avoided any H_2S loss related to volatilization. Furthermore, blood-borne H_2S is rapidly bound and/or metabolized: using a polarographic sensor with a detection limit for H_2S gas corresponding to 100 nM total sulfide in blood at pH 7.4, a 10 μM sodium sulfide spike only transiently increased sulfide from undetectable levels to about 0.5 μM [18]. Finally, the odor threshold of H_2S is 0.01 to 0.3 ppm [9,10], and simply smelling the blood allows one to verify that plasma H_2S concentrations are at, or below, 1 μM : in a phosphate buffer, the human nose can detect as little as 1 μM H_2S [9,19]. The gas/water coefficient of distribution for H_2S is 0.39 [9,10]. Assuming that only 20% of the dissolved gas (that is, 4 to 10% of the total free sulfide) disappears from the blood sample due to volatilization during the 2 or 3 seconds of sniffing [10], a 10 ml blood sample would have a total free sulfide concentration of 20 to 50 μM . This is within the range reported for rat blood, but clearly rat blood does not smell like rotten eggs!

Based on the results of their study, the authors conclude that ‘ H_2S in the blood cannot be used as a marker of hemorrhagic shock’, because any H_2S accumulation resulting from ‘tissue hypoxia must be reconciled with the ability of tissues to oxidize H_2S ’ [1]. Clearly, H_2S consumption via the sulfide quinone reductase system [20] is reduced under hypoxic conditions, which in turn would cause H_2S accumulation: in lung tissue homogenates and pulmonary artery smooth muscle cells [8,9], H_2S consumption dropped at oxygen concentrations of 10 μM (that is, ~8 mmHg) – a 50% drop occurring at oxygen partial pressure values of ~4 mmHg. In the present study, at the end of the hemorrhage the arterial oxygen partial pressure was still normal and, despite a 50% reduction of minute ventilation, the arterial carbon dioxide partial pressure was reduced to 31 mmHg. The hypoxia was probably therefore not sufficiently low to significantly impair H_2S oxidation. Unfortunately, the authors only reported lactatemia, and not pH, so the severity of lactic acidosis cannot be definitively estimated. Finally, core temperature was maintained at 35 to 36°C. Mild hypothermia of 34°C is well established to attenuate lactic acidosis and better preserve tissue oxygenation. Consequently, albeit ultimately lethal, the authors’ model might have been unable to detect the hypoxia threshold necessary to cause H_2S accumulation. In this context, it is

tempting to speculate whether any local H_2S accumulation might decrease energy expenditure and thereby oxygen consumption due to inhibition of cytochrome c oxidase, which in turn would restore the oxygen partial pressure and thereby at least partially resume H_2S oxidation. Furthermore, it is noteworthy that H_2S can even sustain ATP synthesis during mild hypoxia [21], and thus presumably attenuate tissue lactic acidosis.

Van de Louw and Haouzi meticulously conducted a study in a “hot” field. The authors have the merit of re-adjusting the data available on blood H_2S concentrations. Moreover, lacking any observable increase in H_2S levels during hemorrhage coupled with the currently available knowledge on the role of cystathione- γ -lyase for oxygen sensing and vascular homeostasis [8,9,22], blocking endogenous H_2S production most probably has little therapeutic benefit and may actually prove to be contraindicated. Additional studies are mandatory on the exogenous H_2S supplementation during hemorrhagic shock, in order to answer the question of whether H_2S is an innocent bystander or a central player under these conditions.

Abbreviations

H_2S , hydrogen sulfide.

Competing interests

PR received research grants from Ikaria Inc. The remaining authors declare that they have no competing interests.

Acknowledgements

Supported by the Deutsche Forschungsgemeinschaft (Klinische Forschergruppe 200 ‘Die Entzündungsantwort nach Muskulo-Skeletalem Trauma’) and by National Science Foundation Grant Number IOS-1051627.

Author details

¹Sektion Anästhesiologische Pathophysiologie und Verfahrensentwicklung, Klinik für Anästhesiologie, Universitätsklinikum Ulm, Helmholzstrasse 8-1, D-89081 Ulm, Germany. ²Department of Physiology, Indiana University School of Medicine, 1234 Notre Dame Avenue, South Bend, IN 46617, USA.

Published: 29 November 2012

References

1. Van de Louw A, Haouzi P: Oxygen deficit and H_2S in hemorrhagic shock in rats. *Crit Care* 2012, **16**:R178.
2. Morrison ML, Blackwood JE, Lockett SL, Iwata A, Winn RK, Roth MB: Surviving blood loss using hydrogen sulfide. *J Trauma* 2008, **65**:183-188.
3. Ganster F, Burban M, de La Bourdonnaye M, Fizanne L, Douay O, Loufrani L, Mercat A, Calès P, Radermacher P, Henrion D, Asfar P, Meziari F: Effects of hydrogen sulfide on hemodynamics, inflammatory response and oxidative stress during resuscitated hemorrhagic shock in rats. *Crit Care* 2010, **14**:R165.
4. Bracht H, Scheuerle A, Gröger M, Hauser B, Matallo J, McCook O, Seifritz A, Wachter U, Vogt JA, Asfar P, Matejovic M, Möller P, Calzia E, Szabó C, Stahl W, Hoppe K, Stahl B, Lampl L, Georgieff M, Wagner F, Radermacher P, Simon F: Effects of intravenous sulfide during resuscitated porcine hemorrhagic shock. *Crit Care Med* 2012, **40**:2157-2167.
5. Drabek T, Kochanek PM, Stezoski J, Wu X, Bayir H, Morhard RC, Stezoski SW, Tisherman SA: Intravenous hydrogen sulfide does not induce hypothermia or improve survival from hemorrhagic shock in pigs. *Shock* 2011, **35**:67-73.
6. Mok YY, Atan MS, Yoke Ping C, Zhong Jing W, Bhatia M, Moochhala S, Moore PK: Role of hydrogen sulphide in haemorrhagic shock in the rat: protective effect of inhibitors of hydrogen sulphide biosynthesis. *Br J Pharmacol* 2004,

- 143:881-889.
7. Mok YY, Moore PK: **Hydrogen sulphide is pro-inflammatory in haemorrhagic shock.** *Inflamm Res* 2008, **57**:512-518.
 8. Olson KR, Whitfield NL: **Hydrogen sulfide and oxygen sensing in the cardiovascular system.** *Antioxid Redox Signal* 2010, **12**:1219-1234.
 9. Olson KR: **The therapeutic potential of hydrogen sulfide: separating hype from hope.** *Am J Physiol Regul Integr Comp Physiol* 2011, **301**:R297-R312.
 10. Olson KR: **A practical look at the chemistry and biology of hydrogen sulfide.** *Antioxid Redox Signal* 2012, **17**:32-44.
 11. Li L, Whiteman M, Guan YY, Neo KL, Cheng Y, Lee SW, Zhao Y, Baskar R, Tan CH, Moore PK: **Characterization of a novel, water-soluble hydrogen sulfide-releasing molecule (GY4137). New insights into the biology of hydrogen sulfide.** *Circulation* 2008, **117**:2351-2360.
 12. Li L, Whiteman M, Moore PK: **Dexamethasone inhibits lipopolysaccharide-induced hydrogen sulphide biosynthesis in intact cells and in an animal model of endotoxic shock.** *J Cell Mol Med* 2009, **13**:2684-2692.
 13. Wintner EA, Deckwerth TL, Langston W, Bengtsson A, Leviten D, Hill P, Insko MA, Dumpit R, VandenEkart E, Toombs CF, Szabó C: **A monobromobimane-based assay to measure the pharmacokinetic profile of reactive sulphide species in blood.** *Br J Pharmacol* 2010, **60**:941-957.
 14. Tokuda K, Kida K, Marutani E, Crimi E, Bougaki M, Khatri A, Kimura H, Ichinose F: **Inhaled hydrogen sulfide prevents endotoxin-induced systemic inflammation and improves survival by altering sulfide metabolism in mice.** *Antioxid Redox Signal* 2012, **17**:11-21.
 15. Li L, Bhatia M, Zhu YZ, Zhu YC, Ramnath RD, Wang ZJ, Anuar FBM, Whiteman M, Salto-Tellez M, Moore PK: **Hydrogen sulfide is a novel mediator of lipopolysaccharide-induced inflammation in the mouse.** *FASEB J* 2005, **19**:1196-1198.
 16. Baumgart K, Wagner F, Gröger M, Weber S, Barth E, Vogt JA, Wachter U, Huber-Lang M, Knöferl MW, Albuszies G, Georgieff M, Asfar P, Szabó C, Calzia E, Radermacher P, Simkova S: **Cardiac and metabolic effects of hypothermia and inhaled H₂S in anesthetized and ventilated mice.** *Crit Care Med* 2010, **38**:588-595.
 17. Wagner F, Scheuerle A, Weber S, Stahl B, McCook O, Knöferl MW, Huber-Lang M, Thomas J, Asfar P, Szabó C, Möller P, Gebhard F, Georgieff M, Calzia E, Radermacher P, Wagner K: **Cardiopulmonary, histological and inflammatory effects of intravenous H₂S after blunt chest trauma-induced lung contusion in mice.** *J Trauma* 2011, **71**:1659-1667.
 18. Whitfield NL, Kreimier EL, Verdial FC, Skovgaard N, Olson KR: **Reappraisal of H₂S/sulfide concentration in vertebrate blood and its potential significance ischemic preconditioning and vascular signaling.** *Am J Physiol Regul Integr Comp Physiol* 2008, **294**:R1930-R1937.
 19. Furne J, Saeed A, Levitt MD: **Whole tissue hydrogen sulfide concentrations are orders of magnitude lower than presently accepted values.** *Am J Physiol Regul Integr Comp Physiol* 2008, **295**:R1479-R1485.
 20. Bouillaud F, Blachier F: **Mitochondria and sulfide: a very old story of poisoning, feeding, and signaling?** *Antioxid Redox Signal* 2011, **15**:379-391.
 21. Fu M, Zhang W, Wu L, Yang G, Li H, Wang R: **Hydrogen sulfide (H₂S) metabolism in mitochondria and its regulatory role in energy production.** *Proc Natl Acad Sci U S A* 2012, **109**:2943-2948.
 22. Wang R: **Physiological implications of hydrogen sulfide: a whiff exploration that blossomed.** *Physiol Rev* 2012, **92**:791-896.

doi:10.1186/cc11833

Cite this article as: Calzia E, et al.: Endogenous H₂S in hemorrhagic shock: innocent bystander or central player? *Critical Care* 2012, **16**:183.