

Commentary: Macular hole surgery – A tweak here and there

Macular hole surgery is one of the most hotly debated topics in retina circles, with many groups demonstrating innovative surgical approaches with promising results. The standard technique of vitrectomy with internal limiting membrane (ILM) peeling remains the gold standard for small to medium-sized macular holes.

When it comes to large macular holes things get a little tough. Large macular holes have been classified by Gass^[1] as those with a minimum linear diameter (MLD) of 400 μ or more. The Manchester eye study^[2] described excellent success rates for macular holes with size less than 600 μ with even the conventional ILM peeling techniques.

Michalewska *et al.*^[3] described the technique of inverted ILM peeling for large macular holes (>400 μ), where instead of completely removing the ILM, Part of it attached to the margins of the hole was left in place. This ILM remnant was then inverted upside down to cover the macular hole. They achieved 98% closure rates with this technique compared to 88% closure rates with conventional surgery. With this technique, Mahalingam and Sambhav^[4] successfully achieved 100% closure rate and improved visual acuity in macular holes with a mean MLD of 811.4 μ . Khodani *et al.*^[5] achieved type 1 closure in four out of five patients with a mean visual improvement of approximately three lines in patients with a macular hole base diameter ranging from 1280 to 1480 μ with this technique. Kuriyama *et al.*^[6] found this technique to be a good adjuvant in macular hole retinal detachments as well.

For the cases where ILM around the macular hole has already been peeled and for refractory macular hole cases, the technique of autologous ILM transplantation^[7] or ILM translocation^[8] where a free ILM graft is taken from peripheral

retina and transplanted at macular hole has shown encouraging anatomical and visual outcomes. A direct comparison between the techniques inverted ILM peel and ILM transplant by Park *et al.*^[9] found out that both the techniques were equally effective for closure of macular holes, but for recovery of photoreceptor layers and ultimately visual outcomes, the inverted ILM peel technique provided better results.

Another technique of macular hole hydrodissection is described where soft tip extrusion cannula is used to reflux balanced salt solution (BSS) into the macular holes, lysing adhesions and helping in closure. This technique was also found to be useful for anatomical and functional outcomes in persistent/refractory/large macular holes.^[10,11] Subretinal injection of BSS was also proven to be of value in achieving improved anatomical and functional outcomes in refractory macular holes.^[12]

Lens capsule and autologous blood have also been used to stuff recurrent/refractory macular holes.^[13,14]

Grewal and Mahmoud first described the neurosensory retinal graft technique in closing a refractory MH.^[15] Since then multicenter international studies have proven the efficacy and safety of this procedure in refractory macular holes.^[16,17]

Our experience of inverted ILM peel and neurosensory graft for large and refractory macular holes has provided encouraging results.

The technique of ILM tailoring described by the author in this article seems like a logical, thoughtful and a comparatively safe approach for treating large macular holes although its comparison with the inverted flap technique with adequate sample size would be of great help to understand the outcome differences/success rates of the two techniques.^[18]

Even though multiple techniques are available, in our opinion operating surgeons need to choose the technique that works best in their hands considering available facilities and patient factors for the best outcomes in their patients.

**Giriraj Vibhute, Shrinivas Joshi,
Guruprasad Ayachit, Apoorva Ayachit**

Department of Vitreoretina, M M Joshi Eye Institute,
Hubballi, Karnataka, India

Correspondence to: Dr. Apoorva Ayachit,
Department of Vitreoretina, M M Joshi Eye Institute,
Hubballi, Karnataka, India
E-mail: apoorva.ag@gmail.com

References

- Gass JD. Reappraisal of biomicroscopic classification of stages of development of a macular hole. *Am J Ophthalmol* 1995;119:752-9.
- Ch'ng SW, Patton N, Ahmed M, Ivanova T, Baumann C, Charles S, *et al.* The Manchester large macular hole study: Is it time to reclassify large macular holes? *Am J Ophthalmol* 2018;195:36-42.
- Michalewska Z, Michalewski J, Adelman RA, Nawrocki J. Inverted internal limiting membrane flap technique for large macular holes. *Ophthalmology* 2010;117:2018-25.
- Mahalingam P, Sambhav K. Surgical outcomes of inverted internal limiting membrane flap technique for large macular hole. *Indian J Ophthalmol* 2013;61:601-3.
- Khodani M, Bansal P, Narayanan R, Chhablani J. Inverted internal limiting membrane flap technique for very large macular hole. *Int J Ophthalmol* 2016;9:1230-2.
- Kuriyama S, Hayashi H, Jingami Y, Kuramoto N, Akita J, Matsumoto M. Efficacy of inverted internal limiting membrane flap technique for the treatment of macular hole in high myopia. *Am J Ophthalmol* 2013;156:125-31.e1.
- Morizane Y, Shiraga F, Kimura S, Hosokawa M, Shiode Y, Kawata T, *et al.* Autologous transplantation of the internal limiting membrane for refractory macular holes. *Am J Ophthalmol* 2014;157:861-9.e1.
- Pires J, Nadal J, Gomes NL. Internal limiting membrane translocation for refractory macular holes. *Br J Ophthalmol* 2017;101:377-82.
- Park JH, Lee SM, Park SW, Lee JE, Byon IS. Comparative analysis of large macular hole surgery using an internal limiting membrane insertion versus inverted flap technique. *Br J Ophthalmol* 2019;103:245-50.
- Felfeli T, Mandelcorn ED. MACULAR HOLE HYDRODISSECTION: Surgical technique for the treatment of persistent, chronic, and large macular holes. *Retina* 2019;39:743-52.
- Khan TH, Rizvi SF, Mahmood SA, Feroz L. Treatment of chronic large and persistent macular hole by a new technique in a tertiary care hospital. *Pak J Med Sci* 2021;37:979-82.
- Gilani F, Sziagiato A-A, Walsh M, Muni RH. Subretinal BSS injection: A technique for the treatment of chronic persistent macular holes. *Invest Ophthalmol Vis Sci* 2015;56:5076.
- Chen S-N, Yang C-M. Lens capsular flap transplantation in the management of refractory macular hole from multiple etiologies. *Retina Phila Pa* 2016;36:163-70.
- Peng J, Chen C, Jin H, Zhang H, Zhao P. Autologous lens capsular flap transplantation combined with autologous blood application in the management of refractory macular hole. *Retina Phila Pa* 2018;38:2177-83.
- Grewal DS, Mahmoud TH. Autologous neurosensory retinal free flap for closure of refractory myopic macular holes. *JAMA Ophthalmol* 2016;134:229-30.
- Grewal DS, Charles S, Parolini B, Kadonosono K, Mahmoud TH. Autologous retinal transplant for refractory macular holes: Multicenter international collaborative study group. *Ophthalmology* 2019;126:1399-408.
- Robles-Holmes HK, Staropoli PC, Yannuzzi N, Sridhar J. Management of large or recurrent macular holes. *Curr Ophthalmol Rep* 2020;8:62-8.
- Shukla D, Kalliath J. Internal limiting membrane peeling for large macular hole: Tailoring the rhexis to the shape of the hole. *Indian J Ophthalmol* 2022;70:182-6..

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.ijo.in
	DOI: 10.4103/ijo.IJO_2368_21

Cite this article as: Vibhute G, Joshi S, Ayachit G, Ayachit A. Commentary: Macular hole surgery – A tweak here and there. *Indian J Ophthalmol* 2022;70:186-7.