

## Article

# $\alpha$ -Amyrin and $\beta$ -Amyrin Isolated from *Celastrus hindsii* Leaves and Their Antioxidant, Anti-Xanthine Oxidase, and Anti-Tyrosinase Potentials

Tran Duc Viet, Tran Dang Xuan \*  and La Hoang Anh 

Graduate School for International Development and Cooperation, Hiroshima University, Higashi-Hiroshima 739-8529, Japan; viettran1609@gmail.com (T.D.V.); hoanganh6920@gmail.com (L.H.A.)

\* Correspondence: tdxuan@hiroshima-u.ac.jp; Tel.: +81-82-424-6927

**Abstract:** *Celastrus hindsii* is a popular medicinal plant in Vietnam and Southeast Asian countries as well as in South America. In this study, an amount of 12.05 g of an  $\alpha$ -amyrin and  $\beta$ -amyrin mixture was isolated from *C. hindsii* (10.75 g/kg dry weight) by column chromatography applying different solvent systems to obtain maximum efficiency.  $\alpha$ -Amyrin and  $\beta$ -amyrin were then confirmed by gas chromatography-mass spectrometry (GC-MS), electrospray ionization-mass spectrometry (ESI-MS), and nuclear magnetic resonance (NMR). The antioxidant activities of the  $\alpha$ -amyrin and  $\beta$ -amyrin mixture were determined via 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,20-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assays with  $IC_{50}$  of 125.55 and 155.28  $\mu\text{g}/\text{mL}$ , respectively. The mixture exhibited a high potential for preventing gout by inhibiting a relevant key enzyme, xanthine oxidase (XO) ( $IC_{50} = 258.22 \mu\text{g}/\text{mL}$ ). Additionally, an important enzyme in skin hyperpigmentation, tyrosinase, was suppressed by the  $\alpha$ -amyrin and  $\beta$ -amyrin mixture ( $IC_{50} = 178.85 \mu\text{g}/\text{mL}$ ). This study showed that *C. hindsii* is an abundant source for the isolation of  $\alpha$ -amyrin and  $\beta$ -amyrin. Furthermore, this was the first study indicating that  $\alpha$ -amyrin and  $\beta$ -amyrin mixture are promising in future therapies for gout and skin hyperpigmentation.

**Keywords:**  $\alpha$ -amyrin;  $\beta$ -amyrin; GC-MS; ESI-MS; NMR; chromatography; antioxidant; anti-xanthine oxidase; anti-tyrosinase



**Citation:** Viet, T.D.; Xuan, T.D.; Anh, L.H.  $\alpha$ -Amyrin and  $\beta$ -Amyrin Isolated from *Celastrus hindsii* Leaves and Their Antioxidant, Anti-Xanthine Oxidase, and Anti-Tyrosinase Potentials. *Molecules* **2021**, *26*, 7248. <https://doi.org/10.3390/molecules26237248>

Academic Editors: Maria Fernanda Taviano, Rosaria Acquaviva and Giuseppe Antonio Malfa

Received: 8 November 2021  
Accepted: 23 November 2021  
Published: 29 November 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

*Celastrus hindsii* Benth. belongs to the Celastraceae family, which is found in many countries such as Laos, Thailand, Vietnam, and South America [1]. In Vietnam, since ancient times, *C. hindsii* has been used as a traditional medicine to treat numerous diseases such as cancer, ulcers, and inflammation [2]. Previous studies have shown that *C. hindsii* contains many precious compounds such as rosmarinic acid [1], terpenoids [3], alkaloids [4], phenolics, and flavonoids [2]. However, the use of this plant as folk medicine was mainly based on human experience over its long history. Its therapeutic mechanism has not been verified, and the major compounds and their actual contribution to the biological properties of *C. hindsii* have not been clearly investigated.

$\alpha$ -Amyrin and  $\beta$ -amyrin are widely known as a natural mixture of triterpenes. Their pharmacological potentials have been reported in previous studies [5], including antitumor [6], anti-inflammatory [7], anxiolytic [8], and hepatoprotective [9] effects.  $\alpha$ -Amyrin and  $\beta$ -amyrin can be found in numerous plant species such as *Solanum lycopersicum*, *Liriodendron tulipifera*, *Olea europaea*, *Aesculus hippocastanum*, *Aloe vera*, *Betula alba*, *Calendula officinalis*, *Coffea arabica*, *Malus domestica*, *Viscum album*, *Syzygium aromaticum*, *Pisum sativum*, *Brassica oleracea*, *Protium kleinii*, *Symplocos cochinchinensis*, *Alstonia boonei*, *Swertia longifolia*, *Canarium tramdenum*, *Byrsonima crassifolia*, etc. [10,11]. However, the isolation and accurate identification as well as quantification of these compounds in plant sources have been limited. The separation of these compounds for the purpose of treating diseases has not

been specifically proven. On the other hand, *C. hindsii* has been used as a medicinal plant for a long time [1–3]. The distribution area of this plant is also wide, contributing an abundant source of raw materials [2]. Additionally, we have found that  $\alpha$ -amyrin and  $\beta$ -amyrin are the dominant compounds in *C. hindsii*. Therefore, the isolation and purification of  $\alpha$ -amyrin and  $\beta$ -amyrin from *C. hindsii* are significantly needed. In the current research, we aimed to obtain the maximum amount of these two compounds from this species. In our study, column chromatography supported by gas chromatography-mass spectrometry (GC-MS), electrospray ionization mass spectrometry (ESI-MS), and nuclear magnetic resonance (NMR) were used to isolate and purify  $\alpha$ -amyrin and  $\beta$ -amyrin from *C. hindsii*. In addition, the potentials of  $\alpha$ -amyrin and  $\beta$ -amyrin for antioxidant, antigout, and anti-skin hyperpigmentation were investigated for the first time via in vitro assays.

In humans, oxidative stress is an important physiological process involved in many chronic diseases such as diabetes and cancer. Some studies have indicated the existences and contributions of oxidative stress in several disorders via inflammation [12]. Oxidative stress can induce an inflammatory process and vice versa in chronic disorders [13]. Therefore, screening the antioxidant capacities of target compounds was the first important step of our tests for biological activities. Among chronic diseases, gout is a serious disorder caused by increasing uric acid in the blood. This disease seriously affects the populations of many countries in the world [14]. In terms of the underlying mechanism, xanthine oxidase (XO) plays an important role in the elevation of uric acid in the blood, leading to hyperuricemia, followed by the development of gout [15]. Until now, only allopurinol and febuxostat have been acknowledged for their role in inhibiting XO in the treatment of hyperuricemia. However, they also have many undesirable effects such as hypersensitivity syndrome, hepatitis nephropathy, eosinophilia, fever, vasculitis, and skin rash; therefore, further research into XO inhibition compounds is necessary [16,17]. Additionally, other potential compounds inhibiting the activity of xanthine oxidase could be promising source in the development of novel drugs for gout.

Another chronic syndrome, as skin hyperpigmentation, causes changes in skin color, loss of aesthetics, and an increase in the risk of skin cancer [18]. In the formation of skin hyperpigmentation, tyrosinase is a key regulatory enzyme in the proliferation of browning pigments. It plays an important role in controlling the amount of melanin, which is crucial to the health of the skin [19]. Thus, tyrosinase inhibitors are often used in the development of medicines and cosmetics. Furthermore, the synergic inhibition of oxidative stress, xanthine oxidase, and tyrosinase could be a potential approach to the generation of alternative drugs, which can prevent multiple human health problems [20].

For this purpose, many natural and semisynthetic inhibitors have been developed by various methods. The aforementioned rationales prompted us to conduct research focusing on the isolation of  $\alpha$ -amyrin and  $\beta$ -amyrin from *C. hindsii* and examine their antioxidant, anti-xanthine oxidase, and anti-tyrosinase activities. This study contributes to the development of natural-based products for preventing chronic disorders including gout and skin hyperpigmentation.


## 2. Results

### 2.1. Isolation of $\alpha$ -Amyrin and $\beta$ -Amyrin from Leaves of *C. hindsii*

A dried powder of *C. hindsii* (1.12 kg) was extracted by methanol to collect crude extract (260.80 g). Then, it was partitioned with three solvents, including water, hexane, and ethyl acetate. Next, a preliminary activity assessment and thin-layer chromatography were conducted. The results showed that the ethyl acetate mixture had the most potential, and it was used for column chromatography separation. After checking by thin-layer chromatography, the ethyl acetate extract was the most active for the target compounds of  $\alpha$ -amyrin and  $\beta$ -amyrin. This extract was then used to run column chromatography. The results showed that nine fractions (C1–C9) contained the  $\alpha$ -amyrin and  $\beta$ -amyrin. These fractions were combined to run a second-time column chromatography. Finally, 12.05 g of an  $\alpha$ - and  $\beta$ -amyrin mixture was obtained.

## 2.2. Identification of $\alpha$ -Amyrin and $\beta$ -Amyrin by GC-MS and ESI-MS

Based on the GC-MS results (Figure 1), the presence of  $\alpha$ -amyrin and  $\beta$ -amyrin was confirmed. The retention times of  $\alpha$ -amyrin and  $\beta$ -amyrin were 29.53 and 28.92 min, respectively (Table 1).



**Figure 1.** GC-MS chromatogram (a) and mass spectra (b) of  $\alpha$ -amyrin and  $\beta$ -amyrin.

**Table 1.**  $\alpha$ -Amyrin and  $\beta$ -amyrin identified by GC-MS.

Time (min)	Peak Area (% of Total)	Compounds	Formula	Molecular Weight (g/mol)	Chemical Class
28.92	40.61	$\beta$ -Amyrin	$C_{30}H_{50}O$	426.70	Triterpene
29.53	59.10	$\alpha$ -Amyrin	$C_{30}H_{50}O$	426.70	Triterpene

The results of ESI-MS showed the molecular weight and formula of  $\alpha$ -amyrin and  $\beta$ -amyrin, which were verified as dimensionless at  $449.33 + 1 m/z$  and formula  $[C_{30}H_{50}O + Na]^+$  (Figure 2). These results referred to the studies of Banerjee et al. [21] and Pinto et al. [22] to support the results of the GC-MS in which two compounds were isolated:  $\alpha$ -amyrin and  $\beta$ -amyrin.

## 2.3. Elucidation of $\alpha$ -Amyrin and $\beta$ -Amyrin by NMR

Figure 3 shows the  $^{13}C$  NMR spectrum of the mixture of  $\alpha$ -amyrin and  $\beta$ -amyrin. The signals at 139.60 ppm and 145.20 corresponded to  $C_{13}$ , whereas the signals at 124.40 and 121.70 ppm corresponded to  $C_{12}$  of  $\alpha$ -amyrin and  $\beta$ -amyrin, respectively. The other signals were assigned as shown in (Table 2), which are in agreement with the published literature [23,24].

The  $^1H$  NMR spectrum (Figure 4) illustrates that the two signals at 5.18 ppm (H-12, 1H) and 5.13 ppm (H-12, 1H) were assigned to  $\beta$ -amyrin and  $\alpha$ -amyrin, respectively. The ratio of  $\alpha$ -amyrin and  $\beta$ -amyrin was (2.29:1), which was calculated based on the integration value of these two signals [23,24].

Based on the chemical structures of  $\alpha$ -amyrin and  $\beta$ -amyrin in Figure 5, these two compounds share the same molecular weight and chemical formula. The difference only comes from the methyl group ( $CH_3$ ) at  $C_{30}$  and  $C_{29}$  carbon positions. In  $\alpha$ -amyrin structure, the methyl groups of  $C_{29}$  and  $C_{30}$  are attached at two separate positions. One is directed forward and one to back. While in  $\beta$ -amyrin, the methyl groups of  $C_{29}$  and  $C_{30}$  are attached to the same position and are directed forward. This is also the basis for distinguishing the two substances as well as identifying these two compounds.

#### 2.4. Quantitative Analysis of $\alpha$ -Amyrin and $\beta$ -Amyrin in *C. hindsii* Leaves

A total of 12.05 g of  $\alpha$ -amyrin and  $\beta$ -amyrin mixture was isolated from 1.12 kg of *C. hindsii* dried leaves. The ratio of  $\alpha$ -amyrin and  $\beta$ -amyrin in the mixture was determined as (2.29:1) based on the NMR results (Figure 4). The content of  $\alpha$ -amyrin and  $\beta$ -amyrin mixture in leaves of *C. hindsii* was quantified by the method of Shantos et al. [24] as 10.75 g/kg dry weight (DW), and the ratio of  $\alpha$ -amyrin and  $\beta$ -amyrin in the mixture was 69.60% and 30.40%, respectively, equivalent to 7.48 g/kg DW of  $\alpha$ -amyrin and 3.27 g/kg DW of  $\beta$ -amyrin (Table 3).




Figure 2. Identification of  $\alpha$ -amyrin and  $\beta$ -amyrin by ESI-MS.






Figure 3. <sup>13</sup>C-NMR spectrum of the mixture of  $\alpha$ -amyrin and  $\beta$ -amyrin measured in CDCl<sub>3</sub> (125 MHz).

**Table 2.**  $^{13}\text{C}$ -NMR data for  $\alpha$ -amyrin and  $\beta$ -amyrin in  $\text{CDCl}_3$ .

Position	$\beta$ -Amyrin	$\alpha$ -Amyrin	Position	$\beta$ -Amyrin	$\alpha$ -Amyrin
1	38.60	38.80	16	26.20	26.60
2	27.20	27.30	17	32.70	33.80
3	79.00	79.10	18	47.20	59.10
4	39.80	38.80	19	46.80	39.70
5	55.20	55.20	20	31.10	39.60
6	18.40	18.40	21	34.70	31.30
7	32.50	32.90	22	37.10	40.00
8	41.70	40.00	23	28.10	28.10
9	47.60	47.70	24	15.60	15.70
10	37.00	36.90	25	15.50	15.60
11	23.70	23.30	26	16.80	17.40
12	121.70	124.40	27	26.00	23.40
13	145.20	139.60	28	28.40	28.80
14	42.80	41.50	29	33.30	16.90
15	26.90	28.10	30	23.50	21.40

**Figure 4.**  $^1\text{H}$ -NMR spectrum of the mixture of  $\alpha$ -amyrin and  $\beta$ -amyrin measured in  $\text{CDCl}_3$  (400 MHz).**Figure 5.** Chemical structures of  $\alpha$ -amyrin and  $\beta$ -amyrin identified in *C. hindsii*.

**Table 3.** Quantification of  $\alpha$ -amyrin and  $\beta$ -amyrin in *C. hindsii* leaves.

Content of Mixture (g/kg Dry Weight)	Ratio (%)		Contents (g/kg Dry Weight)	
	$\alpha$ -Amyrin	$\beta$ -Amyrin	$\alpha$ -Amyrin	$\beta$ -Amyrin
10.75	69.60	30.40	7.48	3.27

### 2.5. Antioxidant, Anti-Xanthine Oxidase, and Anti-Tyrosinase of $\alpha$ -Amyrin and $\beta$ -Amyrin from *C. hindsii*

The antioxidant activities of  $\alpha$ -amyrin and  $\beta$ -amyrin were determined via DPPH and ABTS free-radical scavenging assays. BHT was used as a standard inhibitor. The results were summarized in Table 4. The IC<sub>50</sub> values of  $\alpha$ -amyrin and  $\beta$ -amyrin were 125.55 and 155.28  $\mu$ g/mL for DPPH and ABTS assays, respectively. For XO inhibitory potential,  $\alpha$ -amyrin and  $\beta$ -amyrin mixture exhibited strong activity, with an IC<sub>50</sub> value of 258.22  $\mu$ g/mL. For suppression of tyrosinase, the mixture had an IC<sub>50</sub> value of 178.85  $\mu$ g/mL. Allopurinol and kojic acid were used as standard inhibitors for XO (IC<sub>50</sub> = 7.58  $\mu$ g/mL) and tyrosinase (IC<sub>50</sub> = 15.55  $\mu$ g/mL), respectively.

**Table 4.** Antioxidant, XO, and tyrosinase inhibitory capacities of  $\alpha$ -amyrin and  $\beta$ -amyrin isolated from *C. hindsii*.

Sample	Antioxidant Activity		XO IC <sub>50</sub> ( $\mu$ g/mL)	Tyrosinase IC <sub>50</sub> ( $\mu$ g/mL)
	DPPH IC <sub>50</sub> ( $\mu$ g/mL)	ABTS IC <sub>50</sub> ( $\mu$ g/mL)		
$\alpha$ -Amyrin and $\beta$ -Amyrin	125.55 $\pm$ 0.98 <sup>a</sup>	155.28 $\pm$ 1.01 <sup>a</sup>	258.22 $\pm$ 2.28 <sup>a</sup>	178.85 $\pm$ 3.28 <sup>a</sup>
Allopurinol	-	-	7.58 $\pm$ 1.29 <sup>b</sup>	-
Kojic acid	-	-	-	15.55 $\pm$ 2.28 <sup>b</sup>
BHT	8.22 $\pm$ 0.89 <sup>b</sup>	53.40 $\pm$ 1.52 <sup>b</sup>	-	-

Values represent means  $\pm$  SD (standard deviation). Values with similar superscript letter in each column are not significantly different ( $p < 0.05$ ) ( $n = 3$ ). -: measurement was not conducted; BHT: butylated hydroxytoluene; DPPH: 2,2-diphenyl-1-picrylhydrazyl; ABTS: 2,20-azinobis (3-ethylbenzothiazoline-6-sulfonic acid); and XO: xanthine oxidase.

### 3. Discussion

*Celastrus hindsii* has shown numerous biological activities in previous research including preventing inflammation, antitumor effects, inhibiting HIV replication, etc. [25,26]. *C. hindsii* has been reported to contain many valuable chemical components, including sesquiterpenes, triterpenes, alkaloids, and flavonoids [27,28]. *C. hindsii* is now attracting greater international research attention [29]. In this study, *C. hindsii* was used to isolate and purify  $\alpha$ -amyrin and  $\beta$ -amyrin. These compounds are classified as triterpenes for which anti-obesity [30], anxiolytic [31], antidepressant and antiplatelet [32], antinociceptive [33], liver-protective [34], gastroprotective [35], anti-inflammatory [36], antiulcer, antihyperlipidemic, and antitumor [37] properties have been reported. However, the potential for these compounds to inhibit gout and skin hyperpigmentation was investigated for the first time in this study.

For the isolation and purification of  $\alpha$ -amyrin and  $\beta$ -amyrin, column chromatography was used in this study with different extracting solvents. This is a popular technique, which is widely used to isolate individual substances of interest in mixtures from large to small sizes and can be applied both at the laboratory scale and in industrial production [38]. To support the isolation and purification, GC-MS, ESI-MS, and NMR were performed. Gas chromatography-mass spectrometry (GC-MS) is an analytical method that involves the combination of gas chromatography and mass spectrometry to identify different components within a test sample [39]. In addition, this technology allows the identification of trace elements, even in tiny amounts, and composition of compounds present in an entity

after it has been decomposed [40]. This technology allows both qualitative and quantitative conversions to the analytical mode in the form of ions [41]. Electrospray ionization mass spectrometry (ESI-MS) was developed as a significant technique to provide the sensitive, robust, and reliable tool for research, conducted at atmospheric pressure and at a moderate temperature, using non-volatile and thermally labile biomolecules that are not amenable to analysis by other conventional techniques [42]. Nuclear magnetic resonance (NMR) is a technology used to determine the structure of molecules, both crystalline and noncrystalline based on results from the specific properties of certain atomic nuclei of hydrogen and carbon.  $^1\text{H}$  spin 1/2 occurs in many biological systems. The NMR proton system produces the displacement of hydrogen in a narrow range with a sharp, signal thereby giving fast quantum results in a short time [43]. The  $^{13}\text{C}$  spin 1/2 method has been used widely in NMR conduction. Although it occurs at a small proportion in nature (approximately 1.1%), it is associated with nuclear fission. Therefore, the sensitivity is low, the output signal is sharp, the effects of spin are prevented, and the mass spectrometry is not disturbed [44]. As a result, a total of 12.05 g of  $\alpha$ -amyrin and  $\beta$ -amyrin were isolated and purified from *C. hindsii* leaves and were able to be confirmed based on the GC-MS, ESI-MS, and NMR results.

In other studies,  $\alpha$ -amyrin and  $\beta$ -amyrin have been found in peas (*Pisum sativum*) [45] and cabbages (*Brassica oleracea*) [46]. Especially, an  $\alpha$ -amyrin and  $\beta$ -amyrin mixture (2.40 g/kg DW) was isolated from *Protium kleinii* [32].  $\beta$ -Amyrin was determined in *Symplocos cochinchinensis* [47], *Alstonia boonei*, and *Swertia longifolia* at the amounts of 1.70; 8.00; and 2.00 g/kg DW, respectively [48]. In the case of  $\alpha$ -amyrin, the compound was quantified in *Melastoma malabathricum* [49] and *Swertia longifolia* [50], in which the contents were 0.60 and 1.00 g/kg, respectively. In *Canarium tramdenum* and *Byrsonima crassifolia*,  $\alpha$ -amyrin and  $\beta$ -amyrin mixtures were determined at 1.52 and 9.00 g/kg DW [51,52].  $\alpha$ -Amyrin was studied in plant cell cultures of *Centella asiatica* [53]. In this study, a total of 12.05 g of  $\alpha$ -amyrin and  $\beta$ -amyrin mixture was obtained from 1.12 kg of *C. hindsii* dried leaves, equivalent to 10.75 g/kg DW (Table 3). The contents of  $\alpha$ -amyrin and  $\beta$ -amyrin were determined based on NMR results (Table 2, Figures 3 and 4).  $\alpha$ -Amyrin and  $\beta$ -amyrin accounted for 7.48 and 3.27 g/kg DW, respectively (Table 3), which are significantly higher than those found in other plant species. This demonstrated that *C. hindsii* is a potential source to isolate  $\alpha$ -amyrin and  $\beta$ -amyrin. A further step in our study was to evaluate the antioxidant, anti-xanthine oxidase, and anti-tyrosinase properties of  $\alpha$ -amyrin and  $\beta$ -amyrin, which is discussed as follows.

Oxidative stress is an imbalance between reactive oxygen/nitrogen species (ROS/RNS), leading to the loss of antioxidant capacity. Oxidative stress has been linked to many diseases such as aging, cardiovascular disease, and cancer [54]. Free-radical-induced damage in oxidative stress has been confirmed as a contributor to the pathogenesis of many chronic health problems such as neurodegenerative, emphysema, cardiovascular and inflammatory diseases, cataracts, and cancer [55]. Oxidative stress is implicated in many diseases [56]. An irreversible progression of oxidative decay caused by reactive oxygen species also exerts its negative influence on the biology of aging, leading to the impairment of physiological functions, promoting disease incidence, and reducing life span [57].  $\alpha$ -Amyrin and  $\beta$ -amyrin isolated from *Myrcianthes pungens* leaves were reported as having antioxidant potential [58]. The antioxidant agent of  $\alpha$ -amyrin and  $\beta$ -amyrin has been shown to combat acute pancreatitis [59]. Oxidative stress is closely related to inflammatory diseases [12,13]. Many studies have shown the strong anti-inflammatory ability of  $\alpha$  and  $\beta$ -amyryns [5,45]. Thus, this assessment of  $\alpha$ - and  $\beta$ -amyrin's antioxidant abilities was consistent with previous findings. Both *C. hindsii*  $\alpha$  amyryn and  $\beta$ -amyryn showed antioxidant capacity [2,51]. The results of this evaluation of  $\alpha$ - and  $\beta$ -amyryn's antioxidant potential by two methods, DPPH and ABTS (Table 4), are important additional evidence of the value of  $\alpha$  and  $\beta$ -amyryn and *C. hindsii* in term of antioxidant capacity.

Gout is a chronic disease caused by the deposition of monosodium urate crystals [60]. The changes of urate transport in the kidneys and gut play an important role in elevation

of uric acid in the blood, which is the cause of gout [61]. Xanthine oxidase is an enzyme that converts hypoxanthine into uric acid, and the strong activity of this enzyme leads to increased uric acid concentration, causing the deposition of monosodium urate crystals that cause gout [62]. This syndrome is closely related to inflammatory reactions; therefore, gout often manifests as monoarthritic of the extremities, so anti-inflammatory agents are widely used to treat gout [63,64]. This was the basis for the evaluation of the anti-gout ability of  $\alpha$  and  $\beta$ -amyrin as they are anti-inflammatory compounds [65]. The results of  $\alpha$ - and  $\beta$ -amyrin's anti-xanthine oxidase potential were shown in (Table 4). The value of this mixture was confirmed. In addition, this evaluation further emphasized the utility of these compounds in the treatment of gout.

Tyrosinase is an enzyme that catalyzes the synthesis of melanin [5]. Tyrosinase has been studied extensively in the treatment of malignant cancers [66]. Anti-tyrosinase has been studied in antibodies in melanoma and vitiligo [67]. Tyrosinases are exploited for a variety of biotechnological and environmental applications. Their potential has been applied in food, pharmaceutical, and environmental sectors [68]. Tyrosinase is involved in neurodegenerative disorders, such as Parkinson's disease, and in melanin-browning reactions important to the cosmetics and food industries [69]. Tyrosinase attracted the interest of immunologists because it was known that melanoma patients could form an immune response to antigens related to melanogenesis [70]. This is the first study of  $\alpha$ -amyrin and  $\beta$ -amyrin to consider anti-tyrosinase potential, and our results support the development of  $\alpha$ -amyrin and  $\beta$ -amyrin as therapeutics for skin hyperpigmentation.

## 4. Materials and Methods

### 4.1. Materials

The plant materials of *C. hindsii* were collected in Cao Duong commune, Luong Son district, Hoa Binh province, Vietnam. Fifty mature plants were randomly selected for leaf harvest. The collected *C. hindsii* plants were in the age range of over 5 years. They have grown naturally. *C. hindsii* plants were healthy and free from pests and diseases, without the use of fertilizers, pesticides, and stimulants. The materials were authenticated and sterilized. Specimens (No. PPBC170506) were preserved before further experiments were carried out. The study was conducted from March to December in 2020.

### 4.2. Preparation of Plant Extract

A total of 1.12 kg of *C. hindsii* dried leaves was used for the experiment. The material was redried for 2 days at 40 °C. After that, the obtained sample was soaked in 10 L of MeOH for 30 days at room temperature. It was then filtered and concentrated by a rotary evaporator (SB-350-EYELA, Tokyo Rikakikai Co., Ltd., Tokyo, Japan) to obtain 260.80 g of crude extract. This crude extract was separated by three solvents (water, hexane, and ethyl acetate).

### 4.3. Fractionation of Ethyl Acetate Extract

The ethyl acetate extract showed the strongest inhibition by thin-layer chromatography test, was subjected to a normal phase of column chromatography (20 mm diameter  $\times$  500 mm height, Climbing G2, Tokyo, Japan) over silica gel (size A 60, 200–400 mesh particle size) (Sigma Aldrich, Tokyo, Japan). This process yielded 9 fractions with following eluents: C1 in CHCl<sub>3</sub>, C2 in CHCl<sub>3</sub>:MeOH (99:1) flask 1–20, C3 in CHCl<sub>3</sub>:MeOH (99:1) flask 21–40, C4 in CHCl<sub>3</sub>:MeOH (99:1) flask 41–60, C5 in CHCl<sub>3</sub>:MeOH (99:1) flask 61–80, C6 in CHCl<sub>3</sub>:MeOH (99:1) flask 81–100, C7 in CHCl<sub>3</sub>:MeOH (97:3), C8 in CHCl<sub>3</sub>:MeOH (95:5), and C9 in CHCl<sub>3</sub>:MeOH (9:1). Then, these fractions were combined to run column chromatography for the second time.

### 4.4. Identification $\alpha$ -Amyrin and $\beta$ -Amyrin by Gas Chromatography-Mass Spectrometry (GC-MS)

The composition of substances in *C. hindsii* was determined by the GC-MS system. An aliquot of 1  $\mu$ L sample dissolved in chloroform was added to the GC-MS system (JMS-T100



GCV, JEOL Ltd., Tokyo, Japan). The column was DB-5MS, 30 mm length, 0.25 mm internal diameter, and 0.25  $\mu\text{m}$  in thickness (Agilent Technologies, JW Scientific Products, Folsom, CA, USA). Helium was used as carrier gas, and ratio was 5:1. The method to operate the GC oven temperatures was conducted, the temperature was 50  $^{\circ}\text{C}$ , and program rate was 10  $^{\circ}\text{C}$  up to 300  $^{\circ}\text{C}$  with holding time of 20 min. The injector and detector temperatures were 300–320  $^{\circ}\text{C}$ , respectively. The mass range scanned from 28 to 800 amu. The control of GC-MS system and data processing were identified following JEOL's GC-MS Mass Center System Version 2.55a (JEOL Ltd., Tokyo, Japan). The peak areas of compounds (>3%) analysis were identified as principal substances [71].

#### 4.5. Electrospray Ionization-Mass Spectrometry (ESI-MS) Analysis

ESI-MS analysis was carried out in negative and positive ion mode. The capillary temperature was conducted at 140  $^{\circ}\text{C}$  (12  $^{\circ}\text{C}$  for S2), and the spray voltage was 3.0 KV (2.7 KV for S2). The measurements were calculated in the positive mode with an ion spray voltage of 3000 V and capillary temperature of 350  $^{\circ}\text{C}$ . Data peaks were estimated within the range of 280–1000  $m/z$  [21].

#### 4.6. Nuclear Magnetic Resonance (NMR) Data of $\alpha$ -Amyrin and $\beta$ -Amyrin

$\alpha$ -Amyrin:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ): 3.24 (dd,  $J = 10.9$  and  $5.1$  Hz, H-3), 5.12 (t,  $J = 3.6$  Hz, H-12).  $\delta$   $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ): 38.8 (C-1), 27.3 (C-2), 79.1 (C-3), 38.8 (C-4), 55.2 (C-5), 18.4 (C-6), 32.9 (C-7), 40.0 (C-8), 47.7 (C-9), 36.9 (C-10), 23.3 (C-11), 124.4 (C-12), 139.6 (C-13), 41.5 (C-14), 28.1 (C-15), 26.6 (C-16), 33.8 (C-17), 59.1 (C-18), 39.7 (C-19), 39.6 (C-20), 31.3 (C-21), 40.0 (C-22), 28.1 (C-23), 15.7 (C-24), 15.6 (C-25), 17.4 (C-26), 23.4 (C-27), 28.8 (C-28), 16.9 (C-29), 21.4 (C-30) [23,24].

$\beta$ -amyrin:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ): 3.24 (dd,  $J = 10.9$  and  $5.1$  Hz, H-3), 5.18 (t,  $J = 3.6$  Hz, H-12).  $\delta$   $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ): 38.6 (C-1), 27.2 (C-2), 79.0 (C-3), 39.8 (C-4), 55.2 (C-5), 18.4 (C-6), 32.5 (C-7), 41.7 (C-8), 47.6 (C-9), 37.0 (C-10), 23.7 (C-11), 121.7 (C-12), 145.2 (C-13), 42.1 (C-14), 26.9 (C-15), 26.2 (C-16), 32.7 (C-17), 47.2 (C-18), 46.8 (C-19), 31.1 (C-20), 34.7 (C-21), 37.1 (C-22), 28.1 (C-23), 15.6 (C-24), 15.5 (C-25), 16.8 (C-26), 26.0 (C-27), 28.4 (C-28), 33.3 (C-29), 23.5 (C-30) [23,24].

#### 4.7. Xanthine Oxidase (XO) Inhibitory Activity

The XO activity was assessed spectrophotometrically, which was based on the research of Andriana et al. [72] with some modifications. In brief, a mixture of 20  $\mu\text{L}$  of sample and 40  $\mu\text{L}$  of 70 mM phosphate buffer (pH = 7.5) and 40  $\mu\text{L}$  of enzyme solution (0.01 units/mL in buffer) was incubated for 8 min at room temperature. After that, 60  $\mu\text{L}$  of xanthine substrate (300  $\mu\text{M}$  in buffer) was injected. The mixture was incubated at room temperature. The reaction was stopped by adding 20  $\mu\text{L}$  of HCl 1M. The result was read at 290 nm by a microplate reader (Multiskan<sup>TM</sup> microplate spectrophotometer, Fisher Scientific, Osaka, Japan). XO inhibitory activity was used according to formula below:

$$\% \text{ Inhibition} = \left\{ \frac{(E - C) - (T - W)}{E - C} \right\} \times 100 \quad (1)$$

where E is the activity of the enzyme without test sample. C is the control of experiment without test sample.

T is the activity of test sample. W is the blank (without test). Allopurinol (10–100  $\mu\text{g}/\text{mL}$ ) was used as a positive control.  $\text{IC}_{50}$  values were calculated from the mean values of the percentage inhibition data.

#### 4.8. 2,2-Diphenyl-1-picrylhydrazyl (DPPH) Free Radical Scavenging Activity

The antioxidant activity of the  $\alpha$ -amyrin and  $\beta$ -amyrin mixture was estimated via DPPH free radical scavenging method described by Quy et al. [73]. In details, a mixture containing 50  $\mu\text{L}$  of  $\alpha$ -amyrin and  $\beta$ -amyrin with 0.25  $\mu\text{L}$  of 0.5 mM DPPH solution and 0.5  $\mu\text{L}$  of 0.1 M acetate buffer (pH 5.5) was prepared and placed in the dark for 30 min

under ambient conditions. The absorbance of the reaction was recorded at 517 nm using a microplate reader (Multiskan™ Microplate Spectrophotometer, Thermo Fisher Scientific, Osaka, Japan). Methanol and BHT (10–50 ppm) were used as negative and positive controls, respectively. The antioxidant capacity of the tested samples was calculated using the following equation:

$$\text{DPPH radical scavenging activity (\%)} = [(C - S)/C]/100 \quad (2)$$

where S is the absorbance of sample. C is the absorbance of the reaction without the sample. The result was expressed as IC<sub>50</sub> value, which determined the concentration of the sample required to scavenge 50% of DPPH.

#### 4.9. 2,20-Azinobis (3-Ethylbenzothiazoline-6-sulfonic Acid) (ABTS) Cation Scavenging Activity

The ABTS method was used to evaluate the antioxidant capacity of *C. hindsii* [73]. The ABTS solution was created by adding 7 mM ABTS to 2.45 mM potassium persulfate solution. After that, the mixture was incubated in the dark at room temperature for 16 h.

MeOH was pushed to achieve an absorbance of  $0.68 \pm 0.05$  at 734 nm. A volume of 24  $\mu\text{L}$  of each sample was added to 120  $\mu\text{L}$  of ABTS solution, and then the mixture was incubated in the dark under ambient condition for 30 min. The absorbance was measured at 734 nm by a microplate reader. BHT (5–125  $\mu\text{g}/\text{mL}$ ) was considered as a standard inhibitor, and MeOH was a negative control. The ABTS radical scavenging activity was evaluated with the same method as presented for DPPH assay.

#### 4.10. Tyrosinase Inhibition

The prepared components included 20 mM potassium phosphate buffer (pH = 6.8) and tyrosinase solution (500 units/mL in buffer). The substrate solution was L-tyrosine (2 mM in distilled water). The sample was dissolved in DMSO. For the reaction, the combination of 20  $\mu\text{L}$  of test sample, 20  $\mu\text{L}$  of enzyme, and 120  $\mu\text{L}$  of buffer was incubated for 5 min at 25 °C. The final solution's absorbance was calculated at 470 nm. Kojic acid was used as a standard inhibitor, and DMSO was used as a negative control. The inhibition potential was evaluated by the following formula:

$$\% \text{ Inhibition} = (C - S)/C \times 100 \quad (3)$$

C: Absorbance of the reaction with DMSO after the substrate control's blank; S: Absorbance of the reaction with sample (without enzymatic reaction on substrate). IC<sub>50</sub> results of the sample against tyrosinase were calculated with the same method of antiradical assays [20].

#### 4.11. Statistical Analysis

The statistical analysis was utilized by using one-way ANOVA in the Minitab software ver. 16.2.3 (©2012 Minitab Inc.; Philadelphia, PA, USA). Tukey's test was used to evaluate the significant differences ( $p < 0.05$ ) among the test sample. The results were expressed as mean values and standard deviations.

## 5. Conclusions

This study observed that *C. hindsii* is an abundant source of  $\alpha$ -amyrin and  $\beta$ -amyrin. The use of column chromatography, GC-MS, ESI-MS, and NMR optimized the effectiveness of extraction and identification this mixture. Especially, the use of NMR for quantification was impressing to determine the high content of  $\alpha$ -amyrin and  $\beta$ -amyrin in *C. hindsii*. Moreover, the anti-xanthine oxidase and anti-tyrosinase potentials of  $\alpha$ -amyrin and  $\beta$ -amyrin were reported for the first time and could contribute to the development of novel drugs for preventing gout and skin hyperpigmentation.

**Author Contributions:** Conceptualization, T.D.V.; methodology, T.D.V.; formal analysis, T.D.V.; investigation, T.D.V. and L.H.A.; data curation, T.D.V.; writing—original draft preparation, review and editing, T.D.X. and L.H.A. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** All data are presented in the article.

**Acknowledgments:** The authors thank the assistance of Yusuf Andriana, Nguyen Van Quan, and Nguyen Hai Dang given to this manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Sample Availability:** Samples of the compounds are available from the authors. Samples was obtained through the process of conducting experiments.

## References

1. Ly, T.N.; Shimoyamada, M.; Yamauchi, R. Isolation and characterization of rosmarinic acid oligomers in *Celastrus hindsii* Benth leaves and their antioxidative activity. *J. Agric. Food Chem.* **2006**, *54*, 3786–3793. [\[CrossRef\]](#)
2. Viet, T.D.; Xuan, T.D.; Van, T.M.; Andriana, Y.; Rayee, R.; Tran, H.D. Comprehensive fractionation of antioxidants and GC-MS and ESI-MS fingerprints of *Celastrus hindsii* leaves. *Medicines* **2019**, *6*, 64. [\[CrossRef\]](#)
3. Kuo, Y.H.; Chen, C.F.; Kuo, L.M.Y.; King, M.L.; Chen, C.F.; Lee, K.H. Celahinine A, a new sesquiterpene pyridine alkaloid from *Celastrus hindsii*. *J. Nat. Prod.* **1995**, *58*, 1735–1738. [\[CrossRef\]](#)
4. Pham, D.C.; Nguyen, H.C.; Nguyen, T.H.L.; Ho, H.L.; Trinh, T.K.; Riyaphan, J.; Weng, C.F. Optimization of ultrasound-assisted extraction of flavonoids from *Celastrus hindsii* leaves using response surface methodology and evaluation of their antioxidant and antitumor activities. *BioMed Res. Int.* **2020**, *2020*, 3497107. [\[CrossRef\]](#)
5. Da Silva Júnior, W.F.; Bezerra de Menezes, D.L.; De Oliveira, L.C.; Koester, L.S.; Oliveira de Almeida, P.D.; Lima, E.S.; Neves de Lima, Á.A. Inclusion complexes of  $\beta$  and HP $\beta$ -cyclodextrin with  $\alpha$ ,  $\beta$  amyryn and in vitro anti-inflammatory activity. *Biomolecules* **2019**, *9*, 241. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Barros, F.W.A.; Bandeira, P.N.; Lima, D.J.B.; Meira, A.S.; de Farias, S.S.; Albuquerque, M.R.J.R.; dos Santos, H.S.; Lemos, T.L.G.; de Moraes, M.O.; Costa-Lotufu, L.V. Amyryn esters induce cell death by apoptosis in HL-60 leukemia cells. *Bioorg. Med. Chem.* **2011**, *19*, 1268–1276. [\[CrossRef\]](#)
7. Lima, E.M.; Nascimento, A.M.; Lenz, D.; Scherer, R.; Meyrelles, S.S.; Boëchat, G.A.P.; Andrade, T.U.; Endringer, D.C. Triterpenes from the *Protium heptaphyllum* resin-chemical composition and cytotoxicity. *Rev. Bras. Farmacogn.* **2014**, *24*, 399–407. [\[CrossRef\]](#)
8. Aragão, G.F.; Carneiro, L.M.V.; Junior, A.P.F.; Vieira, L.C.; Bandeira, P.N.; Lemos, T.L.G.; Viana, G.S.D.B. A possible mechanism for anxiolytic and antidepressant effects of alpha- and beta-amyryn from *Protium heptaphyllum* (Aubl.) March. *Pharmacol. Biochem. Behav.* **2006**, *85*, 827–834. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Oliveira, F.A.; Vieira-Júnior, G.M.; Chaves, M.H.; Almeida, F.R.C.; Florêncio, M.G.; Lima, R.C.P.; Silva, R.M.; Santos, F.A.; Rao, V.S.N. Gastroprotective and anti-inflammatory effects of resin from *Protium paraphyllum* in mice and rats. *Pharmacol. Res.* **2004**, *49*, 105–111. [\[CrossRef\]](#)
10. Yasumoto, S.; Seki, H.; Shimizu, Y.; Fukushima, E.O.; Muranaka, T. Functional characterization of CYP716 family P450 enzymes in triterpenoid biosynthesis in tomato. *Front. Plant Sci.* **2017**, *8*, 21. [\[CrossRef\]](#)
11. Jäger, S.; Trojan, H.; Kopp, T.; Laszczyk, M.N.; Scheffler, A. Pentacyclic triterpene distribution in various plants—rich sources for a new group of multi-potent plant extracts. *Molecules* **2009**, *14*, 2016–2031. [\[CrossRef\]](#)
12. Cachofeiro, V.; Goicochea, M.; de Vinuesa, S.D.; Oubina, P.; Lahera, V.; Luno, J. Oxidative stress and inflammation, a link between chronic kidney disease and cardiovascular disease. *Kidney Int. Suppl.* **2008**, *111*, S4–S9. [\[CrossRef\]](#)
13. Tucker, P.S.; Scanlan, A.T.; Dalbo, V.J. Chronic kidney disease influences multiple systems: Describing the relationship between oxidative stress, inflammation, kidney damage, and concomitant disease. *Oxidative Med. Cell. Longev.* **2015**, *2015*, 806358. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Gliozzi, M.; Malara, N.; Muscoli, S.; Mollace, V. The treatment of hyperuricemia. *Int. J. Cardiol.* **2016**, *213*, 23–27. [\[CrossRef\]](#)
15. Abu Bakar, F.I.; Abu Bakar, M.F.; Rahmat, A.; Abdullah, N.; Sabran, S.F.; Endrini, S. Anti-gout potential of Malaysian medicinal plants. *Front. Pharmacol.* **2018**, *9*, 261. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Liu, G.; Lu, X.; Zhao, J.; Li, X. Sunflower head enzymatic hydrolysate relieves hyperuricemia by inhibiting crucial proteins (xanthine oxidase, adenosine deaminase, uric acid transporter1) and restoring gut microbiota in mice. *J. Funct. Foods.* **2020**, *72*, 104055. [\[CrossRef\]](#)
17. Chinchilla, S.P.; Urionaguena, I.; Perez-Ruiz, F. Febuxostat for the chronic management of hyperuricemia in patients with gout. *Expert Rev. Clin. Pharmacol.* **2016**, *9*, 665–673. [\[CrossRef\]](#)

18. Gupta, A.K.; Mausumi, B.; Ravi, M. Skin cancer concerns in people of color: Risk factors and prevention. *Asian Pac. J. Cancer Prev.* **2016**, *17*, 5257. [[PubMed](#)]
19. Mukherjee, P.K.; Biswas, R.; Sharma, A.; Banerjee, S.; Biswas, S.; Katiyar, C.K. Validation of medicinal herbs for anti-tyrosinase potential. *J. Herb. Med.* **2018**, *14*, 1–16. [[CrossRef](#)]
20. Anh, L.H.; Quan, N.V.; Lam, V.Q.; Iuchi, Y.; Takami, A.; Teschke, R.; Xuan, T.D. Antioxidant, anti-tyrosinase, anti- $\alpha$ -amylase, and cytotoxic potentials of the invasive weed *Andropogon virginicus*. *Plants* **2021**, *10*, 69. [[CrossRef](#)]
21. Banerjee, S.; Mazumdar, S. Electrospray ionization mass spectrometry: A technique to access the information beyond the molecular weight of the analyte. *Int. J. Anal. Chem.* **2012**, *2012*, 282574. [[CrossRef](#)] [[PubMed](#)]
22. Pinto, S.A.H.; Pinto, L.M.S.; Cunha, G.M.A.; Chaves, M.H.; Santos, F.A.; Rao, V.S. Anti-inflammatory effect of  $\alpha$ ,  $\beta$ -amyrin, a pentacyclic triterpene from *Protium heptaphyllum* in rat model of acute periodontitis. *Inflammopharmacology* **2008**, *16*, 48–52. [[CrossRef](#)]
23. Mahato, S.B.; Kundu, A.P.  $^{13}\text{C}$  NMR spectra of pentacyclic triterpenoids—A compilation and some salient features. *Phytochemistry* **1994**, *37*, 1517–1575. [[CrossRef](#)]
24. Shantos, K.B.; Raja, R. Quantitative  $^1\text{H}$  NMR spectroscopy. *TrAC-Trends Anal. Chem.* **2012**, *35*, 5–26.
25. Pham, T.L.; Ha, T.T.T.; Hoang, T.L.T. Comparative proteomic analysis of *Celastrus hindsii* Benth. phenotypes reveal an intraspecific variation. *Plant Biotechnol. J.* **2020**, *47*, 273–282.
26. Kuo, Y.H.; Kuo, L.M.Y. Antitumor and anti-AIDS triterpenes from *Celastrus hindsii*. *Phytochemistry* **1997**, *44*, 1275–1281. [[PubMed](#)]
27. Hu, X.Q.; Han, W.; Han, Z.Z.; Liu, Q.X.; Xu, X.K.; Fu, P.; Li, H.L. Three new diphenylpropanes from *Celastrus hindsii*. *Arch. Pharm. Res.* **2014**, *37*, 1411–1415. [[CrossRef](#)]
28. Sung, T.V.; Thuy, T.T.; Ninh, P.T.; Nhung, L.T.H. Isolation and structural characterization of phenolic glycoside and triterpenes in *Celastrus hindsii* Benth. *New J. Chem.* **2008**, *46*, 224–228.
29. Nicholson, J.K.; Foxall, P.J.; Spraul, M.; Farrant, R.D.; Lindon, J.C. 750 MHz  $^1\text{H}$  and  $^1\text{H}$ - $^{13}\text{C}$  NMR spectroscopy of human blood plasma. *Anal. Chem.* **1995**, *67*, 793–811. [[CrossRef](#)]
30. De Melo, K.M.; De Oliveira, F.T.B.; Silva, R.A.C.; Quinderé, A.L.G.; Marinho Filho, J.D.B.; Araújo, A.J.; Santos, F.A.  $\alpha$ ,  $\beta$ -Amyrin, a pentacyclic triterpenoid from *Protium heptaphyllum* suppresses adipocyte differentiation accompanied by down regulation of PPAR $\gamma$  and C/EBP $\alpha$  in 3T3-L1 cells. *Biomed. Pharmacother.* **2019**, *109*, 1860–1866. [[CrossRef](#)]
31. Ching, J.; Chua, T.K.; Chin, L.C.; Lau, A.J.; Pang, Y.K.; Jaya, J.M.; Koh, H.L. Beta-amyrin from *Ardisia elliptica* Thunb. is more potent than aspirin in inhibiting collagen-induced platelet aggregation. *Indian J. Exp. Biol.* **2010**, *48*, 275–279.
32. Otuki, M.F.; Ferreira, J.; Lima, F.V.; Meyre-Silva, C.; Malheiros, A.; Muller, L.A.; Calixto, J.B. Antinociceptive properties of mixture of  $\alpha$ -amyrin and  $\beta$ -amyrin triterpenes: Evidence for participation of protein kinase C and protein kinase A pathways. *J. Pharmacol. Exp. Ther.* **2005**, *313*, 310–318. [[CrossRef](#)]
33. Oliveira, F.A.; Chaves, M.H.; Almeida, F.R.; Lima, R.C., Jr.; Silva, R.M.; Maia, J.L.; Rao, V.S. Protective effect of  $\alpha$ - and  $\beta$ -amyrin, a triterpene mixture from *Protium heptaphyllum* (Aubl.) March. trunk wood resin, against acetaminophen-induced liver injury in mice. *J. Ethnopharmacol.* **2005**, *98*, 103–108. [[CrossRef](#)] [[PubMed](#)]
34. Oliveira, F.A.; Vieira-Júnior, G.M.; Chaves, M.H.; Almeida, F.R.; Santos, K.A.; Martins, F.S.; Rao, V.S. Gastroprotective effect of the mixture of  $\alpha$ - and  $\beta$ -amyrin from *Protium paraphyllum*: Role of capsaicin-sensitive primary afferent neurons. *Planta Med.* **2004**, *70*, 780–782. [[CrossRef](#)]
35. Akihisa, T.; Yasukawa, K.; Oinuma, H.; Kasahara, Y.; Yamanouchi, S.; Takido, M.; Tamura, T. Triterpene alcohols from the flowers of compositae and their anti-inflammatory effects. *Phytochemistry* **1996**, *43*, 1255–1260. [[CrossRef](#)]
36. Melo, C.M.; Morais, T.C.; Tomé, A.R.; Brito, G.A.C.; Chaves, M.H.; Rao, V.S.; Santos, F.A. Anti-inflammatory effect of  $\alpha$ ,  $\beta$ -amyrin, a triterpene from *Protium heptaphyllum*, on cerulein-induced acute pancreatitis in mice. *Inflamm. Res.* **2011**, *60*, 673–681. [[CrossRef](#)]
37. Ching, J.; Lin, H.S.; Tan, C.H.; Koh, H.L. Quantification of  $\alpha$ - and  $\beta$ -amyrin in rat plasma by gas chromatography–mass spectrometry: Application to preclinical pharmacokinetic study. *J. Mass Spectrom.* **2011**, *46*, 457–464. [[CrossRef](#)] [[PubMed](#)]
38. Akash, M.S.H.; Rehman, K. Column chromatography. In *Essentials of Pharmaceutical Analysis*; Springer: Singapore, 2020; pp. 167–174.
39. Lisec, J.; Schauer, N.; Kopka, J.; Willmitzer, L.; Fernie, A.R. Gas chromatography mass spectrometry–based metabolite profiling in plants. *Nat. Protoc.* **2006**, *1*, 387–396. [[CrossRef](#)] [[PubMed](#)]
40. De Lacy Costello, B.P.J.; Evans, P.; Ewen, R.J.; Gunson, H.E.; Jones, P.R.; Ratcliffe, N.M.; Spencer-Phillips, P.T. Gas chromatography–mass spectrometry analyses of volatile organic compounds from potato tubers inoculated with *Phytophthora* infesting or *Fusarium coeruleus*. *Plant Pathol. J.* **2001**, *50*, 489–496. [[CrossRef](#)]
41. Ho, C.S.; Lam, C.W.K.; Chan, M.H.M.; Cheung, R.C.K.; Law, L.K.; Lit, L.C.W.; Ng, K.F.; Suen, M.W.M.; Tai, H.L. Electrospray ionization mass spectrometry: Principles and clinical applications. *Clin. Biochem. Rev.* **2003**, *24*, 3.
42. Bruins, A.P. Mechanistic aspects of electrospray ionization. *J. Chromatogr. A* **1998**, *794*, 345–357. [[CrossRef](#)]
43. Hoult, D.I.; Bhakar, B. NMR signal reception: Virtual photons and coherent spontaneous emission. *Concepts Magn. Reson. A Bridg. Educ. Res.* **1997**, *9*, 277–297. [[CrossRef](#)]
44. Sternberg, U.; Witter, R.; Kuprov, I.; Lamley, J.M.; Oss, A.; Lewandowski, J.R.; Samoson, A.  $^1\text{H}$  line width dependence on MAS speed in solid state NMR—comparison of experiment and simulation. *J. Magn. Reson.* **2018**, *291*, 32–39. [[CrossRef](#)]

45. Morita, M.; Shibuya, M.; Kushiro, T.; Masuda, K.; Ebizuka, Y. Molecular cloning and functional expression of triterpene synthases from pea (*Pisum sativum*) New  $\alpha$ -amyrin-producing enzyme is a multifunctional triterpene synthase. *Eur. J. Biochem.* **2000**, *267*, 3453–3460. [[CrossRef](#)]
46. Martelanc, M.; Vovk, I.; Simonovska, B. Determination of three major triterpenoids in epicuticular wax of cabbage (*Brassica oleracea* L.) by high-performance liquid chromatography with UV and mass spectrometric detection. *J. Chromatogr. A* **2007**, *1164*, 145–152. [[CrossRef](#)]
47. Sunil, C.; Irudayaraj, S.S.; Duraipandiyar, V.; Al-Dhabi, N.A.; Agastian, P.; Ignacimuthu, S. Antioxidant and free radical scavenging effects of  $\beta$ -amyrin isolated from *Symplocos cochinchinensis* Moore. leaves. *Ind. Crops Prod.* **2014**, *61*, 510–516. [[CrossRef](#)]
48. Okoye, N.N.; Ajaghaku, D.L.; Okeke, H.N.; Ilodigwe, E.E.; Nworu, C.S.; Okoye, F.B.C. Beta-amyrin and alpha-amyrin acetate isolated from the stem bark of *Alstonia boonei* display profound anti-inflammatory activity. *Pharm. Biol.* **2014**, *52*, 1478–1486. [[CrossRef](#)]
49. Sirat, H.M.; Susanti, D.; Ahmad, F.; Takayama, H.; Kitajima, M. Amides, triterpene, and flavonoids from the leaves of *Melastoma malabathricum* L. *J. Nat. Med.* **2010**, *64*, 492–495. [[CrossRef](#)]
50. Saeidnia, S.; Ara, L.; Hajimehdipour, H.; Read, R.W.; Arshadi, S.; Nikan, M. Chemical constituents of *Swertia longifolia* Boiss. with  $\alpha$ -amylase inhibitory activity. *Res. Pharm. Sci.* **2016**, *11*, 23–32.
51. Quan, N.V.; Xuan, T.D.; Tran, H.D.; Thuy, N.T.D.; Trang, L.T.; Huong, C.T.T.; Tuyen, P.T. Antioxidant,  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory activities and potential constituents of *Canarium tramdenum* bark. *Molecules* **2019**, *24*, 605. [[CrossRef](#)]
52. Higuchi, C.T.; Pavan, F.R.; Leite, C.Q.F.; Sannomiya, M.; Vilegas, W.; Leite, S.R.D.A.; Sato, D.N. Triterpenes and antitubercular activity of *Byrsonima crassa*. *Quim. Nova* **2008**, *31*, 1719–1721. [[CrossRef](#)]
53. Hernandez-Vazquez, L.; Bonfill, M.; Moyano, E.; Cusido, R.M.; Navarro-Ocana, A.; Palazon, J. Conversion of  $\alpha$ -amyrin into centellosides by plant cell cultures of *Centella asiatica*. *Biotechnol. Lett.* **2010**, *32*, 315–319. [[CrossRef](#)]
54. Persson, T.; Popescu, B.O.; Cedazo-Minguez, A. Oxidative stress in Alzheimer's disease: Why did antioxidant therapy fail? *Oxidative Med. Cell. Longev.* **2014**, 427318. [[CrossRef](#)]
55. López-Alarcón, C.; Denicola, A. Evaluating the antioxidant capacity of natural products: A review on chemical and cellular-based assays. *Anal. Chim. Acta* **2013**, *763*, 1–10. [[CrossRef](#)]
56. Gutteridge, J.M.; Halliwell, B. Invited review free radicals in disease processes: A compilation of cause and consequence. *Free Radic. Res. Commun.* **1993**, *19*, 141–158. [[CrossRef](#)] [[PubMed](#)]
57. Halliwell, B.A.R.R.Y.; Gutteridge, J.M.; Cross, C.E. Free radicals, antioxidants, and human disease: Where are we now? *J. Lab. Clin. Med.* **1992**, *119*, 598–620. [[PubMed](#)]
58. Karen Cardoso, B.; Line Marko de Oliveira, H.; Zonta Melo, U.; Mariano Fernandez, C.M.; Franco de Araújo Almeida Campo, C.; Gonçalves, J.E.; Cristiani Gazim, Z. Antioxidant activity of  $\alpha$  and  $\beta$ -amyrin isolated from *Myrcianthes pungens* leaves. *Nat. Prod. Res.* **2020**, *34*, 1777–1781. [[CrossRef](#)] [[PubMed](#)]
59. Melo, C.M.; Carvalho, K.M.M.B.; De Sousa Neves, J.C.; Morais, T.C.; Rao, V.S.; Santos, F.A.; De Castro, G.A.; Chaves, M.H.  $\alpha$ ,  $\beta$ -amyrin, a natural triterpenoid ameliorates L-arginine-induced acute pancreatitis in rats. *World J. Gastroenterol.* **2010**, *16*, 4272–4280. [[CrossRef](#)]
60. Shoji, A.; Yamanaka, H.; Kamatani, N. A retrospective study of the relationship between serum urate level and recurrent attacks of gouty arthritis: Evidence for reduction of recurrent gouty arthritis with antihyperuricemic therapy. *Arthritis Care Res.* **2004**, *51*, 321–325. [[CrossRef](#)]
61. Yamanaka, H.; Metabolism, T.G. Essence of the revised guideline for the management of hyperuricemia and gout. *Jpn. Med. Assoc. J.* **2012**, *55*, 324–329.
62. Koto, R.; Nakajima, A.; Horiuchi, H.; Yamanaka, H. Real-world treatment of gout and asymptomatic hyperuricemia: A cross-sectional study of Japanese health insurance claims data. *Mod. Rheumatol.* **2021**, *31*, 261–269. [[CrossRef](#)] [[PubMed](#)]
63. Dalbeth, N.; Haskard, D.O. Mechanisms of inflammation in gout. *Rheumatology* **2005**, *44*, 1090–1096. [[CrossRef](#)]
64. Kuo, C.F.; Grainge, M.J.; Mallen, C.; Zhang, W.; Doherty, M. Rising burden of gout in the UK but continuing suboptimal management: A nationwide population study. *Ann. Rheum. Dis.* **2015**, *74*, 661–667. [[CrossRef](#)] [[PubMed](#)]
65. Nguyen, T.D.; Thuong, P.T.; Hwang, I.H.; Hoang, T.K.H.; Nguyen, M.K.; Nguyen, H.A.; Na, M. Anti-hyperuricemic, anti-inflammatory and analgesic effects of *Siegesbeckia orientalis* L. Resulting from the fraction with high phenolic content. *BMC Complement. Med. Ther.* **2017**, *17*, 191. [[CrossRef](#)]
66. Korner, A.; Pawelek, J. Mammalian tyrosinase catalyzes three reactions in the biosynthesis of melanin. *Science* **1982**, *217*, 1163–1165. [[CrossRef](#)] [[PubMed](#)]
67. Boyle, J.L.; Haupt, H.M.; Stern, J.B.; Mulhaupt, H.A. Tyrosinase expression in malignant melanoma, desmoplastic melanoma, and peripheral nerve tumors: An immunohistochemical study. *Arch. Pathol. Lab. Med.* **2002**, *126*, 816–822. [[CrossRef](#)] [[PubMed](#)]
68. Baharav, E.; Merimski, O.; Shoenfeld, Y.; Zigelman, R.; Gilbrud, B.; Yecheskel, G.; Fishman, P. Tyrosinase as an autoantigen in patients with vitiligo. *Clin. Exp. Immunol.* **1996**, *105*, 84–88. [[CrossRef](#)]
69. Patel, A.R.; Sun, W.Q.; Payne, G.F. Tyrosinase reaction/chitosan adsorption: Potential for removing polymerization storage inhibitors. *Ind. Eng. Chem. Res.* **1994**, *33*, 2168–2173. [[CrossRef](#)]

70. Nichols, L.A.; Chen, Y.; Colella, T.A.; Bennett, C.L.; Clausen, B.E.; Engelhard, V.H. Deletional self-tolerance to a melanocyte/melanoma antigen derived from tyrosinase is mediated by a radio-resistant cell in peripheral and mesenteric lymph nodes. *J. Immunol.* **2007**, *179*, 993–1003. [[CrossRef](#)]
71. Högnadóttir, Á.; Rouseff, R.L. Identification of aroma active compounds in orange essence oil using gas chromatography olfactometry and gas chromatography mass spectrometry. *J. Chromatogr. A* **2003**, *998*, 201–211. [[CrossRef](#)]
72. Andriana, Y.; Xuan, T.D.; Quy, T.N.; Minh, T.N.; Van, T.M.; Viet, T.D. Antihyperuricemia, antioxidant, and antibacterial activities of *Tridax procumbens* L. *Foods* **2019**, *8*, 21. [[CrossRef](#)] [[PubMed](#)]
73. Quy, T.N.; Xuan, T.D. Xanthine oxidase inhibitory potential, antioxidant and antibacterial activities of *Cordyceps militaris* (L.) Link fruiting body. *Medicines* **2019**, *6*, 20. [[CrossRef](#)] [[PubMed](#)]