



Voluntary sustainability standards could significantly reduce detrimental impacts of global agriculture

W. K. Smith^{a,b}, E. Nelson^c, J. A. Johnson^b, S. Polasky^{d,e,1}, J. C. Milder^{f,g}, J. S. Gerber^b, P. C. West^b, S. Siebert^h, K. A. Brauman^b, K. M. Carlsonⁱ, M. Arbutnot^j, J. P. Rozza^k, and D. N. Pennington^{b,j,1}

^aSchool of Natural Resources and the Environment, University of Arizona, Tucson, AZ 85721; ^bInstitute on the Environment, University of Minnesota, St. Paul, MN 55108; ^cDepartment of Economics, Bowdoin College, Brunswick, ME 04011; ^dDepartment of Applied Economics, University of Minnesota, St. Paul, MN 55108; ^eDepartment of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108; ^fEvaluation and Research Program, Rainforest Alliance, New York, NY 10279; ^gDepartment of Natural Resources, Cornell University, Ithaca, NY 14853; ^hDepartment of Crop Sciences, University of Göttingen, 37075 Göttingen, Germany; ⁱDepartment of Natural Resources and Environmental Management, University of Hawaii at Mānoa, Honolulu, HI 96822; ^jConservation Science Program, World Wildlife Fund, Washington, DC 20037; and ^kEnvironmental, Health, and Safety, The Coca-Cola Company, Atlanta, GA 30313

Contributed by S. Polasky, July 12, 2018 (sent for review May 11, 2017; reviewed by Daniel C. Nepstad and Vivek Anand Voora)

Voluntary sustainability standards (VSS) are stakeholder-derived principles with measurable and enforceable criteria to promote sustainable production outcomes. While institutional commitments to use VSS to meet sustainable procurement policies have grown rapidly over the past decade, we still have relatively little understanding of the (i) direct environmental benefits of large-scale VSS adoption; (ii) potential perverse indirect impacts of adoption; and (iii) implementation pathways. Here, we illustrate and address these knowledge gaps using an ecosystem service modeling and scenario analysis of Bonsucro, the leading VSS for sugarcane. We find that global compliance with the Bonsucro environmental standards would reduce current sugarcane production area (–24%), net tonnage (–11%), irrigation water use (–65%), nutrient loading (–34%), and greenhouse gas emissions from cultivation (–51%). Under a scenario of doubled global sugarcane production, Bonsucro adoption would further limit water use and greenhouse gas emissions by preventing sugarcane expansion into water-stressed and high-carbon stock ecosystems. This outcome was achieved via expansion largely on existing agricultural lands. However, displacement of other crops could drive detrimental impacts from indirect land use. We find that over half of the potential direct environmental benefits of Bonsucro standards under the doubling scenario could be achieved by targeting adoption in just 10% of global sugarcane production areas. However, designing policy that generates the most environmentally beneficial Bonsucro adoption pathway requires a better understanding of the economic and social costs of VSS adoption. Finally, we suggest research directions to advance sustainable consumption and production.

sustainability standards | ecosystem services | agriculture | environmental policy | land use change

Increasing agricultural output to meet growing global demand for food, fiber, and fuel is a major driver of ecosystem degradation and biodiversity loss (1–4). Reducing detrimental environmental impacts while maintaining needed agricultural production can be achieved by increasing yields, improving input use efficiency, and targeting agricultural production to areas with lower environmental impact (1, 2, 5–10). While some opportunities for jointly increasing agricultural production and environmental outcomes exist (11), large-scale reductions in the environmental impact of agricultural production will likely require additional incentives for producers to adopt environmentally benign practices (12, 13). Such incentives may include social pressures (i.e., peer pressure to adopt certain practices), price premiums, market access, and/or government-promulgated rewards for adopting certain practices or penalties for not abiding by environmental laws and regulations.

Public policy has traditionally played a key role in driving large-scale transitions toward more sustainable land use (14). Governments have used both involuntary regulatory approaches

and voluntary incentive-based regulatory programs to encourage more sustainable agriculture production. Some governments have used mandates via laws to regulate the environmental impact of commodity production (15, 16). However, involuntary mandates tend to be unpopular with agricultural producers, can be economically inefficient, and can be environmentally ineffective where government institutions are weak or political will is insufficient to enact and enforce legal regulations (16–18). For example, the 2012 Brazilian Forest Code seeks to legally protect the conversion of forests and savannas on Brazil’s 394 Mha of privately owned lands. However, a recent analysis found that full compliance with the Forest Code offered few economic benefits to landowners (19). Full compliance with the law requires high restoration costs and large opportunity costs of foregone production. In addition, expected landowner costs are low for noncompliance due to poor code enforcement. While registered properties initially showed lower deforestation rates than unregistered ones, these differences diminished over time; moreover, only 6% of registered landowners reported taking steps to restore illegally cleared areas on their properties (19).

Significance

Voluntary sustainability standards (VSS) may be an effective way to reduce the negative impacts of agriculture at regional to global scales. Here, we present an approach that highlights the potential of VSS to reduce some of the negative externalities associated with agriculture production. To illustrate this potential, we show that VSS could reduce the global environmental impacts from growing sugarcane. Further, most of this environmental benefit comes from targeting just 10% of production area. To realize these environmental gains, incentives for VSS adoption need to be sufficient to cover the costs of criteria compliance. Determining these costs and public and private-sector mechanisms for efficiently transferring VSS-adoption subsidies to farmers and millers are key future research needs.

Author contributions: W.K.S., E.N., J.A.J., S.P., J.C.M., J.S.G., P.C.W., S.S., K.A.B., K.M.C., M.A., J.P.R., and D.N.P. designed research; W.K.S. and D.N.P. performed research; W.K.S. analyzed data; and W.K.S., E.N., J.A.J., S.P., J.C.M., J.S.G., P.C.W., S.S., K.A.B., K.M.C., M.A., J.P.R., and D.N.P. wrote the paper.

Reviewers: D.C.N., Earth Innovation Institute; and V.A.V., Central European University.

The authors declare no conflict of interest.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: polasky@umn.edu or derrick@umn.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1707812116/-DCSupplemental.

Published online January 22, 2019.

Table 1. Bonsucro VSS environmental requirements considered in the study and their expected environmental outcomes

Indicator	Standard	Impact on efficiency	Impact on expansion
Yield	>50th percentile of estimated yields in the year 2000 by climate zone	Increased sugarcane production per unit area	Reduced need for expansion to meet demand
Water quality (eutrophication potential)	<1.05 actual fertilizer inputs per estimated optimum fertilizer inputs*	Increased sugarcane production per unit eutrophication	NA
Water use	>90 kg sugarcane per mm irrigation water applied	Increased sugarcane production per unit irrigation	Reduced expansion into areas where irrigation water demands are high
GHG _{CULT}	<40 kg CO ₂ eq per metric ton sugarcane	Increased sugarcane production per unit GHG emission	NA
GHG _{LUC}	<50 kg CO ₂ eq per metric ton sugarcane	NA	Reduced expansion into areas of high carbon stocks

NA, not applicable.

*Fertilizer inputs defined to include nitrogen and phosphorus inputs.

gap approach (5, 6), we evaluated production and environmental impacts from the following VSS-adoption scenario comparisons: (i) universal adoption of Bonsucro environmental criteria over all current sugarcane production areas versus current sugarcane production; (ii) doubling of global sugarcane production under universal adoption of Bonsucro versus a business-as-usual (BAU) doubling scenario where there is no deliberate attempt to make new production Bonsucro compliant; and (iii) incremental adoption of the Bonsucro VSS to maximize its near-term global environmental benefits. We also explored the potential indirect effects, such as the displacement of existing land use, of large-scale uptake of Bonsucro VSS.

Results

Global Sugarcane Production Efficiency and Current Compliance with Bonsucro VSS.

First, we assessed the extent of existing sugarcane cultivation that does not comply with Bonsucro's criteria for minimum yield (yield indicator; Table 1), maximum nutrient use (water quality and GHG_{CULT} indicators; Table 1), and water use (water use indicator; Table 1). We found that 63% of global sugarcane production areas are currently noncompliant with at least one of these criteria (Fig. 1 and *SI Appendix, Table S1*): 40% of global production areas are noncompliant with the yield standard, 44% are noncompliant with the water use standard, and 23% are noncompliant with the nutrient use standard. Patterns of compliance and noncompliance were clustered regionally (Fig. 1 and *SI Appendix, Table S1*). For instance, 52% of rain-fed sugarcane production in South America—a region accounting for 57% (26.7%) of global rain-fed (total) sugarcane production—are compliant with at least two of the three standards (Fig. 1 and *SI Appendix, Fig. S5*). By contrast, only 21% of sugarcane production in South Asia—a region accounting for 50% (26.9%) of global irrigated (total) sugarcane production—are compliant with at least two of the three standards (Fig. 1 and *SI Appendix, Fig. S5*).

Our models suggest that universal adoption of all Bonsucro environmental criteria would mean that 24.3% of current production area would need to be abandoned, resulting in a 10.6% reduction in sugarcane production, 65.3% reduction in water use, 33.4% reduction in nutrient loading, and 51.2% reduction in GHG_{CULT} emissions relative to current conditions (*SI Appendix, Table S1*). These potential impacts are clustered regionally, with the largest potential water use and nutrient loading reductions found in South Asia (Fig. 1 and *SI Appendix, Table S1*); this region, currently the second largest in global sugarcane production, also shows the largest reductions in existing production area and production tonnage. By contrast, other regions, such as Central Africa, could potentially see large increases in production tonnage (+219.0%) by increasing nutrient application rates according to the

Bonsucro environmental criteria, with the trade-off of potential increases in eutrophication potential (+104.9%) and GHG_{CULT} emissions (+29.3%). This finding is generalizable to many regions in Africa that appear to be currently low yielding due to under-application of nutrients N and P (*SI Appendix, Figs. S3 and S4*).

Global Sugarcane Expansion Potential and Compliance with Bonsucro VSS.

We estimated the amount of current forest, savanna, grassland, and shrubland in climates suitable for sugarcane production (“available natural land”) (*Methods*) that would be compliant with Bonsucro environmental criteria if converted to sugarcane. The relevant indicators in this case were total greenhouse gas emissions from land clearing (GHG_{LUC} indicator; Table 1) and irrigation




Fig. 1. Compliance with Bonsucro VSS environmental indicators affecting sugarcane production efficiency. Areas currently noncompliant with different combinations of yield (Y), water use (W), and nutrient application (N), as well as areas identified as fully compliant with these criteria (CMLPNT) (Table 1). (A) Spatial pattern of compliance across major rain-fed production areas in and around South America (SAM). (B) Spatial pattern of compliance across major irrigated production areas in and around South Asia (SAS). (C) Regional summary of compliance partitioned by irrigated and rain-fed production. Roughly 63% of global sugarcane production is noncompliant with at least one of the indicators considered. AUS, Australia; CAF, Central Africa; CAM, Central America; CAR, Caribbean; EAF, East Africa; EAS, East Asia; NAF, North Africa; NAM, North America; SAF, South Africa; SAM, South America; SAS, South Asia; SEA, Southeast Asia; WAF, West Africa.




Fig. 3. The potential impact of global uptake of Bonsucro VSS environmental indicators under future doubled sugarcane production. The spatial difference between the Bonsucro (BON) and business-as-usual (BAU) scenarios for (A) global and (B) Southeast Asia production areas. Values indicate increased (green) or decreased (purple) sugarcane production areas under BON relative to BAU. (C) The relative change across environmental indicators under BON relative to BAU. Reported are the mean values across an ensemble of model runs that account for uncertainty in yield, nutrient demand, water demand, GHG emissions, and land use change. GHGE_{CULT}, greenhouse gas emissions associated with cultivation; GHGE_{LUC}, greenhouse gas emissions associated with land use change; LUC_{MAN}, conversion of managed land to sugarcane; LUC_{NAT}, conversion of natural land to sugarcane; PRD_{EXT}, extensified production; PRD_{INT}, intensified production; WTR_{QUAL}, water quality; WTR_{USE}, water use. Error bars represent 2 SDs of the mean.

environmental benefits strategy, converting only ~10% of global sugarcane production area is required to achieve more than 50% of the potential environmental benefits realized under the full BON scenario (Fig. 4A). The large difference in incremental efficacy of these two targeting approaches demonstrates the importance of objective definition—maximizing compliant area versus maximizing environmental benefits per unit area—when targeting future VSS adoption.

Discussion

We find significant potential for environmental and production efficiency improvements across the global sugarcane sector. We also demonstrate that adoption of the environmental criteria in the Bonsucro VSS, whether incrementally targeted or universal, would greatly reduce the direct environmental damage caused by sugarcane production. We found that the potential environmental benefits of the Bonsucro VSS are driven largely through two major changes in current sugarcane production patterns. First, Bonsucro VSS adoption shifts production away from arid ecosystems where annual freshwater use exceeds the water use indicator (Fig. 3). The environmental benefit of this shift is most notable in areas identified as high to severely water-stressed (39).

In these areas, under the doubling of sugarcane production scenario analysis, we found that Bonsucro VSS compliance (the BON scenario) reduces total annual water use by sugarcane by $66 \pm 8.6\%$ relative to BAU (*SI Appendix*, Fig. S8). Indeed, a third of the current sugarcane production area in India, the second largest producer of sugarcane globally, cannot comply with the water use indicator. Such shifts away from current major production areas highlights a significant challenge for those seeking to meet VSS commitments via 100% compliance with Bonsucro. Second, any level of sugarcane expansion under Bonsucro VSS adoption shifts expansion toward managed lands, thus sparing the direct conversion of high carbon density natural ecosystems (Fig. 3).

That expanded production under the Bonsucro VSS (the BON scenario) prevents the direct conversion of natural lands, including forest and savanna ecosystems, is especially important given the growing trend of “deforestation-free” or “land conversion-free” commitments by the private sector. In addition to meeting “conversion-free” policy goals, incremental or universal Bonsucro VSS adoption also promotes production intensification and improves water and climate mitigation outcomes, thus highlighting the value of multicriteria standards to help deliver multiple outcomes and the potential shortcomings of developing and

area is a critical insight for those stakeholders who support adoption of VSS as an effective way to reduce the impact of agricultural production on the environment versus those who only base VSS impacts on market adoption. Many VSS supporters track progress based on the total area of production enrolled under VSS rather than a quality adjusted measure of uptake. For example, World Wildlife Fund for Nature (WWF), a founding member of Bonsucro, established a target whereby 25% of global sugarcane production area is enrolled under Bonsucro by 2020 (www.panda.org/our_work/markets/mti_solutions/certification/agriculture/sugarcane2/bonsucro/). Similar targets have been set by major corporate buyers of sugarcane. However, such production-based targets ignore the tendency for most early adopters to be in places where VSS compliance is most readily achieved, not necessarily those that can generate the most environmental benefit per area. Since ~37% of global sugarcane production is currently compliant with Bonsucro environmental criteria (Fig. 1), 37% adoption of Bonsucro is needed before any additional environmental benefits are realized (Fig. 4). In other words, focusing on quantity of uptake rather than quality of uptake could mean much less new environmental benefit from incremental adoption of the standards. Therefore, environmentally effective incremental adoption of VSS requires targeting of production areas that are underperforming relative to the VSS criteria (33). To do this, incentives for VSS adoption (e.g., price premiums, public and private financial funds, technical information, market access) need to be sufficient to overcome the costs of compliance on land furthest from complying with Bonsucro VSS criteria (42–44). Determining these costs is a key future research priority that has implications for VSS and land-use policy, generally.

Finally, our analysis also highlights the uneven impact that widespread adoption of Bonsucro VSS could have on sugarcane producers' economic livelihoods. For example, we found large regional differences in the ability to comply with Bonsucro VSS criteria (Fig. 3). Reducing or eliminating sugarcane production in such noncomplaint regions (e.g., water-stressed areas in South Asia) is almost certain to cause socioeconomic hardships for people dependent on the regional sugarcane industry. How to meet environmental standards while not imposing undue social and economic burdens remains an important open question.

Conclusions

We demonstrate the potential of Bonsucro—one of the largest and fastest growing global voluntary sustainability standards (VSS)—to reduce eutrophication, water use, greenhouse gas emissions, and natural ecosystem conversion. A better understanding of the potential large-scale sustainability outcomes associated with VSS adoption is vital to determining the optimal contribution that VSS, as part of a portfolio of sustainability policy mechanisms, can make to national and global sustainability commitments and development goals (e.g., United Nation's Sustainable Development Goals) (45). VSS or other sustainability policies for commodity production can use the analytical framework presented here for design and assessment of specific indicators, identification of leverage points, exploration of incentives for widespread adoption, and insight into potential leakage and other indirect market effects. For example, this framework could be adapted to define goals for how much adoption or market penetration is needed to have impact beyond the local level. This approach also provides a logical framework for integrating the assessment of environmental standards with social and economic factors, a key next step and priority research area in sustainability science.

Our framework also presents the opportunity for the agriculture VSS community to rigorously reassess standard design in a more explicit outcome-based and forward-looking way. It is important to separate standards from their implementing mechanism. The standards (i.e., the principles, criteria, and indicators)

define the “rules of the game” for producers. The implementation mechanism can be demand-led or market or governmental regulation. Therefore, standard design is paramount to defining what outcomes can be expected from a VSS certification or a law. Such information could improve standard efficacy and illustrate the potential value from supporting its adoption.

The Bonsucro VSS is unique among existing agriculture VSS schemes in that it uses a performance or metric-based approach to standard design that leverages basic agronomic and conservation principles that could be readily adapted to many commodity sectors. A more common approach to standard design, often described as practice-based, tends to have less explicit, more relative (to baseline conditions) and directional criteria and associated indicators. Moreover, Bonsucro strives to enhance both production intensification and reduced extensification by setting standards to promote production efficiency and reduce natural land conversion. Better understanding of how these design choices affect the environment and adoption patterns can inform the broader private and public-sector debate on the utility of VSS in driving the achievement of sustainable development goals (45).

VSS of the Bonsucro design potentially offer an effective mechanism to manage commodity supply reliability risks, reduce environmental impacts, and meet the growing consumer preference for sustainable products regardless of state actions to increase product sustainability. The results of this research inform long-term commodity sourcing strategies by assessing regional differences in producers' basic ability to meet VSS. These strategies can be adjusted to avoid increasing procurement from areas where complying with VSS will remain a challenge until best practices can be scaled and innovations introduced such that compliance is achievable. Commodity sourcing strategies can also seek to increase procurement in areas where compliance is readily achievable and may also drive greater productivity with a smaller environmental footprint. Moreover, we illustrate the need for VSS, or any land-use policy, to be evaluated within the context of the larger physical and policy landscape. VSS can ultimately only be effective if they complement other landscape-wide efforts.

Our results suggest that the realization of the potential environmental benefits of VSS schemes will depend on (i) accounting for and limiting indirect effects that could undermine direct environmental benefits and (ii) effective intervention to bring current production areas that are underperforming relative to the Bonsucro VSS environmental criteria. Thorough assessment of the costs of VSS adoption and the incentives is needed to drive adoption for farmers, buyers, and customers.

Future research is needed to improve understanding of the explicit economic cost (e.g., management related) and related barriers (e.g., market access) of targeting adoption of VSS in underperforming areas, which can increase the incremental environmental benefits of VSS. To attain such targeting may require leveraging diverse policy and incentive instruments, including public and private financing mechanisms, regulatory controls, and institutions to mitigate up-front economic barriers to compliance and increase benefits to producers and value chain actors of certification in these areas. This analysis provides a first step toward a more comprehensive understanding of the role of VSS in driving a global transition toward sustainable agriculture.

Methods

The spatially explicit sugarcane yield and extent data used in this analysis are originally based on data from ref. 46 and are available at www.earthstat.org/ as the “Harvested Area and Yield for 175 Crops” dataset. These data were computed with national, state, and county level agricultural census statistics from the period 1997 to 2003 and distributed to a 5-arc minute spatial resolution based on satellite observations of global cropland area. We used the reference time period of 1997 to 2003 since it predates the initiation of

the Bonsucro sustainability standard and thus allowed an unbiased evaluation of the potential environmental benefits of standard implementation. Our yield maps give a multiyear average to account for differences between higher yielding first and lower yielding ratoon crop cycles (36). The multiyear average also reduces the impact of year-to-year variation due to weather. Yield data were partitioned into irrigated and rain-fed based on output from the Global Crop Water Model (GCWM) (47). These are the same yield data used by Bonsucro to define yield targets for global sugarcane production and thus enable a focus on the impacts of the policy (SI Appendix, Text S7).

Empirically derived yield models were developed for rain-fed and irrigated sugarcane production using a nonlinear least squares algorithm to fit climate-response curves to yield distributions by climate zone (SI Appendix, Text S2). We next developed nutrient-response curves that explicitly consider soil quality and current rates of nitrogen and phosphorus fertilization using a combination of the best available data to calculate actual, optimum, and excess nutrient application rates (SI Appendix, Texts S3 and S4). We finally utilized the Liebig law of the minimum to estimate the combined effects of water, nitrogen, and phosphorus inputs on yield (SI Appendix, Texts S3 and S4).

We utilized outputs from our global modeling to calculate a global map of sugarcane production in terms of percent deviation from compliance across the full set of environmental criteria considered (SI Appendix, Text S5). We evaluated the potential impacts of the Bonsucro sustainability standard by considering a hypothetical future of doubled sugarcane production. We

then compared a business-as-usual (BAU) scenario against a Bonsucro (BON) scenario in which all sugarcane production shifts to compliant with the full Bonsucro environmental criteria considered (Table 1), allowing the evaluation of the direct potential environmental benefits of large-scale uptake of Bonsucro sustainability standards (SI Appendix, Text S6). We quantified the sensitivity of our global model to four key model inputs: (i) estimated sugarcane yields, (ii) estimated nutrient demand, (iii) estimated irrigation demand, and (iv) estimated carbon stocks. We assumed that these spatially explicit model inputs had a variation of $\pm 10\%$, and their respective uncertainties were propagated by iteratively running the model for all combinations of minimum and maximum input values, resulting in 16 total model runs (SI Appendix, Table S3). We report the mean across the ensemble of model runs and represent uncertainty as 2 SDs of the mean (SI Appendix, Text S7).

ACKNOWLEDGMENTS. We gratefully acknowledge support for this research from the Luc Hoffmann Institute, The World Wildlife Fund, and The Coca-Cola Company Partnership, University of Minnesota's Carlson School of Management, the Belmont Forum/FACCE-JPI/NSF funded DEVIL project (Grant NE/M021327/1), NSF Award 1540195, the Gordon and Betty Moore Foundation, and the US Department of Agriculture National Institute of Food and Agriculture Hatch Project HAW01136-H managed by the College of Tropical Agriculture and Human Resources. The funders had no role in study design, data collection, or analysis, decision to publish, or preparation of the manuscript.

- Foley JA, et al. (2011) Solutions for a cultivated planet. *Nature* 478:337–342.
- West PC, et al. (2014) Leverage points for improving global food security and the environment. *Science* 345:325–328.
- Rockström J, et al. (2009) A safe operating space for humanity. *Nature* 461:472–475.
- Millennium Ecosystem Assessment (2005) *Ecosystems and Human Well-Being: Synthesis* (Island Press, Washington, DC).
- Mueller ND, et al. (2012) Closing yield gaps through nutrient and water management. *Nature* 490:254–257.
- Johnson JA, Runge CF, Senauer B, Foley J, Polasky S (2014) Global agriculture and carbon trade-offs. *Proc Natl Acad Sci USA* 111:12342–12347.
- Erb K-H, et al. (2016) Exploring the biophysical option space for feeding the world without deforestation. *Nat Commun* 7:11382.
- Paustian K, et al. (2016) Climate-smart soils. *Nature* 532:49–57.
- Flynn HC, et al. (2012) Quantifying global greenhouse gas emissions from land-use change for crop production. *Glob Chang Biol* 18:1622–1635.
- Smith P, et al. (2013) How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? *Glob Chang Biol* 19:2285–2302.
- Garbach K, et al. (2017) Examining multi-functionality for crop yield and ecosystem services in five systems of agroecological intensification. *Int J Agric Sustain* 15:11–28.
- Milder JC, et al. (2015) An agenda for assessing and improving conservation impacts of sustainability standards in tropical agriculture. *Conserv Biol* 29:309–320.
- Potts J, et al. (2014) *The State of Sustainability Initiatives Review 2014* (International Institute for Sustainable Development, Winnipeg, MB; International Institute for Environment and Development, London).
- Kamal S, et al. (2015) Conservation on private land: A review of global strategies with a proposed classification system. *J Environ Plann Manage* 58:576–597.
- Barnes AP, et al. (2009) Farmer perspectives and practices regarding water pollution control programmes in Scotland. *Agric Water Manage* 96:1715–1722.
- Barnes AP, et al. (2013) Comparing a 'budger' to a 'nudge': Farmer responses to voluntary and compulsory compliance in a water quality management regime. *J Rural Stud* 32:448–459.
- Christen B, et al. (2015) Can fuzzy cognitive mapping help in agricultural policy design and communication? *Land Use Policy* 45:64–75.
- Popp J, et al. (2007) The role of stakeholders' perceptions in addressing water quality disputes in an embattled watershed. *J Environ Monit Restor* 3:225–263.
- Azevedo AA, et al. (2017) Limits of Brazil's forest code as a means to end illegal deforestation. *Proc Natl Acad Sci USA* 114:7653–7658.
- Morefield PE, et al. (2016) Grasslands, wetlands, and agriculture: The fate of land expiring from the conservation reserve program in the Midwestern United States. *Environ Res Lett* 11:094005.
- Johnson KA, et al. (2016) Conservation Reserve Program (CRP) lands provide ecosystem service benefits that exceed land rental payment costs. *Ecosyst Serv* 18:175–185.
- Bennett MT (2008) China's sloping land conversion program: Institutional innovation or business as usual? *Ecol Econ* 65:699–711.
- Liu J, Li S, Ouyang Z, Tam C, Chen X (2008) Ecological and socioeconomic effects of China's policies for ecosystem services. *Proc Natl Acad Sci USA* 105:9477–9482.
- Pe'er G, et al. (2014) Agriculture policy. EU agricultural reform fails on biodiversity. *Science* 344:1090–1092.
- Banana AY, Sembajjiwe WG (2000) Successful forestry management: The importance of security of tenure and rule enforcement in Ugandan forests. *People and Forest: Communities, Institutions and Governance*, eds Gibson CC, McKean MA, Ostrom E (MIT Press, Cambridge, MA), pp 87–98.
- Agrawal A, et al. (2013) *Economic Contributions of Forests*. United Nations Forum on Forests, Background Paper No. 1, Tenth session, April 8–19, 2013. Instabul, Turkey. Available at citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.361.8278&rep=rep1&type=pdf. Accessed April 12, 2017.
- Wiyu AW, et al. (2014) Solving deforestation, protecting and managing key water catchments in Malawi using smart public and private partnerships. *J Sustain Dev* 8: 251–261.
- Millennium Challenge Account (MCA) (2012) *Malawi Assessment of the Energy Sector Report*. Millennium Challenge Account (Malawi Chapter) (Malawi Government, Lilongwe, Malawi).
- Namirembe S, et al. (2014) Co-investment paradigms as alternatives to payments for tree-based ecosystem services. *Curr Opin Environ Sustain* 6:89–97.
- Biermann F, et al. (2012) Transforming governance and institutions for global sustainability: Key insights from the Earth System Governance Project. *Curr Opin Environ Sustain* 4:51–60.
- Garnett T, et al. (2013) Agriculture. Sustainable intensification in agriculture: Premises and policies. *Science* 341:33–34.
- Phalan B, et al. (2016) CONSERVATION ECOLOGY. How can higher-yield farming help to spare nature? *Science* 351:450–451.
- Lambin EF, et al. (2014) Effectiveness and synergies of policy instruments for land use governance in tropical regions. *Glob Environ Chang* 28:129–140.
- Rueda X, Lambin E (2013) Responding to globalization: Impacts of certification on Colombian small-scale coffee growers. *Ecol Soc* 18:21.
- Brauman KA, Viart N (2016) Development of a regionally sensitive water-productivity indicator to identify sustainable practices for sugarcane growers. *Integr Environ Assess Manag* 12:811–820.
- Bonsucro (2014) *Bonsucro Production Standard Including Bonsucro EU Production Standard* (Bonsucro, London), Version 4.01.
- ISEAL (2014) *Setting Social and Environmental Standards* (ISEAL Alliance, London), Version 6.
- Tilman D, Balzer C, Hill J, Befort BL (2011) Global food demand and the sustainable intensification of agriculture. *Proc Natl Acad Sci USA* 108:20260–20264.
- Gassert FM, Reig P, Shiao T, Landis M, Luck M (2013) *Aqueduct Global Maps 2.0* (World Resources Institute, Washington, DC), Working paper.
- Olson DM, Dinerstein E (2002) The global 200: Priority ecoregions for global conservation. *Ann Mo Bot Gard* 89:199–224.
- Garrett RD, Carlson KM, Rueda X, Noojipady P (2016) Assessing the potential additional certification by the round table on responsible soybeans and the round table on sustainable palm oil. *Environ Res Lett* 11:45003.
- Blackman A, Rivera J (2011) Producer-level benefits of sustainability certification. *Conserv Biol* 25:1176–1185.
- Hatanaka M, Bain C, Busch L (2005) Third-party certification in the global agrifood system. *Food Policy* 30:354–369.
- Auld G, Gulbrandsen LH, McDermott CL (2008) Certification schemes and the impacts on forests and forestry. *Annu Rev Environ Resour* 33:187–211.
- Sachs JD (2012) From millennium development goals to sustainable development goals. *Lancet* 379:2206–2211.
- Monfreda C, Ramankutty N, Foley JA (2008) Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Global Biogeochem Cycles* 22:1–19.
- Siebert S, Döll P (2010) Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation. *J Hydrol (Amst)* 384:198–217.