

Neuroprotective effects of *Rosa damascena* extract on learning and memory in a rat model of amyloid- β -induced Alzheimer's disease

Ebrahim Esfandiary, Mohammad Karimipour, Mohammad Mardani, Mustafa Ghanadian¹, Hojjat Allah Alaei², Daryoush Mohammadnejad³, Abolghasem Esmaeili⁴

Department of Anatomical Sciences and Molecular Biology, ¹Department of Physiology, ²Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Isfahan University of Medical Sciences, Isfahan, ³Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, ⁴Department of Biology, Cells, Molecular Biology and Biochemistry Division, Faculty of Sciences, University of Isfahan, Isfahan, Iran

Abstract

Background: Alzheimer's disease (AD) is an age-related progressive neurodegenerative disease, which is characterized clinically by serious impairment in memory and cognition. Current medications only slow down the dementia progression and the present treatment one-drug one-target paradigm for anti-AD treatment appears to be clinically unsuccessful. Therefore, alternative therapeutic strategies are urgently needed. With respect to multifunctional and multitargeted characteristics of *Rosa damascena* via its effective flavonoids, we investigated the effects of *R. damascena* extract on behavioral functions in a rat model of amyloid- β (A- β)-induced Alzheimer's disease.

Materials and Methods: After preparation of the methanolic extract of the *R. damascena*, HPLC analysis and toxicity studies, median lethal dose (LD50) and dose levels were determined. For evaluation of baseline training behavioral performance, Morris water maze and passive avoidance tests were used. A- β was injected bilaterally into CA1 area of the hippocampus. Twenty-one days after injection of A- β , the first probe trial of the behavioral tests were used to confirm learning and memory impairment. To examine the potential effects of the extract on behavioral tasks, the second probe trials were performed after one month administration of *R. damascena* extract.

Results: Results showed that the *R. damascena* extract significantly improved the spatial and long-term memories in the extract-treated groups in a dose-dependent manner, as in the middle and high doses it had significant effect.

Conclusion: According to these results, we concluded that *R. damascena* can reverse behavioral deficits caused by A- β , and may provide a new potential option for prevention and treatment of the cognitive dysfunction in Alzheimer's disease.

Key Words: Alzheimer's disease, learning and spatial memory, long-term memory, *Rosa damascena*

Access this article online

Quick Response Code:



Website:

www.advbiores.net

DOI:

10.4103/2277-9175.161512

Address for correspondence:

Dr. Mohammad Karimipour, Department of Anatomical Sciences and Molecular Biology, Isfahan University of Medical Sciences, Isfahan, Iran.
E-mail: m_karimipour@resident.mui.ac.ir

Received: 22.01.2014, Accepted: 12.05.2014

INTRODUCTION

Alzheimer's disease (AD) is one of the most serious neurodegenerative diseases and the most common

Copyright: © 2015 Esfandiary. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Esfandiary E, Karimipour M, Mardani M, Ghanadian M, Alaei HA, Mohammadnejad D, et al. Neuroprotective effects of *Rosa damascena* extract on learning and memory in a rat model of amyloid- β -induced Alzheimer's disease. Adv Biomed Res 2015;4:131.

cause of dementia in aging population affecting approximately 26 million people worldwide, and whose prevalence has been calculated to triplicate by 2050.^[1,2] AD is characterized clinically by progressive impairment in memory, cognition, and behavioral functions.^[3] It is believed that AD initiates with synaptic impairment, neuronal loss, microglial cell proliferation, and is followed by inflammation.^[4,5] The neuropathological feature of AD is complex, including cerebral accumulation of extracellular amyloid- β (A- β) plaques and intraneuronal neurofibrillary tau tangles composed of dystrophic neuritis and hyperphosphorylated tau protein.^[6] According to the "amyloid hypothesis," the soluble A- β in the brain, plays an important role in the development of AD.^[7] Despite catastrophic increase of dementia patients worldwide, no effective treatment is available yet, although several acetylcholine esterase inhibitors such as donepezil, rivastigmine, and galantamine usually were used, but these medicines can only slow down the progression of dementia, rather than restoring brain function in a real clinical situation. Therefore, alternative and multitargeted therapeutic strategies are urgently needed to prevent or treat Alzheimer's disease.^[8,9] Because AD arises via multiple pathological or neurotoxic pathways, herbal medicines as a result of multifunction, multitarget characteristics have potential of optimum pharmaceutical and nutraceutical effects on AD patients.^[10]

Medicinal herbs-derived agents have different mechanisms from conventional drugs, which can be effective in clinical treatment. Herbs contain some of the most powerful natural antioxidants and bioactive secondary metabolites such as flavonoids, phenols, or phenolic components that make them valuable in their antioxidant and antiaging effects.^[11] *Rosa damascena* is a plant that belongs to genus *Rosa* and family Rosaceae. Rosaceae are well known as ornamental plants and have referred to the king of flowers. Some members of the Rosaceae family have long been used for food and medical purposes. Most of the studies showed that *R. damascena* as a natural plant with high source of flavonoids such as quercetin, kaempferol, myricetin, gallic acid, and their glycoside derivatives may have curable effect on treatment of diseases.^[11,12] It has been shown that essential oil of *R. damascena* retards the development of behavioral seizures in amygdala electrical kindling and possesses ability of contract kindling acquisition.^[13] There is considerable evidence suggesting that extract of the *R. damascena* significantly induces the neurite outgrowth and inhibits the A- β fibrilization and deposition in brain.^[14] Also it is shown that the *R. damascena* oil can relieve human stress and depression.^[15] Flavonoids perform a multiplicity of neuroprotective

functions within the brain, including neural protection against neurotoxin injuries, neuroinflammatory suppression and promotion of memory, learning, and cognitive function. These effects could be mediated by flavonoids that inhibit apoptosis, promote neuronal survival and synaptic plasticity. They can also promote cerebrovascular blood flow, angiogenesis, neurogenesis, and neuronal morphology.^[16] The aim of the current study was to determine the effects of methanolic extract of *R. damascena* on behavioral functions in a rat model of amyloid- β (1-42) induced Alzheimer's disease.

MATERIALS AND METHODS

Methanolic extraction of *R. damascena*

Dried fresh flowers of *R. damascena* Mill was purchased from Isfahan Pharmaceutical Sciences Research Center (Iran). It was ground to a fine powder (2 kg) and then macerated with methanol (10 L). Extraction procedure was performed three times and each time for 4 consecutive days with intermittent stirring (1 h stirring at every 12 h interval) using magnetic stirrer until the extract was light colored. The combined extracts were filtered and concentrated under reduced pressure in a rotary vacuum evaporator (200 mbar) in a water bath at 40°C. Dried gummy extract (380 g) was stored in 4°C until use.^[11]

HPLC analysis and extract standardization

High-performance liquid chromatographic (HPLC) analysis was performed on a Waters system, equipped with 515 HPLC pump, UV-Visible detector (2487 dual absorbance) operated at 365 nm, and millennium software for the determination of quercetin in the extract. After acid hydrolysis of 100 mg of the extract (1 h in 2N HCl, at 95°C), the hydrolyzed flavonoids were extracted through ethyl acetate to 5 mL. Then 20 μ L of the sample injected into a Nova-Pak C18, 3.9 \times 150 mm (Waters, Milford, MA, USA) using H₃PO₄ 10 mM in water (solvent A) and acetonitrile (solvent B) with gradient elution at a flow rate of 0.8 mL/min. A standard calibration curve in the range of 100–1000 μ g/mL for quantitative analysis was prepared using different concentrations of quercetin (Sigma Aldrich, USA) as standard material (100, 200, 500, and 1000 μ g/mL). The relationship between the concentration and peak-area of standard was measured using the minimum square method (R² value).^[17]

Acute toxicity studies and determination of median lethal dose (LD50)

Toxicity is defined as "the potential of a substance to exert a harmful effect on humans or animals, and a description of the effect and the conditions or concentration under which the effect takes place."^[18]

Acute toxicity is involved in the estimation of LD50 (the dose which has proved to be lethal (causing death) to 50% of the tested group of animals). Determination of acute oral toxicity is usually an initial screening step in the assessment and evaluation of the toxic characteristics of all compounds.^[19,20]

Assignment and housing of the animals for toxicity studies

Each animal was assigned a unique identification number. Thirty-six male Wistar rats (200–250 g) were housed six per cage. Six rats in each group at each dose level were chosen. In the present study, rats were gavaged via a gastric tube with increasing six series doses of *R. damascena* extract to get 50% lethality according to table 1. One dose being used per group. The animals were observed in several times up to 72 h for survival and morbidity.^[21,22] Median lethal dose (LD50) was calculated as given in table 1.

Dose levels and dose selection

This is based on the results of (LD50) finding test. At least three dose levels were used, spaced appropriately to produce test groups. For dose selection, 20% of (LD50) 6000 mg/kg was determined as the highest dose (1200 mg/kg) and decreasing spaced doses of 600 and 300 mg/kg were considered as middle and low doses, respectively, in our experimental study.^[23]

Animals and stereotaxic surgery

Male Wistar rats (200–250 g) were housed four per cage and maintained on a 12 h light–dark cycle in an air-conditioned constant temperature ($23 \pm 1^\circ\text{C}$) room, with food and water made available *ad libitum*.

The Ethics Committee for Animal Experiments at Isfahan University of Medical Sciences approved the study, and all experiments were conducted in accordance with the international guiding principles for biomedical research involving animals, revised in 1985. Initially, all animals were randomly divided into 6 groups ($n = 10$) for evaluation of baseline training performance in Morris water maze and passive avoidance tests. After spatial acquisition phase of Morris water maze and training (learning) phase of the passive avoidance test, animals were grouped as following: First group = control, Second group = sham, Third group = Alzheimer's disease + Normal saline, Fourth group = Alzheimer's disease + 300 mg of *R. damascena* extract, Fifth group = Alzheimer's disease + 600 mg of *R. damascena* extract and sixth group = Alzheimer's disease + 1200 mg of *R. damascena* extract. The protocol of experimental design is summarized in Figure 1.

Rats (groups 3rd to 6th) were anesthetized by intraperitoneal injection of chloral hydrate (350 mg/kg) and placed into stereotaxic device (Stoelting, Kiel, WI, USA). A heating pad was used to maintain body temperature at $36.5 \pm 0.5^\circ\text{C}$. An incision was made along the midline, the scalp was retracted, and the related area was cleaned and dried. In addition, lidocaine and epinephrine solution (0.4 mL) were injected in several locations surrounding the incision. In order to make minimum brain trauma during infusion, injections were performed very slowly (1 $\mu\text{L}/\text{min}$) using a microinjection pump. Amyloid- β (1-42) (Sigma) was dissolved in phosphate buffered saline (PBS 0.1 M) and aliquots at a concentration of 1 $\mu\text{g}/\mu\text{L}$ were stored at -20°C until used. For induction of aggregation, aliquots of amyloid- β (1-42) were incubated for 5 days at 37°C before administration. Then 3 $\mu\text{g}/3 \mu\text{L}$ of fibrillar A- β (1–42) or vehicle (PBS 0.1) were injected into the CA1 area of hippocampus at the following coordinates: Incisor bar -3.3 mm , -3.6 mm anterior-posterior to the bregma, 2.4 mm lateral to the sagittal suture, 2.8 mm dorsoventral from top of the skull bilaterally according to the atlas by Paxinos and Watson.^[24] The needle was kept in place for 5 min to allow the injected solutions and tissue to equilibrate and avoid the possible reflux

Table 1: Determination of (LD50)

Group N=6	Dose (mg/kg)	No. of animals dead in following times (h)						
		1	2	6	12	24	48	72
1	1000	0	0	0	0	0	0	0
2	2000	0	0	0	0	0	0	0
3	3000	0	0	0	0	0	0	0
4	4000	0	0	0	0	0	0	1
5	5000	0	0	0	0	0	1	1
6	6000	0	0	0	0	1	1	1




Figure 1: Experimental schedule of the research design during the course of the study

through the needle track. Incisions were ligated with silk thread. In sham group, same surgery was performed except that PBS was injected in CA1. No surgery was done in the control group.

Morris water maze

The circular tank (180 cm in diameter) was filled with water ($22 \pm 2^\circ\text{C}$) made opaque and was surrounded by a variety of extramaze cues. The tank was divided into four equal quadrants (northeast, northwest, southwest, and southeast) and four start positions were located at the intersection of the quadrants [Figure 2a]. A platform (12.5 cm in diameter) was placed in the northeast (the target quadrant) and submerged 2.0 cm below the water surface where it remained for all spatial trials.^[25-27] Twenty-four hours before water maze training, all rats were habituated to the water and apparatus. In the spatial acquisition phase, the rats learned to find a submerged platform using extramaze cues. Each rat participated in eight trials that were organized into two blocks of four trials (one trial/start position within a block). Each block was considered a separate test session and blocks were separated by 30 min. For each trial, the rat was given a maximum time of 60 s to locate the platform. After mounting the platform, the animals were allowed to remain there for 30 s, and then were placed in a holding cage for 30 s until the start of next trial. If the rat did not locate the platform within 60 s, it was guided to it by the experimenter. After completion of spatial acquisition phase, the animals were returned to their home cages until the initiation of probe trials in the test days. During the different phases of the maze, the animals were monitored by digital camera fixed 2 m above the maze and different parameters were analyzed using computer-based software (Radiab 1). Swim time (in s), swim distance (in cm), and swim speed (in cm/s) were recorded.

The first and second probe trials (spatial retention) 21 days after surgery and following extract treatment were performed, respectively. In the probe trials, the hidden platform was removed and animals were allowed to swim freely for 60 s. During the probe trials, the percentage time in target quadrant was calculated. Cued acquisition task conducted 20 min after completion of the probe trials in order to test their nonmnemonic aspects of water maze performance such as swimming ability, motivation, and visual ability. In this phase, the platform was elevated above the water surface and was placed in different positions, and rats were allowed to swim freely toward visible platform for 60 s. Twenty-one days after operation the 1st probe trial of Morris water maze and passive avoidance tests were performed and memory impairment was confirmed in groups 3–6.

Then dried *R. damascena* extract was dissolved in normal saline 0.9% +1% Tween 20, and then was administrated orally at 300, 600, 1200 mg/kg/day for 1 month via gavage in groups 4–6. Sham and 3rd groups received only normal saline at the same time. After *R. damascena* extract treatment 2nd probe trial of Morris water maze and passive avoidance tests were performed to evaluate the effects of extract on learning and spatial and long-term memories.

Passive avoidance test

The apparatus for the step-through passive avoidance test is an automated shuttle-box divided into an illuminated chamber and a dark chamber of the same size (20 × 25 × 30 cm each) by a wall with a guillotine door^[28-30] [Figure 2b]. In the present study, passive avoidance test was used as an index of long-term memory.

In the adaptation trial, each rat was trained to adapt to the step-through passive avoidance apparatus. The animal was placed into the illuminated chamber, facing away from the dark chamber. After 60 s, the door between these two boxes was opened and the rat was allowed to move into the dark chamber. The training trial was performed 24 h after the adaptation trial. In training trial, each rat was placed individually in the illuminated chamber, and once it entered the dark chamber an electric shock (40 V, 0.3 A, 2 s) was delivered to its feet through the floor grid. The rat was immediately removed and returned to cage. The first probe trial (retention trial) was conducted 21 days after surgery, in which rats were placed again in illuminated chamber and the interval (step-through latency) between placement in illuminated chamber and entry to the dark one was recorded. If the animal did not enter the dark chamber within 5 min, the test was terminated and the step through latency was recorded as 300 s. Memory impairment was confirmed in groups 3–6. Then *R. damascena* extract was applied as described previously. After *R. damascena* extract treatment, 2nd probe trial of passive avoidance was performed to evaluate the effects of extract on long-term memory.




Figure 2: (a) Tank of the Morris water maze and site of the platform. (b) Schematic representation of the passive avoidance test apparatus

Statistical analysis

Data were expressed as mean \pm SEM (standard error of mean) and were analyzed using either SPSS version 19 or GraphPad Prism[®] 5.0. software. Statistical analysis was performed using the following tests. The escape latencies, swim distance, and swim speed in the water maze were analyzed by two-way analysis of variance (ANOVA) for between-subject differences between groups and repeated measures (within subjects) for effects across block interval 1–2 (“BLOCK” effect). The probe trials data for percentage of time in target quadrant were analyzed by multivariate ANOVA followed by Tukey's honestly significant difference (HSD) test as a *post hoc* analysis. One-way ANOVA followed by Tukey's HSD multiple comparison test as a *post hoc* analysis were performed for evaluation of training trial and retention trials of passive avoidance test. A value of $P < 0.05$ was considered statistically significant.

RESULTS

HPLC standardization of the extract

Methanolic extraction of the *R. damascena* was performed and standardized using quercetin as the bioactive marker for standardization of the extract. Quercetin peak of extract appeared at a retention time of 13.48 min and using calibration curve, the extract

was standardized to contain 548.89 ± 20.23 mg/100 g of the total quercetin (0.55% w/w) [Figure 3a and b]. By using of millennium processing software, the calibration curve was determined by linear regression in the range of 100–1000 μ g/mL. The regression equation was $y = 86419x + 794975$, where x was the concentration of total quercetin in the extract (μ g/mL) with the correlation co-factor (R²) of 0.998 [Figure 3c].^[14]

Evaluation of the spatial acquisition phase of the Morris water maze before animal model induction

There were no significant differences in the mean time latency ($F(5, 54) = 0.041, P = 1 > 0.05$), mean swimming distance ($F(5, 54) = 0.128, P = 0.9 > 0.05$), and mean swimming speed ($F(5, 54) = 0.426, P = 0.8 > 0.05$) in blocks 1 and 2 of the spatial acquisition among all groups before application of A- β [Figure 4a–c, blocks 1 and 2], but repeated measure test showed a significant difference in escape latency, swimming distance, and swimming speed ($F(1, 54) = 74.96, P < 0.001$); ($F(1, 54) = 397.22, P < 0.001$); ($F(1, 54) = 20.14, P < 0.001$ respectively) between blocks 1 and block 2 were observed [Figure 4a–c, blocks 1 and 2] to locate the platform across blocks of trials, indicating spatial acquisition performance in all groups improved significantly during the training days and were able to learn the task.




Figure 3: (a) HPLC chromatogram of quercetin standard at 365 nm; 20 μ L of the quercetin standard (Sigma Aldrich, USA) in the range of 100–1000 μ g/mL was injected into a Nova-Pak C18, 3.9 \times 150 mm (Waters, Milford, MA, USA) using H_3PO_4 10 mM in water (solvent A) and acetonitrile (solvent B) with gradient elution at a flow rate of 0.8 mL/min. Quercetin peak appeared at a retention time of 13.64 min. (b) HPLC chromatogram of *Rosa damascena* extract at 365 nm. After acid hydrolysis of 100 mg of the extract (1 h in 2N HCl, at 95°C), the hydrolyzed flavonoids were extracted through ethyl acetate to 5 mL. Then 20 μ L of the sample was injected into a Nova-Pak C18, 3.9 \times 150 mm (Waters, Milford, MA, USA) using H_3PO_4 10 mM in water (solvent A) and acetonitrile (solvent B) with gradient elution at a flow rate of 0.8 ml/min. Quercetin peak of the extract appeared at a retention time of 13.48 min. (c): Calibration curve of quercetin using HPLC method and acetonitrile/water as mobile phase with pH adjusted to 2.3 at 365 nm. Using the millennium processing software, the calibration curve was determined by linear regression in the range of 100–1000 μ g/mL. The regression equation was $y = 86,419x + 7,94,975$

Evaluation of the training trial of the passive avoidance test before animal model induction

One-way ANOVA showed that the step-through latency did not differ in the training trial among the six groups before operation ($F(5, 54) = 0.069$, $P = 0.9 > 0.05$) [Figure 5].

R. damascena extract improved the spatial learning and memory parameters in a rat model of Alzheimer's disease

Multivariate ANOVA analysis showed significant differences in 1st probe trial among groups ($P < 0.001$) [Figure 6]. *Post hoc* Tukey's HSD analysis showed that, the mean percentage of time spent in target quadrant in A- β -injected groups (3rd to 6th) after 21 days surgery was significantly decreased in comparison to control and sham groups ($P < 0.001$), but no significant differences were demonstrated in mean percentage of time spent in target quadrant between control and sham groups, ($P = 0.9 > 0.05$). These results showed that A- β -induced learning and memory impairment in the rats. Multivariate ANOVA analysis showed that mean percentage of time spent in target quadrant in 2nd probe trial after one month *R. damascena* extract treatment was significantly increased among groups ($P < 0.001$) [Figure 6]. *Post hoc* Tukey's HSD analysis showed that, mean percentage of time spent in target quadrant was significantly increased in extract-treated groups (5th and 6th groups) relative to 3rd group, which had received normal saline ($P = 0.006 < 0.05$, $P < 0.001$, respectively), but mean percentage of time in target quadrant in 4th group did not significantly increase relative to 3rd group ($P = 0.3 > 0.05$). Also mean percentage of

time in target quadrant in 6th group was significantly increased in comparison to 4th ($P = 0.002 < 0.05$). Taken together, these results showed that *R. damascena* extract improved spatial learning and memory in a dose-dependent manner.

R. damascena extract improved the long-term memory parameters in a rat model of Alzheimer's disease

One-way ANOVA showed that significant differences in mean step-through latency among groups were observed following surgery in the 1st probe trial ($F(5, 54) = 275.94$, $P < 0.001$) [Figure 7]. *Post hoc* Tukey's HSD analysis showed that mean step-through latency reduced in A- β -treated groups (3rd to 6th) compared with control and sham groups ($F(5, 54) = 119.24$, $P < 0.001$). One-way ANOVA demonstrated that following extract treatment, step-through latency significantly increased among groups ($P < 0.001$) [Figure 7]. The mean step-through latency in groups 5 and 6 was significantly increased in comparison to 3rd group ($P < 0.001$). Mean step-through latency in 4th group did not significantly increase relative to 3rd group ($P = 0.07$). This parameter in 5th and 6th groups significantly increased when compared with 4th group ($P < 0.001$). All together these data show that *R. damascena* extract in medium (600 mg/kg) and high dose (1200 mg/kg) improved long-term memory in a dose-dependent manner.

DISCUSSION

The goodness of usual anti-AD drugs, is just a temporary relief of clinical symptoms and signs of dementia, but the efficacy of these cholinesterase inhibitors in AD treatment is not supported by any high-quality and long-lasting clinical trials.^[9] These medications cannot cure Alzheimer's dementia or stop the disease progression.^[31,32] Because AD is a multifactorial




Figure 4: Behavioral assessment. In spatial acquisition phase there were no significant differences in block 1 and 2 among the six groups before application of amyloid- β , but Repeated measure test showed a significant reduction in (a) escape latency ($P < 0.001$), (b) swimming distance ($P < 0.001$), and (c) swimming speed ($P < 0.001$) in 2nd block compared with 1st block to locate the platform across blocks of trials




Figure 5: In training trial of the Passive avoidance test, there were no significant differences in step-through latency among the six groups ($P = 0.9$)




Figure 6: Effects of the amyloid- β and *Rosa damascene* extract on spatial memory during the first (A, white color) and second (A, black color) probe trials, respectively, as measured by mean percentage time in target quadrant. ** $P < 0.01$; *** $P < 0.001$ different from the control and sham groups. ## $P < 0.01$, ### $P < 0.001$ different from the A β -injected group (Alz + NS), @ $P < 0.05$, @@@ $P < 0.001$ different from the A β -injected group (Alz + 300)

disease, using multifunctions and multitargeted agents have been considered extensively.^[33,34] Enormous medical combination therapies have been proposed to cure Alzheimer's disease.^[35] Herbal medicines as a result of multifunction, multitarget characteristics have potential of optimum pharmaceuticals and nutraceuticals effects on AD patients.^[10] Preclinical and epidemiological studies suggest that herbal drugs might be effective at reversing neurodegenerative pathology and age-related declines in neurocognitive performance.^[36] Herbal drugs may act to protect the brain in a number of ways, including by protection of vulnerable neurons, promotion of peripheral and cerebral vascular system, enhancement of existing neuronal function or by stimulating neuronal regeneration and inhibition of neuroinflammation.^[37] Moreover, medicinal herbs-derived drugs are natural and hence safer than synthetic drugs, and that a complex mixture of herbs can effectively treat complex diseases. These advantages may account for the sudden increase in herbal use in the last decade.^[38] In addition, some of the experimental reports suggest that some herbs may have neuroprotective effects against amyloid- β .^[39,40]

The present study examined the effects of a methanolic extract of *R. damascene* on behavioral deficits in a rat model of amyloid- β (A β 1-42)-induced Alzheimer's disease. Morris water maze and passive avoidance tests were used for examination of behavioral changes caused by amyloid- β and possible effect of *R. damascene* extract on learning, spatial memory, and long-term memories.^[29,30,41,42] The memory impairment induced by intra-CA1 injection of A β in our study was in agreement with Yamada *et al.*^[43] One month after *R. damascene* extract treatment, spatial memory




Figure 7: Effects of the amyloid- β and *Rosa damascene* extract on long-term memory during the first (B, white color) and second (B, black color) probe trials, respectively, as measured by mean time step-through latency. ** $P < 0.01$; *** $P < 0.001$ different from the control and sham groups. ## $P < 0.01$, ### $P < 0.001$ different from the A β -injected group (Alz + NS), @ $P < 0.05$, @@@ $P < 0.001$ different from the A β -injected group (Alz + 300)

and long-term memory were improved, whereas cognitive decline was reversed in a dose-dependent manner. The neuroprotective effect of *R. damascene* against cytotoxicity of A- β was investigated by Awale *et al.* *in vitro*. They showed that *R. damascene* extract significantly causes neurite outgrowth and suppresses the A β - induced atrophy and cell death. Moreover, the length of dendrite in the cells treated with *R. damascene* extract was similar to those of nerve growth factor (NGF)-treated cells.^[14] Significant effect of *R. damascene* on A β -injected groups in our study could be because of its multifactorial effects of bioactive flavonoids, polyphenols, and secondary active metabolites. Kumar *et al.* had detected and quantified some of polyphenols including gallic acid, rutin, quercitrin, myricetin, quercetin, and kaempferol, in methanolic extract of *R. damascene* by using improved HPLC analysis method.^[17] In agreement with Kumar *et al.*, our HPLC analysis showed that quercetin as bioactive compound exists in the *R. damascene* extract. Flavonoids comprise the most common group of polyphenolic compounds in the human diet and are found ubiquitously in plants.^[44,45] The neuroprotective effects of *R. damascene* were mediated by flavonoids that induce memory and cognitive improvement function via scavenging free radicals, and releasing neurotransmitter (acetylcholine). Other multifunctional effects of flavonoids include antioxidant, antiapoptotic, anti-inflammatory, anticholinesterase, antidepressant, and modulation of signaling pathway cascade^[44-48] and cerebrovascular blood flow improvement.^[49] Cellular and molecular mechanisms of flavonoids effects may carry out via protein kinase activation and phosphatase inhibition

that could lead to activation of cyclic AMP-response element-binding protein (CREB). CREB could increase blood flow, angiogenesis, neurogenesis, dendritic spine growth, neuronal communication, and synaptic plasticity via production and activation of neurotrophic factors such as brain-derived growth factor (BDNF), nerve growth factor (NGF), vascular endothelial growth factor (VEGF), and transform growth factor- β (TGF- β). Flavonoids prevent neurodegeneration and brain aging via reduction of induced nitric oxide synthase (iNOS), nitric oxide releasing, suppression inflammatory cytokines, such as TNF- α , IL-1 β .^[50,51] Quercetin as a flavonoid which is present in *R. damascena* attenuates microglia- and/or astrocyte-mediated neuroinflammation.^[52] Many neurodegenerative diseases pathogenesis including Alzheimer's disease correlate to oxidative stress enhancement in brain.^[16] Brain biological functions of flavonoids attribute to their antioxidant functions.^[53] Phenolic compounds are effective hydrogen donors, and because of this they are known as good antioxidants.^[54] Their high antioxidant potential is attributed to their capacity of scavenging reactive oxygen species (ROS), inhibiting lipid peroxidation, chelating metal ions, and other free radicals, which are originated from various cellular activities and lead to oxidative stress.^[55] The cholinergic system is involved in many physiological processes, including synaptic plasticity and learning and memory.^[56,57] Quercetin and rutin, natural compounds widely found in *R. damascena* and diet have anticholinesterase inhibition effects without severe side effects.^[58] Recent studies have focused on amyloid precursor protein (APP) proteolysis and A β -generation as potential targets for AD therapy.^[59] A β is generated from the APP by beta-site APP cleaving enzyme-1 (BACE-1, β -secretase) and γ -secretases through pro-amyloidogenic processing pathway, but α -secretase via nonamyloidogenic processing pathway inhibits A β production and promote A β degradation.^[60] Myricetin, one of the natural flavonoids has been found in *R. damascena* is a strong antioxidant and has a dual neuroprotective effect by activation of α -secretase and direct inhibition of β -secretase (BACE-1). As myricetin-bound BACE-1 does not hydrate APP at the β -cleavage site, A β production would be reduced. It has three hydrogen-binding groups, which stabilize BACE-1 binding for concerning AD drug therapy.^[60,61] Myricetin binds also to A β through its hydroxyl arms and traps A β hydrogen bond, which is necessary to form β -structures or oligomers.^[62] Myricetin via this mechanism prevents A β structural alteration from a random coil to β -sheet and inhibits A β fibril generation. The β -structure of A β forms soluble oligomers that are considered to be AD cytotoxic component.^[7,63] It is demonstrated that quercetin also disrupts the β -sheet structure to form random coils.^[61,64] Therefore

R. damascena with high concentration of quercetin, myricetin, and other flavonoids decreases level of A β (1–42) in brain and finally improves learning and memory.

CONCLUSION

In summary, *R. damascena* reversed the A β -associated memory abnormalities in a rat model of amyloid- β -induced Alzheimer's disease. Taken together, the diverse actions of individual natural components of the *R. damascena* on learning and memory represent a new promising way for generation of memory-enhancing drugs, although complementary experiments are still to be down for better characterizing this valuable effect and its underlying mechanisms, in order to provide a basis for proper clinical trials.

ACKNOWLEDGMENTS

This work was supported by Isfahan University of Medical Sciences. The authors thank Dr. Parham Reisi of Department of Physiology for his excellent technical assistance. None of the authors claim any conflict of interest.

REFERENCES

- Selkoe DJ. The molecular pathology of Alzheimer's disease. *Neuron* 1991;6:487.
- Lee JK, Jin HK, Endo S, Schuchman EH, Carter JE, Bae JS. Intracerebral transplantation of bone marrow-derived mesenchymal stem cells reduces amyloid-beta deposition and rescues memory deficits in Alzheimer's disease mice by modulation of immune responses. *Stem Cells* 2010;28:329-43.
- Selkoe DJ. Alzheimer's disease: Genes, proteins, and therapy. *Physiol Rev* 2001;81:741-66.
- Rogers J, Webster S, Lue LF, Brachova L, Civin WH, Emmerling M, et al. Inflammation and Alzheimer's disease pathogenesis. *Neurobiol Aging* 1996;17:681-6.
- Scheff SW, Price DA, Schmitt FA, DeKosky ST, Mufson EJ. Synaptic alterations in CA1 in mild Alzheimer disease and mild cognitive impairment. *Neurology* 2007;68:1501-8.
- Huang HC, Jiang ZF. Accumulated amyloid-beta peptide and hyperphosphorylated tau protein: Relationship and links in Alzheimer's disease. *J Alzheimers Dis* 2009;16:15-27.
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* 2002;297:353-6.
- Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med* 2010;362:329-44.
- Salloway S, Ferris S, Kluger A, Goldman R, Griesing T, Kumar D, et al. Efficacy of donepezil in mild cognitive impairment: A randomized placebo-controlled trial. *Neurology* 2004;63:651-7.
- Kim HG, Oh MS. Herbal medicines for the prevention and treatment of Alzheimer's disease. *Curr Pharm Des* 2012;18:57-75.
- Kalim MD, Bhattacharyya D, Banerjee A, Chattopadhyay S. Oxidative DNA damage preventive activity and antioxidant potential of plants used in Unani system of medicine. *BMC Complement Altern Med* 2010;10:77.
- Schiber A, Mihalev K, Berardini N, Mollov P, Carle R. Flavonol glycosides from distilled petals of *Rosa damascena* Mill. *Z Naturforsch C* 2005;60:379-84.
- Ramezani R, Moghimi A, Rakhshandeh H, Eftehadi H, Kheirabadi M. The effect of *Rosa damascena* essential oil on the amygdala electrical kindling seizures in rat. *Pak J Biol Sci* 2008;11:746-51.

14. Awale S, Tohda C, Tezuka Y, Miyazaki M, Kadota S. Protective Effects of *Rosa damascena* and its Active Constituent on A β (25-35)-induced Neuritic Atrophy. *Evid Based Complement Alternat Med* 2011;2011:131042.
15. Hongratanaworakit T. Relaxing effect of rose oil on humans. *Nat Prod Commun* 2009;4:291-6.
16. Spencer JP. The impact of fruit flavonoids on memory and cognition. *Br J Nutr* 2010;104:S40-7.
17. Kumar N, Bhandari P, Singh B, Gupta AP, Kaul VK. Reversed phase-HPLC for rapid determination of polyphenols in flowers of rose species. *J Sep Sci* 2008;31:262-7.
18. Yu KH, Nation RL, Dooley MJ. Multiplicity of medication safety terms, definitions and functional meanings: When is enough enough? *Qual Saf Health Care* 2005;14:358-63.
19. Akhila J, Shetty, Deepa Shyamjith, Alwar M. C. Acute toxicity studies and determination of median lethal dose. *Curr Sci* 2007;93:917-20.
20. Poole AF, Leslie GB. *A practical approach to toxicological investigations*. Cambridge, England: Cambridge University Press;1989.
21. Javan M, Ahmadiani A, Semnani S, Kamalinejad M. Antinociceptive effects of *Trigonella foenum-graecum* leaves extract. *J Ethnopharmacol* 1997;58:125-9.
22. Pant J, Deshpande SB. Acute toxicity of Bisphenol A in rats. *Indian J Exp Biol* 2012;50:425-9.
23. Jagetia G, Baliga M, Malagi K, Sethukumar Kamath M. The evaluation of the radioprotective effect of Triphala (an ayurvedic rejuvenating drug) in the mice exposed to γ -radiation. *Phytomedicine* 2002;9:99-108.
24. Paxinos G, Watson C. *The Rat Brain in Stereotaxic Coordinates: Hard Cover Edition*. United States: Academic press; 2007.
25. Morris RG, Garrud P, Rawlins JN, O'Keefe J. Place navigation impaired in rats with hippocampal lesions. *Nature* 1982;297:681-3.
26. de Quervain DJ, Roozendaal B, McGaugh JL. Stress and glucocorticoids impair retrieval of long-term spatial memory. *Nature* 1998;394:787-90.
27. Frick KM, Kim JJ, Baxter MG. Effects of complete immunotoxin lesions of the cholinergic basal forebrain on fear conditioning and spatial learning. *Hippocampus* 2004;14:244-54.
28. Venault P, Chapouthier G, de Carvalho LP, Simiand J, Morre M, Dodd RH, *et al.* Benzodiazepine impairs and beta-carboline enhances performance in learning and memory tasks. *Nature* 1986;321:864-6.
29. Mazzola C, Micale V, Drago F. Amnesia induced by beta-amyloid fragments is counteracted by cannabinoid CB1 receptor blockade. *Eur J Pharmacol* 2003;477:219-25.
30. Abdel-Aal RA, Assi AA, Kostandy BB. Rivastigmine reverses aluminum-induced behavioral changes in rats. *Eur J Pharmacol* 2011;659:169-76.
31. Scarpini E, Scheltens P, Feldman H. Treatment of Alzheimer's disease: Current status and new perspectives. *Lancet Neurol* 2003;2:539-47.
32. Cummings JL. Alzheimer's disease. *N Engl J Med* 2004;351:56-67.
33. Youdim MB, Buccafusco JJ. Multi-functional drugs for various CNS targets in the treatment of neurodegenerative disorders. *Trends Pharmacol Sci* 2005;26:27-35.
34. Pimplikar SW. Reassessing the amyloid cascade hypothesis of Alzheimer's disease. *Int J Biochem Cell Biol* 2009;41:1261-8.
35. Klafki HW, Staufienbiel M, Kornhuber J, Wilfgang J. Therapeutic approaches to Alzheimer's disease. *Brain* 2006;129:2840-55.
36. Manach C, Scalbert A, Morand C, Remesy C, Jimenez L. Polyphenols: Food sources and bioavailability. *Am J Clin Nutr* 2004;79:727-47.
37. Youdim KA, Joseph JA. A possible emerging role of phytochemicals in improving age-related neurological dysfunctions: A multiplicity of effects. *Free Radic Biol Med* 2001;30:583-94.
38. Raskin I, Ribnicky DM, Komarnytsky S, Ilic N, Poulev A, Borisjuk N, *et al.* Plants and human health in the twenty-first century. *Trends Biotechnol* 2002;20:522-31.
39. Kim DS, Kim JY, Han YS. Alzheimer's disease drug discovery from herbs: Neuroprotectivity from beta-amyloid (1-42) insult. *J Altern Complement Med* 2007;13:333-40.
40. Park SY, Kim DS. Discovery of natural products from *Curcuma longa* that protect cells from beta-amyloid insult: A drug discovery effort against Alzheimer's disease. *J Nat Prod* 2002;65:1227-31.
41. Jarrard LE. On the role of the hippocampus in learning and memory in the rat. *Behav Neural Biol* 1993;60:9-26.
42. Jhoo JH, Kim HC, Nabeshima T, Yamada K, Shin EJ, Jhoo WK, *et al.* Beta-amyloid (1-42)-induced learning and memory deficits in mice: Involvement of oxidative burdens in the hippocampus and cerebral cortex. *Behav Brain Res* 2004;155:185-96.
43. Yamada K, Takayanagi M, Kamei H, Nagai T, Dohniwa M, Kobayashi K, *et al.* Effects of memantine and donepezil on amyloid beta-induced memory impairment in a delayed-matching to position task in rats. *Behav Brain Res* 2005;162:191-9.
44. Joseph JA, Shukitt-Hale B, Denisova NA, Prior RL, Cao G, Martin A, *et al.* Long-term dietary strawberry, spinach, or vitamin E supplementation retards the onset of age-related neuronal signal-transduction and cognitive behavioral deficits. *J Neurosci* 1998;18:8047-55.
45. Joseph JA, Shukitt-Hale B, Denisova NA, Bielinski D, Martin A, McEwen JJ, *et al.* Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. *J Neurosci* 1999;19:8114-21.
46. Casadesus G, Shukitt-Hale B, Stellwagen HM, Zhu X, Lee HG, Smith MA, *et al.* Modulation of hippocampal plasticity and cognitive behavior by short-term blueberry supplementation in aged rats. *Nutr Neurosci* 2004;7:309-16.
47. Goyarzu P, Malin DH, Lau FC, Tagliatalata G, Moon WD, Jennings R, *et al.* Blueberry supplemented diet: Effects on object recognition memory and nuclear factor-kappa B levels in aged rats. *Nutr Neurosci* 2004;7:75-83.
48. Joseph JA, Shukitt-Hale B, Casadesus G. Reversing the deleterious effects of aging on neuronal communication and behavior: Beneficial properties of fruit polyphenolic compounds. *Am J Clin Nutr* 2005;81:313S-6.
49. Schroeter H, Heiss C, Balzer J, Kleinbongard P, Keen CL, Hollenberg NK, *et al.* (-)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proc Natl Acad Sci U S A* 2006;103:1024-9.
50. Spencer JP. The interactions of flavonoids within neuronal signalling pathways. *Genes Nutr* 2007;2:257-73.
51. Spencer JP. The impact of flavonoids on memory: Physiological and molecular considerations. *Chem Soc Rev* 2009;38:1152-61.
52. Chen JC, Ho FM, Pei-Dawn Lee C, Chen CP, Jeng KC, Hsu HB, *et al.* Inhibition of iNOS gene expression by quercetin is mediated by the inhibition of I κ B kinase, nuclear factor-kappa B and STAT1, and depends on heme oxygenase-1 induction in mouse BV-2 microglia. *Eur J Pharmacol* 2005;521:9-20.
53. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med* 1996;20:933-56.
54. Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic Res* 1995;22:375-83.
55. Bors W, Heller W, Michel C, Saran M. Flavonoids as antioxidants: Determination of radical-scavenging efficiencies. *Methods Enzymol* 1990;186:343-55.
56. Flood JF, Landry DW, Jarvik ME. Cholinergic receptor interactions and their effects on long-term memory processing. *Brain Res* 1981;215:177-85.
57. Power AE, Vazdarjanova A, McGaugh JL. Muscarinic cholinergic influences in memory consolidation. *Neurobiol Learn Mem* 2003;80:178-93.
58. Ahmed T, Gilani AH. Inhibitory effect of curcuminoids on acetylcholinesterase activity and attenuation of scopolamine-induced amnesia may explain medicinal use of turmeric in Alzheimer's disease. *Pharmacol Biochem Behav* 2009;91:554-9.
59. Haass C. Take five-BACE and the gamma-secretase quartet conduct Alzheimer's amyloid beta-peptide generation. *EMBO J* 2004;23:483-8.
60. Guo T, Hobbs DW. Development of BACE1 inhibitors for Alzheimer's disease. *Curr Med Chem* 2006;13:1811-29.
61. Shimmyo Y, Kihara T, Akaike A, Niidome T, Sugimoto H. Multifunction of myricetin on A β : Neuroprotection via a conformational change of A β

- and reduction of A beta via the interference of secretases. *J Neurosci Res* 2008;86:368-77.
62. McLaurin J, Kierstead ME, Brown ME, Hawkes CA, Lambermon MH, Phinney AL, *et al.* Cyclohexanehexol inhibitors of Abeta aggregation prevent and reverse Alzheimer phenotype in a mouse model. *Nat Med* 2006;12:801-8.
63. Mattson MP. Pathways towards and away from Alzheimer's disease. *Nature* 2004;430:631-9.
64. Shimmyo Y, Kihara T, Akaike A, Niidome T, Sugimoto H. Flavonols and flavones as BACE-1 inhibitors: Structure-activity relationship in cell-free, cell-based and in silico studies reveal novel pharmacophore features. *Biochim Biophys Acta* 2008;1780:819-25.

Source of Support: This study was supported by Isfahan University of Medical Sciences, **Conflict of Interest:** None declared.