

Editorial

Lessons from the results of three national antistigma programmes

Introduction

This supplement brings together results of evaluations of three national antistigma programmes: Time to Change in England (1–5), Opening Minds in Canada (6, 7) and ‘Hjärnkoll’ in Sweden (8). Started within a few years of each other, these programmes share several common features. They make extensive use of contact with people with mental health problems as an evidence-based stigma-reduction method, whether that contact is direct (face to face), or indirect (virtual such as through videos). All have focused on one or more target groups, whether these be young people (Koller in this supplement), employers (8, 9), police (10), health professionals (6) or medical students (11). All have undergone evaluation by academic researchers. The two European programmes use a public health approach to defining stigma, namely in terms of problems of knowledge, attitudes and behaviour, and in addition to target groups include the general population as a target (Hansson; Henderson in this supplement). However, ‘Hjärnkoll’ was initially delivered much more intensely in three regions of Sweden before the rest of the country was included, while in England, the aim from the start has been to target the general population as evenly as possible throughout the country. In Canada, the programme uses Link’s definition of stigma as the co-occurrence of its components: labelling, stereotyping, separation, status loss and discrimination (12), and selected four specific target groups (media, workers, youth and healthcare providers). Following this definition, the Canadian programme also emphasises structural change as a stigma-reduction strategy. To reach members of the target groups, the programme has worked with large numbers of community partners who deliver interventions in

communities across the country. Although all programmes are delivering contact-based education, there is considerable heterogeneity in delivery, which the evaluation team has exploited to identify the active ingredients of interventions, thus paving the way for fidelity criteria for future use (13) as well as toolkits to support a national scale-up. These differences among programmes mean that consideration of all three collectively allows for greater learning compared to consideration of each individually. In this supplement, we present papers from each evaluation and use this editorial to draw lessons across the results.

Both population and target group delivery can be effective at the national level

The positive results from all three evaluations demonstrate emphatically that there is more than one way to deliver an effective national antistigma programme. For Sweden and England, which have used mass media social marketing campaigns (8, 14) in addition to local initiatives and work with target groups, it is not possible to disentangle the different influences of these different components on the general population. However, it is possible that many people have been exposed only to the social marketing campaign and that the population level changes in the outcomes at least in part reflect this exposure. In England, stigma-related knowledge was relatively slow to emerge as a positive outcome (3) while in Sweden this was not the case (8). This may reflect differences in programme content, especially that of the social marketing campaign. After several years, however, it is clear that positive change has been sustained in all outcomes in both countries. In Canada, work with specific target groups has been delivered across the country in preparation for a national scale-up, which is currently underway for youth groups through national summits. Population-level change is not expected to occur until full scale-up has occurred. At this point, the evaluation has contributed considerably to the evidence base for

Editorial

contact-based education, in terms of measurement (15); identification of the key aspects of stigma to be targeted within specific populations (6); and identification of key effective programme ingredients (13).

There may be gender differences in responses to antistigma programmes

Within the general population in England, the lack of change among men compared to women in terms of reported contact with people with mental health problems is striking (3). It suggests that while there may be an impact on men's behaviour of the campaign, this is not detectable to other men; for example, they may disclose only to female friends or partners, so that only women report an increase in contact. Another possibility is that women have increased in their ability to recognise others' mental health problems, leading to greater reported contact, while this has not occurred for men. Among Canadian youth, a positive response to the contact-based intervention among males was highly influenced by whether they self-reported experience of a mental health problem, whereas this was not an influential factor among females (7). Consistent with the population in England, young males were significantly less likely than females to report prior contact with a person with a mental illness. As familiarity is a consistent predictor of positive outcomes (14, 16, 17), it seems possible that lack of contact could hinder further positive change among men in terms of any or all of stigma-related knowledge, attitudes and desire for social distance. These findings suggest that gender-based approaches to stigma reduction may be necessary.

Campaigns and their evaluation must attend to structural discrimination

The positive change in public attitudes mirrors the changes in mental health service users' experiences of discrimination on the part of people with whom they have informal relationships, as friends, family members and when dating (1). These changes contrast in reported experiences of discrimination in several important domains including physical health care, welfare benefits (2) and housing. While education, both with and without contact (13, 18), can be effective at least in the short term with respect to health professionals' attitudes, other studies in health care settings suggest organisational level changes are needed (19–22). Likewise, positive changes in the

way editors and journalists cover mental health topics are small and do not yet show a consistent pattern. Within coverage broadly categorised as stigmatising, they may be replacing one form of negative coverage for another by shifting their focus from violence to focussing on people with mental health problems as 'hopeless victims'. This stereotype risks fuelling within the public the same kind of therapeutic pessimism shown to be a strong component of health professionals' stigma (6, 18). Service users' experiences when seeking or in work lie somewhere between these patterns of positive and lack of change. After an initial improvement (23), the changes in the domain of employment became non-significant, re-emerging later (1). In the UK, other evidence suggests that employers are increasingly aware of the need to consider employees' mental health and to comply with the Equality Act, for example with respect to providing reasonable adjustments (9). However, it is also possible that people with mental health problems were particularly adversely affected by the economic recession in the UK and the stagnation that followed (24). Taken together, the results of our qualitative and quantitative work with mental health service users suggest that organisations that provide services need to consider discrimination both as it affects those of their service users with mental health problems and those of their employees.

Acknowledgements

During Phase 1 of Time to Change, CH was funded in relation to a National Institute for Health Research (NIHR) Applied Programme grant awarded to the South London and Maudsley NHS Foundation Trust (PI Graham Thornicroft). CH was supported during phases 1 and 2 by the grants to Time to Change from Big Lottery, the UK Department of Health and Comic Relief. CH was also funded by grants from the NIHR, Maudsley Charity and Guy's and St Thomas's Charity.

Funding

Time to Change programme, grants from UK Department of Health and Comic Relief.

Declaration of interest

CH has received a consulting fee from Lundbeck. MK and HS have received funding from the Mental Health Commission of Canada.

C. Henderson¹, H. Stuart², L. Hansson³

¹Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK ²Centre for Health Services and Policy Research,

Abramsky Hall, Queen's University, Kingston, ON, Canada and
³*Department of Health Sciences, Lund University, Lund, Sweden*
E-mail: Claire.I.henderson@kcl.ac.uk

References

1. CORKER E, HAMILTON S, ROBINSON E et al. Viewpoint survey of mental health service users' experiences of discrimination in England 2008–14. *Acta Psychiatr Scand* 2016;**134**:6–13.
2. HAMILTON S, PINFOLD V, COTNEY J et al. Qualitative analysis of mental health service users' reported experiences of discrimination. *Acta Psychiatr Scand* 2016;**134**:14–22.
3. HENDERSON C, ROBINSON E, EVANS-LACKO S et al. Public knowledge, attitudes, social distance and reported contact regarding people with mental illness 2009–2015. *Acta Psychiatr Scand* 2016;**134**:23–33.
4. OSUMILI B, HENDERSON C, CORKER E et al. The economic costs of mental health-related discrimination. *Acta Psychiatr Scand* 2016;**134**:34–44.
5. RHYDDERCH D, KROOUPA A-M, SHEFER G et al. Changes in newspaper coverage of mental illness from 2008 to 2014 in England. *Acta Psychiatr Scand* 2016;**134**:45–52.
6. KNAAK S, PATTEN S. A grounded theory model for reducing stigma in health professionals in Canada. *Acta Psychiatr Scand* 2016;**134**:53–62.
7. KOLLER M, STUART H. Reducing stigma in high school youth. *Acta Psychiatr Scand* 2016;**134**:63–70.
8. HANSSON L, STJERNSWÄRD S, SVENSSON B. Changes in attitudes, intended behaviour and mental health literacy in the Swedish population 2009–2014. An evaluation of a national anti-stigma programme. *Acta Psychiatr Scand* 2016;**134**:71–79.
9. HENDERSON C, WILLIAMS P, LITTLE K, THORNICROFT G. Mental health problems in the workplace: changes in employers' knowledge, attitudes and practices in England 2006–2010. *Br J Psychiatry* 2013;**202**(Suppl 55):s70–s76.
10. HANSSON L, MARKSTROM U. The effectiveness of an anti-stigma intervention in a basic police officer training programme: a controlled study. *BMC Psychiatry* 2014;**14**:55.
11. FRIEDRICH B, EVANS-LACKO S, LONDON J, RHYDDERCH D, HENDERSON C, THORNICROFT G. Anti-stigma training for medical students – the education not discrimination project. *Br J Psychiatry* 2013;**202**(Suppl 55):s89–s94.
12. LINK BG, PHELAN JC. Conceptualizing stigma. *Annu Rev Sociol* 2001;**27**:363–385.
13. KNAAK S, MODGILL G, PATTEN SB. Key ingredients of anti-stigma programs for health care providers: a data synthesis of evaluative studies. *Can J Psychiatry* 2014;**59** (10 Suppl 1):S19–S26.
14. EVANS-LACKO S, MALCOLM E, WEST K et al. Influence of Time to Change's social marketing interventions on stigma in England 2009–2011. *Br J Psychiatry* 2013;**202**:s77–s88.
15. MODGILL G, PATTEN SM, KNAAK S, KASSAM A, SZETO AC. Opening minds stigma scale for health care providers (OMS-HC): examination of psychometric properties and responsiveness. *BMC Psychiatry* 2014;**14**:120.
16. EVANS-LACKO S, HENDERSON C, THORNICROFT G. Public knowledge, attitudes and behaviour regarding people with mental illness in England 2009–2012. *Br J Psychiatry* 2013;**202**:s51–s57.
17. ILIC N, HENDERSON H, HENDERSON C, EVANS-LACKO S, THORNICROFT G. Attitudes towards mental illness. In: CRAIG R, FULLER E, MINDELL J, eds. *Health survey for England 2014: health, social care and lifestyles*. London: Health and Social Care Information Centre, 2016:1–15.
18. HENDERSON C, NOBLETT J, PARKE H et al. Mental health-related stigma in health care and mental health-care settings. *Lancet Psychiatry* 2014;**1**:467–482.
19. MOLL S, EAKIN JM, FRANCHE RL, STRIKE C. When health care workers experience mental ill health: institutional practices of silence. *Qual Health Res* 2013;**23**:167–179.
20. SHEFER G, HENDERSON C, HOWARD LM, MURRAY J, THORNICROFT G. Diagnostic overshadowing and other challenges involved in the diagnostic process of patients with mental illness who present in emergency departments with physical symptoms—a qualitative study. *PLoS One* 2014;**9**:e111682.
21. SHEFER G, CROSS S, HOWARD LM, MURRAY J, THORNICROFT G, HENDERSON C. Improving the diagnosis of physical illness in patients with mental illness who present in emergency departments: consensus study. *J Psychosom Res* 2015;**78**:346–351.
22. van NIEUWENHUIZEN A, HENDERSON C, KASSAM A et al. Emergency Department staff views and experiences on diagnostic overshadowing related to people with mental illness. *Epidemiol Psychiatr Sci* 2013;**22**:255–262.
23. HENDERSON C, CORKER E, LEWIS-HOLMES E, HAMILTON S, FLACH C, ROSE D. Reducing mental health related stigma and discrimination in England: one year outcomes of the Time to Change Programme for service user-rated experiences of discrimination. *Psychiatr Serv* 2012;**63**:451–457.
24. EVANS-LACKO S, KNAPP M, McCRONE P, THORNICROFT G, MOITABAI R. The mental health consequences of the recession: economic hardship and employment of people with mental health problems in 27 European countries. *PLoS One* 2013;**8**:e69792.