

Analysis of model replication origins in *Drosophila* reveals new aspects of the chromatin landscape and its relationship to origin activity and the prereplicative complex

Jun Liu, Kristopher McConnell, Michael Dixon, and Brian R. Calvi

Department of Biology, Indiana University, Bloomington, IN 47405

ABSTRACT Epigenetic regulation exerts a major influence on origins of DNA replication during development. The mechanisms for this regulation, however, are poorly defined. We showed previously that acetylation of nucleosomes regulates the origins that mediate developmental gene amplification during *Drosophila* oogenesis. Here we show that developmental activation of these origins is associated with acetylation of multiple histone lysines. Although these modifications are not unique to origin loci, we find that the level of acetylation is higher at the active origins and quantitatively correlated with the number of times these origins initiate replication. All of these acetylation marks were developmentally dynamic, rapidly increasing with origin activation and rapidly declining when the origins shut off and neighboring promoters turn on. Fine-scale analysis of the origins revealed that both hyperacetylation of nucleosomes and binding of the origin recognition complex (ORC) occur in a broad domain and that acetylation is highest on nucleosomes adjacent to one side of the major site of replication initiation. It was surprising to find that acetylation of some lysines depends on binding of ORC to the origin, suggesting that multiple histone acetyltransferases may be recruited during origin licensing. Our results reveal new insights into the origin epigenetic landscape and lead us to propose a chromatin switch model to explain the coordination of origin and promoter activity during development.

Monitoring Editor
Yixian Zheng
Carnegie Institution

Received: May 10, 2011

Revised: Oct 26, 2011

Accepted: Oct 27, 2011

INTRODUCTION

Efficient duplication of large eukaryotic genomes requires that DNA replication initiate from multiple origins. In multicellular eukaryotes, however, it remains largely unknown how certain genomic loci are selected to be active origins of DNA replication; a DNA consensus for origins has yet to emerge. Moreover, the selection of origin loci and their time of initiation during S phase change during develop-

ment (Mechali, 2010). Current evidence suggests that chromatin modifications play a major role in the developmental regulation of origins. Here we investigate the epigenetic regulation of the well-defined model origins that mediate developmental gene amplification during *Drosophila* oogenesis.

The proteins and mechanisms that regulate origins during the cell cycle are conserved in eukaryotes (Remus and Diffley, 2009). During early G1 phase, a prereplicative complex (preRC) assembles onto origins, preparing them for replication (Diffley *et al.*, 1995). Assembly of the preRC begins with the binding of the origin recognition complex (ORC) to DNA, which recruits Cdc6 and Cdt1, both of which are required for loading of the minichromosome maintenance (MCM) helicase complex onto origin DNA (Yan *et al.*, 1991; Bell *et al.*, 1993; Chong *et al.*, 1995; Cocker *et al.*, 1996; Maiorano *et al.*, 2000; Nishitani *et al.*, 2000). Once the preRC is assembled, the origin is "licensed" but has yet to initiate DNA replication. During different times of S phase, subsets of these preRCs are activated by cyclin-dependent kinase (CDK) and Cdc7 kinase, which results in the association of other proteins to the origin, the initiation of DNA replication, and departure of preRC proteins from the origin (Labib, 2010). The preRC is prevented from reassembling onto origins until

This article was published online ahead of print in MBoC in Press (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E11-05-0409>) on November 2, 2011.

The authors declare that they have no conflict of interest.

Address correspondence to: Brian R. Calvi (bcalvi@indiana.edu).

Abbreviations used: ACE1, amplification control element on chromosome X; ACE3, amplification control element on chromosome 3; BrdU, bromodeoxyuridine; CDK, cyclin-dependent kinase; ChIP, chromatin immunoprecipitation; cp, chorion protein; DAFC, *Drosophila* Amplicon in Follicle Cells; GFP, green fluorescent protein; HAT, histone acetyltransferase; HDAC, histone deacetylase; MCM, minichromosome maintenance; ORC, origin recognition complex; preRC, prereplicative complex.

© 2012 Liu *et al.* This article is distributed by The American Society for Cell Biology under license from the author(s). Two months after publication it is available to the public under an Attribution–Noncommercial–Share Alike 3.0 Unported Creative Commons License (<http://creativecommons.org/licenses/by-nc-sa/3.0>). "ASCB," "The American Society for Cell Biology," and "Molecular Biology of the Cell" are registered trademarks of The American Society of Cell Biology.

the next cell cycle, thereby ensuring that the genome duplicates once per cell division (Arias and Walter, 2007). Defects in preRC regulation can result in DNA damage, genome instability, human developmental abnormalities, and cancer (Lengronne and Schwob, 2002; Hook et al., 2007; Liontos et al., 2007; Mehrotra et al., 2008; Green et al., 2010; Bicknell et al., 2011; Guernsey et al., 2011).

Despite the conserved mechanism of origin regulation, it remains unclear in multicellular eukaryotes how certain regions of the genome are selected to be sites of preRC assembly and activation (Mechali, 2010). Although only a handful of metazoan origins have been analyzed both genetically and molecularly, the emerging theme is that most are large and modular, with functional DNA elements spread over several to tens of kilobases. Genomic approaches have mapped thousands of preRC-binding sites and active origins within several multicellular genomes, including *Drosophila* and human (Cadoret, 2008; Sequeira-Mendes, 2009; Gilbert, 2010; Hansen et al., 2010; Karnani et al., 2010; MacAlpine et al., 2010). Nevertheless, a strict DNA consensus for origins has yet to emerge (Aladjem, 2007). In fact, ORC does not display sequence specificity for DNA binding in vitro, although it does prefer AT-rich sequences and negatively supercoiled DNA (Bielinsky et al., 2001; Chesnokov et al., 2001; Vashee et al., 2003; Remus et al., 2004). In contrast, there are thousands of preferred sites for ORC binding and replication initiation on native chromosomes, whose location and S phase timing change during development (Huberman, 1968; Lima-de-Faria and Jaworska, 1968; Mechali, 2010). An important remaining question, therefore, is what specifies genomic loci to be preferred sites for preRC assembly and activation in different cells.

We have been studying developmental gene amplification in the follicle cells of the *Drosophila* ovary as a model for origin structure and regulation in a developmental context. Amplification is a local increase in gene copy number due to site-specific rereplication from origins at two loci that encode eggshell (chorion) proteins on the X (*Drosophila* Amplicon in Follicle Cells-7F, DAFC-7F) and third chromosome (DAFC-66D) and at four other, recently identified loci (DAFC-22B, DAFC-30B, DAFC-34B, and DAFC-62D), some of which encode proteins that assist vitelline membrane and eggshell synthesis (Spradling, 1981; Calvi et al., 1998; Calvi, 2006; Claycomb et al., 2004; Claycomb and Orr-Weaver, 2005; Kim et al., 2011). The amplicon origins are bound by a preRC and regulated by both CDK2 and CDC7 kinases (Calvi et al., 1998; Austin et al., 1999; Landis and Tower, 1999; Whittaker et al., 2000; Schwed et al., 2002). The amplicon origins therefore share many attributes with origins that govern normal genomic replication, and the analysis of these model origins has provided new insights into origin structure and regulation (Claycomb and Orr-Weaver, 2005; Calvi, 2006).

We previously showed that nucleosome acetylation at the amplicon origins contributes to their developmental specificity and efficiency (Aggarwal and Calvi, 2004), consistent with results from others (Lewis et al., 2004; Hartl et al., 2007). A general relationship between active origins and histone acetylation has been supported by genome-wide mapping and the detailed analysis of a few origins in multiple organisms, although this correlation is not perfect (Kim et al., 2003; Danis et al., 2004; Cadoret, 2008; Hiratani et al., 2008, 2010; Schwaiger et al., 2009, 2010; Bell et al., 2010; Eaton et al., 2010a; Gilbert, 2010; MacAlpine et al., 2010). Experiments from yeast to human have also identified some of the histone acetyltransferases (HATs) and histone deacetylases (HDACs) that can influence preRC assembly and the time of origin activation in S phase (Iizuka and Stillman, 1999; Iizuka et al., 2009; Vogelauer et al., 2002; Aggarwal and Calvi, 2004; Stedman et al., 2004; Jorgensen et al., 2007; Crampton et al., 2008; Fox and Weinreich, 2008; Wu and Liu,

2008; Schwaiger et al., 2009; Miotto and Struhl, 2010; Wong et al., 2010). Another attribute of origins in eukaryotes is that they coincide with “nucleosome free regions,” which are now known to represent domains of dynamic nucleosome–DNA association (Simpson, 1990; Lipford and Bell, 2001; Urnov et al., 2002; Mito et al., 2007; Eaton et al., 2010a, 2010b; MacAlpine et al., 2010; Muller et al., 2010). Current evidence suggests, therefore, that both nucleosome modification and position locally influence origin activity. Despite these advances, the molecular mechanisms by which chromatin influences origin function remain poorly defined.

Here we extend our analysis of the epigenetic regulation of the model amplification origins. Our results reveal new aspects of the origin epigenetic landscape and the relationship connecting histone acetylation, ORC binding, and origin activity.

RESULTS

Dynamic acetylation of multiple histone lysine residues is associated with DAFC-66D developmental timing

Site-specific rereplication at six origin loci results in developmental gene amplification during *Drosophila* oogenesis. These origins become active in somatic follicle cells at precisely stage 10B of oogenesis, a time when other origins are not active and genomic replication has ceased, and therefore represents an extreme form of origin developmental specificity (Calvi et al., 1998; Claycomb and Orr-Weaver, 2005). Previous analysis indicated that the N-terminal tails of histones H3 and H4 are hyperacetylated at the origins, and this hyperacetylation stimulates origin activity (Aggarwal and Calvi, 2004; Hartl et al., 2007). These experiments used antibodies against polyacetylated H3 and H4, as well as those raised against specific acetylated lysine residues. It is now widely appreciated that some commercially available antibodies are not as specific as once believed (Egelhofer et al., 2011). Therefore, to further explore the mechanism of origin epigenetic regulation, we used antibodies that have been recently validated by the modENCODE research consortium and, where possible, used multiple antibodies from different suppliers against the same modification (Egelhofer et al., 2011).

To determine whether these antibodies label active amplicon origins, we used them for immunofluorescence labeling of fixed ovaries. These ovaries were first incubated in bromodeoxyuridine (BrdU) to detect exclusive replication from the amplicon origins, which appears as distinct foci of BrdU incorporation in follicle cell nuclei beginning in stage 10B (Calvi et al., 1998; Calvi and Spradling, 2001). The acetylated histone antibodies displayed a general labeling throughout the nucleus up until stage 10A, just before the start of amplification. Figure 1 shows the results for anti-H4K16Ac and anti-H4K12Ac labeling (Figure 1, A–A’ and D–D’). From stage 10B to stage 11/12, however, many of the acetylated histone antibodies strongly labeled the amplicon BrdU foci, in addition to there being a lower level of nucleus-wide labeling (Table 1 and Figure 1, B–B’ and E–E’). Although multiple amplicon BrdU foci were often labeled, the most prominent labeling corresponded to the DAFC-66D locus, which amplifies to the highest DNA copy number (~64-fold) and is the largest BrdU focus (Figure 1, B–B’ and E–E’; Calvi et al., 1998). Consistent with our previous results, the labeling of amplicon foci by acetylated histone antibodies often appeared as a bar in the center of a larger bar of BrdU, which represents waves of replication forks emanating bidirectionally outward from the origins (Figure 1, B–B’; Aggarwal and Calvi, 2004). This pattern suggests that acetylation occurs near the center of the replicon and does not simply represent acetylation of newly deposited nucleosomes behind the




FIGURE 1: Hyperacetylation of H4K16 and H4K12 at active amplification origins. Follicle cells were colabeled with α -BrdU (A'–F') and α -H4K16ac (A–C) or α -H4K12ac (D–F) from oogenesis stages 10A (A–A'' and D–D'), 10B (B–B'' and E–E''), and 12 (C–C'' and F–F''). Merged images are shown in A''–F''. Arrows in B–B'' indicate a single amplicon focus corresponding to DAFC-66D, which is enlarged in the insets. Scale bars, 10 μ m; inset scale bars, 1 μ m.

replication forks. Although other genomic loci in stage 10B/11 follicle cells were labeled by acetylated histone antibodies, the relative level of fluorescence labeling at amplicon loci was much greater, which we previously showed is not an artifact of increased DNA copy number during amplification (Aggarwal and Calvi, 2004). Labeling for these different types of histone acetylation rapidly disappeared at DAFC-66D in early stage 12, a time when the origin shuts off, ORC departs, but forks continue to migrate outward (Figure 1, C–C'' and F–F''; Royzman *et al.*, 1999; Aggarwal and Calvi, 2004). This further argues that this acetylated histone labeling does not correspond to chromatin assembly behind the replication fork. Overall, the immunofluorescence results suggest that nucleosomes near the origins are hyperacetylated on multiple lysine residues during developmental stages when the origin is active.

We next turned to chromatin immunoprecipitation (ChIP) to analyze chromatin modification at the origins, a method that is of higher resolution and more sensitive than immunofluorescence labeling. It has the added advantage that the signal is normalized to input DNA, and, therefore, acetylation per chromatin fiber can be compared among amplicon loci that amplify to different DNA copy numbers. We also quantified the relative level of acetylation at the origins by normalizing the ChIP–real-time quantitative PCR (qPCR) signal to different nonorigin control loci measured in parallel (Supplemental Figure S1). To evaluate the developmental dynamics of chromatin modification, thousands of egg chambers from different stages of oogenesis were mass isolated, followed by hand selection of specific developmental stages and purification of follicle cell nuclei. We initially focused on DAFC-66D because it is the best-characterized amplicon origin. At this origin, the 320–base pair ACE3 and the 840–base pair Ori- β are both necessary and sufficient for origin function and contain preferred binding sites for the ORC (Figure 2A; Orr-Weaver *et al.*, 1989; Austin *et al.*, 1999; Zhang and Tower, 2004; Calvi, 2006). Although both of these elements bind ORC, Ori- β is the preferred site of replication initiation (~80% of the time; Delidakis and Kafatos, 1989; Heck and Spradling, 1990; Zhang and Tower, 2004). Each of these origin elements is immediately upstream of chorion protein genes (cp18 and cp15) but can be functionally separated from its

promoters (Figure 2A; (Orr-Weaver and Spradling, 1986). Initiation from the DAFC-66D origin occurs from stage 10B to early stage 12 (~7 h), after which the origin shuts off and nearby promoters turn on in stage 12/13 (Griffin-Shea *et al.*, 1982; Orr *et al.*, 1984; Royzman *et al.*, 1999).

ChIP-qPCR showed that multiple lysine residues are hyperacetylated at DAFC-66D, confirming the results of immunofluorescence labeling. Similar to the immunofluorescence results, other loci in follicle cells were immunoreactive to the acetylated histone antibodies in ChIP assays. The level of hyperacetylation at DAFC-66D, however, was ~20–60 times greater relative to two other nonorigin loci, using antibodies against different acetylated lysines in the tail of histone H4 (K5, K8, K12, and K16; Table 1, Figure 2, B–E, and Supplemental Figure S1). We previously reported hyperacetylation on lysines in the histone H3 tails at the amplicon and confirmed this result using antibodies against H3K9/14Ac (Aggarwal and Calvi, 2004; Table 1). We also obtained evidence for acetylation on lysine 56 within the core of histone H3, which was the most enriched of any modification in stage 10 (Figure 2F). Control ChIP experiments with antibodies against total histone H3 did not give evidence for enrichment at the amplicons (Supplemental Figure S2). ChIP with antibodies against marks associated with repressive chromatin (e.g., H3K27me3) also did not give evidence for enrichment at the DAFC-66D origin (Supplemental Figure S3A). Analysis of stages before, during, or after the origin is active showed that all the acetylation marks had similar developmental timing. Acetylation on all residues was low in early developmental stages (stage \leq 8) when the origin is inactive, peaked in stage 10/11 when the origin was active, and then rapidly declined in stages 12 and 13 when ORC departs and the origin shuts off (Figure 2, B and F). These results reveal that multiple types of nucleosome hyperacetylation are temporally correlated with developmental origin activity during oogenesis.

Antibody	Immunostaining ^a	ChIP ^b
H4K5ac	+	+
H4K8ac	+	+
H4K12ac (Active Motif)	+	+
H4K12ac (Millipore)	+	+
H4K16ac (Active Motif)	+	+
H4K16ac (Millipore)	f	+
H3K9/14ac	n.d.	+
H3K56ac	+	+
H3K4me3	f	–
H3K27me3	–	–

^a+, BrdU foci labeled; –, BrdU foci not labeled; f, antibody failed to label fixed tissue; n.d.: not determined.

^b+, enriched over control loci; –, not enriched.

^cFrom Aggarwal and Calvi (2004).

TABLE 1: Summary of immunostaining and ChIP results for histone modifications at DAFC-66D in stage 10B follicle cells.




FIGURE 2: Developmentally dynamic acetylation of nucleosomes on multiple lysine residues correlates with activity of the DAFC-66D origin. (A) Diagram of DAFC-66D locus. Hatched boxes represent essential elements ACE3 and Ori- β (middle); the location of ChIP-

Nucleosome acetylation extended across the DAFC-66D origin, with a peak adjacent to the major site of replication initiation

We had used multiple primer pairs for ChIP-qPCR to define the profile of acetylation at DAFC-66D and its relationship to functional origin regions. Hyperacetylation of multiple lysines was not restricted to the essential 320-base pair ACE3 and 840-base pair Ori- β but was at least five- to 15-fold enriched for the different antibodies over a 7-kb region encompassing the four chorion protein genes (Figure 2). Although nucleosome acetylation was significantly enriched across this domain, a distinct peak of acetylation on multiple lysines was detected between ACE3 and Ori- β in the body of the cp18 gene (Figure 2). This peak of acetylation is not an artifact of copy number or ChIP method, because all signals are normalized to input and a nonorigin control locus. The data suggest that nucleosomes are hyperacetylated in a domain at the DAFC-66D locus and that within the population of follicle cells acetylation is most frequent on nucleosomes adjacent to one side of the major site of replication initiation in Ori- β .

Nucleosome acetylation is a general property of amplicon origins and quantitatively correlates with origin activity

To determine whether hyperacetylation of multiple histone lysine residues is a general attribute of amplicon origins, we analyzed the other amplicons by ChIP-qPCR. We first analyzed DAFC-7F on the X chromosome, which encodes five chorion proteins and amplifies to ~16-fold (Figure 3A; Spradling, 1981). Similar to DAFC-66D, nucleosomes were hyperacetylated on multiple lysines extending across a domain at the DAFC-7F origin. There were two peaks of acetylation—one in the body of the cp36 gene and the other in cp38 (Figure 3). The latter peak corresponds to ACE1, a region that has been shown to bind ORC and that is required in *cis* for origin function (Spradling *et al.*, 1987; Austin *et al.*, 1999). The developmental timing of acetylation at DAFC-7F was also similar to that at DAFC-66D. Acetylation was low in early stages of oogenesis, greatly increased when the origin is active in stage 10, and declined later when the origin shuts off (Figure 3, B–F).

We further analyzed four other loci that have been reported to amplify to low levels in follicle cells in late oogenesis (DAFC-22B, DAFC-30B, DAFC-34B, and DAFC-62D; Claycomb *et al.*, 2004; Kim *et al.*, 2011). We found that nucleosomes at these other amplicons were also dynamically acetylated on multiple lysine residues when these origins became active in stage 10 of oogenesis (Figure 4). The one exception was DAFC-22B, which was not significantly hyperacetylated relative to the control locus (Figure 4). Although at first this seemed to challenge the generality of nucleosome hyperacetylation at active amplicon origins, the Orr-Weaver lab recently reported that the DAFC-22B origin is not active in the wild-type strain that we and the modENCODE project used for ChIP, Oregon-R^{modENCODE} (Kim *et al.*, 2011). These results indicate, therefore, that acetylation on multiple histone lysine residues is a general attribute of amplicon origins when they are developmentally activated in late oogenesis.

qPCR products is shown above, and the position of four chorion protein (cp) genes and a flanking gene is shown below. (B–F) ChIP-qPCR analysis with the indicated antibodies using follicle cells from oogenesis stages 8 and earlier (\blacktriangle), stage 10 (\blacksquare), and stage 12 (\circ). The positions of the PCR products are shown on the x-axis and aligned with the map in A; the y-axis represents the fold enrichment of ChIP-qPCR signal over a control region in the genome (at cytogenetic region 64A), and the error bars represent the range of data from two or three biological replicates.




FIGURE 3: Nucleosomes are dynamically acetylated on multiple lysine residues at the DAFC-7F origin. (A) Diagram of DAFC-7F locus and qPCR probes. The hatched box represents the origin ACE1 element. (B–F) ChIP-qPCR analysis using the indicated antibodies on




FIGURE 4: Histone acetylation levels correlate with amplicon origin efficiency. ChIP-qPCR results for six amplicon loci (DAFC-22B, -34B, -62D, -30B, -7F, and -66D) from stage 10 follicle cells using the antibodies indicated. The error bars represent the range of data from two or three biological replicates.

Another relationship between acetylation and origin activity emerged from comparison of the different amplicons; the level of hyperacetylation was greater for loci that amplify to higher final DNA copy number. The fold enrichment of all acetylation marks was higher for DAFC-66D (~64-fold amplified) than DAFC-7F (~16-fold amplified; Figure 4; Spradling, 1981). This trend continued, with acetylation levels being even lower for DAFC-62D (~3- to 6-fold amplified) and DAFC-30B (approximately fourfold amplified; Figure 4; Claycomb *et al.*, 2004). The extreme is DAFC-22B, which was not acetylated or amplified in the Oregon-R^{mod}ENCODE strain (Figure 4). Again, this trend is not an artifact of different DNA copy number, because all ChIP-qPCR is normalized to input DNA, and antibodies against unmodified histone H3 did not give evidence for enrichment (Supplemental Figures S1 and S2). Overall, the results indicated that, whereas acetylation is not unique to the origin loci, the level of acetylation on multiple lysines is much higher at the active origins and quantitatively correlates with the number of times different origins initiate replication.

Histone acetylation is not dependent on origin activation

The foregoing results suggested that multiple lysine residues are rapidly acetylated when amplicon origins become active and then are rapidly deacetylated when the origin shuts off and neighboring promoters turn on. ChIP for a marker of active transcription, H3K4me3, confirmed that neighboring promoters were not active during stages when the origin was active and highly acetylated (Supplemental Figure S3B). This suggests that acetylation at the origins does not represent activation of adjacent promoters. A possible caveat, however, is that hyperacetylation may simply represent modification on newly deposited nucleosomes behind the fork and therefore is a consequence, but not cause, of origin activity. Arguing against this interpretation is the observation that acetylation rapidly declined at DAFC-66D in stages 12 and 13, a time when the origin is quiescent but replication forks continue to migrate (Figure 1, C–C' and F and F', Figure 2, and data not shown; Claycomb *et al.*, 2002; Aggarwal and Calvi, 2004).

follicle cells from oogenesis stages 8 and earlier (▲), stage 10 (■), and stage 12 (○). PCR products are shown below each panel and aligned with A. The error bars represent the range of data from two or three biological replicates.




FIGURE 5: Acetylation is not dependent on replication initiation. Acetylation level at DAFC-66D was compared between wild-type follicle cells and those in which initiation was inhibited by expression of *UAS:dap*. (A–C) ChIP-qPCR results using the indicated antibodies on stage 10 follicle cells from wild-type Oregon R (■) and *c323GAL4/+;UAS:dap/+* (●) flies. The Oregon R data from Figure 2 were graphed for comparison. Drawn to scale for the DAFC-66D locus shown below. The error bars represent the range of data from two or three biological replicates.

To test rigorously whether acetylation simply represents nucleosome deposition behind the fork, we blocked the initiation step of DNA replication and measured acetylation at the DAFC-66D amplicon by ChIP-qPCR. We previously showed that overexpression of the cyclin E/CDK2 inhibitor *dacapo* (*dap*) severely inhibits gene amplification, resulting in a thin eggshell phenotype (Calvi *et al.*, 1998). We overexpressed *UAS:dap* specifically in late-stage follicle cells using the *c323GAL4* driver, which resulted in reduced amplification that was undetectable by BrdU incorporation in all but a few follicle cell nuclei (Calvi *et al.*, 1998). Quantification of DNA copy number at DAFC-66D and DAFC-7F in stage 10 and stage 12 egg chambers by qPCR also indicated that amplicon origin activity was severely inhibited (Supplemental Figure S4). Nonetheless, the level of acetylation on H4K12, H4K16, and H3K56 at DAFC-66D was similar between the *UAS:dap*-expressing and wild-type control cells (Figure 5). These results suggest that the majority of acetylation does not depend on origin activa-




FIGURE 6: ORC binding and histone acetylation occur in a similar extended domain around the DAFC-66D origin. ChIP-qPCR analysis of Orc2 binding (red square, right y-axis) over an ~20-kb region surrounding the DAFC-66D origin in follicle cells from stage 10. H4K5ac (black triangle, left y-axis) and H4K8ac (blue circle, left y-axis) data for the 7 kb around DAFC-66D were taken from Figure 2. Other types of acetylation gave a similar profile over this 20-kb domain but are not shown for simplicity. The error bars represent the range of data from two or three biological replicates.

tion and therefore does not correspond to nucleosome assembly behind the replication fork.

ORC binds in an extended domain at DAFC-66D with a profile that resembles acetylation

We next determined the relationship between acetylation and binding of the ORC to origin DNA, a prerequisite for subsequent assembly of the preRC. It was previously reported that Ori- β and ACE3 are preferred binding sites for the ORC *in vitro* and *in vivo* (Austin *et al.*, 1999). ChIP-qPCR with antibodies against the Orc2 subunit, however, revealed that ORC is significantly enriched in an extended domain around the DAFC-66D locus (Figure 6). We extended the analysis of ORC binding and acetylation by using primer pairs -10 and +10 kb from ACE3, which indicated that both Orc2 and acetylation of multiple lysine residues are at least twofold enriched at these sites over a control locus (Figure 6). The profile of ORC binding in this 20-kb domain was strikingly similar to that of acetylation, except that ORC occupancy did not display a sharp peak of enrichment adjacent to Ori- β . These data indicate that ORC binds and nucleosomes are acetylated in an extended domain around DAFC-66D *in vivo* and are not restricted to the essential origin regions ACE3 and Ori- β .

Acetylation on some histone lysine residues depends on ORC

The similarity in the profile for nucleosome hyperacetylation and ORC binding raised the possibility that they may have a functional relationship to one another. In fact, previous data showed that altering acetylation levels influences the selection of ORC binding sites and active origins in follicle cells (Aggarwal and Calvi, 2004; Lewis *et al.*, 2004). We could not easily test how mutation of different histone lysines affected preRC assembly and origin activity because there are hundreds of copies of histone genes in *Drosophila*, making such an analysis difficult. We could address, however, whether preRC assembly affected acetylation by using genetics to block ORC binding *in vivo* and then analyzing acetylation by ChIP-qPCR. Two mutations in the Orc6 subunit, Orc6^{K76A} and Orc6^{S72A}, impair




FIGURE 7: Acetylation of H4K12 and H3K56 depends on ORC binding. (A) ChIP-qPCR results for DAFC-66D using α -Orc2 antibodies on stage 10 follicle cells from wild-type Oregon R (■) and *c323GAL4/+;UAS:orc6^{S72A}/+* (●) flies. (B–D) ChIP-qPCR results using the indicated histone acetylation antibodies on stage 10 follicle cells from wild-type Oregon R (■) and *c323GAL4/+;UAS:orc6^{K76A}/+* (●) flies. The Oregon R data from Figure 2 are graphed in B–D for comparison. Drawn to scale for the DAFC-66D map shown below. The error bars represent the range of data from two or three biological replicates.

binding of the ORC to ACE3 and Ori- β DNA in vitro (Balasov *et al.*, 2007). We found that expression of *UAS:GFP:orc6^{K76A}* or *UAS:GFP:orc6^{S72A}* (flies provided by I. Chesnokov) using the *c323GAL4* driver partially inhibited amplification. Although most follicle cells had detectable BrdU foci, the fluorescence intensity of these foci was diminished, and females produced eggs with thin shells (data not shown). Quantification of DNA copy number by qPCR in stage 10 and stage 12 follicle cells also showed that amplification was inhibited in *UAS:GFP:orc6^{K76A}*- and *UAS:GFP:orc6^{S72A}*-

expressing flies (Supplemental Figure S4). To evaluate whether the inhibition of amplification was due to reduced ORC binding to DNA in vivo, we conducted ChIP with anti-Orc2 antibodies (Gossen *et al.*, 1995; Royzman *et al.*, 1999). In *UAS:GFP:orc6^{S72A}*-expressing follicle cells, ORC binding at DAFC-66D was not detectable across the entire locus (Figure 7A). As a control, we also expressed a wild-type *UAS:GFP:orc6*. To our surprise, this also inhibited ORC binding and amplification at DAFC-66D (Supplemental Figures S4 and S5). Although we do not understand the molecular basis for this, one possibility is that the green fluorescent protein (GFP) fusions on all these Orc6 proteins poisons the six-subunit ORC and disrupts origin binding. Nevertheless, we can use these transgenes as a tool to disrupt ORC binding to DNA and evaluate its effect on nucleosome acetylation.

Analysis of acetylation in *orc6^{K76A}*-expressing cells indicated that both H4K12ac and H3K56ac levels were reduced at DAFC-66D (Figure 7, B and D). The variance for H4K16ac in wild type, however, was too high for us to confidently conclude how this modification was affected by ORC binding (Figure 7C). These results suggest that at least H4K12ac and H3K56ac partially depend on ORC binding to DNA.

We next blocked the downstream preRC assembly step of MCM helicase loading by expressing geminin, an inhibitor of Cdt1 (Wohlschlegel *et al.*, 2000; Quinn *et al.*, 2001; Tada *et al.*, 2001). Expression of *UAS:gem* in follicle cells severely reduced BrdU incorporation at amplification foci, and females produced eggs with thin shells. BrdU incorporation was mosaic in egg chambers, with only $18 \pm 9\%$ of cells ($n = 225$) having detectable BrdU incorporation. qPCR quantification of amplicon copy number in stages 10 and 12 also showed that *UAS:gem* inhibited amplification (Supplemental Figure S4). Unlike the *UAS:orc6* cells, *UAS:gem* cells had normal levels of H4K12ac (Figure 8A). This suggests that H4K12ac depends on ORC binding but is independent of the downstream MCM helicase-loading step. Again, the variance for H4K16ac was too high among replicates to confidently conclude how MCM loading influences this modification (Figure 8B). Parallel ChIP of the same samples with antibodies against H3K56ac indicated that this modification was increased in the *UAS:gem*-expressing cells (Figure 8C). Combined, the data indicate that H4K12ac is dependent on ORC but independent of the downstream step of MCM loading. In contrast, although H3K56ac also depends on ORC binding, it becomes hyperacetylated when MCM loading is impaired.

DISCUSSION

We used the gene amplification model system to investigate the epigenetic regulation of origins during development. Previous data suggested that nucleosome acetylation contributes to specifying active amplicon origins in follicle cells. Our present data indicate that multiple histone lysines are hyperacetylated when the origin is activated and that all these acetylation marks rapidly decline later when the origin shuts off and nearby promoters turn on. Although nonorigin loci had similar types of acetylation, our data reveal a quantitative relationship between the level of acetylation and origin activity. Together with previous results, the data suggest that hyperacetylation on multiple histone lysines may contribute to origin locus specificity and developmental timing. Our data also provide a higher-resolution picture of the origin epigenetic landscape and show that nucleosomes are hyperacetylated and ORC binds in an extended domain around DAFC-66D, with a peak of acetylation adjacent to Ori- β , the major site of initiation. The most surprising finding, however, was that acetylation on some lysines




FIGURE 8: Inhibition of MCM loading does not affect acetylation of H4K12 but increases acetylation on H3K56. Comparison of acetylation level at DAFC-66D between wild-type and *UAS:gem*-expressing follicle cells. (A–C) ChIP-qPCR results using the indicated antibodies on stage 10 follicle cells from wild-type Oregon R (■) and *c323GAL4/+;UAS:gem/+* (●) flies. The Oregon R data from Figure 2 were graphed for comparison. Drawn to scale for the DAFC-66D locus shown below. The error bars represent the range of data from two or three biological replicates.

depends on ORC binding to the origin, suggesting that multiple HATs may be recruited during origin licensing. These results have important and general implications for understanding origin regulation and the coordination of origin and promoter activity during development.

An epigenetic switch model for developmental gene amplification

Our results provide new insights into the relationship between histone acetylation and the activity of amplicon origins. Previous data indicated that mutation of the general HDAC Rpd3 results in genome-wide hyperacetylation, ORC binding, and genomic replication in stage 10B follicle cells rather than the normal site-specific amplification (Aggarwal and Calvi, 2004; Lewis et al., 2004). That result suggested that acetylation of nucleosomes contributes to origin specification in follicle cells. We also developed a DAFC-66D origin reporter and used it to show that tethering the *Drosophila*

orthologues of the HAT HBO1 or HDAC Rpd3 to the origin in vivo increased or decreased amplification level, respectively (Aggarwal and Calvi, 2004). Recent results from the Orr-Weaver lab using this reporter confirm that tethering the Rpd3 HDAC represses the origin and further show that acetylation of H4K8 in the reporter is decreased (Kim et al., 2011). Results from multiple labs therefore suggest that nucleosome acetylation contributes to amplicon origin efficiency and locus specificity (Hartl et al., 2007). It was not previously known, however, what spectrum of acetylation types occurs at the origin, nor how these different nucleosome modifications coincided with temporal activation and repression of the origin. Our present data with validated antibodies show that nucleosome hyperacetylation at the origins occurs on multiple histone lysines. From our immunofluorescence and ChIP data, however, it is clear that acetylation on these same lysines occurs at some nonorigin loci in stage 10 follicle cells. This suggests that nucleosome acetylation is not sufficient to specify active origins. We found, however, that the relative level of acetylation is much higher at origin loci and that there is a quantitative correlation between the level of hyperacetylation and the number of times different amplicon origins initiate replication. Recent follicle cell array data from the Orr-Weaver lab also show that histone acetylation is not unique to the active amplicons but that there is a quantitative relationship between the level of hyperacetylation and amplicon origin activity (Kim et al., 2011). The current evidence therefore fails to support a qualitative histone code for amplicon origins, but instead reveals a quantitative correlation between the level of nucleosome hyperacetylation and origin activity.

Our data show a temporal correlation between hyperacetylation on multiple lysines and amplicon origin activity during oogenesis. Multiple acetylation marks rapidly increase when the origins become active and then rapidly decline when the origin shuts off. Rapid decline of acetylation at the DAFC-66D origin in stage 12 correlates with the departure of ORC from the origin and is followed by activation of nearby chorion protein promoters (Shea et al., 1990; Austin et al., 1999; Rozyman et al., 1999; Aggarwal and Calvi, 2004). Together with our previous evidence, our data lead us to propose an epigenetic switch model in which dynamic acetylation–deacetylation on multiple lysines participates in the locus specificity, efficiency, and developmental timing of amplicon origins and coordinates sequential origin and promoter activity (Figure 9). Although the mechanistic details of this model have yet to be worked out, it makes testable predictions about the epigenetic regulation of origins and the coordination of origin and promoter activity.

Acetylation depends on preRC assembly

Our most surprising result was that acetylation on different histone lysines depends on ORC. Evidence from yeast and metazoa supports the notion that chromatin can influence both the site of preRC assembly and the time of preRC activation during S phase (Mechali, 2010). Our results, however, indicate that the cause and effect can operate in the opposite direction as well, and that preRC assembly influences chromatin modification. Although nucleosome acetylation was not affected when origin activation was inhibited by *dacapo*, acetylation was reduced on H4K12 and H3K56 when ORC function was disrupted. Our data showed that reduced ORC binding to the origin correlated with a local reduction in nucleosome acetylation in three genotypes and six biological replicates. It remains possible, however, that the reduced acetylation is caused by an unknown indirect effect of ORC overexpression rather than a disruption of ORC binding per se. Nonetheless, H4K12ac was clearly dependent on ORC, but it was not changed when MCM loading was




FIGURE 9: An epigenetic switch model for nucleosome acetylation and origin activity at the DAFC-66D origin. This model illustrates steps of origin specification (initial ORC binding), licensing (loading of MCMs), and origin activation by CDK2 and CDC7 at DAFC-66D. Other replication proteins that associate with the origin are not shown. Genetic manipulations that we used to inhibit these three steps are shown in red italics between the boxes. The drawing shows only one ORC bound to Ori-β and the nucleosome next to it for visual clarity, but our data indicate that both ORC binds and nucleosomes are acetylated in an extended domain that encompasses the origin. Multiple lysines are acetylated, suggesting that multiple HATs participate in origin regulation. H4K12ac and H3K56ac were decreased when ORC binding was inhibited, suggesting that the HATs responsible for these modifications may be recruited or activated, depending on ORC. H3K56 acetylation increased when origin licensing was inhibited, suggesting a possible negative feedback loop between MCMs and the H3K56 HAT (black crossbar in box 3). The last step represents rapid deacetylation of the origin by unknown HDACs in stage 12, which is associated with the departure of ORC, origin silencing, and activation of nearby chorion protein (cp) promoters by transcription factors (TF). See the text for further details.

inhibited. This suggests that the HAT responsible for H4K12ac may be recruited or activated after ORC binding but before the MCM loading step of preRC assembly. H3K56ac was also dependent on ORC but was hyperacetylated when the MCM loading step was inhibited. One possibility is that the HAT for H3K56 is also recruited by ORC to promote loading of MCM helicase rings onto DNA and that a negative feedback mechanism inhibits this HAT or activates an HDAC responsible for regulating acetylation of H3K56 once sufficient MCM helicase complexes are loaded. An important prediction from these results is that multiple HATs may be recruited or activated during different steps of preRC assembly and may contribute to origin licensing. The activation and repression of origins therefore may be analogous to that of promoters, in which codependent recruitment of multiple chromatin modifiers and transcription factors determines the developmental specificity of transcription.

HATs and HDACs in origin regulation

Our results have parallels to growing evidence from yeast to humans that suggests that HATs and HDACs influence origin identity, efficiency, and timing during genomic DNA replication. We previously showed that GAL4 fusions of the fly orthologue of HBO1 could stimulate the DAFC-66D origin when tethered to it (Aggarwal and Calvi, 2004). In human cells, HBO1 is recruited to origins by Cdt1 and is required for MCM loading (Miotto and Struhl, 2010). Evidence suggests that HBO1 acetylates H4K5, H4K8, and H4K12, all modifications that we find at the amplicon origins, but cannot explain the acetylation of H3 tails, H4K16, or H3K56 (Doyon et al., 2006; Miotto and Struhl, 2010). Two candidate HATs for these modifications are CBP, which has been shown to acetylate H3K56 in multicellular eukaryotes, and MOF, which acetylates H4K16 to increase transcription from the *Drosophila* male X chromosome during dosage compensation (Kelley et al., 1999; Das et al., 2009; Gelbart and Kuroda, 2009; Kharchenko et al., 2011). In *Drosophila* tissue culture cells, acetylation of H4K16 by MOF confers early replication to the male X chromosome, and genomic sites of ORC binding and active origins correlate with enrichment for H4K16Ac and other acetylation marks (Schwaiger et al., 2009; Eaton et al., 2010b). Our data are also consistent with a report that nucleosomes at origins in *Saccharomyces cerevisiae* are acetylated on multiple lysine residues, including those that we found at the amplicon origins (Unnikrishnan et al., 2010). Finding multiple, similar acetylation marks at yeast and *Drosophila* origins suggests that the regulation of DNA replication by multiple HATs may be conserved in eukaryotes.

The rapid decline in acetylation in stage 12 suggests that the action of multiple HATs at the origin may be counteracted by multiple HDACs. Our previous data showed that Rpd3 is required to repress nonamplicon origins in stage 10B follicle cells (Aggarwal and Calvi, 2004). In yeast, the HDAC Rpd3 influences the time of origin activation, whereas the HDAC Sir2 can regulate the location of preRC assembly (Vogelauer et al., 2002; Crampton et al., 2008; Fox and Weinreich, 2008). *Drosophila* Sir2 can deacetylate H3K56, and therefore our results raise the possibility that this HDAC also regulates origins in metazoa (Das et al., 2009). It was also recently shown that HDAC11 counteracts acetylation by HBO1 to regulate DNA replication in Chinese hamster ovary cells (Wong et al., 2010). An important future goal, therefore, is to further define how these and other HATs and HDACs are recruited to and directly regulate amplicon and other origins.

The origin epigenetic landscape

A number of origins in metazoa have been analyzed molecularly and genetically, but the picture for origin anatomy and function

remains fuzzy at best. Our analysis of DAFC-66D locus has begun to provide a fine-scale picture of the epigenetic landscape of this well-characterized model origin. Previous reports indicated that the ACE3 and Ori- β regions of DAFC-66D are preferred ORC binding sites, are evolutionarily conserved, and are required in *cis* for amplification (Delidakis and Kafatos, 1989; Orr-Weaver *et al.*, 1989; Austin *et al.*, 1999; Remus *et al.*, 2004; Zhang and Tower, 2004; Claycomb and Orr-Weaver, 2005; Calvi, 2006; Calvi *et al.*, 2007). We found that nucleosome hyperacetylation was not restricted to ACE3 and Ori- β but was spread over the entire 20 kb that we analyzed. Of importance, ORC occupancy was also enriched across the entire hyperacetylated region with a similar profile. Recent array data from the Orr-Weaver lab also showed that H4K8 is hyperacetylated and ORC binds in an extended genomic domain of ~30 kb around DAFC-66D and in smaller domains around other amplicons, consistent with evidence from Mike Botchan's laboratory (Kim *et al.*, 2011; M. Botchan, personal communication). The concordance between nucleosome acetylation and ORC binding profiles is consistent with a functional relationship between them. Our genetic data showing that at least some acetylation marks depend on ORC binding opens the possibility that binding of multiple ORC complexes may promote an extended domain of nucleosome acetylation. At present we do not know whether this extended ORC domain represents binding of ORC to different sites on different DNA strands or binding of multiple ORCs on the same DNA strand. The possibility that multiple ORCs bind a single DNA strand is consistent with previous evidence for increased ORC occupancy on longer origin fragments *in vitro* and the binding of multiple ORCs at some origins in yeast and metazoa *in vivo* (Bielinsky *et al.*, 2001; Takahashi *et al.*, 2003; Bolon and Bielinsky, 2006; Aladjem, 2007). However, the functional significance of multiple ORCs for origin activity or initiation site selection remains unclear. One possibility is that binding of ORC to its preferred sites in ACE3 and/or Ori- β promotes subsequent recruitment and spreading of additional ORCs, analogous to spreading of multiple DnaA complexes at Ori-C in *Escherichia coli* (Clarey *et al.*, 2006; Erzberger *et al.*, 2006; Remus and Diffley, 2009).

Another fine-scale attribute of the DAFC-66D epigenetic landscape was that all acetylation marks that we examined were highest to one side of Ori- β , in the 3' end of the cp18 gene, which is not expressed when the origin is active. This suggests that among the population of chromatin fibers, acetylation occurs more frequently on nucleosomes to one side of Ori- β , the site where replication initiates 80% of the time (Delidakis and Kafatos, 1989; Heck and Spradling, 1990). A peak of acetylation was also observed at DAFC-7F within ACE1, a region that binds ORC and is essential for origin function. In *S. cerevisiae*, nucleosomes positioned adjacent to some origins promote ORC binding, likely through interaction with the bromo adjacent homology (BAH) domain of Orc1, a domain that is also important for the binding of human Orc1 to chromatin and mutation of which causes the developmental abnormality Meier-Gorlin syndrome (Lipford and Bell, 2001; Noguchi *et al.*, 2006; Muller *et al.*, 2010; Bicknell *et al.*, 2011; Guernsey *et al.*, 2011). The peak of acetylation adjacent to the major initiation site in Ori- β may therefore reflect a positioned nucleosome that promotes initial ORC binding or other steps of origin licensing and activation. Further analysis of nucleosome position, nucleosome modification, and preRC occupancy at DAFC-66D will permit us to test this model and dissect origin anatomy and function with high resolution.

MATERIALS AND METHODS

Drosophila strains

The Oregon $R^{modENCODE}$ strain was used as the reference wild-type strain. C323:GAL4 was used to drive expression of $P\{w^+UAS:GFP:orc6^{K76A}\}$, $P\{w^+UAS:GFP:orc6^{S72A}\}$, and $P\{w^+UAS:GFP:orc6\}$ (gifts from Igor Chesnokov, University of Alabama at Birmingham, Birmingham, AL; Balasov *et al.*, 2007), $P\{w^+UAS:dacapo\}$ (Lane *et al.*, 1996), and $P\{w^+UAS:geminin\}$ (this study).

Antibodies

α -BrdU (mouse monoclonal, BDB347580; BD Biosciences, San Diego, CA), α -H4K5ac (07-327; Upstate, Millipore, Billerica, MA), α -H4K8ac (07-328; Upstate), α -H4K12ac (39165; Active Motif, Carlsbad, CA), α -H4K16ac (39167; Active Motif), α -H3K56ac (07-677; Upstate), α -histone H3 C-terminal (39163; Active Motif), H3K4me3 (39159; Active Motif), α -H3K27me3 (07-449; Upstate), α -H3K9/14ac (06-599; Upstate), α -Orc2 (gift of Stephen Bell, MIT, Cambridge, MA; Austin *et al.*, 1999), and α -Orc2 (gift of Michael Botchan, University of California, Berkeley, CA). Except where stated, all antibodies were rabbit polyclonals.

Immunostaining and microscopy

BrdU and antibody labeling of *Drosophila* ovaries was as previously described (Calvi *et al.*, 1998; Calvi and Lilly, 2004). Antibody concentrations were as follows: α -BrdU (1:20), α -H4K12ac (1:100), and α -H4K16ac (1:100). Images are Z-stack projections taken with a Leica (Wetzlar, Germany) SP5 scanning confocal microscope.

Follicle cell nuclear preparation for ChIP

Follicle cell nuclei were purified from different stage egg chambers based on a modification of several protocols (Petri *et al.*, 1976; Woll *et al.*, 1981). Females were conditioned with males on wet yeast for 3 d and then blended by short pulses in cold phosphate-buffered saline buffer in 0.02% Tween-20 in a household blender. Released egg chambers were enriched by serial filtration through 250- to 70- μ m meshes and repeated resuspension-resettling in cold buffer. Eggs were then fixed for 15 min at room temperature in 2% paraformaldehyde solution, followed by fix quenching with 125 mM glycine and washing with cold Dulbecco's phosphate-buffered saline. Stage 10, stage 12, and stage ≤ 8 egg chambers were then further manually separated and stored at -80°C prior to nuclear preparation. Approximately 4000 stage 10 and ~3000 stage 12 egg chambers were used for nuclear preparation. The number of stage ≤ 8 egg chambers was not counted, but a tissue volume similar to that of the other stages was used. Frozen eggs were thawed on ice, resuspended in mHB buffer (0.34 M sucrose, 15 mM NaCl, 60 mM KCl, 0.2 mM EDTA, 0.2 mM ethylene glycol tetraacetic acid, 0.15 mM spermine, 0.15 mM spermidine in 15 mM Tris-HCl, pH 8.0) supplemented with 0.5% NP-40 and transferred to a Kontes 2-ml douncer. Fifteen strokes with a type A pestle were applied, and the content was filtrated with a 15- μ m Nitex nylon membrane to remove the larger nurse cell nuclei. The filtrate was spun 3 min at 500 \times g in a microcentrifuge to pellet follicle nuclei.

Chromatin immunoprecipitation

The ChIP protocol was modified from previous methods (17-295; Millipore) and entailed at least two biological replicates from separate isolations of follicle cells. In brief, prepared follicle nuclei were resuspended in nuclear lysis buffer (1% SDS, 1 mM EDTA in 50 mM Tris-Cl, pH 8.0) and subjected to sonication (Fisher dismembrator model 100 [Thermo Fisher Scientific, Waltham, MA], tip probe, setting 2, five rounds of 20-s sonication on ice) to a modal size of 450

base pairs. One-tenth of the sample was saved as input, and the remainder was aliquoted and diluted at least fivefold into binding buffer (0.01% SDS, 1.1% Triton-X 100, 1.1 mM EDTA, 167 mM NaCl in 20 mM Tris-Cl, pH 8.0) to a final volume of 1 ml. Immunoprecipitation was then performed in parallel by adding 2 μ l (~2 μ g) of antibodies, followed by nutation at 4°C overnight. The next day 30 μ l of 50% protein A agarose beads (15918-014; Invitrogen) pretreated with sheared salmon sperm DNA was added and incubated 1 h at 4°C and precipitated by brief centrifugation. The beads were washed two times each with low-salt, high-salt, LiCl, and Tris-EDTA buffers. Elution and reversal of cross-linking was done in nuclear lysis buffer with heat at 65°C overnight. Both the input and eluate were treated with 1 μ g of RNaseA (11119915001; Roche, Indianapolis, IN) and 50 μ g of proteinase K (P8102S; New England BioLabs, Ipswich, MA). DNA was then purified with standard phenol/chloroform procedure and used for qPCR analysis.

qPCR

Analysis was done on a Stratagene (Santa Clara, CA) Mx3005P machine with SYBR Green Master Mix (600843; Agilent, Santa Clara, CA). Forty cycles of PCR were run with a two-step protocol (denaturation at 95°C for 15 s and annealing/extension at 62°C for 30 s). For ChIP-qPCR experiments, the amount of DNA in the pellet was expressed as percentage of input DNA estimated by a standard curve generated from a serial dilution of the input. The values were then normalized to a control, nonorigin locus at cytogenetic position 64A. Normalization to another, nonamplified region at 93E/F gave similar results. To measure developmentally amplified DNA copy number in wild-type and transgene-expressing flies, the relative copy numbers at different loci were calculated by Δ Ct from a reference locus at cytogenetic region 93E/F. PCR primer sequences are available upon request.

ACKNOWLEDGMENTS

We thank I. Chesnokov and the Bloomington *Drosophila* Stock Center (Bloomington, IN) for flies, S. Bell and M. Botchan for α -Orc2 antibodies, J. Powers of the Indiana Light Microscopy Imaging Center (Bloomington, IN) for helpful advice, and M. Lilly and members of the Calvi lab for advice and comments on the manuscript. We thank M. Botchan, T. Orr-Weaver, and M. Kuroda for communicating results before publication. This work was supported by National Research Service Award Training Grant F32 GM080089 to K.H.M. and by National Institutes of Health Grants R01GM061290-10 and R01GM061290-10S1 to B.R.C.

REFERENCES

Aggarwal BD, Calvi BR (2004). Chromatin regulates origin activity in *Drosophila* follicle cells. *Nature* 430, 372–376.

Aladjem MI (2007). Replication in context: dynamic regulation of DNA replication patterns in metazoans. *Nat Rev Genet* 8, 588–600.

Arias EE, Walter JC (2007). Strength in numbers: preventing rereplication via multiple mechanisms in eukaryotic cells. *Genes Dev* 21, 497–518.

Austin RJ, Orr-Weaver TL, Bell SP (1999). *Drosophila* ORC specifically binds to ACE3, an origin of DNA replication control element. *Genes Dev* 13, 2639–2649.

Balaso M, Huijbregts RP, Chesnokov I (2007). Role of the Orc6 protein in origin recognition complex-dependent DNA binding and replication in *Drosophila melanogaster*. *Mol Cell Biol* 27, 3143–3153.

Bell O, Schwaiger M, Oakeley EJ, Lienert F, Beisel C, Stadler MB, Schubeler D (2010). Accessibility of the *Drosophila* genome discriminates PcG repression, H4K16 acetylation and replication timing. *Nat Struct Mol Biol* 17, 894–900.

Bell S, Kobayashi R, Stillman B (1993). Yeast origin recognition complex functions in transcription silencing and DNA replication. *Science* 262, 1844–1849.

Bicknell LS et al. (2011). Mutations in the pre-replication complex cause Meier-Gorlin syndrome. *Nat Genet* 43, 356–359.

Bielinsky A.-K., Blitzblau H, Beall EL, Ezrokhi M, Smith HS, Botchan MR, Gerbi SA (2001). Origin recognition complex binding to a metazoan replication origin. *Curr Biol* 11, 1427–1431.

Bolon YT, Bielinsky AK (2006). The spatial arrangement of ORC binding modules determines the functionality of replication origins in budding yeast. *Nucleic Acids Res* 34, 5069–5080.

Cadoret JC (2008). Genome-wide studies highlight indirect links between human replication origins and gene regulation. *Proc Natl Acad Sci USA* 105, 15837–15842.

Calvi BR (2006). Developmental DNA amplification. In: *DNA Replication and Human Disease*, ed. ML DePamphilis, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 233–255.

Calvi BR, Byrnes BA, Kolpakas AJ (2007). Conservation of epigenetic regulation, ORC binding and developmental timing of DNA replication origins in the genus *Drosophila*. *Genetics* 177, 1291–1301.

Calvi BR, Lilly MA (2004). Henderson D (2004). BrdU labeling and nuclear flow sorting of the *Drosophila* ovary. *Drosophila Cytogenetics Protocols*, Totowa, NJ: Humana Press, 203–213.

Calvi BR, Lilly MA, Spradling AC (1998). Cell cycle control of chorion gene amplification. *Genes Dev* 12, 734–744.

Calvi BR, Spradling AC (2001). The nuclear location and chromatin organization of active chorion amplification origins. *Chromosoma* 110, 159–172.

Chesnokov I, Remus D, Botchan M (2001). Functional analysis of mutant and wild-type *Drosophila* origin recognition complex. *Proc Natl Acad Sci USA* 98, 11997–12002.

Chong J, Mahbubani H, Khoo C, Blow J (1995). Purification of an MCM-containing complex as a component of the DNA replication licensing system. *Nature* 375, 418–421.

Clarey MG, Erzberger JP, Grob P, Leschziner AE, Berger JM, Nogales E, Botchan M (2006). Nucleotide-dependent conformational changes in the DnaA-like core of the origin recognition complex. *Nat Struct Mol Biol* 13, 684–690.

Claycomb JM, Benasutti M, Bosco G, Fenger DD, Orr-Weaver TL (2004). Gene amplification as a developmental strategy: isolation of two developmental amplicons in *Drosophila*. *Dev Cell* 6, 145–155.

Claycomb JM, MacAlpine DM, Evans JG, Bell SP, Orr-Weaver TL (2002). Visualization of replication initiation and elongation in *Drosophila*. *J Cell Biol* 159, 225–236.

Claycomb JM, Orr-Weaver TL (2005). Developmental gene amplification: insights into DNA replication and gene expression. *Trends Genet* 21, 149–162.

Cocker J, Piatti S, Santocanale C, Nasmyth K, Diffley J (1996). An essential role for the Cdc6 protein in forming the pre-replicative complexes of budding yeast. *Nature* 379, 180–182.

Crampton A, Chang F, Pappas DL Jr, Frisch RL, Weinreich M (2008). An ARS element inhibits DNA replication through a SIR2-dependent mechanism. *Mol Cell* 30, 156–166.

Danis E, Brodolin K, Menu S, Maiorano D, Girard-Reydet C, Mechali M (2004). Specification of a DNA replication origin by a transcription complex. *Nat Cell Biol* 6, 721–730.

Das C, Lucia MS, Hansen KC, Tyler JK (2009). CBP/p300-mediated acetylation of histone H3 on lysine 56. *Nature* 459, 113–117.

Delidakis C, Kafatos F (1989). Amplification enhancers and replication origins in the autosomal chorion gene cluster of *Drosophila*. *EMBO J* 8, 891–901.

Diffley J, Cocker J, Dowell S, Harwood J, Rowley A (1995). Stepwise assembly of initiation complexes at budding yeast replication origins during the cell cycle. *J Cell Sci Suppl* 19, 67–72.

Doyon Y, Cayrou C, Ullah M, Landry AJ, Cote V, Selleck W, Lane WS, Tan S, Yang XJ, Cote J (2006). ING tumor suppressor proteins are critical regulators of chromatin acetylation required for genome expression and perpetuation. *Mol Cell* 21, 51–64.

Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM (2010a). Conserved nucleosome positioning defines replication origins. *Genes Dev* 24, 748–753.

Eaton ML, Prinz JA, Macalpine HK, Tretyakov G, Kharchenko PV, Macalpine DM (2010b). Chromatin signatures of the *Drosophila* replication program. *Genome Res*.

Egelhofer TA et al. (2011). An assessment of histone-modification antibody quality. *Nat Struct Mol Biol* 18, 91–93.

Erzberger JP, Mott ML, Berger JM (2006). Structural basis for ATP-dependent DnaA assembly and replication-origin remodeling. *Nat Struct Mol Biol* 13, 676–683.

- Fox CA, Weinreich M (2008). Beyond heterochromatin: SIR2 inhibits the initiation of DNA replication. *Cell Cycle* 7, 3330–3334.
- Gelbart ME, Kuroda MI (2009). *Drosophila* dosage compensation: a complex voyage to the X chromosome. *Development* 136, 1399–1410.
- Gilbert DM (2010). Evaluating genome-scale approaches to eukaryotic DNA replication. *Nat Rev Genet* 11, 673–684.
- Gossen M, Pak D, Hansen S, Acharya J, Botchan M (1995). A *Drosophila* homolog of the yeast origin recognition complex. *Science* 270, 1674–1677.
- Green BM, Finn KJ, Li JJ (2010). Loss of DNA replication control is a potent inducer of gene amplification. *Science* 329, 943–946.
- Griffin-Shea R, Thireos G, Kafatos F (1982). Organization of a cluster of four chorion genes in *Drosophila* and its relationship to developmental expression and amplification. *Dev Biol* 91, 325–336.
- Guernsey DL et al. (2011). Mutations in origin recognition complex gene ORC4 cause Meier-Gorlin syndrome. *Nat Genet* 43, 360–364.
- Hansen RS, Thomas S, Sandstrom R, Canfield TK, Thurman RE, Weaver M, Dorschner MO, Gartler SM, Stamatoyannopoulos JA (2010). Sequencing newly replicated DNA reveals widespread plasticity in human replication timing. *Proc Natl Acad Sci USA* 107, 139–144.
- Hartl T, Boswell C, Orr-Weaver TL, Bosco G (2007). Developmentally regulated histone modifications in *Drosophila* follicle cells: initiation of gene amplification is associated with histone H3 and H4 hyperacetylation and H1 phosphorylation. *Chromosoma* 116, 197–214.
- Heck M, Spradling A (1990). Multiple replication origins are used during *Drosophila* chorion gene amplification. *J Cell Biol* 110, 903–914.
- Hiratani I et al. (2010). Genome-wide dynamics of replication timing revealed by in vitro models of mouse embryogenesis. *Genome Res* 20, 155–169.
- Hiratani I, Ryba T, Itoh M, Yokochi T, Schwaiger M, Chang CW, Lyou Y, Townes TM, Schubeler D, Gilbert DM (2008). Global reorganization of replication domains during embryonic stem cell differentiation. *PLoS Biol* 6, e245.
- Hook SS, Lin JJ, Dutta A (2007). Mechanisms to control rereplication and implications for cancer. *Curr Opin Cell Biol* 19, 663–671.
- Huberman JA (1968). Visualization of replicating mammalian and T4 bacteriophage DNA. *Cold Spring Harb Symp Quant Biol* 33, 509–524.
- Iizuka M, Stillman B (1999). Histone acetyltransferase HBO1 interacts with the ORC1 subunit of the human initiator protein. *J Biol Chem* 274, 23027–23034.
- Iizuka M, Takahashi Y, Mizzen CA, Cook RG, Fujita M, Allis CD, Frierson HF Jr, Fukusato T, Smith MM (2009). Histone acetyltransferase Hbo1: catalytic activity, cellular abundance, and links to primary cancers. *Gene* 436, 108–114.
- Jorgensen HF, Azuara V, Amoils S, Spivakov M, Terry A, Nesterova T, Cobb BS, Ramsahoye B, Merckenschlager M, Fisher AG (2007). The impact of chromatin modifiers on the timing of locus replication in mouse embryonic stem cells. *Genome Biol* 8, R169.
- Karnani N, Taylor CM, Malhotra A, Dutta A (2010). Genomic study of replication initiation in human chromosomes reveals the influence of transcription regulation and chromatin structure on origin selection. *Mol Biol Cell* 21, 393–404.
- Kelley RL, Meller VH, Gordadze PR, Roman G, Davis RL, Kuroda MI (1999). Epigenetic spreading of the *Drosophila* dosage compensation complex from roX RNA genes into flanking chromatin. *Cell* 98, 513–522.
- Kharchenko PV et al. (2011). Comprehensive analysis of the chromatin landscape in *Drosophila melanogaster*. *Nature* 471, 480–485.
- Kim JC, Nordman J, Xie F, Kashevsky H, Eng T, Li S, MacAlpine DM, Orr-Weaver TL (2011). Integrative analysis of gene amplification in *Drosophila* follicle cells: parameters of origin activation and repression. *Genes Dev* 25, 1384–1398.
- Kim SM, Dubey DD, Huberman JA (2003). Early-replicating heterochromatin. *Genes Dev* 17, 330–335.
- Labib K (2010). How do Cdc7 and cyclin-dependent kinases trigger the initiation of chromosome replication in eukaryotic cells. *Genes Dev* 24, 1208–1219.
- Landis G, Tower J (1999). The *Drosophila* chifon gene is required for chorion gene amplification, and is related to the yeast Dbf4 regulator of DNA replication and cell cycle. *Development* 126, 4281–4293.
- Lane M, Sauer K, Wallace K, Jan Y, Lehner C, Vaessin H (1996). Dacapo, a cyclin-dependent kinase inhibitor, stops cell proliferation during *Drosophila* development. *Cell* 87, 1225–1235.
- Lengronne A, Schwob E (2002). The yeast CDK inhibitor Sic1 prevents genomic instability by promoting replication origin licensing in late G(1). *Mol Cell* 9, 1067–1078.
- Lewis PW, Beall EL, Fleischer TC, Georgette D, Link AJ, Botchan MR (2004). Identification of a *Drosophila* Myb-E2F2/RBF transcriptional repressor complex. *Genes Dev* 18, 2929–2940.
- Lima-de-Faria A, Jaworska H (1968). Late DNA synthesis in heterochromatin. *Nature* 217, 138–142.
- Liontos M et al. (2007). Deregulated overexpression of hCdt1 and hCdc6 promotes malignant behavior. *Cancer Res* 67, 10899–10909.
- Lipford JR, Bell SP (2001). Nucleosomes positioned by ORC facilitate the initiation of DNA replication. *Mol Cell* 7, 21–30.
- MacAlpine HK, Gordan R, Powell SK, Hartemink AJ, MacAlpine DM (2010). *Drosophila* ORC localizes to open chromatin and marks sites of cohesin complex loading. *Genome Res* 20, 201–211.
- Maiorano D, Moreau J, Mechali M (2000). XCDT1 is required for the assembly of pre-replicative complexes in *Xenopus laevis*. *Nature* 404, 622–625.
- Mechali M (2010). Eukaryotic DNA replication origins: many choices for appropriate answers. *Nat Rev Mol Cell Biol* 11, 728–738.
- Mehrotra S, Maqbool SB, Kolpakas A, Murnen K, Calvi BR (2008). Endocycling cells do not apoptose in response to DNA rereplication genotoxic stress. *Genes Dev* 22, 3158–3171.
- Miotto B, Struhl K (2010). HBO1 histone acetylase activity is essential for DNA replication licensing and inhibited by Geminin. *Mol Cell* 37, 57–66.
- Mito Y, Henikoff JG, Henikoff S (2007). Histone replacement marks the boundaries of cis-regulatory domains. *Science* 315, 1408–1411.
- Muller P, Park S, Shor E, Huebert DJ, Warren CL, Ansari AZ, Weinreich M, Eaton ML, MacAlpine DM, Fox CA (2010). The conserved bromo-adjacent homology domain of yeast Orc1 functions in the selection of DNA replication origins within chromatin. *Genes Dev* 24, 1418–1433.
- Nishitani H, Lygerou Z, Nishimoto T, Nurse P (2000). The Cdt1 protein is required to license DNA for replication in fission yeast. *Nature* 404, 625–628.
- Noguchi K, Vassilev A, Ghosh S, Yates JL, DePamphilis ML (2006). The BAH domain facilitates the ability of human Orc1 protein to activate replication origins in vivo. *EMBO J* 25, 5372–5382.
- Orr W, Komitopoulou K, Kafatos F (1984). Mutants suppressing in trans chorion gene amplification in *Drosophila*. *Proc Natl Acad Sci USA* 81, 3773–3777.
- Orr-Weaver T, Johnston C, Spradling A (1989). The role of ACE3 in *Drosophila* chorion gene amplification. *EMBO J* 8, 4153–4162.
- Orr-Weaver T, Spradling A (1986). *Drosophila* chorion gene amplification requires an upstream region regulating s18 transcription. *Mol Cell Biol* 6, 4624–4633.
- Petri WH, Wyman AR, Kafatos FC (1976). Specific protein synthesis in cellular differentiation. III. The eggshell proteins of *Drosophila melanogaster* and their program of synthesis. *Dev Biol* 49, 185–199.
- Quinn LM, Herr A, McGarry TJ, Richardson H (2001). The *Drosophila* Geminin homolog: roles for Geminin in limiting DNA replication, in anaphase and in neurogenesis. *Genes Dev* 15, 2741–2754.
- Remus D, Beall EL, Botchan MR (2004). DNA topology, not DNA sequence, is a critical determinant for *Drosophila* ORC-DNA binding. *EMBO J* 23, 897–907.
- Remus D, Diffley JF (2009). Eukaryotic DNA replication control: lock and load, then fire. *Curr Opin Cell Biol* 21, 771–777.
- Royzman I, Austin RJ, Bosco G, Bell SP, Orr-Weaver TL (1999). ORC localization in *Drosophila* follicle cells and the effects of mutations in *dE2F* and *dDP*. *Genes Dev* 13, 827–840.
- Schwaiger M, Kohler H, Oakeley EJ, Stadler MB, Schubeler D (2010). Heterochromatin protein 1 (HP1) modulates replication timing of the *Drosophila* genome. *Genome Res* 20, 771–780.
- Schwaiger M, Stadler MB, Bell O, Kohler H, Oakeley EJ, Schubeler D (2009). Chromatin state marks cell-type- and gender-specific replication of the *Drosophila* genome. *Genes Dev* 23, 589–601.
- Schwed G, May N, Pechersky Y, Calvi BR (2002). *Drosophila* minichromosome maintenance 6 is required for chorion gene amplification and genomic replication. *Mol Biol Cell* 13, 607–620.
- Sequeira-Mendes J (2009). Transcription initiation activity sets replication origin efficiency in mammalian cells. *PLoS Genet* 5, e1000446.
- Shea M, King D, Conboy M, Mariani B, Kafatos F (1990). Proteins that bind to chorion cis-regulatory elements: A new C2H2 protein and a C2C2 steroid receptor-like component. *Genes Dev* 4, 1128–1140.
- Simpson RT (1990). Nucleosome positioning can affect the function of a cis-acting DNA element in vivo. *Nature* 343, 387–389.
- Spradling A (1981). The organization and amplification of two chromosomal domains containing *Drosophila* chorion genes. *Cell* 27, 193–201.

- Spradling A, de Cicco DV, Wakimoto B, Levine J, Kalfayan L, Cooley L (1987). Amplification of the X-linked *Drosophila* chorion gene cluster requires a region upstream from the s38 chorion gene. *EMBO J* 6, 1045–1053.
- Stedman W, Deng Z, Lu F, Lieberman PM (2004). ORC, MCM, and histone hyperacetylation at the Kaposi's sarcoma-associated herpesvirus latent replication origin. *J Virol* 78, 12566–12575.
- Tada S, Li A, Maiorano D, Mechali M, Blow JJ (2001). Repression of origin assembly in metaphase depends on inhibition of RLF-B/Cdt1 by geminin. *Nat Cell Biol* 3, 107–113.
- Takahashi T, Ohara E, Nishitani H, Masukata H (2003). Multiple ORC-binding sites are required for efficient MCM loading and origin firing in fission yeast. *EMBO J* 22, 964–974.
- Unnikrishnan A, Gafken PR, Tsukiyama T (2010). Dynamic changes in histone acetylation regulate origins of DNA replication. *Nat Struct Mol Biol* 17, 430–437.
- Urnov FD, Liang C, Blitzblau HG, Smith HS, Gerbi SA (2002). A DNase I hypersensitive site flanks an origin of DNA replication and amplification in *Sciara*. *Chromosoma* 111, 291–303.
- Vashee S, Cvetic C, Lu W, Simancek P, Kelly TJ, Walter JC (2003). Sequence-independent DNA binding and replication initiation by the human origin recognition complex. *Genes Dev* 17, 1894–1908.
- Vogelauer M, Rubbi L, Lucas I, Brewer BJ, Grunstein M (2002). Histone acetylation regulates the time of replication origin firing. *Mol Cell* 10, 1223–1233.
- Whittaker AJ, Rozyman I, Orr-Weaver TL (2000). *Drosophila* double parked: a conserved, essential replication protein that colocalizes with the origin recognition complex and links DNA replication with mitosis and the down-regulation of S phase transcripts. *Genes Dev* 14, 1765–1776.
- Wohlschlegel JA, Dwyer BT, Dhar SK, Cvetic C, Walter JC, Dutta A (2000). Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Science* 290, 2309–2312.
- Woll WW, Duffy JJ, Giese NA, Lindell TJ (1981). Nuclear isolation by a modified method of Hewish and Burgoyne: implications for the study of nuclear enzymology. *Life Sci* 29, 2709–2719.
- Wong PG, Glozak MA, Cao TV, Vaziri C, Seto E, Alexandrow M (2010). Chromatin unfolding by Cdt1 regulates MCM loading via opposing functions of HBO1 and HDAC11-geminin. *Cell Cycle* 9, 4351–4363.
- Wu ZQ, Liu X (2008). Role for Plk1 phosphorylation of Hbo1 in regulation of replication licensing. *Proc Natl Acad Sci USA* 105, 1919–1924.
- Yan H, Gibson S, Tye BK (1991). Mcm2 and Mcm3, two proteins important for ARS activity, are related in structure and function. *Genes Dev* 5, 944–957.
- Zhang H, Tower J (2004). Sequence requirements for function of the *Drosophila* chorion gene locus ACE3 replicator and Ori-beta origin elements. *Development* 131, 2089–2099.