

## Synthetic biology and therapeutic strategies for the degenerating brain

Synthetic biology approaches can transform classical cell and gene therapies, to provide new cures for neurodegenerative diseases

Carmen Agustín-Pavón and Mark Isalan\*

Synthetic biology is an emerging engineering discipline that attempts to design and rewire biological components, so as to achieve new functions in a robust and predictable manner. The new tools and strategies provided by synthetic biology have the potential to improve therapeutics for neurodegenerative diseases. In particular, synthetic biology will help design small molecules, proteins, gene networks, and vectors to target disease-related genes. Ultimately, new intelligent delivery systems will provide targeted and sustained therapeutic benefits. New treatments will arise from combining protect and repair strategies: the use of drug treatments, the promotion of neurotrophic factor synthesis, and gene targeting. Going beyond RNAi and artificial transcription factors, site-specific genome modification is likely to play an increasing role, especially with newly available gene editing tools such as CRISPR/Cas9 systems. Taken together, these advances will help develop safe and long-term therapies for many brain diseases in human patients.

### Keywords:

artificial cell systems; genome editing; neurodegeneration; synthetic proteins

DOI 10.1002/bies.201400094

Department of Life Sciences, Imperial College London, London, UK

### \*Corresponding author:

Mark Isalan  
E-mail: m.isalan@imperial.ac.uk

### Abbreviations:

**AAV**, adeno-associated virus; **AD**, Alzheimer's disease; **ALS**, amyotrophic lateral sclerosis; **GDNF**, glial-derived neurotrophic factor; **HD**, Huntington's disease; **PD**, Parkinson's disease; **ZF**, zinc finger.


### Introduction

'Our pleasures, joys, laughter, and jests arise from no other source than the brain; and so do our pains, grief, anxieties, and tears.' Two millennia after Hippocrates first acknowledged this simple truth, understanding how the brain works is still one of our major challenges. Indeed, two ambitious initiatives are currently devising tools for imaging and controlling brain activity, ultimately to create a working computational model of the entire brain [1]. Although considerable scientific efforts are being made towards making the 21st century the age of the brain, we still seem to have caught just a glimpse of the function of the most complex organ in the human body.

Understanding the healthy brain is one goal, working out how to treat it when it becomes diseased is quite another. Although human lifespan has increased as standards of living have risen, the incidence of brain diseases has increased as a consequence [2, 3]. Brain pathologies include neuropsychiatric conditions, such as depression, as well as neurodegenerative diseases – the former being a risk factor for the latter [4]. Taking both types of disorders together, up to one third of the European population suffers from brain diseases every year [5].

Treating brain degeneration is now more necessary than ever, but it is especially difficult for a number of reasons. Putting aside our limited understanding of neural circuit functions, the intricate mixture of cell types populating the brain, the difficulty of targeted delivery, and the problem of designing drugs capable of crossing the blood brain barrier effectively (Fig. 1), we still face the daunting problem of rewiring and restoring the neural circuits when some of their components are lost as a result of degeneration. To circumvent this problem, it is critical to make the earliest possible interventions, ideally before neurodegeneration takes place. In other words, the focus should be in targeting pathophysiology rather than pathogenesis [6].

Discovering early biomarkers of disease is also essential, so as to improve diagnostic tools and to identify early



**Figure 1.** Overcoming the blood brain barrier for therapy. The sketch shows the cellular components of the blood brain barrier (BBB), which isolates and protects the neural tissue. The barrier prevents most drugs and therapeutic molecules from reaching their target sites in the central nervous system, when administered peripherally. Several strategies can be used to overcome the BBB, although these vary in their invasiveness. Recent developments include the use of nanoparticles and ‘Trojan peptides’ that naturally pass through lipid membranes to deliver their cargo. For a recent review of advances in delivery systems to the central nervous system, see [99].

symptoms. For example, in most common neurodegenerative conditions, early psychiatric symptoms such as depression and anxiety may precede the onset of neuronal loss by one or two decades. Some of these early symptoms are already being tested in the clinic for diagnostic potential [7, 8]. Metabolic and molecular biomarkers could also enable accurate predictions before the clinical onset of the disease, but they need extensive validation, and they raise ethical issues because of the unavoidable uncertainties associated with prediction [9].

Even if efficient early diagnosis were possible, we currently lack effective treatments for the degenerating brain. Most available drugs merely alleviate symptoms, without stopping disease progression or treating the underlying causes. It is therefore essential to devise novel, efficient therapeutic approaches to overcome these challenges; and, fortunately, some new potentially game-changing ideas are in the pipeline. These include the tools of synthetic biology, such as protein and gene network engineering [10, 11], gene targeting [12, 13] and genome editing [14–16].


In this review, we will look at the application of these tools to advance gene and cell therapy for neurodegenerative diseases. The discipline of synthetic biology is emerging with the promise of building standard, programmable and reusable parts for engineering biological systems [17]. It is beyond our

scope here to review the entire field, which is diverse. Very briefly, synthetic biology includes topics such as: engineering synthetic transcription, signalling and patterning networks [18, 19]; engineering metabolic networks to produce useful biomolecules (e.g. the malarial drug artemisinin [20]); engineering multicellular systems (e.g. cellular logic gates and computers [21]); building synthetic biomachines (e.g. swimming ‘jellyfish’ made from heart cells [22]).

Synthetic biology can contribute to gene and cell therapy for neurodegenerative diseases in several ways (Fig. 2). First, it can be used to modify classical gene therapy vectors for avoiding the immune system and yet for expressing therapeutic transgenes at the appropriate time, location, and quantity, to achieve long term therapeutic benefit. Second, synthetic biologists can design new molecules to target and modify expression in a tightly controlled manner. Pathogenic genes can be targeted, while taking into account their interactions in a given gene network. Genome engineering can be performed *ex vivo* in cells from different sources – including patient-specific cells and induced-pluripotent stem cells – or *in vivo*. Third, synthetic biomolecules can be designed to interact and clear aberrant proteins. Finally, it is possible to rewire genetic circuits in cells to build robust artificial systems that perform tasks of interest. These can then be delivered to the patient in the form of controllable microencapsulated cells. By combining some of these new approaches with classical gene therapy, we may obtain some urgently needed solutions for the problems of neurodegenerative diseases.

## How will gene therapy conquer neurodegenerative diseases?

The idea of gene therapy is beautifully simple: by repairing, replacing or modulating the activity of a diseased gene, you



**Figure 2.** Overview of synthetic biology approaches for tackling neurodegenerative diseases. The sketch shows a marble bust of Hippocrates, who recorded some of the earliest insights about the functions of the brain. Note that many of the approaches can be used in combination.

should be able to target the underlying causes of the pathology directly. Gene therapy approaches typically deliver genes or gene modulating factors (e.g. RNAi [23]) with viral vectors. Standard vectors for use in the brain are adeno-associated viruses (AAVs) [24] and lentiviruses [25]; these are modified to infect cell subtypes efficiently while addressing safety concerns by being non-replicating. Single gene diseases

are easier to target and, in the context of neurological diseases, these include the metabolic disease of metachromatic leukodystrophy [26] and Huntington's disease (HD) [27]. However, even single gene diseases are not always straightforward to tackle, because we often have a limited understanding of the multiple interactions of single genes embedded within larger disease-causing networks.

Neurodegenerative diseases such as Parkinson's (PD), Alzheimer's disease (AD), or amyotrophic lateral sclerosis (ALS), are generally complex (Box 1), hence making it difficult to choose a single gene to target, – although several trials are in development (Table 1). The preferred approach so far has therefore been that of neuroprotection, via the administration of neurotrophic factors (factors that promote the growth,

**Box 1****Common and 'rare' neurodegenerative diseases**

Parkinson's and Alzheimer's disease are two of the most common neurodegenerative diseases of the brain. Both can be idiopathic or familial, and a number of different genes have been related to each condition. For AD, these include amyloid- $\beta$ , presenilins, tau, and apolipoprotein E. For PD, genes include  $\alpha$ -synuclein, parkin, or Leucine-rich repeat kinase 2. The multifactorial nature of these diseases makes them intrinsically more difficult to understand and to treat.

By contrast, HD has a single cause, but occurs more rarely with a frequency of approximately 1:10,000 people. HD belongs to a family of nine neurodegenerative disorders known as polyglutamine diseases, caused by the expansion of glutamine stretches coded within several unrelated genes. HD shares some features with PD and AD, including onset in late adulthood (for shorter polyglutamine expansions), selective neuronal vulnerability to disease-related proteins and abnormal protein processing and aggregation. Thus, HD is gaining importance as a model disease for developing therapeutic strategies that might translate across to the more common neurodegenerative diseases of diverse aetiology [27].

Similarly, whereas most of the cases of the motor neuron disease ALS are sporadic, the rare disease spinal muscular atrophy (SMA) is monogenic, making it an ideal candidate for gene therapy studies [100]. SMA has a similar prevalence to HD, and can be caused by several mutations in the autosomic Survival Motor Neuron gene (*SMN1*). These include intragenic deletions, nonsense and point mutations, which reduce the production of SMN protein.

survival and maintenance of neurons) or by promoting their synthesis [6]. Cell transplantation of bioengineered cells secreting neurotrophic factors is another promising strategy [28, 29], but requires overcoming problems such as the limited distribution of factors in brain tissue, and the limited controllability of the release (Table 1). Synthetic biology has the potential to improve these approaches both by modifying the gene delivery vectors and by providing new payloads to control gene expression.

**Modifying gene therapy vectors reduces host clearance of therapeutic transgenes**

Before discussing the possible genetic payloads that will be used in the next generation of gene therapy studies, it is important to consider the problem of long-term therapeutic expression of transgenes in the brain and vector design. Dramatic problems, such as the sudden death of a young patient in a gene therapy trial, as a direct consequence of viral vector infusion [30], have driven researchers to improve the safety of the viral vectors, by reducing the immunogenicity of vector capsid proteins [31]. Another well-documented adverse effect has been insertional mutagenesis – the development of cancerous cells after insertion of the vector in the host genome [32]. As a result, researchers have worked to reduce genomic insertion capabilities [33]. Nevertheless, the development of increasingly safe vectors does not necessarily imply successful gene therapy, for it is also necessary to consider the toxicity and immunogenicity of administered genes. A striking example of this issue has been shown recently in a series of papers using rodents and non-human primates, simply using the common reporter gene GFP [34–36].

GFP is ubiquitous in modern biological research, and on the whole no toxicity is reported. However, it has been demonstrated that brain injections of an AAV vector

**Table 1. Recent clinical trials illustrating strategies in gene and cell therapy for neurodegenerative diseases, showing trial outcomes and potential improvements from synthetic biology tools**

Disease	Therapeutic approach	Outcome	Reference	Improvement with synthetic biology tools
AD	Antibody against amyloid $\beta$	Safety	[110]	Improved specificity, tailored antibody engineering
	Encapsulated cells delivering NGF	Safety, low delivery levels after retrieval	[29]	Tightly controlled delivery levels
PD	Transplantation of levodopa-producing cells	No improvement	[111]	Improvement in delivery and distribution
	Gene therapy: administration of neurotrophic factor	Safety	[24]	Improvement in delivery control and distribution
	Gene therapy: administration of rate-limiting enzyme for dopamine	Safety and short term efficacy	[112]	Improvement in delivery and distribution
	Transplant of fetal cells	Long term improvement in two patients	[113]	Synthetic non-fetal stem cells
SMA	Antisense oligonucleotides for RNA editing	Safety and dose-dependent muscle function improvement	[45]	Fully synthetic antisense oligonucleotides

As noted in the main text, the outcome of the treatments might critically depend on early administration, i.e. growth factors might be able to slow disease progression, but they will not rescue neuronal death. AD, Alzheimer's disease; PD, Parkinson's disease.



enhancing the expression of neurotrophic factors to promote neuroprotection and self-recovery. Further to the RNAi-related developments of the last decade, synthetic biology has the potential to engineer transcription factors to control disease-related gene expression.

### 'Killing the messenger': RNA silencing and editing technologies are reaching maturity

One of the most tried and tested gene therapy strategies so far has been the silencing of mutated transcripts by means of antisense and RNA interference (RNAi) molecules. The use of chemically-modified nucleic acids to build stable small hairpin oligonucleotides (shRNAs) or microRNAs (miRNAs) has helped in this direction.

A pre-clinical study that successfully used this approach provided a safe, long-term treatment in several mouse models of HD. Targeting the mutant huntingtin mRNA gene with antisense 2'-O-methoxyethyl-phosphorothioate-modified chimeric oligonucleotides led to its degradation with RNase H [41]. This approach provided significant reductions of the mutant proteins for two months after treatment, in early-onset HD mouse models, and reversal of the motor abnormalities for six months after treatment, in a late-onset mouse model. Importantly, this treatment reduced the mRNA levels of huntingtin in a non-human primate for eight weeks after the termination of the treatment [41], highlighting its potential in translational medicine.

Similar modified oligonucleotides are already being translated to the clinic for spinal muscular atrophy (SMA) (Box 1). The Survival Motor Neuron gene (*SMN1*) is mutated in the disease, but the therapeutic strategy takes advantage of the presence of an almost identical gene variant (*SMN2*). *SMN2* contains an RNA splicing site that excludes exon 7 and produces a non-functional protein [42]. However, antisense oligonucleotides can silence this splice site to promote the inclusion of exon 7, thus rescuing SMA. After successful pre-clinical studies in mouse models [43, 44], several clinical trials are being developed [45] (Table 1).

As well as targeting mRNAs that code for disease-related proteins, recent studies are tackling non-coding RNAs that are also involved in pathogenesis [46]. For example, a GGGGCC expansion is found in the promoter and first intron of the C9ORF72 gene, that makes a toxic non-coding transcript. This mutation is found in both ALS and frontotemporal dementia patients. Targeting the transcript with antisense oligonucleotides thus provides a promising common strategy for two different neurodegenerative diseases [47].

Although generally safe, some of these interference strategies are not devoid of peril for the organism. For example, shRNA has sometimes been found to be toxic in the brain [48–51], most likely because of saturation of the endogenous miRNA machinery, or because of off-target effects. A step forward in this strategy is the use of novel, fully synthetic, genetic polymers [52]. These polymers, based on nucleic acid architectures not found in nature, can have the advantage of not being recognised by cellular enzymes and processing machinery, hence reducing the possibility of rejection, while retaining therapeutic potential.

## Synthetic transcription factors can flexibly regulate disease target genes

Whereas RNAi is generally limited to inhibiting target expression, artificial DNA binding proteins can be targeted to virtually any sequence, and can be fused to either transcription repression or activation domains [53, 54].

For example, we designed and built a long poly-ZF protein able to bind to expanded stretches of CAG repeats, a mutation found in polyglutamine disorders, such as HD (Box 1). We fused our custom-made ZF to a KRAB repressor domain, and this synthetic transcription factor was efficient in repressing mutant huntingtin in human patient cell lines [12]. The ZF was vectorised using AAV, and was delivered to mouse brains in the R6/2 HD model. This resulted in an acute reduction of the levels of mutant RNA and protein, and a concurrent delay in the onset of HD symptoms [12].

A similar approach was used in the neurons of the retina, as a treatment for retinitis pigmentosa. This disease causes blindness because of a mutation in rhodopsin. In a mouse model overexpressing mutated human rhodopsin, retinal injections of a viral vector carrying an engineered ZF, coupled to a transcriptional repressor, reduced transcription levels of mutated rhodopsin and prevented retinal degeneration [55].

Synthetic transcription factors have also been used to upregulate the transcription of endogenous neurotrophic factors. Generally, the virally-derived VP16 or VP64 transcription activation domains can be fused to any DNA binding domain to make activators [56, 57]. In a rat model of PD, a synthetic ZF coupled to the activator domain targeting the endogenous glial-derived neurotrophic factor (GDNF) promoter, was indeed neuroprotective [58]. When following this strategy, it is important that the expression levels of the enhanced gene do not exceed the normal physiological range, since this risks toxicity [59]. The therapeutic potential of artificial gene regulation is great, and, it is complemented by another level of synthetic biology: manipulating the genome at will.

## Genome editing with artificial nucleases: A geneticist's dream

Perhaps the most transformative technology to emerge in the last decade has been the possibility to harness the power of synthetic nucleases to allow genome-editing [16, 60, 61] (Table 2). Nucleases make targeted double-stranded DNA breaks, which naturally induce knockouts (by the process of error-prone non-homologous end-joining) or knock-ins (by homologous recombination). Zinc finger nucleases (ZFNs), Transcription Activator like Effectors (TALENs) and, most recently, RNA-guided Clustered Regularly Interspaced Short Palindromic Repeat systems (CRISPR/Cas9), allow easy and efficient genetic engineering of mammalian cells. In particular, the latest generation of lentiviral Cas9 systems can reach genome modification efficiencies of >90% of targeted cells [14] – this would have seemed like science fiction just a few years ago.

Synthetic nucleases can be used to knock out, mutate, or repair a gene causing a given disease. One of the most promising aspects of this approach relies on the editing of human pluripotent stem cells for autologous transplants [62]. An example is the recent success using ZF nucleases to delete the CCR5 HIV receptor in white blood cells, demonstrated to be a potential functional cure for HIV in patients [63].

Haematopoietic cell transplants, using engineered cells corrected for a mutation, have already shown promise for neurodegenerative diseases arising from metabolic defects, and therefore such diseases could also be treated with the latest generation of synthetic nucleases. Amongst these, lysosomal storage diseases are fatal conditions in which the accumulation of cell by-products in lysosomes leads to neurodegeneration. In metachromatic leukodystrophy, the deficiency of a single enzyme, arylsulfatase A, leads to fatality in infancy, within a few years of disease onset. Recently, a clinical trial has successfully used the autologous transplantation of engineered haematopoietic cells expressing the functional enzyme. These cells can migrate to the brain and clear the toxic accumulations of the enzyme substrate [26]. The use of new genome editing technologies will greatly improve the efficiency and safety of genome editing with such aims, by integrating transgenes at specific loci, using homologous recombination.

The challenge of target specificity of synthetic nucleases is being actively researched and refined [64]. Researchers are devising methods to improve specificity and avoid potential off-target effects, such as using DNA nickases rather than full cleavases [65]. Nickases induce breaks in just one DNA strand, and they enhance homology-directed repair without activating non-homologous end joining repair. Although the CRISPR/Cas9 system is rapidly emerging as the easiest system to apply – simply requiring cloning of a guide RNA sequence to reprogram nuclease specificity – it originally had a weakness in terms of target specificity. Generally, >18 bp are required to specify a unique locus in the human genome. Although the CRISPR/Cas9 recognition complex covers 20 bp, ~6 bp of the DNA recognition site tolerates mismatches, opening up the possibility of off-target effects. By combining two nickases, Ran et al. effectively doubled the specific binding target length, making unwanted off-target effects far less likely [66]. The use of bioinformatic tools to predict off-target effects can also increase target specificity [67].

CRISPR systems are very practical, but some applications will require sustained expression, and *in vivo* studies have not yet investigated the long-term consequences of expressing these bacterial factors in the brain. The prokaryotic TALE effectors have also only been used for short studies: for example, when a light-controlled TALE was injected in mouse brain [68], animals were sacrificed only eight days post injection. By contrast, we have used eukaryotic ZFs to modulate the gene causing HD for up to four weeks in mouse brains [12], and can potentially make the entire protein sequence mouse-like, to avoid the immune system.

While *in vivo* techniques may be essential for complex diseases such as PD, in which discreet populations of neurons are lost, single-gene metabolic neurodegenerative diseases naturally suit the more tractable *ex vivo* approaches. As well as not having to consider immune rejection, one can deliver

genome editing tools in the form of RNA *ex vivo*, hence eliminating the risks of long-term expression or unwanted random integration. Moreover, if stem cells can be modified, they can provide a long-term therapeutic benefit [62, 63]. Pluripotent stem cell transplant technology for the brain is still in its infancy, and it is still not clear how genome-edited neurons would be rewired correctly. While this major hurdle will require further research, genome editing already has other useful applications such as creating disease models (Box 2).

## Engineering approaches improve the clearance of aberrant proteins



One of the common hallmarks of neurodegenerative diseases is the aggregation of mutated proteins that possess seeding activity and can be transmitted in a prion-like fashion [69]. A strategy to prevent aggregation is the administration of antibodies targeting these proteins. The advantage of this approach is that the antibodies can be delivered directly to the brain with minipumps, without the use of viral vectors, and without the need of modifying the genome, hence preventing undesired off-target effects. For example, a recent study in a mouse model of AD showed that selected antibodies administered chronically for three weeks to the brain ventricle decreased aggregation and improved cognitive deficits [70]. These antibodies can also be administered as vaccines against prion-like diseases, and several clinical trials are on-going [71]. Monoclonal antibodies are expensive to generate, however, and their production requires the use of living animals. Synthetic antibodies made by protein engineering offer a good alternative to naturally generated antibodies, with the possibility of improved specificity [72].

Antibodies are not the only peptides that can help clear mutant protein aggregates. Interactions with some natural proteins prevents aggregation and may be used as a therapeutic approach [73]. Bauer et al. [74] tackled mutant Huntingtin protein in HD by designing a polyglutamine-binding peptide, fused to heat shock cognate protein 70 binding motif, such that it would promote degradation of aggregates by chaperone-mediated autophagy. Impressively, AAV delivery of this synthetic protein had a strong therapeutic effect in the R6/2 HD mouse model. Thus, synthetic biology principles can be used for the design and production of engineered proteins that will act with high affinity to target mutated proteins and prevent aggregation.

## Synthetic gene circuits: Creating intelligent genetic therapies

Synthetic biology can go beyond designed therapeutic proteins, expressed by simple promoters, and can achieve the precise control of gene expression by means of new promoter designs, gene switches, and circuit engineering.

Current gene therapy vector constructs utilise a relatively limited range of promoters to drive transgene expression. The

**Box 2****Genome editing for neurodegenerative disease models****1. Microinjection of fertilized monkey eggs with Cas9 and guide RNA****2. Embryonic culture and genetic screening****3. Transfer to surrogate mother****4. Genetically modified non-human primate model of neurodegenerative disease**

Synthetic nucleases have already been used to create brain disease models. For example, using ZF nucleases, PD, and AD model cell lines expressing mutations were created and, conversely, mutations in patient-derived induced pluripotent cells were corrected [101, 102]. Such techniques not only underpin the hope that we will be able to model diseases in a dish [103], building systems at will in order to increase our understanding, but they also pave the way for future therapies based on genome-edited patient stem cell transplants.

Although useful, cell culture approaches unfortunately cannot recapitulate the complex environment of an organism completely; factors such as immunity and cell-cell communication are of extreme importance. Moreover, the common inbred rodent models used are far removed from many aspects of human physiology in terms of both size and genetic diversity. Reflecting the move towards more personalised medicine, synthetic biology tools are helping to develop genetically modified animal models beyond the mouse, more rapidly and more efficiently [104–106]. For example, using three plasmids coding TALENs, the *MECP2* gene was knocked down in monkey embryos with no detectable off-target mutations [107]. *MECP2* deficiency is the cause of Rett syndrome, a severe neurodevelopmental disorder affecting only females (since the mutation is lethal for male embryos). A non-human primate model of this disease will help to shed light on the autism-related phenotype, and will enable more efficient and realistic pre-clinical trials. Potentially, this method could be used to create other neurodegenerative disease models in any non-human primate species, such as marmosets (see picture), an increasingly popular model species for neuroscience research due to its small size, ease of breeding and brain features comparable to humans [108, 109].

combined CMV enhancer chicken  $\beta$ -actin (CAG) promoter and woodchuck hepatitis virus post-transcriptional regulatory element (WPRE) [75] are widely used for strong constitutive expression, whereas cell-type specific promoters are often used to limit transgene expression to subsets of cells [76, 77]. Since quantitative promoter characterisation is a key aspect of synthetic biology [78, 79], new promoters should become available as a result.

Gene switches, based on small molecules or drugs administered to the patient, also have great potential for building conditional expression constructs in the brain. For example, the synthetic steroid hormone mifepristone – commonly used to treat Cushing's syndrome [80] – has been used as a gene switch in the brain. Mifepristone activated the production of GDNF, resulting in a neuroprotective treatment in a rat model of PD [81]. Furthermore, in some cases, the switch molecule can also directly treat disease symptoms, resulting in double treatments [82].

An alternative and increasingly popular genetic switch is light-driven: optogenetics has revolutionized the field of

neuroscience because of its precision and rapid kinetics [83]. Light-controlled systems have successfully modulated neuron specific genes both in primary culture [84] and in the brains of mice: a TALE binding domain fused to plant cryptochrome 2 (sensitive to blue light) thus interacted conditionally with its partner protein CIB1, itself fused to an activator domain [68]. Moreover, the use of fibre optics devices for gene control in the brain is feasible. Implanted electrodes for stimulation are already in use for PD patients [85], and experimental models in which neuronal activity is controlled in vivo by optogenetic techniques is achieving great refinement [86].

Moving from switches to full genetic circuits, perhaps the most exciting developments will come from intelligent constructs that integrate multiple inputs to get conditional outputs. Mammalian gene circuit engineering is still in its infancy, and most examples still focus on systems to make desired patterns of gene expression [19, 87]. However, several research projects are developing genetic logic gates, analogous to the gates used in electronics [10, 11]. For example, AND gates require two simultaneous activation inputs, such as the



binding of heterodimer parts of transcription factors, to achieve conditional expression of a desired output gene (Fig. 1). Based on similar principles, cancer-detection circuits are being developed [13, 88], and it is only a matter of time before these are adapted to gene therapy applications.

## Will artificial cell transplants achieve long-term neuroprotection?

One of the most ambitious aims of synthetic biologists is to create synthetic cells equipped with tightly regulated gene networks to transplant into host organisms. These would release therapeutic compounds in a controlled manner to help organisms to self-repair. Such cells need to be carefully isolated from the host to prevent immune rejection and undesired proliferation. This is achieved by encapsulation with a biocompatible, semipermeable material, such as alginate-poly-(L-lysine)-alginate, which allows the exchange of essential biomolecules and ions, while isolating the cells from the immune system of the host [82, 89, 90].

In recent work following this strategy, a biosensor cell was built that could sense a change in the neurotransmitter dopamine, which is involved in processes such as movement and motivation [91]. The D1 dopamine receptor was coupled to an intracellular signalling cascade to activate the production of atrial natriuretic peptide (ANP), a natural anti-hypertensive molecule. Implanting these genetically engineered encapsulated cells in the peritoneum of hypertensive mice demonstrated that the production of ANP was enhanced in a controllable way by dopamine agonists or sexual arousal. This reduced and controlled the blood pressure of the mice in both situations [89].

Pharmacological and gene therapies can be combined for diseases with multiple symptoms. To treat metabolic syndrome, a chimeric TAAR (trace amine-associated receptor, coupled to G protein) was rewired to activate two genes controlling food intake, namely glucagon-like peptide 1 and leptin. The chimeric receptor responds to guanabenz, a drug clinically prescribed to treat hypertension. Microencapsulated cells bearing this system were implanted in the peritoneal cavity of mice. This double treatment was successful in attenuating several symptoms of metabolic syndrome in obese mice [82].

Encapsulated cells have long been used to release neurotrophic factors in the treatment of neurodegenerative diseases [90], but the main outcomes of those clinical trials have been the general safety of the implants, rather than strong therapeutic results. Problems have included the limited distribution of secreted neurotrophic factors, and fluctuating production and responses, caused by circadian rhythms. Mammalian synthetic biology researchers are building artificial oscillators, inspired by circadian clocks [92], and tunable secretion and cell-cell communication systems [19]. In the future, we may build more complex sender-receiver systems, including feedback elements that enhance the distribution of secreted therapeutic factors.

As noted in the introduction, successful treatments for neurodegenerative diseases require molecular biomarkers for

accurate early prediction. It was recently discovered that the depletion of ten lipids in peripheral blood predicts the occurrence of amnesic mild cognitive impairment or AD within 2–3 years, with over 90% accuracy [93]. Similarly, an auto-antibody against a potassium channel, whose dysfunction is related to neurodegeneration [94], was found in multiple sclerosis patients before symptom onset [95]. If such biomarkers could be linked to biosensor circuits in encapsulated cells [96], for the conditional production of neurotrophic factors, one could envisage predictive and protective strategies for neurodegenerative diseases. These would aim to delay onset and to slow disease progression, and would be targeted to susceptible populations.

Artificial cells can also be used to release specific factors for the promotion of self-repair. Adult neurogenesis is active in some specific niches in the human brain, and this process is dysregulated in neurodegenerative disorders [97]. The recent discovery of adult neurogenesis in the human striatum [98] opens the exciting possibility of investigating the regulatory factors required for cell migration and differentiation. A deeper knowledge of the factors, and their roles in diseases like HD and PD, would allow the design of regulators to enhance these processes, perhaps in combination with neuroprotection. Another issue is that projection neurons (large neurons wiring main circuits) are lost in such diseases, while adult-produced cells only differentiate into interneurons (small neurons that modulate the main circuits) and glia. The challenges are to design regulators that could further influence the fate of the progenitors to differentiate into projection neurons, and to promote rewiring.

## Conclusions and outlook

Synthetic biology can provide many useful approaches for the rational design of therapeutic factors and intelligent systems to treat a variety of diseases, including targeting the most complicated organ in our body, the brain. We can already find examples of the design of factors for genome editing, the regulation of gene expression, protein targeting, and for biosensor cells able to release therapeutic molecules in a tightly regulated manner, both in response to endogenous signals or to drugs. With all these new tools, we will be able to repair or to complement the pathogenic genes in specific conditions, such as those with single-gene metabolic causes. For many diseases, we will need to establish combined protective and restorative actions. Harnessing natural processes to promote self-repair and an improved basic knowledge of the underlying development and circuitry of the brain will be essential for developing new synthetic therapies that act in concert with host physiology. Early detection and intervention is crucial for beating the devastation caused by neurodegeneration. Perhaps the most exciting long-term possibility is to link biomarkers to biosensors, to express effectors conditionally, and to reverse pathogenesis as early as possible. Although the work has begun, we still need a lot more research to design a safe and long-term gene or cell therapy based on synthetic biology principles. Using nature as our inspiration, we will continue to improve our intelligent synthetic designs.

## Acknowledgements

CAP and MI are funded by the European Research Council grant FP7 ERC 201249 ZINC-HUBS. MI is supported by Wellcome Trust UK New Investigator Award WT102944MA. MI is a co-applicant on a patent application related to engineered zinc fingers that bind expanded poly-CAG repeats in Huntington's disease (WO2012/049332A1).

## References

1. Reardon S. 2014. Brain-mapping projects to join forces. *Nat News* doi: 10.1038/nature.2014.14871
2. Evans SJ, Douglas I, Rawlins MD, Wexler NS, et al. 2013. Prevalence of adult Huntington's disease in the UK based on diagnoses recorded in general practice records. *J Neurol Neurosurg Psychiatry* **84**: 1156–60.
3. Thies W, Bleiler L, Alzheimer's A. 2013. 2013 Alzheimer's disease facts and figures. *Alzheimers Dement* **9**: 208–45.
4. Byers AL, Yaffe K. 2011. Depression and risk of developing dementia. *Nat Rev Neurol* **7**: 323–31.
5. Wittchen HU, Jacobi F, Rehm J, Gustavsson A, et al. 2011. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol* **21**: 655–79.
6. Lu B, Nagappan G, Guan X, Nathan PJ, et al. 2013. BDNF-based synaptic repair as a disease-modifying strategy for neurodegenerative diseases. *Nat Rev Neurosci* **14**: 401–16.
7. Julien CL, Thompson JC, Wild S, Yardumian P, et al. 2007. Psychiatric disorders in preclinical Huntington's disease. *J Neurol Neurosurg Psychiatry* **78**: 939–43.
8. Weintraub D, Burn DJ. 2011. Parkinson's disease: the quintessential neuropsychiatric disorder. *Mov Disord* **26**: 1022–31.
9. Menendez-Gonzalez M. 2014. The many questions on the use of biomarkers for neurodegenerative diseases in clinical practice. *Front Aging Neurosci* **6**: 45.
10. Auslander S, Auslander D, Muller M, Wieland M, et al. 2012. Programmable single-cell mammalian biocomputers. *Nature* **487**: 123–7.
11. Moon TS, Lou C, Tamsir A, Stanton BC, et al. 2012. Genetic programs constructed from layered logic gates in single cells. *Nature* **491**: 249–53.
12. Garriga-Canut M, Agustín-Pavón C, Herrmann F, Sanchez A, et al. 2012. Synthetic zinc finger repressors reduce mutant huntingtin expression in the brain of R6/2 mice. *Proc Natl Acad Sci USA* **109**: E3136–45.
13. Xie Z, Wroblewska L, Prochazka L, Weiss R, et al. 2011. Multi-input RNAi-based logic circuit for identification of specific cancer cells. *Science* **333**: 1307–11.
14. Shalem O, Sanjana NE, Hartenian E, Shi X, et al. 2014. Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* **343**: 84–7.
15. Miller JC, Tan S, Qiao G, Barlow KA, et al. 2011. A TALE nuclease architecture for efficient genome editing. *Nat Biotechnol* **29**: 143–8.
16. Urnov FD, Miller JC, Lee YL, Beausejour CM, et al. 2005. Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature* **435**: 646–51.
17. Gardner TS, Hawkins K. 2013. Synthetic biology: evolution or revolution? A co-founder's perspective. *Curr Opin Chem Biol* **17**: 871–7.
18. Basu S, Gerchman Y, Collins CH, Arnold FH, et al. 2005. A synthetic multicellular system for programmed pattern formation. *Nature* **434**: 1130–4.
19. Carvalho A, Menendez DB, Senthivel VR, Zimmermann T, et al. 2014. Genetically encoded sender-receiver system in 3D mammalian cell culture. *ACS Synth Biol* **3**: 264–72.
20. Ro DK, Paradise EM, Ouellet M, Fisher KJ, et al. 2006. Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature* **440**: 940–3.
21. Regot S, Macia J, Conde N, Furukawa K, et al. 2011. Distributed biological computation with multicellular engineered networks. *Nature* **469**: 207–11.
22. Nawroth JC, Lee H, Feinberg AW, Ripplinger CM, et al. 2012. A tissue-engineered jellyfish with biomimetic propulsion. *Nat Biotechnol* **30**: 792–7.
23. Yu D, Pendergraff H, Liu J, Kordasiewicz HB, et al. 2012. Single-stranded RNAs use RNAi to potently and allele-selectively inhibit mutant huntingtin expression. *Cell* **150**: 895–908.
24. Bartus RT, Baumann TL, Siffert J, Herzog CD, et al. 2013. Safety/feasibility of targeting the substantia nigra with AAV2-neurturin in Parkinson patients. *Neurology* **80**: 1698–701.
25. Palfi S, Gurruchaga JM, Ralph GS, Lepetit H, et al. 2014. Long-term safety and tolerability of ProSavin, a lentiviral vector-based gene therapy for Parkinson's disease: a dose escalation, open-label, phase 1/2 trial. *Lancet* **383**: 1138–46.
26. Biffi A, Montini E, Lorioli L, Cesani M, et al. 2013. Lentiviral hematopoietic stem cell gene therapy benefits metachromatic leukodystrophy. *Science* **341**: 1233158.
27. Ross CA, Tabrizi SJ. 2011. Huntington's disease: from molecular pathogenesis to clinical treatment. *Lancet Neurol* **10**: 83–98.
28. Fjord-Larsen L, Kusk P, Emerich DF, Thanos C, et al. 2012. Increased encapsulated cell biodelivery of nerve growth factor in the brain by transposon-mediated gene transfer. *Gene Ther* **19**: 1010–7.
29. Wahlberg LU, Lind G, Almqvist PM, Kusk P, et al. 2012. Targeted delivery of nerve growth factor via encapsulated cell biodelivery in Alzheimer disease: a technology platform for restorative neurosurgery. *J Neurosurg* **117**: 340–7.
30. Stolberg SG. 1999. The biotech death of Jesse Gelsinger. *NY Times Mag* **136–40**: 49–50.
31. Basner-Tschakarjan E, Bijjiga E, Martino AT. 2014. Pre-clinical assessment of immune responses to adeno-associated virus (AAV) vectors. *Front Immunol* **5**: 28.
32. Howe SJ, Mansour MR, Schwarzwaelder K, Bartholomae C, et al. 2008. Insertional mutagenesis combined with acquired somatic mutations causes leukemogenesis following gene therapy of SCID-X1 patients. *J Clin Invest* **118**: 3143–50.
33. Papayannakos C, Daniel R. 2013. Understanding lentiviral vector chromatin targeting: working to reduce insertional mutagenic potential for gene therapy. *Gene Ther* **20**: 581–8.
34. Ciesielska A, Hadaczek P, Mittermeyer G, Zhou S, et al. 2013. Cerebral infusion of AAV9 vector-encoding non-self proteins can elicit cell-mediated immune responses. *Mol Ther* **21**: 158–66.
35. Hadaczek P, Forsayeth J, Mirek H, Munson K, et al. 2009. Transduction of nonhuman primate brain with adeno-associated virus serotype 1: vector trafficking and immune response. *Hum Gene Ther* **20**: 225–37.
36. Samaranch L, San Sebastian W, Kells AP, Salegio EA, et al. 2014. AAV9-mediated expression of a non-self protein in nonhuman primate central nervous system triggers widespread neuroinflammation driven by antigen-presenting cell transduction. *Mol Ther* **22**: 329–37.
37. Parker AS, Zheng W, Griswold KE, Bailey-Kellogg C. 2010. Optimization algorithms for functional deimmunization of therapeutic proteins. *BMC Bioinformatics* **11**: 180.
38. Perez-Pinera P, Ousterout DG, Brunger JM, Farin AM, et al. 2013. Synergistic and tunable human gene activation by combinations of synthetic transcription factors. *Nat Methods* **10**: 239–42.
39. Gilbert LA, Larson MH, Morsut L, Liu Z, et al. 2013. CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* **154**: 442–51.
40. Maeder ML, Linder SJ, Cascio VM, Fu Y, et al. 2013. CRISPR RNA-guided activation of endogenous human genes. *Nat Methods* **10**: 977–9.
41. Kordasiewicz HB, Stanek LM, Wancewicz EV, Mazur C, et al. 2012. Sustained therapeutic reversal of Huntington's disease by transient repression of huntingtin synthesis. *Neuron* **74**: 1031–44.
42. Singh NK, Singh NN, Androphy EJ, Singh RN. 2006. Splicing of a critical exon of human survival motor neuron is regulated by a unique silencer element located in the last intron. *Mol Cell Biol* **26**: 1333–46.
43. Hua Y, Vickers TA, Okunola HL, Bennett CF, et al. 2008. Antisense masking of an hnRNP A1/A2 intronic splicing silencer corrects SMN2 splicing in transgenic mice. *Am J Hum Genet* **82**: 834–48.
44. Williams JH, Schray RC, Patterson CA, Ayitey SO, et al. 2009. Oligonucleotide-mediated survival of motor neuron protein expression in CNS improves phenotype in a mouse model of spinal muscular atrophy. *J Neurosci* **29**: 7633–8.
45. Zanetta C, Nizzardo M, Simone C, Monguzzi E, et al. 2014. Molecular therapeutic strategies for spinal muscular atrophies: current and future clinical trials. *Clin Ther* **36**: 128–40.
46. Vucicevic D, Schrewe H, Orom UA. 2014. Molecular mechanisms of long ncRNAs in neurological disorders. *Front Genet* **5**: 48.
47. Riboldi G, Zanetta C, Ranieri M, Nizzardo M, et al. 2014. Antisense oligonucleotide therapy for the treatment of C9ORF72 ALS/FTD diseases. *Mol Neurobiol* in press, doi: 10.1007/s12035-014-8724-7
48. Martin JN, Wolken N, Brown T, Dauer WT, et al. 2011. Lethal toxicity caused by expression of shRNA in the mouse striatum: implications for therapeutic design. *Gene Ther* **18**: 666–73.

49. **Khodr CE, Sapru MK, Pedapati J, Han Y, et al.** 2011. An alpha-synuclein AAV gene silencing vector ameliorates a behavioral deficit in a rat model of Parkinson's disease, but displays toxicity in dopamine neurons. *Brain Res* **1395**: 94–107.
50. **Ehlert EM, Eggers R, Niclou SP, Verhaagen J.** 2010. Cellular toxicity following application of adeno-associated viral vector-mediated RNA interference in the nervous system. *BMC Neurosci* **11**: 20.
51. **McBride JL, Boudreau RL, Harper SQ, Staber PD, et al.** 2008. Artificial miRNAs mitigate shRNA-mediated toxicity in the brain: implications for the therapeutic development of RNAi. *Proc Natl Acad Sci USA* **105**: 5868–73.
52. **Pinheiro VB, Taylor AI, Cozens C, Abramov M, et al.** 2012. Synthetic genetic polymers capable of heredity and evolution. *Science* **336**: 341–4.
53. **Morbitzer R, Romer P, Boch J, Lahaye T.** 2010. Regulation of selected genome loci using de novo-engineered transcription activator-like effector (TALE)-type transcription factors. *Proc Natl Acad Sci USA* **107**: 21617–22.
54. **Hu J, Lei Y, Wong WK, Liu S, et al.** 2014. Direct activation of human and mouse Oct4 genes using engineered TALE and Cas9 transcription factors. *Nucleic Acids Res* **42**: 4375–90.
55. **Mussolino C, Sanges D, Marrocco E, Bonetti C, et al.** 2011. Zinc-finger-based transcriptional repression of rhodopsin in a model of dominant retinitis pigmentosa. *EMBO Mol Med* **3**: 118–28.
56. **Kliem MA, Heeke BL, Franz CK, Radovitskiy I, et al.** 2011. Intramuscular administration of a VEGF zinc finger transcription factor activator (VEGF-ZFP-TF) improves functional outcomes in SOD1 rats. *Amyotroph Lateral Scler* **12**: 331–9.
57. **Rebar EJ, Huang Y, Hickey R, Nath AK, et al.** 2002. Induction of angiogenesis in a mouse model using engineered transcription factors. *Nat Med* **8**: 1427–32.
58. **Laganieri J, Kells AP, Lai JT, Guschin D, et al.** 2010. An engineered zinc finger protein activator of the endogenous glial cell line-derived neurotrophic factor gene provides functional neuroprotection in a rat model of Parkinson's disease. *J Neurosci* **30**: 16469–74.
59. **Ellison SM, Trabalza A, Tisato V, Pazarentzos E, et al.** 2013. Dose-dependent neuroprotection of VEGF(1)(6)(5) in Huntington's disease striatum. *Mol Ther* **21**: 1862–75.
60. **Christian M, Cermak T, Doyle EL, Schmidt C, et al.** 2010. Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics* **186**: 757–61.
61. **Mali P, Yang L, Esvelt KM, Aach J, et al.** 2013. RNA-guided human genome engineering via Cas9. *Science* **339**: 823–6.
62. **Li M, Suzuki K, Kim NY, Liu GH, et al.** 2014. A cut above the rest: targeted genome editing technologies in human pluripotent stem cells. *J Biol Chem* **289**: 4594–9.
63. **Tebas P, Stein D, Tang WW, Frank I, et al.** 2014. Gene editing of CCR5 in autologous CD4 T cells of persons infected with HIV. *N Engl J Med* **370**: 901–10.
64. **Fu Y, Foden JA, Khayter C, Maeder ML, et al.** 2013. High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. *Nat Biotechnol* **31**: 822–6.
65. **Wang J, Friedman G, Doyon Y, Wang NS, et al.** 2012. Targeted gene addition to a predetermined site in the human genome using a ZFN-based nicking enzyme. *Genome Res* **22**: 1316–26.
66. **Ran FA, Hsu PD, Lin CY, Gootenberg JS, et al.** 2013. Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* **154**: 1380–9.
67. **Hsu PD, Scott DA, Weinstein JA, Ran FA, et al.** 2013. DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol* **31**: 827–32.
68. **Konermann S, Brigham MD, Trevino AE, Hsu PD, et al.** 2013. Optical control of mammalian endogenous transcription and epigenetic states. *Nature* **500**: 472–6.
69. **Guo JL, Lee VM.** 2014. Cell-to-cell transmission of pathogenic proteins in neurodegenerative diseases. *Nat Med* **20**: 130–8.
70. **Yanamura K, Kfoury N, Jiang H, Mahan TE, et al.** 2013. Anti-tau antibodies that block tau aggregate seeding in vitro markedly decrease pathology and improve cognition in vivo. *Neuron* **80**: 402–14.
71. **Panza F, Logroscino G, Imbimbo BP, Solfrizzi V.** 2014. Is there still any hope for amyloid-based immunotherapy for Alzheimer's disease? *Curr Opin Psychiatry* **27**: 128–37.
72. **Adams JJ, Sidhu SS.** 2014. Synthetic antibody technologies. *Curr Opin Struct Biol* **24C**: 1–9.
73. **de Chiara C, Pastore A.** 2014. Kaleidoscopic protein-protein interactions in the life and death of ataxin-1: new strategies against protein aggregation. *Trends Neurosci* **37**: 211–8.
74. **Bauer PO, Goswami A, Wong HK, Okuno M, et al.** 2010. Harnessing chaperone-mediated autophagy for the selective degradation of mutant huntingtin protein. *Nat Biotechnol* **28**: 256–63.
75. **Garg S, Oran AE, Hon H, Jacob J.** 2004. The hybrid cytomegalovirus enhancer/chicken beta-actin promoter along with woodchuck hepatitis virus posttranscriptional regulatory element enhances the protective efficacy of DNA vaccines. *J Immunol* **173**: 550–8.
76. **Delzor A, Dufour N, Petit F, Guillermier M, et al.** 2012. Restricted transgene expression in the brain with cell-type specific neuronal promoters. *Hum Gene Ther Meth* **23**: 242–54.
77. **Portales-Casamar E, Swanson DJ, Liu L, de Leeuw CN, et al.** 2010. A regulatory toolbox of MiniPromoters to drive selective expression in the brain. *Proc Natl Acad Sci USA* **107**: 16589–94.
78. **Kelly JR, Rubin AJ, Davis JH, Ajo-Franklin CM, et al.** 2009. Measuring the activity of BioBrick promoters using an in vivo reference standard. *J Biol Eng* **3**: 4.
79. **Trinklein ND, Aldred SJ, Saldanha AJ, Myers RM.** 2003. Identification and functional analysis of human transcriptional promoters. *Genome Res* **13**: 308–12.
80. **Katznelson L, Loriaux DL, Feldman D, Braunstein GD, et al.** 2014. Global clinical response in Cushing's syndrome patients treated with mifepristone. *Clin Endocrinol (Oxf)* **80**: 562–9.
81. **Maddalena A, Tereshchenko J, Bahr M, Kugler S.** 2013. Adeno-associated virus-mediated, mifepristone-regulated transgene expression in the brain. *Mol Ther Nucleic Acids* **2**: e106.
82. **Ye H, Charpin-El Hamri G, Zwicky K, Christen M, et al.** 2013. Pharmaceutically controlled designer circuit for the treatment of the metabolic syndrome. *Proc Natl Acad Sci USA* **110**: 141–6.
83. **Packer AM, Roska B, Hausser M.** 2013. Targeting neurons and photons for optogenetics. *Nat Neurosci* **16**: 805–15.
84. **Polstein LR, Gersbach CA.** 2012. Light-inducible spatiotemporal control of gene activation by customizable zinc finger transcription factors. *J Am Chem Soc* **134**: 16480–3.
85. **Katzenschlager R.** 2014. Parkinson's disease: recent advances. *J Neurol* **261**: 1031–6.
86. **Wykes RC, Heeroma JH, Mantoan L, Zheng K, et al.** 2012. Optogenetic and potassium channel gene therapy in a rodent model of focal neocortical epilepsy. *Sci Transl Med* **4**: 161ra52.
87. **Greber D, Fussenegger M.** 2010. An engineered mammalian band-pass network. *Nucleic Acids Res* **38**: e174.
88. **Nissim L, Bar-Ziv RH.** 2010. A tunable dual-promoter integrator for targeting of cancer cells. *Mol Syst Biol* **6**: 444.
89. **Rosser K, Charpin-El Hamri G, Fussenegger M.** 2013. Reward-based hypertenor control by a synthetic brain-dopamine interface. *Proc Natl Acad Sci USA* **110**: 18150–5.
90. **Emerich DF, Orive G, Thanos C, Tornøe J, et al.** 2013. Encapsulated cell therapy for neurodegenerative diseases: from promise to product. *Adv Drug Deliv Rev* **67–68**: 131–41.
91. **Salamone JD, Correa M.** 2012. The mysterious motivational functions of mesolimbic dopamine. *Neuron* **76**: 470–85.
92. **Tigges M, Marquez-Lago TT, Stelling J, Fussenegger M.** 2009. A tunable synthetic mammalian oscillator. *Nature* **457**: 309–12.
93. **Mapstone M, Cheema AK, Fiandaca MS, Zhong X, et al.** 2014. Plasma phospholipids identify antecedent memory impairment in older adults. *Nat Med* **20**: 415–8.
94. **Tong X, Ao Y, Faas GC, Nwaobi SE, et al.** 2014. Astrocyte Kir4.1 ion channel deficits contribute to neuronal dysfunction in Huntington's disease model mice. *Nat Neurosci* **17**: 694–703.
95. **Nerrant E, Charif M, Ayrignac X, Carra-Dalliere C, et al.** 2014. Is KIR4.1 antibody a marker of multiple sclerosis? *Neurology* **82**: Supplement P4.130.
96. **Gil B, Kahan-Hanum M, Skirtenko N, Adar R, et al.** 2011. Detection of multiple disease indicators by an autonomous biomolecular computer. *Nano Lett* **11**: 2989–96.
97. **Hamilton LK, Joppe SE, Cochard LM, Fernandes KJ.** 2013. Aging and neurogenesis in the adult forebrain: what we have learned and where we should go from here. *Eur J Neurosci* **37**: 1978–86.
98. **Ernst A, Alkass K, Bernard S, Salehpour M, et al.** 2014. Neurogenesis in the striatum of the adult human brain. *Cell* **156**: 1072–83.
99. **Stockwell J, Abdi N, Lu X, Maheshwari O, et al.** 2014. Novel central nervous system drug delivery systems. *Chem Biol Drug Des* **83**: 507–20.
100. **Mulcahy PJ, Iremonger K, Karyka E, Herranz-Martin S, et al.** 2014. Gene therapy: a promising approach to treating SMA. *Hum Gene Ther* **25**: 575–86.
101. **Soldner F, Laganieri J, Cheng AW, Hockemeyer D, et al.** 2011. Generation of isogenic pluripotent stem cells differing exclusively at two early onset Parkinson point mutations. *Cell* **146**: 318–31.

102. **Fong H, Wang C, Knoferle J, Walker D**, et al. 2013. Genetic correction of tauopathy phenotypes in neurons derived from human induced pluripotent stem cells. *Stem Cell Rep* **1**: 226–34.
103. **Marchetto MC, Gage FH**. 2012. Modeling brain disease in a dish: really? *Cell Stem Cell* **10**: 642–5.
104. **Zhou J, Shen B, Zhang W, Wang J**, et al. 2014. One-step generation of different immunodeficient mice with multiple gene modifications by CRISPR/Cas9 mediated genome engineering. *Int J Biochem Cell Biol* **46**: 49–55.
105. **Hwang WY, Fu Y, Reyon D, Maeder ML**, et al. 2013. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol* **31**: 227–9.
106. **Niu Y, Shen B, Cui Y, Chen Y**, et al. 2014. Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated gene targeting in one-cell embryos. *Cell* **156**: 836–43.
107. **Liu H, Chen Y, Niu Y, Zhang K**, et al. 2014. TALEN-mediated gene mutagenesis in rhesus and cynomolgus monkeys. *Cell Stem Cell* **14**: 323–8.
108. **Agustin-Pavon C, Braesicke K, Shiba Y, Santangelo AM**, et al. 2012. Lesions of ventrolateral prefrontal or anterior orbitofrontal cortex in primates heighten negative emotion. *Biol Psychiatry* **72**: 266–72.
109. **Kishi N, Sato K, Sasaki E, Okano H**. 2014. Common marmoset as a new model animal for neuroscience research and genome editing technology. *Dev Growth Differ* **56**: 53–62.
110. **Adolfsson O, Pihlgren M, Toni N, Varisco Y**, et al. 2012. An effector-reduced anti-beta-amyloid (Abeta) antibody with unique abeta binding properties promotes neuroprotection and glial engulfment of Abeta. *J Neurosci* **32**: 9677–89.
111. **Gross RE, Watts RL, Hauser RA, Bakay RA**, et al. 2011. Intraatrial transplantation of microcarrier-bound human retinal pigment epithelial cells versus sham surgery in patients with advanced Parkinson's disease: a double-blind, randomised, controlled trial. *Lancet Neurol* **10**: 509–19.
112. **Mittermeyer G, Christine CW, Rosenbluth KH, Baker SL**, et al. 2012. Long-term evaluation of a phase 1 study of AADC gene therapy for Parkinson's disease. *Hum Gene Ther* **23**: 377–81.
113. **Kefalopoulou Z, Politis M, Piccini P, Mencacci N**, et al. 2014. Long-term clinical outcome of fetal cell transplantation for Parkinson disease: two case reports. *JAMA Neurol* **71**: 83–7.
114. **Herrmann F, Garriga-Canut M, Baumstark R, Fajardo-Sanchez E**, et al. 2011. p53 Gene repair with zinc finger nucleases optimised by yeast 1-hybrid and validated by Solexa sequencing. *PLoS One* **6**: e20913.
115. **Bibikova M, Carroll D, Segal DJ, Trautman JK**, et al. 2001. Stimulation of homologous recombination through targeted cleavage by chimeric nucleases. *Mol Cell Biol* **21**: 289–97.
116. **Li T, Huang S, Jiang WZ, Wright D**, et al. 2011. TAL nucleases (TALNs): hybrid proteins composed of TAL effectors and FokI DNA-cleavage domain. *Nucleic Acids Res* **39**: 359–72.
117. **Cong L, Ran FA, Cox D, Lin S**, et al. 2013. Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**: 819–23.