

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

#### Table 1. (continued)

SDG	Blue biotechnology solution	
14 UFE BELOW WATER	Sustainable economic models, circular economy, and biobased businesses growth Conservation of natural resources and ecosystem services as economic assets, genomics for assisted evolution to promote more resilient phenotypes, and ecosystem management to promote biodiversity and stock sustainability	
15 LIFE ON LAND		

from the fish processing industry into pricey fashion products [14], fish bones bioprocessed into high-value collagen for cosmeceutical, and fish scales used for bone tissue engineering applications [15]). Finally, at the scientific and technological challenges level, blue biotechnology is driven by innovation. Innovative tools, such as miniature next-generation sequencing (NGS) or lab-on-a-chip and microfluidic devices, are paving the way for further and deeper marine exploration. By facilitating sample collection and allowing onboard and laboratory growth of specimens collected under extreme deep-sea conditions, blue biotechnology continues to unravel the 'blue gold': the hidden chemodiversity of world oceans. The use of biotechnology to solve challenges in aquaculture in aquafeed (e.g., formulations), overfishing, integrated ecosystem management approaches, and the production of new species (namely, microorganisms grown *ex situ*) is now generalized and will continue to expand. Most importantly, the recent range of novel hologenomic approaches and biodata gathering and mining are paving novel ways to interpret and use this information for as-yet-unforeseen applications that are likely to promote disruptive breakthroughs in science and technology. By building bridges between technology, stakeholders, and the economy, blue biotechnology is among the most innovative, yet overlooked,

innovation ecosystems that can contribute to a bluer bioeconomic future.

#### Acknowledgments

We acknowledge the financial support to CESAM (UIDB/50017/2020+UIDP/50017/2020) by FCT/MEC through national funds and the co-funding by FEDER, within the PT2020 Partnership Agreement and Compete 2020. We also acknowledge the financial support to BioISI (UIDB/04046/2020) by FCT/MEC through national funds and the co-funding by FEDER, within the PT2020 Partnership Agreement and Compete 2020 and the EU Commission via ongoing H2020 projects. M.C.L. was supported by the Integrated Programme of SR&TD 'SmartBioR – Smart Valorization of Endogenous Marine Biological Resources Under a Changing Climate' (Centro-01-0145-FEDER-000018), co-funded by the Centro 2020 program, Portugal 2020, EU, through the European Regional Development Fund.

<sup>1</sup>BiolSI – Biosystems and Integrative Sciences Institute, Faculty of Sciences, University of Lisboa, Lisboa, Portugal <sup>2</sup>ECOMARE, Centre for Environmental and Marine Studies (CESAM), Department of Biology, University of Aveiro, Santiago University Campus, 3810-193 Aveiro, Portugal <sup>9</sup>https://www.linkedin.com/in/helenavieirabio/ <sup>4</sup>https://www.linkedin.com/in/niguel-costa-leal-698595a <sup>6</sup>https://www.linkedin.com/in/nicardo-calado-55ba6b8/

\*Correspondence: hmvieira@fc.ul.pt (H. Vieira). https://doi.org/10.1016/j.tibtech.2020.03.011

© 2020 Elsevier Ltd. All rights reserved.

#### References

- 1. Sillanpää, M. et al. (2017) A Sustainable Bioeconomy, Springer
- Bracco, S. *et al.* (2018) Assessing the contribution of bioeconomy to the total economy: a review of national frameworks. *Sustainability* 10, 1698
- Ronzon, T. et al. (2015) The Bioeconomy in the European Union in Numbers. Facts and Figures on Biomass, Turnover and Employment, European Commission, Joint

Research Centre, Institute for Prospective Technological Studies

- Cleveland, B.M. (2019) A perspective of the future value and challenges of genetic engineering in aquaculture. *J. World Aquac. Soc.* 50, 890–893
- Chakravarti, L.J. and van Oppen, M.J. (2018) Experimental evolution in coral photosymbionts as a tool to increase thermal tolerance. *Front. Mar. Sci.* 5, 227
- 6. Macreadie, P.I. *et al.* (2019) The future of blue carbon science. *Nat. Commun.* 10, 3998
- Calado, R. *et al.* (2018) How to succeed in marketing marine natural products for nutraceutical, pharmaceutical and cosmeceutical markets. In *Grand Challenges in Marine Biotechnology*, pp. 317–403, Springer
- German Bioeconomy Council (2018) *Bioeconomy Policy* (*Part III*) – *Update Report of National Strategies around the World*, Office of the Bioeconomy Council
- 9. Ligtvoet, A. et al. (2019) Blue Bioeconomy Forum: Roadmap for the Blue Bioeconomy, European Commission
- Hurst, D. et al. (2016) Marine Biotechnology Strategic Research and Innovation Roadmap: Insights to the Future Direction of European Marine Biotechnology, Marine Biotechnology ERA-NET
- 11. Vasconcelos, V. et al. (2019) Portugal Blue Bioeconomy Roadmap, CIIMAR
- Caporgno, M.P. and Mathys, A. (2018) Trends in microalgae incorporation into innovative food products with potential health benefits. *Front. Nutr.* 5, 58
- Alves, A. et al. (2020) Exploring trade-offs among the multiple benefits of green-blue-grey infrastructure for urban flood mitigation. Sci. Total Environ. 703, 134980
- Mallon, J. (2019) Is fish skin the new frontier for ecofriendly fashion? *Fashion United*. Published online November 5, 2019. https://fashionunited.uk/news/ fashion/is-fish-skin-the-new-frontier-for-eco-friendlyfashion/2019110546042
- Kara, A. et al. (2019) Bioactive fish scale incorporated chitosan biocomposite scaffolds for bone tissue engineering. Int. J. Biol. Macromol. 130, 266–279

## **Science & Society**

Working Hard or Hardly Working? Regulatory Bottlenecks in Developing a COVID-19 Vaccine

Lisette Pregelj <sup>(b)</sup>,<sup>1,5,@</sup> Damian C. Hine <sup>(b)</sup>,<sup>2,5,@</sup> Maria G. Oyola-Lozada <sup>(b)</sup>,<sup>3,@</sup> and Trent P. Munro <sup>(b)</sup>,<sup>4,\*,@</sup>

Vaccine solutions rarely reach the public until after an outbreak



abates; an Ebola vaccine was approved 5 years after peak outbreak and SARS, MERS, and Zika vaccines are still in clinical development. Despite massive leaps forward in rapid science, other regulatory bottlenecks are hamstringing the global effort for pandemic vaccines.

When an international team of researchers led by Jeremy Farrar from the Wellcome Trust published an article last November in Nature calling for theories and practices underpinning infectious disease epidemiology to keep pace with the translational nature of epidemics in the 21st century [1], few thought the next pandemic would come so soon. Yet, when the COVID-19 outbreak was confirmed by Wuhan Municipal Health Commission on 31 December 2019, the scientific community acted swiftly, collaborating intensely across disciplines and sharing new knowledge in a truly open innovation approach to advancing scientific research [2-5].

The combined efforts of the scientific community, mobilization of public health resources, increased public and private investment, and a 'completely new culture of doing research' [6] accelerated the first novel vaccine candidate through discovery, lead-candidate optimization, preclinical development, and into clinical trials within 2 months of the outbreak. Four additional candidates entered the clinic within the next month and 18 more are expected to begin clinical trials by late 2020 [7].

#### Rapid Development

The rapid spread of COVID-19 helps us understand the volume, diversity, and quality of resources mobilized to realize these unprecedented vaccine development speeds, especially when we compare them with the response to other global outbreaks over the last two decades (Figure 1). The first novel vaccine candidate for COVID-19 began clinical trials 1 month faster than the first candidate for the H1N1 swine flu epidemic in 2009 (93 days) and 3 months faster than the first Ebola vaccine candidate in 2014 (167 days) (Table S1 in the supplemental information online). These vaccine candidates also started clinical trials ahead of their respective outbreak peaks (Figure 2).

Yet, only the H1N1 vaccine was approved in time to deliver a preventative solution to the then ongoing public health crisis. Novel vaccines against the viruses responsible for the 2003 severe acute respiratory syndrome (SARS), 2012 Middle East respiratory syndrome (MERS), and 2015 Zika virus outbreaks are still in clinical testing, long after their respective outbreaks peaked and the World Health Organization (WHO) declared the outbreaks over. While there is now an Ebola vaccine, it took 5 years to complete 12 supporting and one pivotal clinical trial, then be reviewed and approved (in 2019) by the European Medicines Agency (EMA) and the FDA Centre for Biological Evaluation and Research (CBER).

The reality is that a 5 year development program for a novel vaccine is fast. The only other novel vaccines approved by CBER in 2019, one for dengue fever (Dengvaxia, Sanofi) and one for smallpox and monkeypox (Jynneos, Bavarian Nordic), were both in clinical trials for approximately 15 years. Five-year clinical development is fast, even for drugs, and generally limited to ground-breaking precision medicines for indications with unmet medical need where the FDA's Centre for Drug Evaluation and Research works intensively with drug developers to impart their knowledge and expertise, built over decades and across thousands of new drug development programs [8,9].

To achieve the Ebola vaccines' 5-year clinical program, global regulators from developed countries combined and leveraged their collective expertise to assist the African Vaccine Regulatory Forum and worked closely with vaccine developer Merck to conduct supporting and pivotal trials when and where the outbreak was underway [10]. This significantly improved the regulatory and ethics timelines in key areas of presubmission discussions with sponsors and manufacturers, authorization of clinical trials and emergency use of investigational vaccines, and implementation of joint reviews for ethical and regulatory approvals [10,11]. The leadership of WHO was the cornerstone of this coordinated regulatory response.


#### **Regulatory Bottlenecks**

While there are ongoing collaborative efforts around the harmonization of regulatory frameworks in pharmaceuticals and medical devices, the ability to coordinate cross-national regulatory responses in times of crises remains a sticking point. For example, while Merck submitted the same evidence dossier for the Ebola vaccine to both the EMA and FDA at the same time, there were different meetings, review and response timelines, and questions from each, and ultimately the vaccine was approved 1 month earlier in Europe than the United States<sup>viii</sup>. Further, regulatory and ethical issues could make it challenging for SARS, MERS, and Zika candidates to even complete clinical testing now that their respective outbreaks are over. For instance, the development of a Zika presents vaccine still unresolved questions around the identification of immune markers likely to predict clinical benefit, the role of flavivirus immunity in vaccine safety, and ethical concerns related to the vaccination of pregnant women [12].

There is ongoing debate about how long a novel COVID-19 vaccine will really take to advance through trials, achieve regulatory







Trends in Biotechnology

Figure 1. Timing and Duration of Key Development Milestones towards a Vaccine in First 365 Days of Recent Major Global Outbreaks. Double lines at 365 days represent ongoing activities. Slanted lines indicate overlap in development milestones; the genomes of Middle East respiratory syndrome (MERS) and severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2; COVID-19) were sequenced and diagnostic tests developed before the World Health Organization (WHO) issued their alert. See Table S1 in the supplemental information online for dates of featured events.

approval across multiple jurisdictions, and be manufactured and distributed at the required scale to impact global immunity and prevent further transmission. Vaccines often present commercial viability challenges due to long, complex development timelines and 'capacity to pay' issues in developing countries. While some that achieve global population health success, such as the human papillomavirus vaccine, can provide substantial return on investment<sup>ix</sup>, these trends have meant that, compared with novel therapeutics, far fewer vaccines have made it through clinical testing. As a result, regulators and vaccine developers lack experience in vaccine development comparable with novel therapeutics [13]. This could become increasingly

problematic if all current COVID-19 clinical and preclinical vaccine candidates are simultaneously vying for regulators' attention and resources to expeditiously review clinical protocols and evidence dossiers. Not to mention disruption to other clinical programs, as regulators are redirected to reviewing the clinical protocols of dozens of COVID-19 vaccine candidates and even more therapeutic candidates. Beyond these bottlenecks, the underlying support required within bioanalytical and pathology providers will also be critical to ensure rapid availability and interpretation of clinical data.


The changing dynamics of the disease as it continues to emerge could also affect the development of vaccines when trial designs deliver solutions at an earlier stage of an

are informed by the epidemiology of the virus. As governments implement lockdowns and nonessential travel and transmission rates decline, clinical trial recruitment will become challenging across the potentially dozens of independent clinical programs striving for patients and healthy volunteers. In addition, patient screening for COVID-19negative status and monitoring of potential virus exposure while on study will create additional challenges and opportunities for early trial endpoints.

#### **Potential Solutions**

If regulatory bottlenecks prevent solutions being brought to the public in time to reduce the impact of an epidemic, then we need to consider alternatives that





#### Trends in Biotechnology

Figure 2. Number of Reported Cases per Month and Key Events in the Development of a Novel Vaccine for Global Epidemics and Pandemics over the Last Two Decades. See Table S1 in the supplemental information online for key events. Data sources: severe acute respiratory syndrome (SARS): World Health Organization (WHO)<sup>1</sup>; H1N1: WHO<sup>11</sup>, FluNet/WHO Global Influenza Surveillance and Response System (GISRS)<sup>111</sup> (H1N1 data comes from surveillance of global circulation of influenza viruses; number of specimens positive for H1N1 reported to GISRS); Middle East respiratory syndrome (MERS): European Centre for Disease Prevention and Control (ECDC)<sup>11</sup>; Ebola: Centers for Disease Control and Prevention (CDC)<sup>12</sup>; Zika: Pan American Health Organization (PAHO)/WHO<sup>11</sup>; COVID19: Johns Hopkins University<sup>11</sup> (COVID-19 cases for month 4 are from 1 April 2020 to 27 April 2020).

epidemic, in the geographical jurisdictions where the outbreak is occurring. Governments can leverage harmonized regulatory standards, mutual recognition, and regulatory reliance mechanisms to enhance existing regulatory pathways and accelerate and coordinate approvals globally<sup>×</sup>. This would be particularly beneficial in countries that traditionally suffer submission and approval lags for novel drugs and vaccines.

With the data at hand from previous outbreaks, we can consider predictive epidemiological models that can guide globally coordinated responses ahead of the curve and take advantage of the availability of patients across what are typically northern and southern hemisphere seasonal outbreaks. Public-private partnerships, such as the Accelerating COVID-19 Therapeutic Interventions and Vaccines partnership led by the National Institutes of Health, and programs for expedited responses from leading regulators such as the FDA's Coronavirus Treatment Acceleration Program and the EMA's Rapid Response team could be preemptively designed and developed for global health emergencies, rather than introduced during an outbreak, and when governments with advanced health systems such as Australia, New Zealand, South Korea, Hong Kong, Taiwan, and Singapore have already flattened their curves. This may point to an expanded leadership role for international organizations like WHO to coordinate regulatory jurisdictions in a more focused and targeted manner. In short, global coordination of regulatory efforts for COVID-19 are still lagging research and clinical development efforts.

We also need to be better prepared in the areas of preclinical science, including animal challenge models, clinical trial pathways from safety to efficacy, and for innovative and adaptive clinical trial designs such as umbrella studies (where multiple candidates are tested under a single protocol) to become the norm in times of epidemics. While some of these trial initiatives have started to emerge, such as the SOLIDARITY trial, they are still few and far between. This is especially problematic considering there are currently over 500 randomized controlled studies for COVID-19 preventatives and treatments, 31 of which have at least two arms and a target sample size of at least 1000 patients (some

with recruitment targets of 40 000–55 000) [14]. Large, randomized and placebocontrolled studies have been associated with longer total development times, as opposed to smaller, nonrandomized singlearm or extension-arm studies [8]. Last, guidance documents for pandemic trials should be available in anticipation of pandemics, ready to be triggered as soon as the WHO declares a public health emergency, and remain in effect after the public health crisis has abated.

While it is clear the scientific community's attention is now shifting towards regulatory solutions and regulators are demonstrating flexible regulatory processes, there is no time for competition and fragmentation among governments and regulators. Regulation requires collaboration, especially as a solution would provide neither prophylactic nor therapeutic benefit to communities, or commercial return to the developer, if marketed after the peak of the outbreak.

#### Acknowledgments

The authors would like to thank the anonymous reviewer and the Editor for comments and suggestions to improve the manuscript. M. Oyola-Lozada would like to acknowledge Peru's National Council for Science, Technology and Technological Innovation (CONCYTEC) for supporting her doctoral research at The University of Queensland.

#### **Disclaimer Statement**

Professor Munro is Project Director of the Coalition for Epidemic Preparedness Innovations (CEPI) funded Vaccine Rapid Response pipeline at the University of Queensland.

#### Supplemental Information

Supplemental information associated with this article can be found online at https://doi.org/10.1016/j. tibtech.2020.06.004.

#### Resources

<sup>i</sup>www.who.int/csr/sars/country/en/

<sup>ii</sup>www.who.int/csr/disease/swineflu/updates/en/

<sup>iii</sup>http://apps.who.int/flumart/Default?ReportNo=10

<sup>iv</sup>www.ecdc.europa.eu/en/publications-data/ communicable-disease-threats-report-31-december-2017-6-january-2018-week-1

<sup>v</sup>www.cdc.gov/vhf/ebola/history/2014-2016outbreak/cumulative-cases-graphs.html#anchor\_ 1515096547931

<sup>vi</sup>www.paho.org/data/index.php/es/?option=com\_ content&view=article&id=529:zika-subregionses&ltemid=353

viihttps://gisanddata.maps.arcgis.com/apps/ opsdashboard/index.html#/

bda7594740fd40299423467b48e9ecf6

viiihttps://cdn.ymaws.com/www.casss.org/resource/ resmgr/wcbp/wcbp\_2020/Duffy\_Kim\_Hassis\_Kim\_ Slides.pdf

<sup>Ix</sup>www.mrknewsroom.com/newsroom/newsreleases/news-details/2020/Merck-Announces-Fourth-Quarter-and-Full-Year-2019-Financial-Results/default.aspx

<sup>x</sup>www.ifpma.org/resource-centre/covid-19biopharmaceutical-industry-regulatory-guidingprinciples/

<sup>1</sup>School of Chemistry and Molecular Biosciences and Australian Institute for Business and Economics, The University of Queensland, St Lucia, Queensland 4072, Australia

<sup>2</sup>Kemmy Business School, University of Limerick, Limerick, V94 T9PX, Ireland

<sup>3</sup>School of Chemistry and Molecular Biosciences and Faculty of Business, Economics and Law, The University of Queensland, St Lucia, Queensland 4072, Australia

<sup>4</sup>Australian Institute for Bioengineering and Nanotechnology, The University of Queensland, St Lucia,

Queensland 4072, Australia <sup>5</sup>These authors contributed equally to this work.

#### \*Correspondence:

t.munro@uq.edu.au(T.P. Munro)

@Twitter: @lisettepregelj (L. Pregelj), @hine\_damian (D.C. Hine), @g\_oyola (M.G. Oyola-Lozada), and @trent\_munro (T.P. Munro). https://doi.org/10.1016/j.tibtech.2020.06.004

© 2020 Elsevier Ltd. All rights reserved.

#### References

Bedford, J. *et al.* (2019) A new twenty-first century science for effective epidemic response. *Nature* 575, 130–136

Dong, E. *et al.* (2020) An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect. Dis.* 20, 533–534

- Xu, B. *et al.* (2020) Open access epidemiological data from the COVID-19 outbreak. *Lancet Infect. Dis.* 20, 534
  Chu, D.K.W. *et al.* (2020) Molecular diagnosis of a novel
- Chit, D.A.W. et al. (2020) Molecular diagnosis of a hover coronavirus (2019-nCoV) causing an outbreak of pneumonia. *Clin. Chem.* 66, 549–555
- Corman, V. *et al.* (2012) Detection of a novel human coronavirus by real-time reverse-transcription polymerase chain reaction. *Eurosurveillance* 17, 20285
- Kupferschmidt, K. (2020) 'A completely new culture of doing research.' Coronavirus outbreak changes how scientists communicate. *Science* Published online February 26, 2020. https://doi.org/10.1126/science.abb4761
- Le, T. et al. (2020) The COVID-19 vaccine development landscape. Nat. Rev. Drug Discov. 19, 305–306
- Pregelj, L. et al. (2018) Precision medicines have faster approvals based on fewer and smaller trials than other medicines. *Health Aff. (Millwood*) 37, 724–731
- Darrow, J.J. et al. (2020) FDA approval and regulation of pharmaceuticals, 1983-2018. JAMA 323, 164–176

- Akanmori, B. *et al.* (2015) The African vaccine regulatory forum (AVAREF): a platform for collaboration in a public health emergency. *WHO Drug Info.* 29, 127–132
- Henao-Restrepo, A.M. et al. (2016) On a path to accelerate access to Ebola vaccines: the WHO's research and development efforts during the 2014–2016 Ebola epidemic in West Africa. Curr. Opin. Virol. 17, 138–144
- Vannice, K.S. et al. (2019) Demonstrating vaccine effectiveness during a waning epidemic: a WHO/NIH meeting report on approaches to development and licensure of Zika vaccine candidates. Vaccine 37, 863–868
- Hodgson, J. (2020) The pandemic pipeline. Nat. Biotechnol. 38, 523–532
- Davis, J.S.Ferreira, D.Denholm, J.T.Tong, S.Y.C. (2020) Clinical trials for the prevention and treatment of coronavirus disease 2019 (COVID-19): The current state of play. Med. J. Aust. Published online: 27 April 2020. https://www.mja.com.au/journal/2020/clinical-trialsprevention-and-treatment-coronavirus-disease-2019covid-19-current

### Forum

## Decellularized Scaffolds for Intervertebral Disc Regeneration

Morena Fiordalisi (D, 1,2,3,4 Ana João Silva (D, 1,2) Mário Barbosa (D, 1,2,3,4) Raquel Gonçalves (D, 1,2,3,4) Joana Caldeira (D 1,2,4,\*)

In the last decade, intervertebral disc (IVD) decellularization has gained significant attention for tissue regenerative purposes as a successful therapeutic alternative for low back pain (LBP). We discuss the recent advances in IVD decellularization, repopulation, and sterilization procedures, highlighting the major challenges that need to be addressed for clinical translation.

# Recapitulating a Healthy Disc Environment to Treat LBP

IVD degeneration is often the cause of LBP, which affects over 70% of the world's population and is the most

## **CIIPTESS** REVIEWS