



ORIGINAL ARTICLE

Prevalence and determinants of depression up to 5 years after colorectal cancer surgery: results from the ColoRECTal Wellbeing (CREW) study

Lynn Calman¹ | Joshua Turner¹ | Deborah Fenlon² | Natalia V. Permyakova³ | Sally Wheelwright¹ | Mubarak Patel⁴ | Amy Din¹ | Jane Winter^{1,5} | Alison Richardson^{1,5} | Peter W. F. Smith⁶ | Claire Foster¹ | members of the CREW Study Advisory Committee†

¹School of Health Sciences, University of Southampton, Southampton, UK

²College of Human and Health Sciences, Swansea University, Swansea, UK

³Faculty of Medicine, University of Southampton, Southampton, UK

⁴Division of Health Sciences, University of Warwick, Warwick, UK

⁵University Hospitals Southampton NHS Trust, Southampton, UK

⁶Social Statistics and Demography, Social Sciences, University of Southampton, Southampton, UK

Correspondence

Claire Foster, Macmillan Survivorship Research Group (MSRG), School of Health Sciences, University of Southampton, Building 67, Southampton SO17 1BJ, UK
Email: c.l.foster@soton.ac.uk

Funding information

The work was supported by Macmillan Cancer Support as part of the Macmillan Survivorship Research Group programme (grant no. 3546834).

Abstract

Aim: Depression experienced by people with colorectal cancer (CRC) is an important clinical problem affecting quality of life. Recognition of depression at key points in the pathway enables timely referral to support. This study aimed to examine depression before and 5 years after surgery to examine its prevalence and identify determinants.

Method: The ColoRECTal Wellbeing (CREW) study is a prospective UK cohort study involving 872 adults with nonmetastatic CRC recruited before surgery with curative intent. Questionnaires completed before surgery and 3, 9, 15, 24, 36, 48 and 60 months after surgery captured socio-demographics and assessed depression (Centre for Epidemiologic Studies Depression Scale, CES-D) and other psychosocial factors. Clinical details were also gathered. We present the prevalence of clinically significant depression (CES-D \geq 20) over time and its predictors assessed before and 2 years after surgery.

Results: Before surgery, 21.0% of the cohort reported CES-D \geq 20 reducing to 14.7% 5 years after surgery. Presurgery risk factors predicting subsequent depression were clinically significant depression and anxiety, previous mental health service use, low self-efficacy, poor health, having neoadjuvant treatment and low social support. Postsurgery risk factors at 2 years predicting subsequent depression were clinically significant depression, negative affect, cognitive dysfunction, accommodation type and poor health.

Conclusion: Depression is highly pervasive in people with CRC, exceeding prevalence in the general population across follow-up. Our findings emphasize the need to screen and treat depression across the pathway. Our novel data highlight key risk factors of later depression at important and opportune time points: before surgery and at the end of routine surveillance. Early recognition and timely referral to appropriate support is vital to improve long-term psychological outcomes.

†Members are listed in the Acknowledgements.

Prior presentations: presented as an oral presentation at the International Psycho-Oncology Society (IPOS) Annual Conference, Hong Kong, 31 October to 2 November 2018 and the British Psycho-Oncology Society (BPOS) Annual Conference, Chester, UK, 28 February to 1 March 2019. Presented as a poster presentation at the National Cancer Research Institute (NCRI) Cancer Conference, Glasgow, UK, 4–6 November 2018.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Colorectal Disease published by John Wiley & Sons Ltd on behalf of Association of Coloproctology of Great Britain and Ireland.

**KEYWORDS**

colorectal cancer, determinants, depression, quality of life, risk factors

INTRODUCTION

Depressive disorders are one of the highest contributors to the global burden of disease, affecting 4% of people [1]. The prevalence of depression is greater in people living with cancer (rates range between 8% and 27% [2]) and thus it is a clinical and policy priority [3]. In people living with colorectal cancer (CRC), the prevalence of depression is also greater when compared with the general population, with rates reaching 37% [4–7]. Depression in people living with cancer is associated with poor quality of life (QoL) [8], reduced adherence to treatment [9] and reduced survival [10] and is associated with an increased risk of suicide [11]. More specifically, people with CRC and depression have poorer QoL, health status and wellbeing after diagnosis and surgery [5,7].

Despite this, depression is often not identified in people with cancer, and few are treated [6], possibly due to a range of factors associated with underreporting of mental health issues (e.g. stigma) [12]. Timely recognition and referral to support and intervention is recommended in international clinical guidelines for depression [13], with UK guidelines recommending targeting screening in people at most risk [14]. However, determinants of depression in people living with cancer are poorly characterized [15] with calls for more evidence in people with CRC [16]. Identification of people most at risk also informs the development of interventions, reduces disease burden and improves planning of psychosocial care resources [17].

Several determinants of depression in people with CRC have been described, including female gender [4,18,19], low socioeconomic status [20], higher stage disease [21], receipt of neoadjuvant and adjuvant treatments [22] and low social support [23,24]. Findings related to age are inconsistent [19,21,25] but may be due to sample characteristics (e.g. recruitment of an older sample [25]). Depressive symptoms in people with CRC are also reported to reduce over time [4,5] but much research is cross-sectional [16]. We conducted a scoping review and found that no published longitudinal studies in people with nonmetastatic CRC have examined presurgery risk factors associated with reporting depression up to 5 years after treatment (Appendix 1).

UK clinical guidance recommends before and after treatment as key time points to appraise psychological outcomes in people living with cancer [26]. Assessment close to diagnosis allows for a better understanding of the impact of cancer and its treatment on depression outcomes [15,26]. Early screening is also encouraged due to its benefits: improving treatment adherence, reducing the burden on health services and patients, enhancing communication between clinical teams and patients and more timely referral to support [17,27]. Addressing psychological outcomes after treatment gives patients the opportunity to reflect on the impact and psychosocial concerns following the intense scheduling of cancer treatment [15,26]. Posttreatment CRC surveillance is valuable to

What does this paper add to the literature?

Depression in people living with colorectal cancer is an important clinical problem. Our study shows that the prevalence of depression exceeds prevalence in the general population over time. It also highlights risk factors for later depression at important time points (before surgery and at the end of routine follow-up) which informs strategies for recognition and referral to appropriate psychosocial support.

provide reassurance, as patients feel greater concern when its frequency decreases [28], possibly due to reduced contact with clinical teams [29], so we highlight this time point as being important. In the UK it is recommended that routine surveillance appointments cease after 2 years [30] with some variation in international guidelines [31].

This paper presents analysis from the ColoRECTal Wellbeing (CREW) study [32], a UK prospective cohort study investigating factors associated with the recovery of health and wellbeing following CRC. Data were collected before and at regular intervals up to 5 years after surgery. Data comprised socio-demographic, clinical information and patient-reported outcomes examining a selection of psychosocial variables (including depression) informed by a conceptual framework of recovery following cancer diagnosis and treatment [33]. The analysis assesses 'clinically significant levels' of depression via self-reporting, and whilst this is not a 'clinical diagnosis' of depression, which requires a comprehensive assessment accounting for contextual factors [14], the cut-off used has high concordance with psychiatric interviews [34] suggesting the experience of high levels of depressive symptomology [35]. In this paper we describe the prevalence of clinically significant levels of depression from before surgery up to 5 years after surgery and, given the levels of depression before and at 2 years after surgery, identify which characteristics are associated with subsequent clinically significant levels of depression up to 5 years after surgery.

METHOD**Study sample**

The CREW study is a prospective cohort study of adults (≥ 18 years) with nonmetastatic CRC (Dukes A–C) treated by surgery with curative intent. Inclusion and exclusion criteria are published elsewhere [32].

Data collection

Details of study procedures have been previously reported [32]. Eligible participants were recruited from 29 UK National Health Service (NHS) centres between November 2010 and March 2012. Participants consented and completed questionnaires before surgery (baseline). Follow-up questionnaires were posted at regular intervals (3, 9, 15, 24 months and then annually up to 5 years after surgery). Clinical and treatment information was gathered from NHS medical databases at participating centres. Ethical approval was granted by the UK NHS NRES Committee South Central – Oxford B (REC ref. 10/H0605/31). Information collected in the study did not inform the care of the participants involved due to the study design and anonymization of the data.

Measures

Patient-reported depression was captured using the 20-item Centre for Epidemiologic Studies Depression Scale (CES-D) [36]. Higher scores indicate greater levels of depression (range 0–60). A recent meta-review demonstrated that CES-D is responsive to change and suitable for screening for depression in people with cancer [37].

A score of ≥ 20 has previously been used in studies involving people with cancer to define a 'clinically significant level' of depression [38] and has been shown to be highly concordant with psychiatric interviews [34]. A recent meta-analysis examining the screening accuracy of CES-D noted the ≥ 20 cut-off to be more appropriate than the standard ≥ 16 cut-off [35]. Thus the ≥ 20 cut-off was selected as an indicator of a clinically significant levels of depression for this study, but this does not constitute a formal diagnosis of clinical depression.

Determinants/covariates

Table 1 lists the validated patient-reported outcome measures, socio-demographic questions and clinical information captured before surgery and 2 years after surgery which were used as covariates in the analyses. Covariates are presented according to the conceptual framework domains [33] and the rationale for each measure is provided elsewhere [32]. Validated measures were repeated at every time point unless otherwise indicated. Selection of covariates, including European Organisation for Research and Treatment of Cancer (EORTC) subscales, was informed by our scoping review (Appendix 1). Alongside depression, accommodation type, health status (EQ-5D), age and ethnicity were found to be significantly associated with participant attrition in the CREW study [39] and were included in the model to account for this.

Statistical analysis

The total CES-D score was summarized at each time point using its median and interquartile range (IQR) to examine changes over time.

The number and proportion of participants reporting clinically significant levels of depression ($\text{CES-D} \geq 20$) were also assessed over time.

Two multivariable logistic regression models were fitted to predict clinically significant levels of depression up to 5 years after surgery (Appendix 2): model 1 included depression together with other covariates collected before surgery (baseline); model 2 included depression together with other covariates collected at 2 years after surgery. Multicollinearity was assessed in each model using the variance inflation factor (VIF). The VIF ranged from 1.05 to 2.18 for model 1 and from 1.08 to 2.51 for model 2. A VIF below 10 indicates that there was no evidence of multicollinearity in our models.

Missing data were imputed according to published guidelines for the measures selected. If unavailable, these were omitted from the final model. The number of comorbidities was first assessed at 3 months but was included in model 1 due to its stability over time [40].

A population-average approach was applied to account for the time-varying nature of the binary outcome, where each model was adjusted for the clustering of observations within the participants [41]. Regression analyses were based on a backwards elimination of statistically nonsignificant predictors. The significance level was fixed at 5% and all analyses were completed in Stata 14.

RESULTS

Sample characteristics

One thousand and eighteen participants were recruited into CREW and 872 consented to questionnaire follow-up. Figure 1 presents the participant flow over follow-up; full details of study recruitment and descriptive statistics are published elsewhere [39,42]. The sample was representative of the eligible patients treated during the recruitment period [39,42]. Table 2 shows the demographic and clinical characteristics of the 741 participants who returned a baseline questionnaire and had completed the CES-D; these participants had a mean age of 67.54 years ($\text{SD} = 10.26$ years). Over 54% of the sample underwent laparoscopic surgery and 40% underwent open surgery for CRC.

Depression over time

At baseline (before surgery), people who were women, single, living in rented accommodation and had previously used mental health services were more likely to report clinically significant CES-D scores (Table 2).

Median scores peaked before surgery at 12.0 (IQR = 11.7) and decreased to 9.5 (IQR = 12.0) at 5 years (Table 3). Similarly, the proportion of participants reporting clinically significant levels of depression also peaked before surgery at 21.0% and reduced to 14.7% at 5 years (Table 3). Overall, 303 participants (34.8%) reported

TABLE 1 CREW study measures presented by conceptual framework domains [33] for regression analysis

Domain	Characteristic of interest	Measure
Preexisting factors (socio-demographics)	Age	
	Gender	
	Ethnicity	
	Employment status ^a	
	Accommodation type ^a	
Clinical factors	Deprivation index	Index of Multiple Deprivation (IMD) [61]
	Tumour site	
	Dukes stage ^a	
	Neoadjuvant treatment ^a	
	Surgery type	
	Adjuvant treatment ^a	
	Stoma status	
	Number of comorbidities ^b	Self-reported measure [40]
	Previous use of mental health services ^c	
	Environmental factors	Domestic status
Life events		List of Threatening Experience Questionnaire (LTE-Q) [62]
Social support		Medical Outcomes Study - Social Support Survey (MOS-SSS) [63]
Personal factors	Self-efficacy	Self-efficacy for Managing Chronic Disease (SEMCD) scale [64] Cancer Survivors' Self-Efficacy Scale (CS-SES) [65]
	Affect	Positive and Negative Affect Schedule Short Form (PANAS-SF) [66]
Psychosocial outcomes	State anxiety ^d	State-Trait Anxiety Inventory, State scale (STAI-S) [67]
	Wellbeing	Personal Wellbeing Index-Adult (PWI-A) [68]
	Health status	EuroQoL, five dimensions three levels (EQ-5D-3L) [69]
	Quality of life (QoL) ^e	Quality of Life in Adult Cancer Survivors (QLACS) scale [70]: Cancer-Specific Summary Score (QLACS-CSS), Benefit of Cancer (QLACS-BC)
	Symptoms and functioning ^b	European Organization for Research and Treatment of Cancer quality of life measure (EORTC QLQ-C30) [71]: function scales: physical, emotional, cognitive, social; symptom scales: fatigue, pain, insomnia, financial worry

^aTo avoid imprecise estimates from the low counts in the regression analyses two or more groups were merged together: unemployed and retired (employment status); renting and other (accommodation type); Stages C1 and C2 (Dukes stage); radiotherapy, chemotherapy and both (neoadjuvant treatment; adjuvant treatment).

^bCollected from 3 months onwards. Selection of EORTC subscales was informed by previous work involving people with CRC [7,22,24,25].

^cSelf-reported at baseline only.

^dWe used a cut-off of ≥ 40 to indicate a clinically significant level of anxiety [72].

^eItems comprising the QLACS-CSS and QLACS-BC were collected from 9 months onwards.

TABLE 2 Sociodemographic and clinical information comparisons of CES-D < 20 and ≥20 reported at baseline (N = 741)

Covariates reported at baseline	n (%)	CES-D < 20, n (%)	CES-D ≥ 20, n (%)	p-value (chi-square test)
Age groups (years)				
50 or younger	47 (6.4%)	32 (68.1%)	15 (31.9%)	0.073
51–60	113 (15.3%)	83 (73.5%)	30 (26.5%)	
61–70	285 (38.6%)	236 (82.8%)	49 (17.2%)	
71–80	217 (29.4%)	173 (79.7%)	44 (20.3%)	
81 or older	77 (10.4%)	58 (75.3%)	19 (24.7%)	
Gender				
Male	440 (59.4%)	373 (84.8%)	67 (15.2%)	<0.001
Female	301 (40.6%)	210 (69.8%)	91 (30.2%)	
Ethnicity				
White British	623 (92.7%)	491 (78.8%)	132 (21.2%)	0.898
Other ethnic group	49 (7.3%)	39 (79.6%)	10 (20.4%)	
Deprivation (IMD) quintile				
1st quintile (least deprived)	146 (20.1%)	123 (84.2%)	23 (15.8%)	0.086
2nd quintile	150 (20.6%)	123 (82%)	27 (18%)	
3rd quintile	142 (19.5%)	113 (79.6%)	29 (20.4%)	
4th quintile	136 (18.7%)	99 (72.8%)	37 (27.2%)	
5th quintile (most deprived)	153 (21%)	114 (74.5%)	39 (25.5%)	
Domestic status				
Married/living with partner	524 (71.1%)	430 (82.1%)	94 (17.9%)	<0.001
Single/widowed/divorced/separated	213 (28.9%)	150 (70.4%)	63 (29.6%)	
Employment status				
Employed	201 (27.3%)	158 (78.6%)	43 (21.4%)	0.980
Unemployed/retired	535 (72.7%)	421 (78.7%)	114 (21.3%)	
Accommodation type				
Owner occupied	589 (79.9%)	473 (80.3%)	116 (19.7%)	0.021
Renting/other ^a	148 (20.1%)	106 (71.6%)	42 (28.4%)	
Previous use of mental health services				
No	670 (94.5%)	536 (80%)	134 (20%)	<0.001
Yes	39 (5.5%)	22 (56.4%)	17 (43.6%)	
Tumour site				
Colon	475 (64.4%)	374 (78.7%)	101 (21.3%)	0.911
Rectum	263 (35.6%)	208 (79.1%)	55 (20.9%)	
Dukes stage				
A	109 (14.7%)	93 (85.3%)	16 (14.7%)	0.335
B	391 (52.8%)	303 (77.5%)	88 (22.5%)	
C (C1 and C2)	229 (30.9%)	178 (77.7%)	51 (22.3%)	
Could not be determined ^b	11 (1.5%)	9 (81.8%)	2 (18.2%)	
Neoadjuvant treatment (any type)				
No	592 (80.7%)	465 (78.5%)	127 (21.5%)	0.649
Yes	142 (19.3%)	114 (80.3%)	28 (19.7%)	
Surgery type ^c				
Laparoscopic	401 (54.3%)	–	–	–
Open	299 (40.5%)	–	–	
Not available	38 (5.2%)	–	–	

(Continues)

TABLE 2 (Continued)

Covariates reported at baseline	n (%)	CES-D < 20, n (%)	CES-D ≥ 20, n (%)	p-value (chi-square test)
Adjuvant treatment (any type) ^c				
No	477 (64.6%)	-	-	-
Yes	261 (35.4%)	-	-	-
Stoma ^c				
No	262 (35.9%)	-	-	-
Yes	468 (64.1%)	-	-	-
Number of comorbidities ^d				
0	168 (27.6%)	143 (85.1%)	25 (14.9%)	.055
1	194 (31.9%)	160 (82.5%)	34 (17.5%)	
2	144 (23.6%)	107 (74.3%)	37 (25.7%)	
3+	103 (16.9%)	78 (75.7%)	25 (24.3%)	

Note: p-values in bold indicate a statistically significant difference at the 5% level.

Abbreviation: IMD, Index of Multiple Deprivation.

^aOther accommodation includes: temporary accommodation, living in residential or nursing home, living with others (e.g. friends or family).

^bDukes stage could not be determined for 11 full consent patients with small tumours following neoadjuvant therapy.

^cCaptured from the medical records after baseline.

^dSelf-reported at 3 months.

TABLE 3 Descriptive statistics for the CES-D score and clinically significant level of depression (CES-D ≥ 20) at each time point from before surgery to 5 years after surgery

	Presurgery	Postsurgery						
	Baseline	3 months	9 months	15 months	24 months	36 months	48 months	60 months
N	741	642	605	534	483	382	369	319
CES-D total, median (IQR)	12.0 (11.7)	11.1 (12.0)	10.0 (13.0)	9.0 (12.0)	9.0 (10.9)	8.0 (11.7)	9.0 (11.0)	9.5 (12.0)
CES-D ≥ 20, n (%)	158 (21.3)	124 (19.3)	106 (17.5)	70 (13.1)	73 (15.1)	49 (12.8)	48 (13.0)	47 (14.7)

Abbreviations: CES-D, Centre for Epidemiologic Studies Depression Scale; IQR, interquartile range

clinically significant levels of depression at 2 years were at greater risk of subsequent depression up to 5 years (OR = 3.14, 95% CI = 1.41–7.04). Those who had higher scores for negative affect were also at greater risk (OR = 1.21, 95% CI = 1.08–1.36).

People reporting problems with cognitive function (OR = 2.21, 95% CI = 1.03–4.77) and poorer wellbeing (OR = 2.40, 95% CI = 1.25–4.61) at 2 years also had higher odds of experiencing clinically significant depression later. Participants who did not own their accommodation were also at greater risk of reporting clinically significant depression (OR = 2.38, 95% CI = 1.23–4.62).

In contrast, the risk of reporting clinically significant levels of depression was lower amongst those who had 'perfect' health status at 2 years (OR = 0.28, 95% CI = 0.12–0.68).

DISCUSSION

This is the first prospective cohort study to examine the prevalence and risk factors associated with clinically significant levels of

depression in people with nonmetastatic CRC assessed before and up to 5 years after surgery. Our results reveal that clinically significant levels of depression remain a long-term problem for a considerable proportion of people, despite median CES-D scores reducing over time from initial diagnosis. These results are consistent with previous findings [4,5]. For example, our prevalence rates across each time point occur within the range observed by cross-sectional studies of people living with CRC (7%–37% [4–7]) and are considerably higher than the median prevalence found in the general population (CES-D ≥ 20; 11.8%) [35].

The novelty of this study is the investigation of risk factors of clinically significant levels of depression at two key time points in the cancer care pathway as recommended by UK clinical guidance [26]: close to diagnosis (before surgery) and when posttreatment routine surveillance ends (2 years after surgery). Identifying risk factors improves planning of psychosocial care and informs the development of interventions [17]. We identified several pre- and postsurgery risk factors of depression consistent with previous work [4,5,7,19,20,22–25].



TABLE 4 Multivariable logistic regression model of clinically significant depression (CES-D \geq 20) up to 5 years after surgery, significant covariates collected before surgery (baseline)

Theme block	Covariates	OR	95% CI	p-value
Socio-demographic factors	Age groups (years)			
	50 or younger	Ref.	-	-
	51-60	0.50	0.22-1.10	0.086
	61-70	0.50	0.26-0.97	0.040
	71-80	0.55	0.27-1.13	0.103
Clinical and treatment factors	81 or older	0.77	0.33-1.80	0.550
	Tumour site			
	Colon	Ref.	-	-
	Rectum	0.55	0.35-0.87	0.011
	Neoadjuvant treatment			
	None	Ref.	-	-
	Yes, any therapy	2.99	1.75-5.09	<0.001
Previous use of mental health services	No	Ref.	-	-
	Yes	3.33	1.81-6.12	<0.001
	Unknown	0.65	0.23-1.81	0.411
Environmental factors	Domestic status			
	Married/living with a partner	Ref.	-	-
	Single/widowed/divorced/separated	2.02	1.32-3.09	0.001
Personal factors	Self-efficacy (SEMCD)			
	Low confidence	Ref.	-	-
	Moderate confidence	0.42	0.24-0.73	0.002
	Confident	0.35	0.20-0.61	<0.001
Psychosocial factors	Very confident	0.18	0.08-0.37	<0.001
	Depression (CES-D)			
	<20	Ref.	-	-
	\geq 20 (clinical level)	3.44	2.18-5.45	<0.000
	Anxiety (STAI-S)			
	<40	Ref.	-	-
	\geq 40 (high level)	1.82	1.15-2.87	0.010
	Social support (MOS-SSS)			
	<100 (not full)	Ref.	-	-
	\geq 100 (full)	0.41	0.23-0.74	0.003
Health status (EQ-5D-3L)	Not perfect health	Ref.	-	-
	Perfect health	0.42	0.24-0.75	0.003

Note: p-values in bold indicate a statistically significant difference at the 5% level. The model controls for the time point of the outcome report (postsurgery 3-60 m), which was statistically significant.

Abbreviations: CES-D, Centre for Epidemiologic Studies Depression Scale; EQ-5D-3L, EuroQoL five dimensions, three levels; MOS-SSS, Medical Outcome Study Social Support Scale; SEMCD, Self-Efficacy for Managing Chronic Disorders Scale; STAI-S, State-Trait Anxiety Inventory, State scale.

Importantly, our findings underscore the need for depression screening close to diagnosis, with clinically significant levels before surgery being identified as a risk for later depression. Early screening

has been shown to have a positive impact on care by leading to more timely referrals for psychological intervention [17,27]. Our analysis at 2 years after surgery also suggests the need for assessment of

Theme block	Covariates	OR	95% CI	p-value
Socio-demographic factors	Accommodation type			
	Owner occupied	Ref.	-	-
	Rented/other	2.38	1.23-4.62	0.010
Personal factors	Affect (PANAS-SF)			
	Negative affect	1.21	1.08-1.36	0.001
Psychosocial factors	Depression (CES-D)			
	<20	Ref.	-	-
	≥20 (clinical level)	3.14	1.41-7.04	0.005
	Health status (EQ-5D-3L)			
	Not perfect health	Ref.	-	-
	Perfect health	0.28	0.12-0.68	0.005
	Wellbeing (PWI-A)			
	≥70 (good)	Ref.	-	-
	<70 (poorer)	2.40	1.25-4.61	0.008
	Cognitive functioning (EORTC QLQ-C30)			
No problem	Ref.	-	-	
Some problem	2.21	1.03-4.77	0.043	

Note: p-values in bold indicate a statistically significant difference at the 5% level.

Abbreviations: CES-D, Centre for Epidemiologic Studies Depression Scale; EORTC-QLQ-C30, European Organisation for Research and Treatment of Cancer Quality of Life Core-30 Questionnaire; EQ-5D-3L, EuroQoL five dimensions, three levels; PANAS-SF, Positive and Negative Effect Schedule Short Form; PWI-A, Personal Wellbeing Index – Adult.

depression and depressive symptomology (negative affect) when posttreatment surveillance ends. Regular appraisal of psychological needs throughout the pathway aligns with recent emphasis on risk stratification in the UK NHS Long Term Plan to inform personalized care for people with cancer and facilitate referral to appropriate levels of care [43]. Psychosocial interventions for people with CRC have been reported to be beneficial in improving symptomology of depression and anxiety, as well as QoL [44,45]. Novel strategies for follow-up have been tested in Australia [46] and Canada [47] and are being considered in the USA [48]. Such strategies can help target specialist resources as these become increasingly scarce [48]. Innovative models of psychological screening and care (e.g. stepped care and nurse-led collaborative interventions) for people with cancer are effective in reducing psychological symptoms, improve QoL for people with a depressive or anxiety disorder and are cost-effective [49,50]. Internationally, variability in models and approaches to survivorship care and complexity in reimbursement for psychosocial and integrated care make implementation a challenge [51].

Our presurgery analysis also highlights at-risk groups to whom we should direct depression screening. People undergoing neoadjuvant treatment commonly face more complex surgery, stoma formation, additional side effects and increased treatment time [22], which can explain our findings and so attention should focus on this group. People with rectal cancer had a lower risk of clinically significant depression over time but no previous CRC studies have reported tumour site as a significant predictor of depression [16]. This

TABLE 5 Multivariable logistic regression model of clinically significant depression (CES-D ≥ 20) up to 5 years after surgery, significant covariates collected at 2 years

relationship was also unexpected, as people with rectal cancer often have complex treatment regimens (including neoadjuvant treatment) [52] which may impact psychological outcomes, particularly those who later have a permanent stoma [53]. One possible explanation could be that a more complex treatment pathway may result in greater contact with clinical teams and this may improve perceptions of support [29,54] that could help to reduce depression symptomology. Nevertheless, this finding requires further investigation.

Our analysis further recommends that depression screening should target people with a history of mental health problems or with psychological comorbidities (e.g. anxiety). This is unsurprising, as the level of anxiety tends to peak close to diagnosis [55] and it commonly co-occurs with depression [8].

The value of assessing self-efficacy and social support early in the pathway was highlighted by our presurgery analysis. This is important given the increasing role of self-management for people with cancer [56]; thus confidence to manage consequences of cancer and its treatment need to be assessed early on. Assessing the level of social support at the point of diagnosis is imperative given its importance for depression outcomes and later QoL [23].

Our analysis at 2 years after surgery highlighted other at-risk groups in whom assessment and support for depression may be helpful. People with cognitive difficulties postsurgery were at greater risk of later depression; this is important as cognitive dysfunction is a commonly reported consequence of CRC treatment [57]. However, caution should be applied as it can be difficult to delineate cognitive dysfunction as a result of cancer treatment or as a symptom



of depression and/or anxiety [58]. Type of accommodation (rented or other) was also highlighted as a risk factor, but this specifically has not been reported previously. It could be used as a descriptor of socio-economic status which has been noted to be a risk factor for anxiety, depression and distress in people with cancer [20,59]. This highlights the need for additional support for this group as low socio-economic status may indicate a low availability of resources important for coping, which may result in poorer psychological outcomes [59].

Study strengths include the scale and representative nature of the CREW sample with over 91% of all eligible patients approached to participate [39,42]. Loss to follow-up is expected in cohort studies but our response rates remained high up to 5 years (Figure 1; 71%). Participants who withdrew by 5 years were more likely to report clinically significant depression, were ≥ 80 years of age, did not own accommodation (renting or other) and were of non-White ethnicity at baseline [39]. Therefore, our findings may underestimate the true prevalence of depression among CRC cancer survivors in the UK. Additionally, our sample represents patients from one type of healthcare system (the UK NHS) whereby access to and provision of specialist services is universal and free at the point of delivery.

Patient-reported depression may not account for contextual factors considered in diagnostic interviews [14]. Nevertheless, the cut-off used suggested clinically significant levels of depressive symptomology [35] and is highly concordant with psychiatric interviews [34]. A high prevalence of depression over time may be attributed to its undertreatment [6]. However, due to a high level of missing data, as a result of poor self-reporting of health service use in CREW, we were unable to examine use of psychological treatment, which may explain our findings. We examined our mental health service use data in a bivariate analysis with CES-D scores for interest (Appendix 3).

Scoping of the literature (Appendix 2) identifies this as one of the first studies to include a presurgery assessment on a range of socio-demographic, psychosocial and clinical factors and the only one to collect data for up to 5 years afterwards to examine risk factors of clinically significant levels of depression. The importance of this work is highlighted by the dearth of evidence examining the long-term psychological impacts in people living with and beyond cancer [3], including people with CRC [16]. The need for research into the short- and long-term psychological impacts of cancer and its treatment has been identified as a Top 10 research priority in the UK [60] and our analysis contributes knowledge to this for two crucial time points in the CRC care pathway.

In summary, our results indicate that depression is an enduring problem in people with nonmetastatic CRC even at 5 years after surgery. Before surgery it affects one in five people and one in seven people at 5 years after surgery, both of which are higher than reported in the general population. Our findings clearly highlight the need for screening for depression across the pathway to improve depression outcomes in the long term. Early screening

should be focused on those with a history of mental health issues, high levels of anxiety, low self-efficacy, poor health status and low levels of support, whilst clinicians should also monitor people who undergo neoadjuvant treatment. The end of routine oncology surveillance is also an opportune time to assess symptoms of depression, especially as the frequency of contact with clinical teams decreases. At this time point, assessment should focus on people with poor health, a lower socio-economic status and problems from treatment (e.g. cognitive dysfunction). Depression in people living with cancer is associated with poor health and wellbeing and has an impact on survival and adherence to treatment, so early recognition and treatment may lead to overall improved outcomes for patients.

ACKNOWLEDGEMENTS

We thank all CREW study participants and recruiting NHS Trusts; Carol Hill, Kerry Coleman, Bjoern Schukowsky, Christine May (study support); Matthew Breckons, Cassandra Powers, Alex Recio-Saucedo, Bina Nausheen, Ikumi Okamoto, Kim-Chivers Seymour, Joanne Haviland (researchers); Jo Clough, Alison Farmer (research partners).

Members of the Study Advisory Committee: Jo Armes, Janis Baird, Andrew Bateman, Nick Beck, Graham Moon, Claire Hulme, Peter Hall, Karen Poole, Susan Restorick-Banks, Paul Roderick, Claire Taylor, Jocelyn Walters, Fran Williams, Lynn Batehup, Jessica Corner and Deborah Fenlon. We would also like to thank Michael Sharpe for his valuable feedback on our manuscript and Angus McNair for his valuable advice.

Professor Alison Richardson is a National Institute for Health Research (NIHR) Senior Investigator. The views expressed in this article are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

CONFLICT OF INTERESTS

LC has received an honorarium for teaching from Boehringer Ingelheim. DF has received an honorarium for teaching from Roche.

AUTHOR CONTRIBUTIONS

CF, DF, PWS conceived and obtained funding for the study. LC, JW, AR and the Study Advisory Committee made substantial contributions to the development and design of this work. NP and MP analysed the data with the support of PWS. JT, LC, DF, NP, PWS, SW, JW and CF contributed to interpretation of data. JT drafted the manuscript with substantial contributions from LC, CF and SW. All authors provided critical comments on drafts of the manuscript and approved the final manuscript.

ETHICAL APPROVAL

The study was approved by the UK National Health Service National Research Ethics Service (REC reference number: 10/H0605/31). Informed consent was obtained from all individual participants involved in the study.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author. Information regarding data access is available at http://horizons-hub.org.uk/access_data.html

REFERENCES

- World Health Organization. Depression and other common mental disorders: global health estimates, 2017.
- Krebber AMH, Buffart LM, Kleijn G, Riepma IC, de Bree R, Leemans CR, et al. Prevalence of depression in cancer patients: a meta-analysis of diagnostic interviews and self-report instruments. *Psycho-Oncology*. 2014;23(2):121–30.
- Niedzwiedz CL, Knifton L, Robb KA, Katikireddi SV, Smith DJ. Depression and anxiety among people living with and beyond cancer: a growing clinical and research priority. *BMC Cancer*. 2019;19(1):1–8.
- Clark C, Fino N, Liang J, Hiller D, Bohl J, Clark CJ, et al. Depressive symptoms in older long-term colorectal cancer survivors: a population-based analysis using the SEER-Medicare healthcare outcomes survey. *Support Care Cancer*. 2016;24(9):3907–14.
- Mols F, Schoormans D, de Hingh I, Oerlemans S, Husson O. Symptoms of anxiety and depression among colorectal cancer survivors from the population-based, longitudinal PROFILES Registry: prevalence, predictors, and impact on quality of life. *Cancer*. 2018;124(12):2621–8.
- Walker J, Hansen CH, Martin P, Symeonides S, Ramessur R, Murray G, et al. Prevalence, associations, and adequacy of treatment of major depression in patients with cancer: a cross-sectional analysis of routinely collected clinical data. *Lancet Psychiat*. 2014;1(5):343–50.
- Tsunoda A, Nakao K, Hiratsuka K, Yasuda N, Shibusawa M, Kusano M. Anxiety, depression and quality of life in colorectal cancer patients. *Int J Clin Oncol*. 2005;10(6):411–7.
- Spiegel D, Giese-Davis J. Depression and cancer: mechanisms and disease progression. *Biol Psychiat*. 2003;54(3):269–82.
- DiMatteo M, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med*. 2000;160(14):2101–7.
- Pinquart M, Duberstein PR. Depression and cancer mortality: a meta-analysis. *Psychol Med*. 2010;40(11):1797–810.
- Henson KE, Brock R, Charnock J, Wickramasinghe B, Will O, Pitman A. Risk of suicide after cancer diagnosis in England. *JAMA Psychiat*. 2019;76(1):51–60.
- Kissane DW. Unrecognised and untreated depression in cancer care. *Lancet Psychiat*. 2014;1(5):320–1.
- Ferenchick EK, Ramanuj P, Pincus HA. Depression in primary care: part 1—screening and diagnosis. *BMJ*. 2019;365:l794.
- National Institute for Health and Clinical Excellence (NICE). Depression in adults: recognition and management (CG90). In: National Institute for Health and Clinical Excellence (NICE), editor. London: National Institute for Health and Clinical Excellence (NICE); 2009.
- Pitman A, Suleman S, Hyde N, Hodgkiss A. Depression and anxiety in patients with cancer. *BMJ*. 2018;361:k1415.
- Mosher CE, Winger JG, Given BA, Helft PR, O'Neil BH. Mental health outcomes during colorectal cancer survivorship: a review of the literature. *Psycho-Oncology*. 2016;25(11):1261–70.
- Mitchell AJ. Screening for cancer-related distress: when is implementation successful and when is it unsuccessful? *Acta Oncol*. 2013;52(2):216–24.
- Braamse AMJ, van Turenhout ST, Terhaar Sive Droste JS, de Groot GH, van der Hulst RWM, Klemm-Kropp M, et al. Factors associated with anxiety and depressive symptoms in colorectal cancer survivors. *Eur J Gastroenterol Hepatol*. 2016;28(7):831–5.
- Powell-Chandler A, Boyce K, James O, Scourfield L, Torkington J, Bisson J, et al. Psychological sequelae of colonic resections. *Colorectal Dis*. 2020;22(8):945–51.
- Dunn J, Ng SK, Holland J, Aitken J, Youl P, Baade PD, et al. Trajectories of psychological distress after colorectal cancer. *Psycho-Oncology*. 2013;22(8):1759–65.
- Xia S, Sun M, Liu X. Major depression but not minor to intermediate depression correlates with unfavorable prognosis in surgical colorectal cancer patients undergoing adjuvant chemotherapy. *Psychol Health Med*. 2020;25(3):309–18.
- Gray NM, Hall SJ, Browne S, Johnston M, Lee AJ, Macleod U, et al. Predictors of anxiety and depression in people with colorectal cancer. *Support Care Cancer*. 2014;22(2):307–14.
- Haviland J, Sodergren S, Calman L, Corner J, Din A, Fenlon D, et al. Social support following diagnosis and treatment for colorectal cancer and associations with health-related quality of life: results from the UK ColoREctal Wellbeing (CREW) cohort study. *Psycho-Oncology*. 2017;26(12):2276–84.
- Gonzalez-Saenz de Tejada M, Bilbao A, Baré M, Briones E, Sarasqueta C, Quintana JM, et al. Association between social support, functional status, and change in health-related quality of life and changes in anxiety and depression in colorectal cancer patients. *Psycho-Oncology*. 2017;26(9):1263–9.
- Deckx L, van Abbema DL, van den Akker M, van den Broeke C, van Driel M, Bulens P, et al. A cohort study on the evolution of psychosocial problems in older patients with breast or colorectal cancer: comparison with younger cancer patients and older primary care patients without cancer. *BMC Geriatr*. 2015;15:79.
- National Institute for Health and Clinical Excellence (NICE). Improving supportive and palliative care for adults with cancer (CSG4). In: National Institute for Health and Clinical Excellence (NICE), editor. London: National Institute for Health and Clinical Excellence (NICE); 2004.
- Carlson LE, Waller A, Mitchell AJ. Screening for distress and unmet needs in patients with cancer: review and recommendations. *J Clin Oncol*. 2012;30(11):1160–77.
- Berian JR, Cuddy A, Francescatti AB, O'Dwyer L, Nancy You Y, Volk RJ, et al. A systematic review of patient perspectives on surveillance after colorectal cancer treatment. *J Cancer Surviv*. 2017;11(5):542–52.
- Arora NK, Finney Rutten LJ, Gustafson DH, Moser R, Hawkins RP. Perceived helpfulness and impact of social support provided by family, friends, and health care providers to women newly diagnosed with breast cancer. *Psycho-Oncology*. 2007;16(5):474–86.
- Cairns SR, Scholefield JH, Steele RJ, Dunlop MG, Thomas HJ, Evans GD, et al. Guidelines for colorectal cancer screening and surveillance in moderate and high risk groups (update from 2002). *Gut*. 2010;59(5):666–89.
- Jorgensen ML, Young JM, Solomon MJ. Optimal delivery of colorectal cancer follow-up care: improving patient outcomes. *Patient Relat Outcome Meas*. 2015;6:127–38.
- Fenlon D, Richardson A, Addington-Hall J, Smith P, Corner J, Winter J, et al. A cohort study of the recovery of health and wellbeing following colorectal cancer (CREW study): protocol paper. *BMC Health Serv Res*. 2012;12(1):90.
- Foster C, Fenlon D. Recovery and self-management support following primary cancer treatment. *Br J Cancer*. 2011;105:S21.
- Agrell B, Dehlin O. Comparison of six depression rating scales in geriatric stroke patients. *Stroke*. 1989;20(9):1190–4.
- Vilagut G, Forero CG, Barbaglia G, Alonso J. Screening for depression in the general population with the Center for Epidemiologic Studies Depression (CES-D). A systematic review with meta-analysis. *PLoS One*. 2016;11(5):e0155431.



36. Radloff LS. The CES-D Scale: a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1(3):385–401.
37. Wakefield CE, Butow PN, Aaronson NA, Hack TF, Hulbert-Williams NJ, Jacobsen PB. Patient-reported depression measures in cancer: a meta-review. *Lancet Psychiat*. 2015;2(7):635–47.
38. Katz MR, Kopek N, Waldron J, Devins GM, Tomlinson G. Screening for depression in head and neck cancer. *Psycho-Oncology*. 2004;13(4):269–80.
39. Wheelwright S, Permyakova NV, Calman L, Din A, Fenlon D, Richardson A, et al. Does quality of life return to pre-treatment levels five years after curative intent surgery for colorectal cancer? Evidence from the ColoREctal Wellbeing (CREW) study. *PLoS One*. 2020;15(4):e0231332.
40. Cummings A, Grimmett C, Calman L, Patel M, Permyakova NV, Winter J, et al. Comorbidities are associated with poorer quality of life and functioning and worse symptoms in the 5 years following colorectal cancer surgery: results from the ColoREctal Well-being (CREW) cohort study. *Psycho-Oncology*. 2018;27(10):2427–35.
41. Hu FB, Goldberg J, Hedeker D, Flay BR, Pentz MA. Comparison of population-averaged and subject-specific approaches for analyzing repeated binary outcomes. *Am J Epidemiol*. 1998;147(7):694–703.
42. Foster C, Haviland J, Winter J, Grimmett C, Chivers Seymour K, Batehup L, et al. Pre-surgery depression and confidence to manage problems predict recovery trajectories of health and wellbeing in the first two years following colorectal cancer: results from the CREW cohort study. *PLoS One*. 2016;11(5):e0155434.
43. NHS England. Universal personalised care: implementing the comprehensive model. NHS England, 2019.
44. Zhang X, Liu J, Zhu H, Zhang X, Jiang Y, Zhang J. Effect of psychological intervention on quality of life and psychological outcomes of colorectal cancer patients. *Psychiatry*. 2020;83(1):58–69.
45. Meng X, Wang X, Dong Z. Impact of non-pharmacological interventions on quality of life, anxiety, and depression scores in patients with colorectal cancer: a systematic review and meta-analysis of randomized controlled trials. *Support Care Cancer*. 2021;29(10):5635–52.
46. Jefford M, Kinnane N, Howell P, Nolte L, Galetakis S, Bruce Mann G, et al. Implementing novel models of posttreatment care for cancer survivors: enablers, challenges and recommendations. *Asia Pac J Clin Oncol*. 2015;11(4):319–27.
47. Mittmann N, Beglaryan H, Liu N, Seung SJ, Rahman F, Gilbert J, et al. Examination of health system resources and costs associated with transitioning cancer survivors to primary care: a propensity-score-matched cohort study. *J Oncol Pract*. 2018;14(11):682.
48. Alfano CM, Mayer DK, Bhatia S, Maher J, Scott JM, Nekhlyudov L, et al. Implementing personalized pathways for cancer follow-up care in the United States: proceedings from an American Cancer Society–American Society of Clinical Oncology summit. *CA Cancer J Clin*. 2019;69(3):234–47.
49. Krebber AMH, Jansen F, Witte BI, Cuijpers P, de Bree R, Becker-Commissaris A, et al. Stepped care targeting psychological distress in head and neck cancer and lung cancer patients: a randomized, controlled trial. *Ann Oncol*. 2016;27(9):1754–60.
50. Sharpe M, Walker J, Hansen CH, Martin P, Symeonides S, Gourley C, et al. Integrated collaborative care for comorbid major depression in patients with cancer (SMART Oncology-2): a multicentre randomised controlled effectiveness trial. *Lancet*. 2014;384(9948):1099–108.
51. Recklitis CJ, Syrjala KL. Provision of integrated psychosocial services for cancer survivors post-treatment. *Lancet Oncol*. 2017;18(1):e39–50.
52. National Institute for Health and Clinical Excellence (NICE). Colorectal cancer: diagnosis and management (CG131). National Institute for Health and Clinical Excellence (NICE), London; 2011.
53. Chongpison Y, Hornbrook MC, Harris RB, Herrinton LJ, Gerald JK, Grant M, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psycho-Oncology*. 2016;25(11):1350–6.
54. Clark L, Holcombe C, Hill J, Downey H, Fisher J, Krespi MR, et al. The perception of support received from breast care nurses by depressed patients following a diagnosis of breast cancer. *Ann R Coll Surg Engl*. 2009;91(1):43–5.
55. Stark DPH, House A. Anxiety in cancer patients. *Br J Cancer*. 2000;83(10):1261.
56. NHS England. The NHS Long Term Plan. NHS England; 2019.
57. Vardy JL, Dhillon HM, Pond GR, Rourke SB, Bekele T, Renton C, et al. Cognitive function in patients with colorectal cancer who do and do not receive chemotherapy: a prospective, longitudinal, controlled study. *J Clin Oncol*. 2015;33(34):4085–92.
58. Yi JC, Syrjala KL. Anxiety and depression in cancer survivors. *Med Clin*. 2017;101(6):1099–113.
59. Andrykowski MA, Aarts MJ, van de Poll-Franse LV, Mols F, Slooter GD, Thong MS. Low socioeconomic status and mental health outcomes in colorectal cancer survivors: disadvantage? advantage? or both? *Psycho-Oncology*. 2013;22(11):2462–9.
60. Li F, Morgan A, McCullagh A, Johnson A, Giles C, Greenfield D, et al. Abstract 3348: Top 10 living with and beyond cancer research priorities. *Can Res*. 2019;79(13 Supplement):3348.
61. Department for Communities and Local Government. The English index of multiple deprivation (IMD) 2015—guidance. Department for Communities and Local Government; 2015.
62. Brugha TS, Cragg D. The list of threatening experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand*. 1990;82(1):77–81.
63. Sherbourne CD, Stewart AL. The MOS social support survey. *Soc Sci Med*. 1991;32(6):705–14.
64. Lorig KR, Ritter P, Stewart AL, Sobel DS, Brown BW, Bandura A, et al. Chronic Disease Self-Management Program: 2-year health status and health care utilization outcomes. *Med Care*. 2001;39(11):1217–23.
65. Foster C, Breckons M, Hankins M, Fenlon D, Cotterell P. Developing a scale to measure self-efficacy to self-manage problems following cancer treatment. *Psycho-Oncology*. 2013;22:1–29.
66. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol*. 1988;54(6):1063.
67. Spielberger C, Gorsuch R, Lushene R, Vagg P, Jacobs G. Manual for the State-Trait Anxiety Inventory. Consulting Psychologists Press; 1970.
68. International Wellbeing Group. Personal Wellbeing Index. Australian Centre on Quality of Life, Deakin University Melbourne; 2006.
69. EuroQol Group. EuroQol—a new facility for the measurement of health-related quality of life. *Health Policy*. 1990;16(3):199.
70. Avis NE, Smith KW, McGraw S, Smith RG, Petronis VM, Carver CS. Assessing quality of life in adult cancer survivors (QLACS). *Qual Life Res*. 2005;14(4):1007–23.
71. Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst*. 1993;85(5):365–76.
72. Knight RG, Waal-Manning HJ, Spears GF. Some norms and reliability data for the State-Trait Anxiety Inventory and the Zung Self-Rating Depression Scale. *Br J Clin Psychol*. 1983;22(4):245–9.
73. Dunn J, Ng SK, Holland J, Aitken J, Youl P, Baade PD, et al. Trajectories of psychological distress after colorectal cancer. *Psycho-Oncology*. 2013;22:1759–65.

74. Hart SL, Charles ST. Age-related patterns in negative affect and appraisals about colorectal cancer over time. *Health Psychol.* 2013;32:302.
75. Gonzalez-Saenz de Tejada M, Bilbao A, Baré M, Briones E, Sarasqueta C, Quintana JM, et al. Association between social support, functional status, and change in health-related quality of life and changes in anxiety and depression in colorectal cancer patients. *Psycho-Oncology.* 2017;26:1263–9.
76. Quintana JM, Antón-Ladislá A, González N, Lázaro S, Baré M, de Larrea NF, et al. Outcomes of open versus laparoscopic surgery in patients with colon cancer. *Eur J Surg Oncol.* 2018;44:1344–53.
77. Mols F, Schoormans D, de Hingh I, Oerlemans S, Husson O. Symptoms of anxiety and depression among colorectal cancer survivors from the population-based, longitudinal PROFILES registry: prevalence, predictors, and impact on quality of life. *Cancer.* 2018;124:2621–8.
78. Dunn J, Ng SK, Holland J, Aitken J, Youl P, Baade PD, et al. Trajectories of psychological distress after colorectal cancer. *Psycho-Oncology.* 2013;22:1759–65.
79. Hart SL, Charles ST. Age-related patterns in negative affect and appraisals about colorectal cancer over time. *Health Psychol.* 2013;32:302.
80. Gonzalez-Saenz de Tejada M, Bilbao A, Baré M, Briones E, Sarasqueta C, Quintana JM, et al. Association between social support, functional status, and change in health-related quality of life and changes in anxiety and depression in colorectal cancer patients. *Psycho-Oncology.* 2017;26:1263–9.
81. Quintana JM, Antón-Ladislá A, González N, Lázaro S, Baré M, de Larrea NF, et al. Outcomes of open versus laparoscopic surgery in patients with colon cancer. *Eur J Surg Oncol.* 2018;44:1344–53.
82. Mols F, Schoormans D, de Hingh I, Oerlemans S, Husson O. Symptoms of anxiety and depression among colorectal cancer survivors from the population-based, longitudinal PROFILES Registry: prevalence, predictors, and impact on quality of life. *Cancer.* 2018;124:2621–8.

How to cite this article: Calman L, Turner J, Fenlon D, Permyakova NV, Wheelwright S, Patel M, et al; members of the CREW Study Advisory Committee. Prevalence and determinants of depression up to 5 years after colorectal cancer surgery: results from the ColoREctal Wellbeing (CREW) study. *Colorectal Dis.* 2021;23:3234–3250. <https://doi.org/10.1111/codi.15949>

APPENDIX 1

Scoping review search strategy and key studies identified which involve people with CRC including at least one follow-up time point

Dates literature searches conducted:

- October 2016
- August 2017
- September 2018
- May 2019
- July 2019
- October 2020
- March 2021

Databases searched:

- CINHALL
- APA PsycINFO
- APA PsycARTICLES
- MEDLINE (EBSCO)
- ISI Web of Science

Limiters:

- Date published: 1 January 2010 to 1 October 2020
- English language
- Human studies
- Peer-reviewed

Search terms:

Subject/MeSH headings used where appropriate

Colorectal neoplasms OR Colorectal cancer

(Colon OR Rectum) AND (neoplasms OR cancer)

Depression OR MM Depression

Anxiety OR MH Anxiety Disorder

'Mental Health'

'Psychological disorder'

'Psychological distress'

'Worry'

MH Stress, Psychological

MH Mental Disorders

MH Fear

Lead author and year	Country of study	Sample	Assessment time points	Depression measure	Key findings	Comparison with the CREW study
Dunn et al., 2013 [1]	Australia	1884 CRC survivors; Stages I-IV	T1: 5 months after diagnosis. Follow-up: 12 (T2), 24 (T3), 36 (T4), 48 (T5) and 60 (T6) months postdiagnosis	BSI-18	<ul style="list-style-type: none"> • Four trajectories of depressive symptoms: constant low levels, constant high levels and people who increase from low and those who reduce from high levels • Men, younger participants, later stage, poor social support and lower education were more likely to experience high levels of depression • 16.1% of participants were in the 'constant high' level trajectory for depressive symptoms (BSI-18 Depression Subscale) 	<ul style="list-style-type: none"> • No presurgery assessment of psychological distress • Recruitment of patients with metastatic CRC
Hart and Charles, 2013 [2]	USA	139 CRC patients (Stages I-IV)	T1: presurgery. Follow-up: T2, 6 months; T3, 12 months; T4, 18 months postsurgery	CES-D	<ul style="list-style-type: none"> • Mean (SD): T1, 10.45 (8.11); T2, 9.33 (7.80); T3, 9.41 (8.74); T4, 9.49 (9.28) • Older adults reported lower levels of depressive symptoms. Men had fewer depressive symptoms than women 	<ul style="list-style-type: none"> • The prevalence of clinical levels of depression was not assessed • No follow-up assessment beyond 18 months postsurgery • Recruitment of patients with metastatic CRC

Lead author and year	Country of study	Sample	Assessment time points	Depression measure	Key findings	Comparison with the CREW study
Gonzalez-Saenz de Tejada et al., 2017 [3]; Quintana et al., 2018 [4]	Spain	972 CRC patients (including patients in relapse)	T1: presurgery. Follow-up: T2, 12 months; T3, 24 months postsurgery	HADS	<ul style="list-style-type: none"> • 19.6% of participants reported depression at T1 • Patients with depression improved less than participants not reporting depression or anxiety in all health-related quality of life domains (EORTC QLQ-C30) • Overall, few differences in depression symptoms in people undergoing either open or laparoscopic surgery • Mean (SD) (laparoscopy vs. open): T1, 4.28 (4.12) vs. 5.33 (4.84); T2, 3.52 (3.85) vs. 4.08 (4.31); T3, 3.50 (3.97) vs. 4.28 (4.38) 	<ul style="list-style-type: none"> • Recruitment of patients with metastatic CRC • Recruitment of patients in relapse (CREW excluded patients with previous cancer diagnosis) • No follow-up assessment beyond 24 months postsurgery
Mols et al., 2018 [5]	Netherlands	315 CRC survivors (Stages I–IV)	Annual follow-up (1–4 years): T1, 2010; T2, 2011; T3, 2012; T4, 2013	HADS	<ul style="list-style-type: none"> • Significantly higher prevalence of depression (19.0%, N = 2625) compared with a matched population (12.8%, N = 315) during their first assessment • Reduction in depression symptoms over time with the largest difference identified when examining the first and fourth assessments (mean change –0.89) • Fewer depressive symptoms were reported in people who were older, low QoL and lower physical, role, cognitive, emotional and social functioning 	<ul style="list-style-type: none"> • Participants recruited 1 to 4 years postdiagnosis • Recruitment of participants with metastatic CRC

Abbreviations: BSI–18, Brief Symptom Inventory–18; CES-D, Centre for Epidemiologic Studies Depression Scale; CRC, colorectal cancer; EORTC QLQ-C30, European Organisation for Research and Treatment of Cancer Quality of Life Core-30 Questionnaire; HADS, Hospital Anxiety and Depression Scale; QoL, quality of life.

APPENDIX 2

Availability of the covariates at two time points of the regression analyses

Thematic block	Topic/measure	Taken time points in separate regression models	
		Baseline (presurgery)	2 years postsurgery
Preexisting factors (socio-demographics)	Age	+	+
	Gender	+ ^a	+ ^a
	Ethnicity	+ ^a	+ ^a
	Employment status	+	+
	Accommodation type	+	+
	Index of multiple deprivation	+ ¹	+ ¹
Clinical factors	Tumour site	+	+
	Dukes stage	+	+
	Neoadjuvant treatment	+	+
	Adjuvant treatment	-	+
	Surgery type	-	+
	Stoma status	-	+
	Number of comorbidities	+ ⁿ	+
	Previous use of mental health services	+	-
Environmental factors	Domestic status	+	+
	Life events	-	+
	Medical Outcome Study Social Support Scale (MOS-SSS)	+	+
Personal factors	Self-Efficacy for Managing Chronic Disease 6-Item Scale (SEMCD)	+	-
	Cancer Survivor Self-Efficacy Scale (CS-SES)	-	+
	Positive and Negative Affect Schedule Short Form (PANAS-SF)	+	+
Psychosocial outcomes	Centre for Epidemiologic Studies Depression Scale (CES-D)	+	+
	Quality of Life in Adult Cancer Survivors (QLACS) scale: QLACS Cancer-Specific Summary Score	-	+
	QLACS Benefit of Cancer subscale	-	+
	State-Trait Anxiety Inventory, State scale (STAI-S)	+	+
	Personal Wellbeing Index - Adult (PWI-A)	+	+
	EQ-5D-3L	+	+
	EORTC-QLQ-C30: physical functioning	-	+
	EORTC-QLQ-C30: emotional functioning	-	+
	EORTC-QLQ-C30: cognitive functioning	-	+
	EORTC-QLQ-C30: social functioning	-	+
	EORTC-QLQ-C30: fatigue	-	+
	EORTC-QLQ-C30: pain	-	+
	EORTC-QLQ-C30: insomnia	-	+
EORTC-QLQ-C30: financial worry	-	+	

Note: data are taken from the same time point unless otherwise annotated: ^adata taken from baseline time point, ^bdata taken from the 3 month follow-up time point. Abbreviations: EORTC QLQ-C30, European Organisation for Research and Treatment of Cancer Quality of Life Core-30 Questionnaire; EQ-5D-3L, EuroQoL five dimensions, three levels. Key: '+' included in regression analysis for the time point; '-' indicates excluded from regression analysis for the time point due to the measure not being assessed at the time point.

APPENDIX 3

Self-reported health service use (Have you used any of the following health and social services in the last 12 months?)

Time point (postsurgery)	24 months		36 months		48 months		60 months	
	≥20, n (%)	<20, n (%)	≥20, n (%)	<20, n (%)	≥20, n (%)	<20, n (%)	≥20, n (%)	<20, n (%)
N	73	410	49	333	48	321	47	272
Mental health services	2 (2.7)	0 (0)	2 (4.1)	1 (0.3)	3 (6.3)	1 (0.3)	1 (2.1)	1 (0.4)
Counselling services	4 (5.5)	2 (0.5)	6 (12.2)	2 (0.6)	2 (4.2)	5 (1.6)	1 (2.1)	2 (0.7)
Psychiatrist	4 (5.5)	2 (0.5)	4 (8.2)	1 (0.3)	3 (6.3)	0 (0)	2 (4.3)	1 (0.4)
Self-help group	5 (6.8)	4 (1.0)	2 (4.1)	5 (1.5)	3 (6.3)	6 (1.9)	3 (6.4)	4 (1.5)

Abbreviation: CES-D, Centre for Epidemiologic Studies Depression Scale.