

Hepatic bile acids and bile acid-related gene expression in pregnant and lactating rats

Qiong N. Zhu¹, Hong M. Xie², Dan Zhang¹, Jie Liu^{1,3} and Yuan F. Lu¹

¹ Department of Pharmacology and Key Lab of Basic Pharmacology of Guizhou, Zunyi Medical College, Zunyi, China

² Department of Gynaecology and Obstetrics, The Third Affiliated Hospital of Zunyi Medical College, Zunyi, China

³ University of Kansas Medical Center, Kansas City, KS, USA

ABSTRACT

Background. Significant physiological changes occur during pregnancy and lactation. Intrahepatic cholestasis of pregnancy (ICP) is a liver disease closely related to disruption of bile acid homeostasis. The objective of this study was to examine the regulation of bile acid synthesis and transport in normal pregnant and lactating rats.

Materials and Methods. Livers from timed pregnant SD rats were collected on gestational days (GD) 10, 14 and 19, and postnatal days (PND) 1, 7, 14 and 21. Total bile acids were determined by the enzymatic method, total RNA was isolated and subjected to real time RT-PCR analysis. Liver protein was extracted for western-blot analysis.

Results. Under physiological conditions hepatic bile acids were not elevated during pregnancy but increased during lactation in rats. Bile acid synthesis rate-limiting enzyme Cyp7a1 was unchanged on gestational days, but increased on PND14 and 21 at mRNA and protein levels. Expression of Cyp8b1, Cyp27a1 and Cyp7b1 was also higher during lactation. The mRNA levels of small heterodimer partner (SHP) and protein levels of farnesoid X receptor (FXR) were increased during pregnancy and lactation. Bile acid transporters Ntcp, Bsep, Mrp3 and Mrp4 were lower at gestation, but increased during lactation. Hepatic Oatp transporters were decreased during pregnancy and lactation.

Conclusion. Hepatic bile acid homeostasis is maintained during normal pregnancy in rats, probably through the FXR-SHP regulation. The expression of bile acid synthesis genes and liver bile acid accumulation were increased during lactation, together with increased expression of bile acid efflux transporter Bsep, Mrp3 and Mrp4.

Subjects Biochemistry, Toxicology, Pharmacology

Keywords Liver bile acids, FXR-SHP, Ntcp and Bsep, Cyp7a1, Pregnant and lactating rats, Oatps

INTRODUCTION

Significant physiological changes occur during pregnancy and lactation to support the nutritional demands of the developing fetus and lactating pups (*Carlin & Alfirevic, 2008; Athippozhy et al., 2011*). Bile acids and cholesterol metabolism are important changes during pregnancy and lactation to support and to protect offspring development

Submitted 24 June 2013
Accepted 5 August 2013
Published 27 August 2013

Corresponding authors
Jie Liu, JLi@kumc.edu
Yuan F. Lu, Yflu@zmc.edu.cn

Academic editor
Rebecca Berdeaux

Additional Information and
Declarations can be found on
page 12

DOI 10.7717/peerj.143

© Copyright
2013 Zhu et al.

Distributed under
Creative Commons CC-BY 3.0

OPEN ACCESS

(Wooton-Kee, Cohen & Vore, 2008; Athipposzhy et al., 2011; Abu-Hayyeh, Papacleovoulou & Williamson, 2013). Such physiological changes would also affect hepatic drug processing genes of phase-1, phase-2 metabolism and transporters (Aleksunes et al., 2012; Shuster et al., 2013). The alteration of bile acid homeostasis during pregnancy could unmask cholestatic disease in genetically predisposed but otherwise asymptomatic individuals (Milona et al., 2010). Recent work suggests that in pregnant mice farnesoid X receptor (FXR)-SHP (small heterodimer partner, NR0B2) regulation could be dysfunctional in its ability to downregulate the rate-limiting bile acid synthetic enzyme Cyp7a1 and 8b1, resulting in bile acids accumulation in the liver of late pregnancy mice (Milona et al., 2010; Aleksunes et al., 2012).

Intrahepatic cholestasis of pregnancy (ICP) is a liver disease which can occur in the third trimester of pregnancy (Abu-Hayyeh, Papacleovoulou & Williamson, 2013). The etiology and pathogenesis of ICP are still not clear, but many studies have related this disease to abnormal bile acid metabolism (Abu-Hayyeh, Papacleovoulou & Williamson, 2013; Floreani et al., 2013). ICP with elevated bile acids in serum and liver is a major cause for premature embryo development and embryonic death (Diken, Usta & Nassar, 2013). Genetic variations or mutations of farnesoid X receptor (FXR) (Van Mil et al., 2007), bile salt export pump (BSEP/ABCB11) (Dixon et al., 2009), and ATP-binding cassette, sub-family B (MDR/TAP), member 4 (ABCB4/MDR3) and ABCB11 (Dixon et al., 2000; Anzivino et al., 2013) contribute to the etiology of ICP. To fully understand bile acid synthesis, transport, and regulation in normal pregnancy would help us to shed light on the pathology of ICP.

Estradiol and/or its metabolites may interfere with FXR activity during pregnancy (Milona et al., 2010; Aleksunes et al., 2012), and a defect in progesterone metabolism is also implicated in the etiology of ICP (Pascual et al., 2002). Estrogen signaling is associated with pregnancy-induced hepatotoxicity and cholestasis in mice (Arrese et al., 2008), and reduced hepatic PPAR- α function in the mouse also appears to be estrogen-dependent (Papacleovoulou, Abu-Hayyeh & Williamson, 2011).

The above scenario has been studied extensively in mice (Milona et al., 2010; Aleksunes et al., 2012; Shuster et al., 2013). Mice and rats are the two most commonly used experimental animals, but some physiological responses are different. For example, in mice, Cyp7a1 and liver bile acid pool were not increased during lactation (Aleksunes et al., 2012), whereas the bile acid synthesis gene Cyp7a1 and hepatic bile acids are increased 2 to 3-fold in lactating rats (Wooton-Kee, Cohen & Vore, 2008; Wooton-Kee et al., 2010). In mice, pregnancy and lactation are associated with decreases in hepatic transporters, including bile acid transporters (Aleksunes et al., 2012), and such a phenomenon should also be characterized in rats. To fully understand bile acid synthesis, transport, and regulation in normal pregnancy would help us to shed light on the pathology of ICP. This study was initiated to investigate bile acid metabolism and transport gene expressions in pregnant and lactating rats, and the results suggest that under physiological conditions, FXR-SHP regulation might play a role in bile acid homeostasis in pregnant and lactating rats.

MATERIALS AND METHODS

Animals

Adult Sprague Dawley (SD) rats (250 g) were purchased from the Experimental Animal Center of Third Military Medical College (Chongqing, China; certificate No. CXX 2007-0005). Rats were kept in a SPF-grade animal facilities (certificate No. SYXX 2011-004) at Zunyi Medical College, with regulated environment ($22 \pm 1^\circ\text{C}$, $50 \pm 2\%$ humidity and a 12 h:12 h light:dark cycle) and free access to purified water and standard rodent chow. Rats were acclimatized for 1 week, and subjected to timely mating overnight. A vaginal plug the next morning was designated as day 0 (GD 0) of gestation. Maternal livers were collected without fasting on GD10, GD14 and GD19, as well as on the postnatal days (PND) 1, 7, 14 and 21. The age-matched virgin rats were used as controls. The experimental design followed similar time points in mice (*Aleksunes et al., 2012*). Livers were weighed, snap frozen in liquid nitrogen, and stored at -80°C until analysis. All animal procedures followed the NIH guide of Humane Use and Care Animals, and were approved by Institutional Animal Use and Care Committee of Zunyi Medical College.

Bile acid determination

Bile acids were extracted from the liver and measured with the “Total” Bile Acid assay (TBA) kit (Nanjing Jian-Cheng Bioengineering Co., China). Briefly, livers were homogenized in physiological saline (1:9, wt:vol), followed by centrifugation at 2500 rpm/min for 10 min. The supernatant (30 μl) was taken for determination of bile acids according to the manufacturer’s protocol.

RNA Isolation and real-time RT-PCR analysis

Total RNA was isolated from frozen liver samples (50–100 mg) using 1 ml TRIzol (Takara, Biotechnology, Dalian, China) and subsequently purified with Total RNA (Mini) Kit (Watson Biotechnology, Shanghai, China). The quality of purified RNA was determined by spectrophotometry with the 260/280 ratio > 1.8 . Purified RNA was reverse transcribed with the High Capacity Reverse Transcriptase Kit (Applied Biosystems, Foster City, CA, USA). The primer pairs were designed with the Primer3 software and listed in [Table S1](#). The Power SYBR Green Master Mix (Applied Biosystems, Foster City, CA, USA) was used for real-time RT-PCR analysis. The cycle time for reaching threshold (Ct) of each target gene was normalized to the housekeeping genes (G3PDH and β -actin), and expressed as % of housekeeping genes.

WESTERN BLOT ANALYSIS

Livers were homogenized in a RIPA lysis buffer (Beyotime, P0013B, Shanghai, China) containing freshly-prepared proteinase inhibitors. The supernatants were centrifuged at 12000 rpm 10 min at 4°C , and protein concentrations were quantified by the BCA assay (Beyotime, P0012, Shanghai, China). Aliquot proteins were denatured with a protein loading buffer (Beyotime, P0015, Shanghai, China), and approximately 50 μg of protein/lane was separated on 10% SDS-PAGE and transferred to PVDF membranes. Membranes




Figure 1 Liver bile acid levels in pregnant and lactating rats. Bile acids were quantified in livers from control and pregnant rats on GD10, 14, and 19 and PND 1, 7, 14, and 21. Dark gray bars represent pregnant rats, and black bars represent lactating rats. Data are presented as mean \pm SEM ($n = 3-6$). Asterisks (*) represent statistically significant difference ($p < 0.05$) compared with the control.

were blocked in 5% non-fat milk in TBST, followed by incubation overnight at 4°C with 1:1000 CYP7A1 (ab65586) and β -actin (Ab8227) from Abcam (Cambridge, MA), or FXR (sc-13063) from Santa Cruz Biotechnology (Santa Cruz, CA) in 1% BSA. After washing with TBST three times, the membranes were incubated with HRP-conjugated anti-rabbit or anti-mouse IgG (Beyotime, A0208 and A0216, Shanghai, China). Protein-antibody complexes were visualized using an enhanced chemiluminescent reagent (ECL-Plus) (Beyotime, P0018, Shanghai, China), and exposed to Gel Imaging (Bio-Rad, ChemiDoc XRS, USA). The intensity of the band was semi-quantified with Quantity One software.

STATISTICAL ANALYSIS

The software SPSS 17.0 was used for statistical analysis. The results were described using mean \pm SEM. The difference between virgin and pregnant rats was determined by two-tailed independent samples test, $P < 0.05$ was considered statistically significant.

RESULTS

Liver bile acid levels in pregnant and lactating rats

Bile acids were quantified in livers from control and pregnant rats at GD10, 14, and 19 and PND 1, 7, 14, and 21. Liver bile acid levels slightly decreased in late pregnancy, especially on GD 10, and 19. After birth, liver bile acid concentrations tended to increase, and there is a significant increase in PND 21 (30% over control) (Fig. 1).




Figure 2 Hepatic mRNA expression of bile acid synthetic pathway genes in pregnant and lactating rats. The expression of bile acid synthetic classic pathway genes Cyp7a1, Cyp8b1 and alternative pathway genes Cyp27a1 and Cyp7b1 was quantified from control and GD10,14 and 19 and PND 1, 7, 14 and 21. Data were normalized to controls (set to 100%) and presented as mean \pm SEM ($n = 3-6$). Dark gray bars represent pregnant rats, and black bars represent lactating rats. Asterisks (*) represent statistically significant difference ($p < 0.05$) compared with the control.

Hepatic mRNA expression of bile acid synthesis genes in pregnant and lactating rats

The expression of the classic pathway bile acid synthetic enzyme genes (Cyp7a1 and 8b1) and alternative pathway (Cyp27a1 and 7b1), is shown in Fig. 2. The expression of rate-limiting Cyp7a1 mRNA was unchanged during pregnancy, and increased on postpartum. Cyp8b1 mRNA decreased in GD10 and GD14, and increased about 2-fold in PND14. The expression of alternative pathway genes Cyp27a1 and Cyp7b1 were unchanged in gestation days and increased in postnatal days.

Hepatic expression of bile acid synthetic rate-limiting protein CYP7A1 in pregnant and lactating rats

Western blots were performed using liver homogenates from control rats, pregnant rats at GD 10, 14, 19 and lactating rats at PND 1, 7, 14 and 21. The expressions of CYP7A1 protein were semi-quantified by band intensity. CYP7A1 protein was basically unchanged during




Figure 3 Hepatic expression of bile acid synthesis rate-limiting protein CYP7A1 in pregnant and lactating rats. Western blots were performed using liver homogenates from control, and pregnant rats in GD 10, 14, 19 and PND 1, 7, 14 and 21. The expression of CYP7A1 was semi-quantified by band intensity. Values are mean \pm SEM ($n = 5$). Dark gray bars represent pregnant rats, and black bars represent lactating rats. The statistically significant difference was confirmed with a two-tailed independent samples test method ($P < 0.05$).

pregnancy, a result similar to Cyp7a1 mRNA expression, but increased on lactation days PND7, 14 and 21 (Fig. 3).

Hepatic mRNA expression of nuclear receptors FXR, SHP, and ESR-1, PPAR- α in pregnant and lactating rats

The expression of bile acid regulation nuclear receptor genes farnesoid X receptor (FXR, NR1H4) did not show significant increases during pregnancy, while FXR gradually increased on postpartum. The small heterodimer partner (SHP; NR0B2) significantly increased in the late gestational days, increased 3-fold on GD 19 compared to controls. FXR plays an important role in bile acid homeostasis by inducing the transcription repressor SHP (Chiang, 2009). Estrogen receptor alpha (ESR-1) decreased to 64.7% and 57.7% on GD10 and GD14. In postnatal days, ESR-1 increased 2.33-fold in PND1 and then decreased to 68% of control on PND21. Proliferator-activated receptor α (PPAR α) increased 3.79-fold compared to controls during lactation (Fig. 4).

Hepatic expression of FXR protein in pregnant and lactating rats

Western blots were performed using liver homogenates from control rats, pregnant rats at GD 10, 14, 19 and lactating rats at PND 1, 7, 14 and 21. The expression of FXR protein was semi-quantified by band intensity. FXR protein was increased during late pregnancy (GD10 to GD19) and early lactation (PND1 to PND7) (Fig. 5).




Figure 4 Hepatic mRNA expression of nuclear receptors SHP, FXR and ESR-1 and PPAR- α in pregnant and lactating rats. The expression of bile acid regulation nuclear receptors genes SHP, FXR and ESR-1, PPAR- α were quantified using total hepatic RNA from control and pregnant mice at gestational days 10, 14, 19 and postnatal days 1, 7, 14, 21. Data were normalized to controls and presented as mean \pm SEM ($n = 3-6$). Dark gray bars represent pregnant rats, and black bars represent lactating rats. Asterisks (*) represent statistically significant differences ($p < 0.05$) compared to the control.

Hepatic mRNA expression of bile acid transporters in pregnant and lactating rats

As illustrated in Fig. 6, the expression of bile acid efflux transporter bile salt export pump (Bsep/ABCB11) was decreased during pregnancy but increased during lactation. The multidrug resistance protein 3 (Mrp3) and Mrp4 showed a similar pattern, with slight increases during lactation. The ATP-binding cassette sub-family G member 2 (Abcg2/BCRP) was also decreased in the gestation and lactation days except PND1.

Hepatic mRNA expression of uptake OATP transporters and Ntcp in pregnant and lactating rats

Figure 7 demonstrates that the expression of canalicular uptake transporter solute carrier organic anion transporter (Oatp1/Slco1a1), solute carrier organic anion transporter




Figure 5 Hepatic expression of nuclear receptor FXR in pregnant and lactating rats. Western blots were performed using liver homogenates from control, pregnant rats in GD 10, 14, 19 and PND 1, 7, 14 and 21. The expression of FXR was semi-quantified by band intensity. Values are mean \pm SEM ($n = 3$). Dark gray bars represent pregnant rats, and black bars represent lactating rats. Statistically significant difference was confirmed with a two-tailed independent samples test method ($P < 0.05$).

(Oatp2/Slco1b2), and organic anion-transporting polypeptide 4 (Oatp4/Slc21a10) were all decreased in the gestation days, on PND1 Oatp1 increased 1.68-fold, then decreased in postnatal days. In comparison, Oatp2 and Oatp4 decreased in the both gestation days and lactation days. The uptake transporter Na^+ -taurocholate co-transporting polypeptide (Ntcp) was also decreased during pregnancy, increased on PND1, but decreased again thereafter during lactation.

DISCUSSION

The present study demonstrates that in pregnant rats, hepatic bile acids were not elevated. Consistent with hepatic bile acid concentrations, bile acid synthesis genes and/or enzymes, i.e., Cyp7a1, Cyp8b1, Cyp27a1 and Cyp7b1 were not increased during pregnancy. Increased FXR protein and SHP mRNA are associated with bile acid homeostasis during pregnancy. In comparison, lactating rats had increased liver bile acid, increased bile acid synthetic enzymes, and increased expression of bile acid efflux transporters. In general, OATP transporters and bile acid uptake transport Ntcp were downregulated during pregnancy and lactation in rats.

ICP is characterized by raised serum bile acid levels and abnormal liver function tests (Geenes & Williamson, 2009; Diken, Usta & Nassar, 2013). However, in normal pregnant women, serum bile acid levels are not necessarily increased during pregnancy, regardless of gestation days (Barth et al., 2005; Egan et al., 2012). In experimental animal studies, a




Figure 6 Hepatic mRNA expression of bile acid transporter in pregnant and lactating rats. The expression of bile acid efflux transporter Bsep, Mrp3, Mrp4 and Abcg2 was quantified using total hepatic RNA from pregnant rats on GD 10, 14 and 19, and postpartum rats on PND 1, 7, 14 and 21. Data were normalized to controls and presented as mean \pm SEM ($n = 3-6$). Dark gray bars represent pregnant rats, and black bars represent lactating rats. Asterisks (*) represent statistically significant differences ($p < 0.05$) compared to the control.

mild increase in liver bile acid levels during normal pregnancy in mice was reported in some studies (Aleksunes *et al.*, 2012), but not in others (Abu-Hayyeh, Papacleovoulou & Williamson, 2013). In the majority of cases, such mild increases do not reach pathological levels and remain below the upper end of the reference range for serum bile acid levels (Abu-Hayyeh, Papacleovoulou & Williamson, 2013). Thus, it is not surprising that in the present study, liver bile acids were not elevated during pregnancy in gestation days (Fig. 1). The expression of bile acid synthesis gene and proteins during the gestation days (Figs. 2 and 3) is in agreement with hepatic bile acid profiles.

ICP has a complex etiology including genetic factors, endocrine factors, and the impact of pregnancy on FXR function (Abu-Hayyeh, Papacleovoulou & Williamson, 2013; Floreani *et al.*, 2013). The present study focused on FXR-SHP regulation under physiological conditions. It is proposed that pregnancy in mice resembles a state of FXR inactivation (Milona *et al.*, 2010; Aleksunes *et al.*, 2012), and attenuated FXR function during mouse




Figure 7 Hepatic mRNA expression of uptake transporter Oatps and Ntcp in pregnant and lactating rats. The expression of hepatic canalicular uptake transporter solute carrier organic anion transporters (Oatps) and Na⁺-taurocholate co-transporting polypeptide (Ntcp) transporters was quantified using total hepatic RNA from pregnant rats on GD10, 14 and 19, and postpartum rats on PND 1, 7, 14 and 21. Data were normalized to controls and presented as mean ± SEM (*n* = 3–6). Dark gray bars represent pregnant rats, and black bars represent lactating rats. Asterisks (*) represent statistically significant differences (*p* < 0.05) compared to the control.

pregnancy has been reported (*Papacleovoulou, Abu-Hayyeh & Williamson, 2011; Aleksunes et al., 2012*) and the 3 β -sulfated progesterone metabolite epiallopregnanolone sulfate was found to inhibit FXR, resulting in reduced FXR-mediated bile acid efflux (*Abu-Hayyeh et al., 2013*). In the present study, the expression of FXR mRNA in rats during pregnancy was basically unchanged. However, the FXR protein and FXR-inducible negative target SHP were markedly increased at the late gestation days and reached approximately 3-fold higher at GD14 and GD19, despite FXR mRNA not increasing. The increases in FXR-SHP may play an important role in maintaining the bile acid homeostasis and preventing the liver bile acids from accumulating to protect the fetus from the bile acid toxicity.

Lactation is a time of a five-fold increase in energy demand, as suckling young requires a proportional adjustment in the ability of the lactating dam to absorb nutrients (*Cripps & Williams, 1975; Vernon et al., 2002*). Lactating rats have a 2 to 3-fold increase in

food consumption to ensure lactating dams absorb nutrients and synthesize critical molecules including bile acids to meet the dietary needs of the offspring and the dam (Vernon *et al.*, 2002). The size and hydrophobicity of the bile acid pool increase during lactation, implying an increased absorption and disposition of lipid, sterols, nutrients, and xenobiotics (Athippozhy *et al.*, 2011). In essence, rats (Wooton-Kee, Cohen & Vore, 2008) are different from mice (Aleksunes *et al.*, 2012) in bile acid homeostasis during lactation. In the present study, hepatic bile acid pool (Fig. 1), mRNA levels of bile acid synthesis gene Cyp7a1, Cyp8b1, Cyp27a1 and Cyp7b1 (Figs. 2 and 3) were all increased during lactation, consistent with this scenario.

The mRNA levels of bile acid transporters Ntcp and Bsep followed a similar pattern. Ntcp is the major bile acid transporter for conjugated bile acid (Csanaky *et al.*, 2011) and Bsep is the major bile acid efflux pump located at the bile canalicular apical domain of hepatocytes (Lam, Soroka & Boyer, 2010). Downregulation of Ntcp and Bsep was observed in pregnant rats (Arrese *et al.*, 2003; Cao *et al.*, 2001), however, they increase during early postpartum, probably under the influence of prolactin (Cao *et al.*, 2001). In the present study, the changes in mRNA levels of Ntcp and Bsep showed a similar pattern, i.e., lower expression during pregnancy but returned to normal and even increased during lactation.

Sulfated progesterone metabolite (P4-S) levels are raised in normal pregnancy and elevated further in ICP, which can cause a competitive inhibition of NTCP-mediated uptake of taurocholate in *Xenopus* oocytes (Abu-Hayyeh *et al.*, 2010), and also can cause inhibition of BSEP (Vallejo *et al.*, 2006). In the present study, mRNA levels of Ntcp and Bsep (Fig. 5) were lower during pregnancy, and Bsep was increased during lactation, consistent with liver bile acid homeostasis profile. Mrp3 and Mrp4 are two major bile acid efflux (Cui *et al.*, 2009; Aleksunes *et al.*, 2012), and mRNA levels of these two genes expression showed the similar pattern (Fig. 5), i.e., lower during the pregnancy and higher during lactation. The pattern of these transporter mRNA levels coincide with FXR-SHP regulation of bile acid homeostasis, and fortifying the concept that under physiological conditions, FXR-SHP regulation of bile acid synthesis could be essential for maintaining the bile acid homeostasis and could prevent the occurrence of ICP, an unusual pathological condition.

One of the major findings in the study is the decreased mRNA levels of Oatp transporters (Fig. 6), and this finding is consistent with that observed in mice (Aleksunes *et al.*, 2012; Shuster *et al.*, 2013). Oatps are important not only for bile acid transport (Zhang *et al.*, 2012), but also for drug and xenobiotic transport (Lu *et al.*, 2008). In pregnant rats, the expression of Oatp2, but not Oatp1, was reported to decrease (Cao *et al.*, 2001; Cao *et al.*, 2002). The generalized downregulation of Oatp transporters could be an adaptive mechanism for the dam to protect developing fetuses and nursing pups from toxicants. Abcg2 is involved in epithelial transport/barrier functions, including bile acid transport (Blazquez *et al.*, 2012). Abcg2 is proposed to play a key role in bile acid transport in placenta, as Bsep does in liver (Blazquez *et al.*, 2012). In the present study, mRNA levels of Abcg 2 was depressed during pregnancy and lactation except for a transient increase at PND1. The pattern of Abcg2 expression is similar to Oatps, and can also be envisioned as an adaptive mechanism during pregnancy and lactation.

In summary, the present study suggests that in pregnant rats, FXR-SHP could regulate bile acid synthesis enzyme genes to prevent the accumulation of bile acids in the liver, together with downregulation of bile acid transporters Ntcp and Bsep. Pregnancy and lactation is associated with a general downregulation of Oatp and Abcg2 in rats. These data would add to our understanding of FXR-SHP regulation of bile acid homeostasis under physiological conditions.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by the Science and Technology Foundation of Guizhou Province (QK2008-002 and 2009-70019) and the Foundation of Zunyi Medical College (QK023). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Science and Technology Foundation of Guizhou Province: QK2008-002, 2009-70019.
Foundation of Zunyi Medical College: QK023.

Competing Interests

Jie Liu is an Academic Editor for PeerJ.

Author Contributions

- Qiong N. Zhu and Hong M. Xie performed the experiments, analyzed the data, wrote the paper.
- Dan Zhang performed the experiments, analyzed the data.
- Jie Liu conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper.
- Yuan F. Lu conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All animal procedures follow the NIH guide of Humane Use and Care Animals, and were approved by Institutional Animal Use and Care Committee of Zunyi Medical College (2012-07).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.143>.

REFERENCES

- Abu-Hayyeh S, Martinez-Becerra P, Sheikh Abdul Kadir SH, Selden C, Romero MR, Rees M, Marschall H-U, Marin JJG, Williamson C. 2010. Inhibition of Na⁺-taurocholate Co-transporting polypeptide-mediated bile acid transport by cholestatic sulfated progesterone metabolites. *The Journal of Biological Chemistry* 285:16504–16512 DOI 10.1074/jbc.M109.072140.
- Abu-Hayyeh S, Papacleovoulou G, Williamson C. 2013. Nuclear receptors, bile acids and cholesterol homeostasis series—bile acids and pregnancy. *Molecular and Cellular Endocrinology* 368:120–128 DOI 10.1016/j.mce.2012.10.027.
- Abu-Hayyeh S, Papacleovoulou G, Lövgren-Sandblom A, Tahir M, Oduwole O, Jamaludin NA, Ravat S, Nikolova V, Chambers J, Selden C, Rees M, Marschall HU, Parker MG, Williamson C. 2013. Intrahepatic cholestasis of pregnancy levels of sulfated progesterone metabolites inhibit farnesoid X receptor resulting in a cholestatic phenotype. *Hepatology* 57:716–726 DOI 10.1002/hep.26055.
- Aleksunes LM, Yeager RL, Wen X, Cui JY, Klaassen CD. 2012. Repression of hepatobiliary transporters and differential regulation of classic and alternative bile acid pathways in mice during pregnancy. *Toxicological Sciences* 130:257–268 DOI 10.1093/toxsci/kfs248.
- Anzivino C, Odoardi MR, Meschiari E, Baldelli E, Facchinetti F, Neri I, Ruggiero G, Zampino R, Bertolotti M, Loria P, Carulli L. 2013. ABCB4 and ABCB11 mutations in intrahepatic cholestasis of pregnancy in an Italian population. *Digestive and Liver Disease* 45:226–232 DOI 10.1016/j.dld.2012.08.011.
- Arrese M, Trauner M, Ananthanarayanan M, Pizarro M, Solís N, Accatino L, Soroka C, Boyer JL, Karpen SJ, Miquel JF, Suchy FJ. 2003. Down-regulation of the Na⁺/taurocholate cotransporting polypeptide during pregnancy in the rat. *Journal of Hepatology* 38:148–155 DOI 10.1016/S0168-8278(02)00379-3.
- Arrese M, Macias RI, Briz O, Perez MJ, Marin JJ. 2008. Molecular pathogenesis of intrahepatic cholestasis of pregnancy. *Expert Reviews in Molecular Medicine* 10:e9 DOI 10.1017/S1462399408000628.
- Athipozhy A, Huang L, Wooton-Kee CR, Zhao T, Jungsuwadee P, Stromberg AJ, Vore M. 2011. Differential gene expression in liver and small intestine from lactating rats compared to age-matched virgin controls detects increased mRNA of cholesterol biosynthetic genes. *BMC Genomics* 12(1):95 DOI 10.1186/1471-2164-12-95.
- Barth A, Rost M, Kindt A, Peiker G. 2005. Serum bile acid profile in women during pregnancy and childbed. *Experimental and Clinical Endocrinology & Diabetes* 113:372–375 DOI 10.1055/s-2005-865719.
- Blazquez AG, Briz O, Romero MR, Rosales R, Monte MJ, Vaquero J, Macias RI, Cassio D, Marin JJ. 2012. Characterization of the role of ABCG2 as a bile acid transporter in liver and placenta. *Molecular Pharmacology* 81:273–283 DOI 10.1124/mol.111.075143.
- Cao J, Huang L, Liu Y, Hoffman T, Stieger B, Meier PJ, Vore M. 2001. Differential regulation of hepatic bile salt and organic anion transporters in pregnant and postpartum rats and the role of prolactin. *Hepatology* 33:140–147 DOI 10.1053/jhep.2001.20895.
- Cao J, Stieger B, Meier PJ, Vore M. 2002. Expression of rat hepatic multidrug resistance-associated proteins and organic anion transporters in pregnancy. *American Journal of Physiology Gastrointestinal and Liver Physiology* 283:G757–G766 DOI 10.1152/ajpgi.00126.2002.
- Carlin A, Alfirevic Z. 2008. Physiological changes of pregnancy and monitoring. Best Practice & research. *Clinical Obstetrics & Gynaecology* 22:801–823 DOI 10.1016/j.bpobgyn.2008.06.005.

- Chiang JY. 2009.** Bile acids: regulation of synthesis. *Journal of Lipid Research* **50**:1955–1966 DOI [10.1194/jlr.R900010-JLR200](https://doi.org/10.1194/jlr.R900010-JLR200).
- Cripps AW, Williams VJ. 1975.** The effect of pregnancy and lactation on food intake, gastrointestinal anatomy and the absorptive capacity of the small intestine in the albino rat. *The British Journal of Nutrition* **33**:17–32 DOI [10.1079/BJN19750005](https://doi.org/10.1079/BJN19750005).
- Cui YJ, Aleksunes LM, Tanaka Y, Goedken MJ, Klaassen CD. 2009.** Compensatory induction of liver efflux transporters in response to ANIT-induced liver injury is impaired in FXR-null mice. *Toxicological Sciences* **110**:47–60 DOI [10.1093/toxsci/kfp094](https://doi.org/10.1093/toxsci/kfp094).
- Csanaky IL, Lu H, Zhang Y, Ogura K, Choudhuri S, Klaassen CD. 2011.** Organic anion-transporting polypeptide 1b2 (Oatp1b2) is important for the hepatic uptake of unconjugated bile acids: studies in Oatp1b2-null mice. *Hepatology* **53**:272–281 DOI [10.1002/hep.23984](https://doi.org/10.1002/hep.23984).
- Diken Z, Usta IM, Nassar AH. 2013.** A clinical approach to intrahepatic cholestasis of pregnancy. *American Journal of Perinatology* Epub ahead of print Jan 28 2013 DOI [10.1055/s-0033-1333673](https://doi.org/10.1055/s-0033-1333673).
- Dixon PH, Weerasekera N, Linton KJ, Donaldson O, Chambers J, Egginton E, Weaver J, Nelson-Piercy C, de Swiet M, Warnes G, Elias E, Higgins CF, Johnston DG, McCarthy MI, Williamson C. 2000.** Heterozygous MDR3 missense mutation associated with intrahepatic cholestasis of pregnancy: evidence for a defect in protein trafficking. *Human Molecular Genetics* **9**:1209–1217 DOI [10.1093/hmg/9.8.1209](https://doi.org/10.1093/hmg/9.8.1209).
- Dixon PH, van Mil SW, Chambers J, Strautnieks S, Thompson RJ, Lammert F, Kubitz R, Keitel V, Glantz A, Mattsson LA, Marschall HU, Molokhia M, Moore GE, Linton KJ, Williamson C. 2009.** Contribution of variant alleles of ABCB11 to susceptibility to intrahepatic cholestasis of pregnancy. *Gut* **58**:537–544 DOI [10.1136/gut.2008.159541](https://doi.org/10.1136/gut.2008.159541).
- Egan N, Bartels A, Khashan AS, Broadhurst DI, Joyce C, O'Mullane J, O'Donoghue K. 2012.** Reference standard for serum bile acids in pregnancy. *BJOG* **119**:493–498 DOI [10.1111/j.1471-0528.2011.03245.x](https://doi.org/10.1111/j.1471-0528.2011.03245.x).
- Floreani A, Caroli D, Lazzari R, Memmo A, Vidali E, Colavito D, D'Arrigo A, Leon A, Romero R, Gervasi M. 2013.** Intrahepatic cholestasis of pregnancy: new insights into its pathogenesis. *The Journal of Maternal-Fetal & Neonatal Medicine* Epub ahead of print Mar 12 2013 DOI [10.3109/14767058.2013.783810](https://doi.org/10.3109/14767058.2013.783810).
- Geenes V, Williamson C. 2009.** Intrahepatic cholestasis of pregnancy. *World Journal of Gastroenterology* **15**:2049–2066 DOI [10.3748/wjg.15.2049](https://doi.org/10.3748/wjg.15.2049).
- Lam P, Soroka CJ, Boyer JL. 2010.** The bile salt export pump: clinical and experimental aspects of genetic and acquired cholestatic liver disease. *Seminars in Liver Diseases* **30**:125–133 DOI [10.1055/s-0030-1253222](https://doi.org/10.1055/s-0030-1253222).
- Lu H, Choudhuri S, Ogura K, Csanaky IL, Lei X, Cheng X, Song PZ, Klaassen CD. 2008.** Characterization of organic anion transporting polypeptide 1b2-null mice: essential role in hepatic uptake/toxicity of phalloidin and microcystin-LR. *Toxicological Sciences* **103**:35–45 DOI [10.1093/toxsci/kfn038](https://doi.org/10.1093/toxsci/kfn038).
- Milona A, Owen BM, Cobbold JF, Willemsen EC, Cox IJ, Boudjelal M, Cairns W, Schoonjans K, Taylor-Robinson SD, Klomp LW, Parker MG, White R, van Mil SW, Williamson C. 2010.** Raised hepatic bile acid concentrations during pregnancy in mice are associated with reduced farnesoid X receptor function. *Hepatology* **52**:1341–1349 DOI [10.1002/hep.23849](https://doi.org/10.1002/hep.23849).
- Papacleovoulou G, Abu-Hayyeh S, Williamson C. 2011.** Nuclear receptor-driven alterations in bile acid and lipid metabolic pathways during gestation. *Biochimica et Biophysica Acta* **1812**:879–887 DOI [10.1016/j.bbadis.2010.11.001](https://doi.org/10.1016/j.bbadis.2010.11.001).

- Pascual MJ, Serrano MA, El-Mir MY, Macias RI, Jiménez F, Marin JGJ. 2002.** Relationship between asymptomatic hypercholanemia of pregnancy and progesterone metabolism. *Clinical Science* 102:587–593 DOI 10.1042/CS20010258.
- Shuster DL, Bammler TK, Beyer RP, Macdonald JW, Tsai JM, Farin FM, Hebert MF, Thummel KE, Mao Q. 2013.** Gestational age-dependent changes in gene expression of metabolic enzymes and transporters in pregnant mice. *Drug Metabolism and Disposition* 41:332–342 DOI 10.1124/dmd.112.049718.
- Vallejo M, Briz O, Serrano MA, Monte MJ, Marin JGJ. 2006.** Potential role of trans-inhibition of the bile salt export pump by progesterone metabolites in the etiopathogenesis of intrahepatic cholestasis of pregnancy. *Journal of Hepatology* 44:1150–1157 DOI 10.1016/j.jhep.2005.09.017.
- van Mil SWC, Milona A, Dixon PH, Mullenbach R, Geenes VL, Chambers J, Shevchuk V, Moore GE, Lammert F, Glantz AG, Mattsson LA, Whittaker J, Parker MG, White R, Williamson C. 2007.** Functional variants of the central bile acid sensor FXR identified in intrahepatic cholestasis of pregnancy. *Gastroenterology* 133:507–516 DOI 10.1053/j.gastro.2007.05.015.
- Vernon RG, Denis RG, Sorensen A, Williams G. 2002.** Leptin and the adaptations of lactation in rodents and ruminants. *Hormone Metabolic Research* 34:678–685 DOI 10.1055/s-2002-38258.
- Wooton-Kee CR, Cohen DE, Vore M. 2008.** Increased cholesterol 7 α -hydroxylase expression and size of the bile acid pool in the lactating rat. *American Journal of Physiology Gastrointestinal and Liver Physiology* 294:G1009–G1016 DOI 10.1152/ajpgi.00017.2008.
- Wooton-Kee CR, Coy DJ, Athipozhy AT, Zhao T, Jones BR, Vore M. 2010.** Mechanisms for increased expression of cholesterol 7 α -hydroxylase (Cyp7a1) in lactating rats. *Hepatology* 51:277–285 DOI 10.1002/hep.23289.
- Zhang Y, Csanaky IL, Cheng X, Lehman-McKeeman LD, Klaassen CD. 2012.** Organic anion transporting polypeptide 1a1 null mice are sensitive to cholestatic liver injury. *Toxicological Sciences* 127:451–462 DOI 10.1093/toxsci/kfs123.