

Original article

Identification of potential mediators of the relationship between body mass index and colorectal cancer: a Mendelian randomization analysis

Emmanouil Bouras ^{1,*}, Dipender Gill ^{2,3}, Verena Zuber ³, Neil Murphy ⁴, Niki Dimou,⁴
Krasimira Aleksandrova ^{5,6}, Sarah J Lewis,^{7,8} Richard M Martin,^{7,8,9} James Yarmolinsky,^{7,8}
Demetrius Albanes,¹⁰ Hermann Brenner,^{11,12,13} Sergi Castellví-Bel,¹⁴ Andrew T Chan,^{15,16,17,18,19,20}
Iona Cheng ²¹, Stephen Gruber,²² Bethany Van Guelpen ^{23,24}, Christopher I Li,²⁵ Loic Le Marchand,²⁶
Polly A Newcomb,^{25,27} Shuji Ogino,^{18,19,28,29} Andrew Pellatt,³⁰ Stephanie L Schmit,^{31,32} Alicja Wolk ³³,
Anna H Wu,³⁴ Ulrike Peters,^{25,27} Marc J Gunter^{3,4} and Konstantinos K Tsilidis^{1,3}

¹Department of Hygiene and Epidemiology, University of Ioannina School of Medicine, Ioannina, Greece, ²Chief Scientific Advisor Office, Research and Early Development, Novo Nordisk, Copenhagen, Denmark, ³Department of Epidemiology and Biostatistics, Imperial College London, School of Public Health, London, UK, ⁴Nutrition and Metabolism Branch, International Agency for Research on Cancer, Lyon, France, ⁵Faculty of Human and Health Sciences, University of Bremen, Bremen, Germany, ⁶Department Epidemiological Methods and Etiological Research, Leibniz Institute for Prevention Research and Epidemiology, Bremen, Germany, ⁷Department of Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK, ⁸MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK, ⁹NIHR Bristol Biomedical Research Centre, University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol, ¹⁰Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA, ¹¹Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany, ¹²Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany, ¹³German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany, ¹⁴Department of Gastroenterology, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBERehd), Hospital Clínic, Barcelona, Spain, ¹⁵Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA, ¹⁶Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA, ¹⁷Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA, ¹⁸Broad Institute of Harvard and MIT, Cambridge, MA, USA, ¹⁹Department of Epidemiology, Harvard TH Chan School of Public Health, Harvard University, Boston, MA, USA, ²⁰Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public Health, Harvard University, Boston, MA, USA, ²¹Department of Epidemiology and Biostatistics, University of California-San Francisco, San Francisco, CA, USA, ²²Department of Medical Oncology & Therapeutics Research and Center for Precision Medicine, City of Hope National Medical Center, Duarte, CA, USA, ²³Department of Radiation Sciences, Oncology Unit, Umeå University, Umeå, Sweden, ²⁴Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden, ²⁵Public Health Sciences Division, Fred Hutchinson Cancer Center, Seattle, Washington, USA, ²⁶University of Hawaii Cancer Center, Honolulu, HI, USA, ²⁷Department of Epidemiology, University of Washington, Seattle, WA, USA, ²⁸Program in MPE Molecular Pathological Epidemiology, Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA, ²⁹Department of Oncologic Pathology, Dana-Farber Cancer Institute, Boston, MA, USA, ³⁰Department of Medicine, University of Utah, Salt Lake City, UT, USA, ³¹Genomic Medicine Institute, Cleveland Clinic, Cleveland, OH, USA, ³²Population and Cancer Prevention Program, Case Comprehensive Cancer Center, Cleveland, OH, USA, ³³Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden and ³⁴University of Southern California, Preventative Medicine, Los Angeles, CA, USA

*Corresponding author. Faculty of Medicine, Department of Hygiene and Epidemiology, University of Ioannina campus, P.C. 45110, Ioannina, Greece.
E-mail: ebouras@uoi.gr

Abstract

Background: Colorectal cancer (CRC) is the third-most-common cancer worldwide and its rates are increasing. Elevated body mass index (BMI) is an established risk factor for CRC, although the molecular mechanisms behind this association remain unclear. Using the Mendelian randomization (MR) framework, we aimed to investigate the mediating effects of putative biomarkers and other CRC risk factors in the association between BMI and CRC.

Methods: We selected as mediators biomarkers of established cancer-related mechanisms and other CRC risk factors for which a plausible association with obesity exists, such as inflammatory biomarkers, glucose homeostasis traits, lipids, adipokines, insulin-like growth factor 1 (IGF1), sex hormones, 25-hydroxy-vitamin D, smoking, physical activity (PA) and alcohol consumption. We used inverse-variance weighted MR in the main univariable analyses and performed sensitivity analyses (weighted-median, MR-Egger, Contamination Mixture). We used multivariable MR for the mediation analyses.

Received: 15 September 2023. Editorial Decision: 25 March 2024. Accepted: 15 April 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of the International Epidemiological Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Results: Genetically predicted BMI was positively associated with CRC risk [odds ratio per SD (5 kg/m^2) = 1.17, 95% CI: 1.08–1.24, P -value = 1.4×10^{-9}] and robustly associated with nearly all potential mediators. Genetically predicted IGF1, fasting insulin, low-density lipoprotein cholesterol, smoking, PA and alcohol were associated with CRC risk. Evidence for attenuation was found for IGF1 [explained 7% (95% CI: 2–13%) of the association], smoking (31%, 4–57%) and PA (7%, 2–11%). There was little evidence for pleiotropy, although smoking was bidirectionally associated with BMI and instruments were weak for PA.

Conclusions: The effect of BMI on CRC risk is possibly partly mediated through plasma IGF1, whereas the attenuation of the BMI–CRC association by smoking and PA may reflect confounding and shared underlying mechanisms rather than mediation.

Keywords: Body mass index, BMI, obesity, mediation analysis, colorectal cancer, CRC, Mendelian randomization.

Key Messages

- Using the Mendelian randomization framework, we investigated the plausibility of putative biomarkers and other colorectal cancer (CRC) risk factors as potential mediators in the association between body mass index (BMI) and CRC.
- The effect of BMI on CRC risk is possibly partly mediated through plasma insulin-like growth factor 1 (IGF1), whereas the attenuation observed by smoking and physical activity may reflect shared underlying mechanisms and confounding rather than mediation.
- Understanding the pathophysiological pathways underlying the association between adiposity and risk of CRC may facilitate the development of targeted interventions for susceptible individuals.

Introduction

One in 10 incident cancers diagnosed in 2020 was colorectal cancer (CRC) and its global rates are increasing.¹ This increase may partly reflect changes in dietary and lifestyle habits, and a rise in the prevalence of excess body weight in the population, commonly captured using body mass index (BMI). Increased BMI is considered an established risk factor for CRC and 12 other cancers.² Based on a World Cancer Research Fund dose–response meta-analysis, a 5-kg/m^2 increase in BMI is associated with a 5% increase in CRC risk.³ Understanding the pathophysiological pathways behind the link between excess adiposity and CRC may inform the development of targeted interventions for susceptible individuals.

Epidemiological and experimental studies suggest that increased adiposity may contribute to elevated CRC risk through effects on inflammation, sex-steroid hormone and adipokine concentrations, hyperglycaemia and hyperinsulinaemia, and other metabolic intermediates.^{4–8} Even though no clear consensus on which mediators explain the obesity–CRC relationship has been reached, specific patterns of mediators have emerged through observational studies. For instance, metabolic intermediates related to glucose metabolism and leptin bioavailability appear to be responsible for mediating the effects of adiposity on CRC risk more strongly than inflammatory- or sex-hormone-related markers.^{5–9}

Most studies to date have used traditional methods for studying mediation in observational settings based on the model proposed by Baron and Kenny, which relies on strong and untestable assumptions (these include absence of unmeasured confounding in the associations between exposure, mediator and outcome, and absence of exposure–mediator interaction).^{4–6,10,11} Furthermore, most of these observational studies do not cover a wide range of plausible molecular pathways, typically include a small number of participants (<1000 cases in most of the studies) and are prone to biases (such as confounding).^{4–10}

The Mendelian randomization (MR) framework that could overcome some of the above limitations can be used in a multivariable (MV) context, using summary genetic association estimates, allowing a mediation analysis to be performed.^{12,13} In the present study, we use the MR framework to: (i) investigate the plausibility of putative biomarkers and

other CRC risk factors as potential mediators in the association between BMI (as a measure of general adiposity) and CRC risk, and (ii) to estimate the proportion of the mediated effect explained by plausible mediators.

Methods

Source of genetic instruments

We selected as potential mediators biomarkers of established cancer-related mechanisms for which a plausible association with BMI exists based on epidemiological and experimental evidence.¹⁴ These potential mediators included inflammation-related biomarkers [C-reactive protein (CRP) and interleukin-6 (IL6)], metabolism-related biomarkers [glucose homeostasis, lipids, adipokines, insulin-like growth factor 1 (IGF1)], sex hormones and 25-hydroxy-vitamin D [25 (OH)D].^{3,14} We also included CRC risk factors that typically act as confounders, such as smoking, physical activity (PA), alcohol consumption and type 2 diabetes, in the analysis, to examine the possibility of mediation vs confounding (Supplementary Figure S1, available as [Supplementary data at IJE online](#)).¹⁵

Details of the genome-wide association studies (GWASs) from which we selected the genetic instruments are presented in the [Supplementary Methods](#), [Supplementary Table S1](#) (available as [Supplementary data at IJE online](#)) and in the originally published studies.

Summary genetic association estimates for CRC, as well as cancers of the colon, proximal and distal colon and rectum, and in men and women separately, were obtained from a GWAS meta-analysis of the Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO), Colorectal Transdisciplinary Study (CORECT) and Colon Cancer Family Registry (CCFR)¹⁶ ([Supplementary Table S2](#), available as [Supplementary data at IJE online](#)).

Instrument selection and MR analyses

In the univariable MR analyses, we selected as instruments single-nucleotide polymorphisms (SNPs) that were associated with the exposure of interest (BMI and the putative mediators) at genome-wide significance ($P < 5 \times 10^{-8}$), allowing a weak linkage disequilibrium ($r^2 < 0.001$) ([Supplementary Table S3](#), available as [Supplementary data at IJE online](#)). The random-




Figure 1. To obtain valid causal estimates for mediation, the three typical Mendelian randomization (MR) assumptions should be satisfied in the context of multivariable MR. These are: (i) relevance—the single-nucleotide polymorphisms (SNPs) (G) should be strongly associated with body mass index (BMI) [given the putative mediator(s) included in the model]; (ii) exchangeability—the SNPs (G) should be independent of all confounders (U) of any of the exposures (in the model) and colorectal cancer (CRC) and (iii) exclusion restriction—there should be no other path (P) through which the SNPs affect CRC but via BMI [or the putative mediator(s) in the model]

effects inverse-variance weighted (IVW) model was used as the main analysis and the MR-Egger, Contamination Mixture and weighted-median MR methods were used as sensitivity analyses.^{17–19} These analyses were used to investigate the association of genetically predicted BMI on CRC, mediators on CRC and BMI on mediators bidirectionally. Bidirectional analyses between BMI and mediators were performed to examine confounding vs mediation and to gain a deeper insight into the main associations. Mediators for which a ‘robust’ association was found in the univariable analysis with BMI (BMI to mediators) and CRC (mediators to CRC) were included in the subsequent mediation analyses ([Supplementary Methods](#), available as [Supplementary data](#) at *IJE* online). For discrepant BMI-to-mediator and mediator-to-CRC associations, clustered heterogeneity in the BMI-to-mediator association was assessed using an expectation-maximization-based model fitting, to identify potentially distinct pathways that would explain the observed discrepancy.²⁰

In the multivariable MR (MVMR) analyses, we used a combined genetic instrument, selecting uncorrelated variants ($r^2 < 0.001$) associated with any of the exposures in the model. The genetic association estimates for CRC were regressed on the genetic association estimates for BMI and mediator(s), weighted for the precision of the genetic association estimates for CRC and with the intercept fixed to zero.²¹ Similarly, we performed MV MR-Egger analyses, allowing an intercept, as an indicator of directional pleiotropy.

Mediation analysis

Univariable MR analyses were used to estimate the total effect of genetically predicted BMI on CRC risk. MVMR analyses were used to assess mediation and correction for horizontal pleiotropy. Using MVMR models, adjusting for one mediator at a time, the direct effects of BMI on CRC risk was estimated (i.e. the effect of BMI on CRC through

pathways other than the mediator in the model). The indirect effect of BMI on outcome was estimated using the difference in coefficients method, which was expressed as the proportion attenuated.^{12,13}

To investigate the validity of the three core MR assumptions ([Figure 1](#)), in the context of MVMR: (i) relevance [the SNPs should be strongly associated with BMI given the mediator(s) included in the model], (ii) exchangeability (the SNPs should be independent of confounders of any of the exposures in the model and CRC) and (iii) exclusion restriction [there should be no other path through which the SNPs affect CRC but via BMI, or the mediator(s) in the model], several approaches were used ([Supplementary Methods](#), available as [Supplementary data](#) at *IJE* online).¹³ Briefly, we used the conditional F-statistic (F_{cond}) as an indicator of instrument strength.²² For mediators with conditionally weak instruments ($F_{\text{cond}} < 10$), we reran the MVMR and mediation analyses using alternative instruments for BMI as a means of enhancing the conditional strength of the mediator(s) in the model. Sensitivity analyses were performed excluding genetic instruments significantly associated with both BMI and mediator in the MV model, as a means of distinguishing the mediation phenomenon from horizontal pleiotropy.¹³ The Phenoscanner database was used to explore the previously reported associations of the selected genetic instruments and identify potentially pleiotropic pathways.

In secondary analyses, we investigated the mediation on the subtypes of CRC, and among men and women separately. All analyses were performed using R (v.4.1.1).²³

Results

Evaluating the effect of BMI on CRC (total effects)

There was consistent evidence across the IVW and sensitivity analyses that higher genetically predicted BMI increased




Figure 2. Mendelian randomization (MR) mediation analysis of genetically predicted body mass index (BMI) on colorectal cancer (CRC) risk. The forest plot (left, top and middle) presents the MR inverse-variance weighted (IVW) estimate (OR, 95% CI) of BMI on CRC risk in the MVMR model, adjusted for one putative mediator at a time (direct effects). Where there was a robust BMI-to-putative-mediator and mediator-to-CRC-risk association, the percent attenuation and 95% CI were estimated and are presented in labels. The forest plot on the right side displays the marginal associations of BMI on putative mediators (the 95% CI of the MR-IVW using a thick grey line) and mediators on CRC risk (the MR-IVW beta and 95% CI). The check marks in the first column ('Robustness') indicate robust marginal associations [i.e. significant associations in the IVW analysis ($P < 0.05$) that were qualitatively consistent in sensitivity analyses], in the second column ['Instrument strength (MVMR)'] indicate conditional F-statistics of >10 [for both predicted body mass index (BMI) and the putative mediator in the model] and in the third column ('Egger P -int. >0.1 ') indicate that the P -value of the intercept in the MV MR-Egger models was >0.1 . The forest plot at the bottom displays the total effects (OR, 95% CI) of BMI on CRC risk in univariable MR-IVW models

overall CRC risk [odds ratio (OR) per SD (4.7 kg/m^2) higher BMI: 1.16, 95% CI: 1.08–1.24, $P = 1.4 \times 10^{-6}$] (Figure 2 and Supplementary Table S4, available as Supplementary data at *IJE* online). A positive effect of BMI remained in the analyses by CRC subtypes and by sex. It was marginally stronger for colon vs rectal cancer ($P_{\text{het}}=0.07$) but similar for proximal vs distal colon cancer ($P_{\text{het}}=0.32$) and by sex ($P_{\text{het}}=0.58$) (Figure 3 and Supplementary Table S4, available as Supplementary data at *IJE* online).

Evaluating the effect of mediators on CRC

There was consistent evidence across the IVW and sensitivity analyses that CRC risk was associated with genetically predicted IGF1 (OR per 1 unit in rank-based inverse normal transformed IGF1, which approximates to 5.7 nmol/L : 1.08, 95% CI: 1.03–1.15), fasting insulin (OR per 1 unit in natural log-transformed pmol/L: 1.58, 95% CI: 1.13–2.22), low-density lipoprotein (LDL) cholesterol (OR per 1 unit increase in rank-based inverse normal transformed LDL cholesterol: 1.08, 95% CI: 1.01–1.16), smoking (OR per 1 unit of lifetime smoking index, which approximates to 20 cigarettes/day for 10 years: 1.24, 95% CI: 1.01–1.53), PA {OR per SD [0.08 m/s^2], which approximates to 3 units of metabolic equivalent task score (METs)-h/day]: 0.70, 95% CI: 0.52–0.93} and alcohol drinking [OR per SD (~ 5.6 drinks/week): 1.25, 95% CI: 1.00–1.57]. Furthermore, T2DM was positively associated with CRC (OR per 1-unit increase in log odds for T2DM liability: 1.04, 95% CI: 1.00–1.07), albeit the association was not consistent in sensitivity analyses (Supplementary Table S5, available as Supplementary data at *IJE* online).

Evaluating the effect of BMI on mediators

Using MR-IVW analyses, we found that genetically predicted BMI was positively associated with HbA1c, fasting glucose and insulin (using the instrument unadjusted for BMI), T2DM liability, triglycerides, bioavailable testosterone, oestradiol among men, IL6, CRP, leptin and smoking, and inversely associated with IGF1, fasting glucose and insulin, and two-h glucose (2hGlu) (using instruments adjusted for BMI, likely due to collider bias), HDL-C, total cholesterol, total testosterone, sex hormone binding globulin (SHBG), 25(OH) D, adiponectin, PA and alcohol drinking. The above associations were largely consistent in the sex-stratified analyses, except for the associations of BMI with total testosterone and bioavailable testosterone, which were inverse in men and positive in women (Supplementary Table S6, available as Supplementary data at *IJE* online).

Evaluating the mediator-to-mediator associations

Several mediator-to-mediator associations, consistent in sensitivity analyses, were found in the univariable MR analyses. Evidence of bidirectionality with BMI was found for fasting insulin and 2hGlu (using instruments adjusted for BMI), total cholesterol and smoking (Supplementary Figure S2 and Supplementary Table S7, available as Supplementary data at *IJE* online).

MR mediation analysis

IGF1 was found to be a plausible mediator (7% of total effects mediated, 95% CI: 2–13%, $P = 0.01$) in the association of genetically predicted BMI with overall CRC risk, with




Figure 3. Mendelian randomization (MR) mediation of genetically predicted body mass index (BMI) on colorectal cancer (CRC) subtypes. The forest plot shows the OR (95% CI) per SD of genetically predicted BMI (5 kg/m²), on CRC subtype in the MR inverse-variance weighted (IVW) models (total effects) and in sensitivity analyses. Where there was a robust BMI-to-putative-mediator and mediator-to-CRC-risk association, the percent attenuation and 95% CI were estimated and are presented in labels

little evidence of residual directional pleiotropy or weak instrument bias ($F_{\text{cond}}=46$) (Figure 2 and Supplementary Table S8, available as Supplementary data at *IJE* online).

Additionally, attenuation of the effects of BMI on CRC risk was found for smoking (31%, 95% CI: 4–57%, $P=0.02$), PA (7%, 2–11%, $P=2 \times 10^{-3}$) and alcohol drinking (1%, 0–2%, $P=0.05$). However, the conditional F-statistics were low, indicating weak instruments (F_{cond} range: 2–7) (Figure 2 and Supplementary Table S8, available as Supplementary data at *IJE* online). When we used alternative instruments for BMI, the results did not change materially (Supplementary Table S9, available as Supplementary data at *IJE* online).

In sensitivity analysis excluding genetic instruments significantly associated with both BMI and the mediator in the model, the results did not change materially (Supplementary Table S10 and Supplementary Figure S3, available as Supplementary data at *IJE* online).

We found no major pleiotropic pathways (other than those included as mediators), apart from a few SNPs previously associated with inflammatory bowel disease (Supplementary Tables S11 and S12, available as Supplementary data at *IJE* online) that are unlikely to have substantially influenced the mediation estimation. Discrepant associations were found for IGF1 and alcohol, with evidence of clustered heterogeneity in the association of BMI with both mediators, supporting the presence of distinct pathways, potentially inferring positive and inverse effects (Supplementary Table S13 and Supplementary Figure S4, available as Supplementary data at *IJE* online).

In the secondary analyses performed by subtype and sex, evidence for mediation was found for IGF1, total cholesterol and smoking in relation to colon cancer; IGF1, total cholesterol and PA for distal colon cancer; and smoking for CRC in men (Figure 3 and Supplementary Table S8, available as Supplementary data at *IJE* online). However, most of these analyses were based on smaller samples and the power was limited.

Discussion

Using MR analyses, we investigated the mediating effects of several biologically plausible intermediates in the association

between BMI and CRC. Our results suggest that some of the effects of genetically predicted BMI on CRC and subtypes is mediated through IGF1. Fasting insulin was also a plausible mediator, robustly associated with BMI and CRC risk, although no mediating effect was found, possibly due to the weak instrument for insulin. Attenuation of the effects of BMI on CRC risk was observed by smoking and PA, although these exposures typically act as confounders or common underlying factors, hence adjustment via MVMR likely provides a reflection of adjustment for horizontal pleiotropy. The bidirectional association of BMI with (smoking) and the shared carcinogenic mechanisms for BMI and PA strengthen the notion that these factors are less likely to act as mediators.

We found that higher genetically predicted BMI was associated with decreased IGF1 concentrations and that IGF1 was positively associated with CRC risk. Such a discrepancy in the direction of the marginal associations might suggest that BMI acts to increase CRC risk via IGF1, via complex modes of effect that potentially involve other intermediates (such as insulin).²⁴ Such a complexity is underscored by clusters of variants influencing exposures and mediators in diverse ways, potentially targeting distinct causal effect parameters, and is partly addressed by using the composite instruments. Nevertheless, such a complexity may not be sufficiently represented by individual biomarkers, but rather require patterns of biomarkers, which, however, were not captured in our analyses, partly due to the limited power that exists with multiple mediators within the MVMR framework. Though not in universal agreement, epidemiological studies support an inverse association of BMI with circulating IGF1 levels.^{25,26} Such an inverse association might be explained by the presence of a negative feedback loop among individuals with a prolonged state of obesity, whereby reductions in IGF binding protein 1 (IGFBP1) and IGF binding protein 2 (IGFBP2), as a result of obesity-associated hyperinsulinemia, may lead to increased negative feedback by free IGF1 (unbound to IGFBPs) on pituitary growth hormone secretion (a major regulator of IGF1 synthesis in the liver).²⁷ Furthermore, time-varying effects of BMI on the

concentrations of IGF1 have been previously reported, which could also explain the observed discrepancy in the direction of the BMI-to-IGF1 association. For instance, a cross-sectional study ($n=4241$ participants from the Long Life Family Study) found evidence of an inverse association among individuals in the lowest age groups (20–66 years old) and a positive association among older individuals (>87 years old).²⁸ Prospective studies have shown that increased concentrations of circulating IGF1 are positively associated with CRC risk—an association that has been replicated in previous MR analyses and is concordant with our results.^{29,30} In support of the observed positive association, a large body of mechanistic evidence has shown that IGF1 has proliferative and anti-apoptotic effects and activates cellular pathways linked to tumorigenesis such as the PI3K-mTOR and RAS pathways.^{31–33} A recent observational study investigating the contribution of metabolic mediators in the association between adult weight gain and CRC found no evidence of mediation for IGF1, but the study was small, including 266 colon and 186 rectal cancer cases.⁵

A relatively strong attenuation of the effects of BMI on CRC risk was observed for smoking. Causal associations of BMI on smoking and of smoking on CRC have been previously reported and confirmed in our analysis.^{34,35} Purported mechanisms through which BMI is linked to smoking are complex and most likely have behavioural and metabolic implications.³⁶ Higher BMI and tobacco smoking genetically share a biological basis for addictive behaviours, such as nicotine addiction and higher energy intake.³⁴ Higher BMI has been related to lower socio-economic status, income and educational attainment, and lower educational attainment may lead to increased smoking, whereas obesity may increase nicotine dependence and affect smoking intensity.^{37,38} Tobacco smoking as a source of a multitude of carcinogenic compounds, likely acting via their immunosuppressive effects, which can play an important tumour-promoting role in the serrated neoplasia pathway (characterized by high-level microsatellite instability and vigorous immune response), has been well established as a CRC risk factor.³⁹ We found a bidirectional (positive) association of genetic predisposition to higher lifetime amount of smoking on BMI, which supports the hypothesis that smoking may act as a confounder. Such a positive association (of smoking on BMI) finds little support in previously published observational or MR analyses.^{36,40}

There is ample epidemiological evidence suggesting that obesity in childhood is associated with reduced PA, but research on adult populations is largely focused on the effectiveness of PA as a means of maintaining a healthy weight.^{41,42} Previous observational and MR analyses provided evidence to support an association between BMI and PA levels, whereas a bidirectional association with BMI has also been reported.^{43–45} We found an inverse association between genetically predicted BMI and device-measured PA, but not vice-versa, which could be attributed to the relatively weak instrument for device-measured PA. Furthermore, higher genetically predicted PA was associated with a 30% decreased CRC risk, in line with previous studies.⁴³ Unhealthy lifestyle behaviours (such as sedentary behaviour, time spent watching television) that accompany obesity might impact an individual's levels of PA.⁴⁶ On the other hand, increased PA may act to reduce CRC risk via multiple biological mechanisms that include improvement in insulin sensitivity and inflammation, digestion stimulation and

transit time stimulation in the intestine.⁴⁷ The fact that BMI and PA share multiple carcinogenic mechanisms questions the notion that PA in fact acts as a mediator in the association between BMI and CRC risk.

In this large-scale MR analysis, we included a wide panel of biologically plausible mediators covering different mechanisms and the largest samples available to date for the exposure, mediators and outcomes. We acknowledge that our study has several limitations that should be considered when interpreting the results. The present analysis is based on BMI as a measure of general adiposity, which does not capture the type of adiposity and the proportion relative to lean mass; nevertheless, BMI is highly correlated with objective measures of adiposity and, considering its wide use in the epidemiological setting (because it is relatively easy and straightforward to measure), large GWAS sample sizes have been achieved and hence power is increased compared with other measures of adiposity (such as radiology-based measures).⁴⁸ Several potentially important pathways or pathway-specific traits, such as detailed inflammation-related intermediates or adipokines, were not included (or inadequately represented in our panel of mediators) due to absence of valid instruments. This also limited our ability to identify patterns of biomarkers, which could elucidate the complex pathways in which higher BMI is associated with increased CRC risk. The total effects of BMI on CRC risk were relatively small, which might have caused the precision in the estimate of the indirect effects and the proportion attenuated to inflate. There was no distinction between childhood and adulthood BMI; however, a recent analysis demonstrated that most of the effects of the childhood BMI on CRC are mediated through adulthood BMI.⁴⁹ Time-varying effects of BMI on some biomarkers (such as IGF1) have been reported, which could be a threat to the validity of the mediating estimates and should be explored in future MR mediation analyses using individual-level data.²⁸ Measurement errors may not be of utmost concern in the present analysis, although we cannot rule out the possibility that intra-individual variation for some of the included traits may have affected the validity of the estimates to some extent.^{50,51} Finally, the analysis was based on individuals of European descent and there was limited power to expand the mediation analysis to populations of different racial and ethnic groups.

Conclusions

In conclusion, the effects of BMI on CRC risk may be partly mediated through IGF1. Future studies using stronger instruments for plausible intermediates are warranted to more comprehensively unveil the mechanisms linking adiposity to CRC risk. The complex nature of the causal associations of the intermediates highlights the need to consider the major risk factors together in the development of CRC prevention strategies.

Disclaimer

Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer/World Health Organization.

Ethics approval

Not applicable.

Data availability

All data used in this work are presented in the Additional files that accompany the manuscript and are available in the original publications.

Supplementary data

Supplementary data are available at *IJE* online.

Author contributions

E.B.: Data curation, Formal analysis, Investigation, Visualization, Writing—Original Draft, Writing—Review & Editing; D.G., V.Z.: Investigation, Methodology, Resources, Validation, Writing—Review & Editing; N.M., N.D., K.A., S.L., R.M., J.Y., D.A., H.B., S.C.B., A.C., I.C., S.G., B.v.G., C.L., L.L.M., P.N., S.O., A.P., S.S., A.W., A.H.W.: Methodology, Writing—Review & Editing; U.P., M.G.: Methodology, Supervision, Writing—Review & Editing; K. T.: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing—Review & Editing. All authors critically revised the manuscript for intellectual content. All authors read and approved the final manuscript.

Funding

This study was funded by Cancer Research UK (PPRCPT \100005) to K.K.T. R.M.M. is a National Institute for Health Research Senior Investigator (NIHR202411). R.M.M. and S. J.L. are supported by a Cancer Research UK 25 (C18281/A29019) programme grant (the Integrative Cancer Epidemiology Programme). R.M.M. is also supported by the NIHR Bristol Biomedical Research Centre, which is funded by the NIHR (BRC-1215-20011) and is a partnership between University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. R.M.M. and S.J.L. are affiliated with the Medical Research Council Integrative Epidemiology Unit at the University of Bristol, which is supported by the Medical Research Council (MC_UU_00011/1, MC_UU_00011/3, MC_UU_00011/6 and MC_UU_00011/4) and the University of Bristol. J.Y. is supported by a Cancer Research UK Population Research Postdoctoral Fellowship (C68933/A28534). Department of Health and Social Care disclaimer: The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. GECCO funding & acknowledgements are described in detail in the [Supplementary notes](#).

Conflict of interest

I have read the Journal's policy and the authors of this manuscript have the following competing interests: D.G. is employed part-time by Novo Nordisk. The remaining authors have no competing interests to declare.

References

1. Ferlay J, Colombet M, Soerjomataram I *et al.* Cancer statistics for the year 2020: an overview. *Int J Cancer* 2021;**149**:778–89.
2. Kyrgiou M, Kalliala I, Markozannes G *et al.* Adiposity and cancer at major anatomical sites: umbrella review of the literature. *BMJ* 2017;**356**:j477.
3. World Cancer Research Fund/American Institute for Cancer Research. *Continuous Update Project Expert Report 2018. Diet, Nutrition, Physical Activity and Colorectal Cancer*. <https://www.wcrf.org/diet-activity-and-cancer/> (5 March 2023, date last accessed).
4. Aleksandrova K, Drogan D, Boeing H *et al.* Adiposity, mediating biomarkers and risk of colon cancer in the European prospective investigation into cancer and nutrition study. *Int J Cancer* 2014;**134**:612–21.
5. Aleksandrova K, Schlesinger S, Fedirko V *et al.* Metabolic mediators of the association between adult weight gain and colorectal cancer: data from the European Prospective Investigation into Cancer and Nutrition (EPIC) Cohort. *Am J Epidemiol* 2017;**185**:751–64.
6. Ho GY, Wang T, Gunter MJ *et al.* Adipokines linking obesity with colorectal cancer risk in postmenopausal women. *Cancer Res* 2012;**72**:3029–37.
7. Petimar J, Tabung FK, Valeri L *et al.* Mediation of associations between adiposity and colorectal cancer risk by inflammatory and metabolic biomarkers. *Int J Cancer* 2019;**144**:2945–53.
8. Dashti SG, Viallon V, Simpson JA *et al.* Explaining the link between adiposity and colorectal cancer risk in men and postmenopausal women in the UK Biobank: A sequential causal mediation analysis. *Int J Cancer* 2020;**147**:1881–94.
9. Fritz J, Bjørge T, Nagel G *et al.* The triglyceride-glucose index as a measure of insulin resistance and risk of obesity-related cancers. *Int J Epidemiol* 2020;**49**:193–204.
10. Aleksandrova K, Boeing H, Jenab M *et al.* Leptin and soluble leptin receptor in risk of colorectal cancer in the European Prospective Investigation into Cancer and Nutrition cohort. *Cancer Res* 2012;**72**:5328–37.
11. VanderWeele TJ. Mediation analysis: a practitioner's guide. *Annu Rev Public Health* 2016;**37**:17–32.
12. Burgess S, Thompson DJ, Rees JMB, Day FR, Perry JR, Ong KK. Dissecting causal pathways using Mendelian randomization with summarized genetic data: application to age at menarche and risk of breast cancer. *Genetics* 2017;**207**:481–7.
13. Carter AR, Sanderson E, Hammerton G *et al.* Mendelian randomisation for mediation analysis: current methods and challenges for implementation. *Eur J Epidemiol* 2021;**36**:465–78.
14. World Cancer Research Fund/American Institute for Cancer Research. *Continuous Update Project Expert Report 2018. The Cancer Process*. <https://www.wcrf.org/diet-activity-and-cancer/> (5 March 2023, date last accessed).
15. MacKinnon DP, Krull JL, Lockwood CM. Equivalence of the mediation, confounding and suppression effect. *Prev Sci* 2000;**1**:173–81.
16. Huyghe JR, Bien SA, Harrison TA *et al.* Discovery of common and rare genetic risk variants for colorectal cancer. *Nat Genet* 2019;**51**:76–87.
17. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* 2013;**37**:658–65.
18. Burgess S, Foley CN, Allara E, Staley JR, Howson JMM. A robust and efficient method for Mendelian randomization with hundreds of genetic variants. *Nat Commun* 2020;**11**:376.
19. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in mendelian randomization with some invalid instruments using a weighted median estimator. *Genet Epidemiol* 2016;**40**:304–14.
20. Foley CN, Mason AM, Kirk PDW, Burgess S. MR-Clust: clustering of genetic variants in Mendelian randomization with similar causal estimates. *Bioinformatics* 2020;**37**:531–41.

21. Burgess S, Thompson SG. Multivariable Mendelian randomization: the use of pleiotropic genetic variants to estimate causal effects. *Am J Epidemiol* 2015;**181**:251–60.
22. Sanderson E, Spiller W, Bowden J. Testing and correcting for weak and pleiotropic instruments in two-sample multivariable Mendelian randomization. *Stat Med* 2021;**40**:5434–52.
23. R Development Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing, 2020.
24. Rajpathak SN, He M, Sun Q *et al.* Insulin-like growth factor axis and risk of type 2 diabetes in women. *Diabetes* 2012;**61**:2248–54.
25. Tin Tin S, Reeves GK, Key TJ. Body size and composition, physical activity and sedentary time in relation to endogenous hormones in premenopausal and postmenopausal women: Findings from the UK Biobank. *Int J Cancer* 2020;**147**:2101–15.
26. Faupel-Badger JM, Berrigan D, Ballard-Barbash R, Potischman N. Anthropometric correlates of insulin-like growth factor 1 (IGF-1) and IGF binding protein-3 (IGFBP-3) levels by race/ethnicity and gender. *Ann Epidemiol* 2009;**19**:841–9.
27. Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 2004;**4**:579–91.
28. Sherlala RA, Kammerer CM, Kuipers AL *et al.* Relationship between serum IGF-1 and BMI differs by age. *J Gerontol A Biol Sci Med Sci* 2021;**76**:1303–8.
29. Rinaldi S, Cleveland R, Norat T *et al.* Serum levels of IGF-I, IGFBP-3 and colorectal cancer risk: results from the EPIC cohort, plus a meta-analysis of prospective studies. *Int J Cancer* 2010;**126**:1702–15.
30. Murphy N, Carreras-Torres R, Song M *et al.* Circulating levels of insulin-like growth factor 1 and insulin-like growth factor binding protein 3 associate with risk of colorectal cancer based on serologic and mendelian randomization analyses. *Gastroenterology* 2020;**158**:1300–12.e20.
31. Samani AA, Yakar S, LeRoith D, Brodt P. The role of the IGF system in cancer growth and metastasis: overview and recent insights. *Endocr Rev* 2007;**28**:20–47.
32. Fukuda R, Hirota K, Fan F, Jung YD, Ellis LM, Semenza GL. Insulin-like growth factor 1 induces hypoxia-inducible factor 1-mediated vascular endothelial growth factor expression, which is dependent on MAP kinase and phosphatidylinositol 3-kinase signaling in colon cancer cells. *J Biol Chem* 2002;**277**:38205–11.
33. Xie L, Wang W. Weight control and cancer preventive mechanisms: role of insulin growth factor-1-mediated signaling pathways. *Exp Biol Med (Maywood)* 2013;**238**:127–32.
34. Carreras-Torres R, Johansson M, Haycock PC *et al.* Role of obesity in smoking behaviour: Mendelian randomisation study in UK Biobank. *BMJ* 2018;**361**:k1767.
35. Dimou N, Yarmolinsky J, Bouras E *et al.* Causal effects of lifetime smoking on breast and colorectal cancer risk: mendelian randomization study. *Cancer Epidemiol Biomarkers Prev* 2021;**30**:953–64.
36. Taylor AE, Richmond RC, Palviainen T *et al.* The effect of body mass index on smoking behaviour and nicotine metabolism: a Mendelian randomization study. *Hum Mol Genet* 2019;**28**:1322–30.
37. Tyrrell J, Jones SE, Beaumont R *et al.* Height, body mass index, and socioeconomic status: Mendelian randomisation study in UK Biobank. *BMJ* 2016;**352**:i582.
38. Gage SH, Bowden J, Davey Smith G, Munafò MR. Investigating causality in associations between education and smoking: a two-sample Mendelian randomization study. *Int J Epidemiol* 2018;**47**:1131–40.
39. Hamada T, Nowak JA, Masugi Y *et al.* Smoking and risk of colorectal cancer sub-classified by tumor-infiltrating T cells. *J Natl Cancer Inst* 2019;**111**:42–51.
40. Graff-Iversen S, Hewitt S, Forsén L, Grøtvedt L, Ariansen I. Associations of tobacco smoking with body mass distribution; a population-based study of 65,875 men and women in midlife. *BMC Public Health* 2019;**19**:1439.
41. Jakicic JM, Powell KE, Campbell WW, 2018 PHYSICAL ACTIVITY GUIDELINES ADVISORY COMMITTEE *et al.* Physical Activity and the Prevention of Weight Gain in Adults: A Systematic Review. *Med Sci Sports Exerc* 2019;**51**:1262–9.
42. Sagelv EH, Ekelund U, Hopstock LA *et al.* The bidirectional associations between leisure time physical activity change and body mass index gain. The Tromsø Study 1974–2016. *Int J Obes (Lond)* 2021;**45**:1830–43.
43. Papadimitriou N, Dimou N, Tsilidis KK *et al.* Physical activity and risks of breast and colorectal cancer: a Mendelian randomisation analysis. *Nat Commun* 2020;**11**:597.
44. Carrasquilla GD, García-Ureña M, Fall T, Sørensen TIA, Kilpeläinen TO. Mendelian randomization suggests a bidirectional, causal relationship between physical inactivity and adiposity. *Elife* 2022;**11**:e70386.
45. Barone Gibbs B, Aaby D, Siddique J *et al.* Bidirectional 10-year associations of accelerometer-measured sedentary behavior and activity categories with weight among middle-aged adults. *Int J Obes (Lond)* 2020;**44**:559–67.
46. Cassidy S, Chau JY, Catt M, Bauman A, Trenell MI. Low physical activity, high television viewing and poor sleep duration cluster in overweight and obese adults; a cross-sectional study of 398,984 participants from the UK Biobank. *Int J Behav Nutr Phys Act* 2017;**14**:57.
47. Hojman P, Gehl J, Christensen JF, Pedersen BK. Molecular Mechanisms Linking Exercise to Cancer Prevention and Treatment. *Cell Metab* 2018;**27**:10–21.
48. Flegal KM, Shepherd JA, Looker AC *et al.* Comparisons of percentage body fat, body mass index, waist circumference, and waist-stature ratio in adults. *Am J Clin Nutr* 2009;**89**:500–8.
49. Papadimitriou N, Bull CJ, Jenab M *et al.* Separating the effects of early and later life adiposity on colorectal cancer risk: a Mendelian randomization study. *BMC Med* 2023;**21**:5.
50. Benjamin W, James Y, Ville K, Kate T, Dipender G. Re-evaluating the robustness of Mendelian randomisation to measurement error. *medRxiv* 2022; 2022.10.02.22280617.
51. Li X, Jansen L, Chang-Claude J, Hoffmeister M, Brenner H. Risk of colorectal cancer associated with lifetime excess weight. *JAMA Oncol* 2022;**8**:730–7.