

## SYSTEMATIC REVIEW

# Effectiveness of outpatient geriatric rehabilitation after inpatient geriatric rehabilitation or hospitalisation: a systematic review and meta-analysis

ASTRID PREITSCHOPF<sup>1,2,3</sup>, MARIJE HOLSTEGE<sup>3,4,5</sup>, ANDREA LIGTHART<sup>3</sup>, WIM GROEN<sup>1,2</sup>,  
GEORGE BURCHELL<sup>7</sup>, MARGRIET POL<sup>1,2,6</sup>, BIANCA BUURMAN<sup>1,2</sup>

<sup>1</sup>Department of Medicine for Older People, Amsterdam UMC, location Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

<sup>2</sup> Amsterdam Public Health, Aging & Later Life, Amsterdam, The Netherlands

<sup>3</sup>Department of Research GRZPLUS; Omring and Zorgcirkel, Hoorn, The Netherlands

<sup>4</sup>Department of Research Omring, Hoorn, The Netherlands

<sup>5</sup>Faculty of Health, Sports and Social Work, Inholland University of Applied Sciences, Amsterdam, Netherlands

<sup>6</sup>Research Group Occupational Therapy: Participation and Environment, Faculty of Health, Center of Expertise Urban Vitality, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands

<sup>7</sup>Medical Library, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

Address correspondence to: Astrid D. Preitschopf, MSc, Department of Medicine for Older People, Amsterdam UMC, location Vrije Universiteit Amsterdam, de Boelelaan 1109, 1081 HV, Amsterdam, The Netherlands. Tel: (+31) 651260776. Email: [a.d.vangestel@amsterdamumc.nl](mailto:a.d.vangestel@amsterdamumc.nl)

---

## Abstract

**Background:** Due to the increasing number of older people with multi-morbidity, the demand for outpatient geriatric rehabilitation (OGR) will also increase.

**Objective:** To assess the effects of OGR on the primary outcome functional performance (FP) and secondary outcomes: length of in-patient stay, re-admission rate, patients' and caregivers' quality of life, mortality and cost-effectiveness. We also aim to describe the organisation and content of OGR.

**Methods:** Systematic review and meta-analysis. Five databases were queried from inception to July 2022. We selected randomised controlled trials written in English, focusing on multidisciplinary interventions related to OGR, included participants aged  $\geq 65$  and reported one of the main outcomes. A meta-analysis was performed on FP, patients' quality of life, length of stay and re-admissions. The structural, procedural and environmental aspects of OGR were systematically mapped.

**Results:** We selected 24 studies involving 3,405 participants. The meta-analysis showed no significant effect on the primary outcome FP (activity). It demonstrated a significant effect of OGR on shortening length of in-patient stay ( $P = 0.03$ , MD =  $-2.41$  days, 95%CI:  $[-4.61—0.22]$ ). Frequently used elements of OGR are: inpatient start of OGR with an interdisciplinary rehabilitation team, close cooperation with primary care, an OGR coordinator, individual goal setting and education for both patient and caregiver.

**Conclusion:** This review showed that OGR is as effective as usual care on FP activity. It shows low certainty of evidence for OGR being effective in reducing the length of inpatient stay. Further research is needed on the various frequently used elements of OGR.

**Keywords:** outpatient geriatric rehabilitation, functional performance, length of stay, re-admission, quality of life, systematic review, older people

### **Key Points**

- Geriatric rehabilitation is more frequently offered at home known as ambulatory or outpatient geriatric rehabilitation (OGR).
- OGR is effective in reducing the length of in-patient stay.
- OGR is as effective as usual care on functional performance activity, patients' quality of life, and re-admission rate.
- Further research is needed on the various frequently used elements of OGR.

---

## **Introduction**

Older people with an acute decline in function or with subacute health decline in chronic diseases should be offered Geriatric Rehabilitation (GR) [1]. According to the European consensus, the definition of GR is 'a multidimensional approach of diagnostic and therapeutic interventions, to optimise functional capacity, promote activity and preserve functional reserve and social participation in older people with disabling impairments' [2]. Currently, GR is mostly provided in inpatient hospitals or skilled nursing homes. However, GR is evolving internationally and is more frequently offered at home, in an outpatient setting or in the patients' residence through a specialised multidisciplinary team (known as ambulatory or outpatient geriatric rehabilitation, OGR) [1].

Over recent years, the general intention of GR professionals and policymakers has been to shorten the inpatient phase of GR and continue rehabilitation in the home or an outpatient setting with the aims of cost-effectiveness and providing a high quality of care [3–5]. Earlier studies have discussed that well-structured OGR, offered in line with inpatient GR, may lead to effective rehabilitation outcomes and early discharge [6]. OGR appears to positively affect mobility, balance, risk of falling, self-employment and general health [7]. Task-oriented training in OGR improves the patient's functional outcome and quality of life [7, 8]. Furthermore, involving close relatives can reduce caregiver burden [9]. According to Nanninga et al. [8], caregivers have more confidence in guiding the patient in their daily functioning when that guidance is practiced at home.

Internationally, OGR is organised in different ways with some countries, e.g. Italy and Greece, not able to provide OGR at all, because of the health care finance system [1, 10]. Moreover, there is no consensus about the inclusion criteria for OGR [1, 10]. Multiple reports have indicated a need for further research into OGR [5, 11]. Unfortunately, evidence on the effectiveness of OGR is scarce and mainly focuses on stroke and hip fractures [1]. Additionally, there are many uncertainties about the various structures, processes and rehabilitation environments that may affect the quality and outcome of OGR. To manage the uncertainties regarding the content and organisation of OGR, the post-acute care (PAC) rehabilitation quality framework [12] can be useful to map

these elements and to report on the quality of rehabilitation. This framework is based on two models that are widely used within healthcare: the Structure, Process, Outcome model of Donabedian [13] and the International classification of functioning disability, and health (ICF) model of the World Health Organisation (WHO) which includes the patient-centred aspect of rehabilitation [14].

Earlier research [1, 3, 6–8, 15, 16] indicated that OGR could have an impact on patients' functional performance (FP), total duration of rehabilitation, re-admission to hospital or a nursing home, patients' and caregivers' quality of life, mortality and cost-effectiveness. However, to our knowledge, there have been no studies that systematically mapped out the effects of OGR. Therefore, the present systematic review and meta-analysis assesses the effects of OGR on the primary outcome FP (activity and participation) and secondary outcomes: (i) length of in-patient stay, (ii) re-admission rate, (iii) patients' and caregivers' quality of life, (iv) mortality and (v) cost-effectiveness in comparison to usual care. Moreover, we aim to describe the organisation and content of OGR.

## **Methods**

We performed a systematic review and meta-analysis according to the PRISMA (Preferred Reporting Items for Systematic reviews and meta-Analysis) statement [17]. This systematic review has been registered in the PROSPERO register of systematic reviews (registration number: CRD42021260264).

### **Search strategy**

We conducted a systematic literature search in PubMed, Embase, CINAHL, Cochrane and Web of Science from their inception to July 2022. The search strategy was conducted by GB and AP. The following terms were used (including synonyms and closely related words): 'ambulatory', 'outpatient', 'geriatric', 'rehabilitation' and 'randomized controlled trial'. Appendix 1 shows the full search strategy. Additional trials were identified by screening the references of selected articles.

### Study selection strategy

After removal of duplicates, title and abstract screening was performed independently by two reviewers (AP, AL). Whenever there was a lack of consensus, a third party was consulted (MP, MH). Next, the full text of all eligible studies was screened by two reviewers (AP, AL). We included studies that met the following criteria: (i) Design: randomised controlled intervention trials (RCTs). (ii) Intervention: related to OGR; rehabilitation at home and/or in an outpatient setting and coordinated by a multidisciplinary team with a duration at least three weeks. (iii) Population: people who have an acute decline in function, suffer from a (complex) disease or multi-morbidity in the sub-acute phase, living at home after an inpatient rehabilitation period. (iv) Outcome: at least one of the outcomes of OGR (FP, patients' and caregivers' quality of life (pQoL/cQoL), length of in-patient stay (i.e. acute care) in hospital and/or a post-acute in-patient setting prior to discharge to OGR (LOS), re-admission to hospital or a nursing home, mortality and cost-effectiveness). (v) Control: usual care; geriatric rehabilitation in an inpatient setting (hospital, stroke unit, skilled nursing home), home-based rehabilitation without multidisciplinary organisation, day rehabilitation, primary health care, discharge with no support or no follow-up. Studies were excluded based on the following criteria: age < 65 years, less than two involved disciplines or patients with chronic diseases in a stable phase.

### Definition of FP

A definition of FP was used according to the ICF model and the WHO disability assessment schedule (WHODAS 2.0). FP can be divided into two categories: activity and participation. It states that 'functional performance activity (FPa) describes the execution of a task or action by an individual. It can be sub-divided into: a) cognition, b) mobility, and c) self-care. Functional performance participation (FPP) describes the involvement in a life situation. It can be sub-divided into: a) getting along, b) life activities, and c) participation' [14, 18].

### Data extraction

Data extraction was completed individually by two reviewers (AP, AL). A data extraction form was developed based on the Cochrane Handbook for Systematic Reviews of Interventions [19]. For each study, the following data were extracted: study characteristics, participant characteristics, setting characteristics and outcome. If data were missing, we contacted the corresponding author by email.

### Quality assessment

Two reviewers (AP, AL) independently assessed the risk of bias (RoB) in the studies using the Cochrane risk of bias tool for RCTs (RoB2) [20]. A third party (MH) was consulted when consensus could not be reached. When the

original article did not contain sufficient information for the appraisal, the authors were contacted. Publication bias was assessed by visual inspection of funnel plots when at least 10 studies were included in the meta-analysis [21]. To determine the certainty of evidence, we applied the Grades of Recommendation, Assessment, Development and Evaluation (GRADE) approach to each meta-analysis (Appendix 5) [22, 23].

### Data syntheses and analysis

Review Manager software (RevMan 5.4) [24] was used to perform the meta-analysis. We selected post-intervention outcome measures reported in at least five of the studies and for which sufficient data were available, such as mean and standard deviation. The embedded Review Manager calculator was used when standard deviations were missing, but *P*-values or 95% confidence intervals were given. Authors were contacted when data were insufficient. We used a random effects model, which assumes heterogeneity and expresses effects as standardised mean differences (SMD) or mean differences (MD) with accompanying confidence intervals. To examine statistical heterogeneity, we used the  $I^2$ , at which 50–90% may represent substantial heterogeneity. To examine study heterogeneity, we performed subgroup analyses based on diagnose-group. A sensitivity analysis was conducted by excluding studies with high RoB. [21]

Data synthesis was conducted in all studies using a harvest plot. For the outcomes FPP, cQoL, mortality and cost-effectiveness, the studies are coded based on whether there was no effect on the intervention, a favourable effect on the intervention or a favourable effect on the control group.

We performed a narrative synthesis [25, 26] to systematically summarise the specific OGR elements using the PAC rehabilitation quality framework [12]. This framework provides an overview of components of a patient-centred approach in PAC rehabilitation and how they interact. The process (patient-care, inter-professional) and immediate-intermediate outcomes (ICF) have an iterative and integrative connection which is influenced by the structure and environmental context elements. All the elements are related to the end goals of rehabilitation [12].

## Results

### Identification of eligible studies

The search identified 8,386 references. Figure 1 shows the Prisma flow chart. After screening by title and abstract, a total of 57 studies were considered for a full text review, where 18 studies were excluded due to various reasons (see Figure 1). A third reviewer was needed in eight cases to achieve consensus during the study selection process. Ultimately, 40 articles describing 24 studies were eligible for inclusion of which 8 were eligible for inclusion for the meta-analysis on FPa, 9 on LOS, 11 on re-admission and 5 on pQoL.




Figure 1. Flow diagram.

### Study characteristics

The study characteristics are shown in Table 1. The number of participants in the 24 included studies totalled 3,405 allocated to the intervention ( $n = 1,777$ ) or control ( $n = 1,628$ ) group. The mean age ranged between 66.5 and 84.0 years. We found 15 studies (62.5%) on stroke [27–51], 5 (20.8%) on general GR [52–57], 3 (12.5%) on hip fractures [58–65] and 1 (4.1%) on COPD [66]. Baseline functioning of the participants was measured in 17 of the studies with the (modified) Barthel Index, whereas most studies included participants with moderate dependency [67].

### Risk of bias assessment

Results of the RoB assessment are documented in Appendix 2. A summary of the overall result is presented in Table 2. In general, there is a low RoB arising from the randomisation process (100%), the missing outcome data (85%,  $n = 34$ ), the measurement of the outcome (90%,  $n = 36$ ) and the selection of the reported result (80%,  $n = 32$ ). The RoB due to adhering to the intervention highlights 12 articles (30%) demonstrating a high RoB. In all of the studies, the participants and personnel were aware of the intervention. In many of the studies, the usual care intervention is not explained in enough detail, so it is not clear if the non-protocol interventions are balanced between the intervention and control group, and 45% of the 40 articles scored an overall high RoB. Visual inspection

of funnel plots showed no indication of publication bias (Appendix 3).

### Meta-analysis

#### Primary outcome

##### FP activity

Twenty-four studies assessed FP. The effect of OGR on the Barthel Index (BI) [67] was assessed in 15 studies of which 8 [33, 39, 43, 44, 47, 49, 51, 64] were included in the meta-analysis (Figure 2A). The analysis was based on 1,038 participants (574 for intervention and 464 for control). There is no significant difference between OGR and usual care ( $P = 0.32$ ). The certainty of evidence is low. There was a negligible effect (SMD) of 0.11 (95% CI: [-0.11–0.34]). There was significant ( $P = 0.003$ ) and substantial heterogeneity ( $I^2 = 67%$ ). When excluding high RoB studies [39, 44, 49], the analysis also demonstrates no significant difference ( $P = 0.72$ ) with a small effect size (SMD) of -0.04 (95% CI: [-0.27–0.19]). The heterogeneity reduced to  $I^2 = 43%$  ( $P = 0.14$ ). The subgroup analyses did not alter the conclusion (Appendix 4).

#### Secondary outcomes

##### Length of in-patient stay

The length of in-patient stay (LOS) in hospital (acute care) and/or in an in-patient setting (PAC) before discharge to

**Table 1.** Characteristics of included studies

First author, year, country	Setting acute → setting outpatient	Diagnosis: No. of patients (I/C)	Age mean (SD): Female % (I/C)	Living alone % (I/C)	Functioning at baseline (I/C): BI, mBI, Katz, FIM, NEADL	Cognitive (I/C) MMSE
Donald, 1995 [52]‡ UK	Hospital → Hospital at home	Geriatric: 30/30	81.6 (5.4): 73.3-84.0 (6.0): 76.7	56.7/80.0	BI: mean (IQR) 15.9 (15-17)/15.7 (15-17)	NI
Rodgers, 1997 [27]‡ McManee, 1998 [28] UK	Hospital → Stroke discharge rehabilitation service	Stroke: 46/46	*73.0 (47-93): 43.0 73.0 (44-91): 48.0	48.0/46.0	BI: median (range): 15.0 (2-20)/13.0 (2-20)	NI
Holmqvist, 1998 [29]‡ Von Koch, 2000 [30]‡ Von Koch, 2001 [31] Thorsén, 2005 [32] Sweden	Hospital → Home rehabilitation	Stroke: 41/40	70.9 (7.6): 46.3-72.7 (8.9): 45.0	26.8/32.5	Katz ADL: 17.0/20.0	MMSE: median (IQR) 27.0 (26-29)/28.0 (26-28)
Basket, 1999 [33]‡ New Zealand	Hospital → Supervised home-based program	Stroke: 50/50	67.8 (11.6): 46.0-71.7 (9.1): 40.0	NI	mBI: mean (SD) 86.8 (15.7)/87.6 (14.2)	NI
Anderson, 2000 [34]‡ New Zealand	Hospital → Early hospital discharge and home-based rehabilitation	Stroke: 42/44	72.0 (11): 38.1 71.0 (11): 50.0	40.5/43.2	mBI: median (IQR) 85.0 (80-97)/86.0 (77-95)	MMSE: median (IQR) 28.0 (25-29)/28.0 (77-95)
Indredavik, 2000 [35]‡ [36], Fjærtroft, 2004 [37] Fjærtroft 2011 [38] Norway	Hospital SU → (rehabilitation clinic → Extended stroke unit service	Stroke: 160/160	¥74.0: 45.0 73.8: 56.0	41.0/43.0	mBI: mean (median) 60.4 (65.0)/58.5 (60.0)	NI
Mayo, 2000 [39]‡ Teng, 2003 [40]‡ Canada	Hospital → Early supported home-discharge	Stroke: 58/56	70.3 (12.7): 63.8-69.6 (12.7): 71.4	NI	mBI: mean (SD) 84.6 (14.4)/82.7 (13.9)	NI
Roderick, 2001 [41]‡ UK	Hospital → Domiciliary rehabilitation	Stroke: 66/74	*78.6 (62-91): 50.0 79.6 (60-95): 56.8	40.9/35.1	BI: mean (SD) 12.6 (4.4)/12.8 (5.0)	NI
Baurz-Holter, 2002 [42]‡ Norway	Hospital SU → Early supported discharge	Stroke: 42/40	*79.5 (69-84): 50.0 78.0 (74-82): 60.0	57.1/62.5	BI: median (IQR) 16.5 (12-19)/14.0 (11-18)	MMSE: median (IQR) 27.5 (24-29)/27.0 (22-30)
Crofty, 2002 [58]‡ Australia	Hospital → Early discharge with home-based rehabilitation	Hip fracture: 34/32	*81.6 (78.2-84.8): 62.0 83.5 (76.6-85.5): 75.0	44.0/34.0	mBI: median (IQR) 85.0 (79-89)/85.0 (77-89)	MMSE: median (IQR) 26.0 (24-29)/28.0 (27-29)
Askim, 2004 [43]‡ Norway	Hospital or rehabilitation clinic → Extended stroke unit service	Stroke: 31/31	¥76.9: 48.4 76.3: 45.2	35.5/48.4	mBI: mean (median) 57.7 (55)/54.0 (55)	NI
Cunliffe, 2004 [53]‡ Miller, 2005 [54] UK	Hospital → Early discharge rehabilitation service	Geriatric: 185/185 Fracture 28% Neurological 26% Cardio-respiratory 14% Musculoskeletal 8% Gastroenterological 5% Infection 4% Peripheral vascular 3% Other medical or surgical condition 2% Non-specific condition 11%	*80.0 (73-85): 71.0 79.0 (72-86): 62.0	69.0/65.0	BI: median (IQR) 15.0 (13-16)/15.0 (13-16)	NI

(Continued)



Table 1. Continued

First author, year, country	Setting acute → setting outpatient	Diagnosis: No. of patients (I/C)	Age mean (SD); Female % (I/C)	Living alone % (I/C)	Functioning at baseline (I/C): BI, mBI, Katz, FIM, NEADL	Cognitive (I/C) MMSE
Caplan, 2006 [68]‡ Australia	Hospital → Home rehabilitation service	Frail older patients (ischaemic heart disease, diabetes, dementia) 64/33	83.9 (7.8); 61.4 84.0 (7.0): 64.7	NI	FIM: mean (SD) 100.3 (16.9)/ 78.9 (16.0)	MMSE: mean (SD) 22.7 (7.3)/ 23.2 (4.9)
Zidén, 2008 [60]‡, Zidén, 2010, [61] Sweden	Hospital → Home rehabilitation	Hip fracture: 48/54	81.2 (5.9); 60.4 82.5 (7.6): 77.8	54.0/72.2	NI	NI
Crotty, 2008 [56]‡ Australia	Hospital → Home rehabilitation	Geriatric: 113/116 Total knee replacement: 19%, Stroke: 44% home General rehabilitation (hip fracture, orthopaedic injury, functional decline) 37%	71.2 (13.4); 53.1 72.2 (14.8): 51.7	46.0/45.0	mBI: mean (SD) 92.2 (6.4)/ 92.5 (6.5)	MMSE: mean (SD) 27.0 (3.0)/ 26.9 (3.1)
Eaton, 2008 [66]‡ New Zealand	Hospital → Early inpatient-outpatient rehabilitation	Exacerbation COPD: 47/50	70.1 (10.3); 55.0 69.7 (9.4): 58.0	NI	NI	NI
Parker, 2011 [57]‡ UK	Hospital → Home-based rehabilitation	Geriatric: 47/42 Stroke 22%	74.0 (11); 45.2 (76.0 (11): 45.0	NI	NEADL: 15.0/16.9	NI
ESD stroke Bergen study: Hofstad, 2014 [44]‡ Gjelsvik, 2014 [45]‡ Taulle, 2015 [46] Norway Rasmussen, 2016 [47]‡ Denmark	Hospital SU or rehabilitation clinic → Early supported discharge day unit and early supported discharge home Hospital → Early home-based rehabilitation before and after discharge	Falls assessment 27%, Mobility assessment 21% Orthopaedic rehabilitation 9% Other reasons 20% Stroke: 207/99	*71.3 (27–92): 43.0 74.2 (32–98): 47.5 *78.0 (72–84): 58.0 79.0 (71–85): 58.0	NI	mBI: median (IQR) 100.0 (50), 92.5 (35)/95.0 (45)	NI
Santana 2017 [48]‡ Portugal	Hospital SU or rehabilitation clinic → Early home supported discharge	Stroke: 95/95	*67.5 (40–84): 51.0 66.5 (35–84): 43.0	NI	FIM: (SD) (min-max); 69.0 (21.3) (30–100)/70.5 (18.7) (24–100)	NI
Karlson, 2016 [62]‡ Berggren, 2019 [63]‡ Karlson, 2020 [64]‡ Karlson, 2020 [65] Sweden Rafsten, 2019 [49]‡ Sweden	Hospital → Geriatric interdisciplinary home rehabilitation Hospital → Very early supported discharge	Hip fracture: 107/98 Stroke: 69/71	83.2 (7.0); 73.8 82.6 (6.4): 69.4 75.5 (11.1); 39.1 72.7 (12.4): 38.0	72.9/69.4 NI	BI: median (IQR) 18.0 (13–20)/ 18.0 (13–20) mBI: median(IQR) 82.5 (65–90)/ 80 (65–90)	MMSE: mean (SD) 17.3 (8.4)/ 17.3 (8.9) NI
Vluggen, 2021 [50] Netherlands	Nursing home rehabilitation SU → Integrated multidisciplinary geriatric rehabilitation program	Stroke: 99/91	78.9 (7.0); 69.7 79.0 (6.5): 51.1	54.0/47.3	Katz-15: 6.0 (4.0)/6.5 (3.3)	NI
Kam Yuet Wong, 2022 [51]‡ China	Hospital → home based transitional care program	Stroke 58/58	66.21 (10.07); 67.00 (8.61)	12.1/10.3	mBI: mean (SE) 83.86 (2.03) 87.93 (1.76)	NI

SU = stroke, unit I/C = intervention/control, NI = no information, geriatric diagnosis = mix of different diagnosis (stroke, orthopaedic, COPD, trauma and other), AMPS = assessment of motor and process skills, (m)BI = (modified) Barthel index, IPA = impact on participation and activity, MMSE = mini mental state examination, mRS = modified Rankin scale, NEADL = Nottingham extended ADL, CAPE = Clifton assessment procedure for the elderly; information/orientation sub-scale score < 8. ‡ = included meta-analysis. \* = SD not available. † = Age median (range).

Table 2. Description of intervention

Author, year (colour indicates Risk of Bias)	Structure			Process				Environment			Outcome
	Type of acute care setting	Type of OGR setting	OGR multi-disciplinary team	Facilities, external health-care	Amount and timing	Coordination of care	Specific intervention	Caregiver	Reported outcome (measurement)		
Donald, 1995	Hospital	OGR starts inpatient	Day care/outpatient clinic	Facilitated transport	< 4 weeks	Coordinator	Weekly team meeting	Education meeting for patient	Favours intervention	No difference	Favours control
Rodgers, 1997	X	X	4	Cooperation with primary care	4-10 weeks	Case manager	Coordinator	Self-training program	Participation and meeting for caregiver	Du (LOS), CE	Rea, Mor,
McManee, 1998	X	X	5	Cooperation with primary care	X	X	X	Home visit during inpatient period	Training/education	Du (LOS), Mor, CE	Rea, Mor,
Holmqvist, 1998 Von Koch, 2000, 2001	X	X	4	Specialised from inpatient members	X	X	X	Task, context-specific training	Individual goal setting	FPa (Kara), FFP (FAU), Du (LOS), CE	Rea, Mor,
Thorsén, 2005	X	X	2	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (BI, nhptet, 10mnus), Du (freq. du), Rea, Mor	Rea, Mor,
Baskett, 1999	X	X	2	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (BI, 10mnus, nhptet, Edtent), Du (du) (LOS), pQoL (HADS), cQoL (CHQ28)	Rea, Mor,
Anderson, 2000	X	X	5	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (mBI), FFP (44Profile), Rea, pQoL (SF-36), cQoL (CSI)	Rea, Mor,
Indredavik, 2000 Fjearroff, 2003	X	X	3	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (BI), FFP (FAU), Du (LOS), cQoL (CSI), Mor	Rea, Mor,
Fjearroff, 2004, 2011	X	X	5	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (OAKS, BI), Mor	Rea, Mor,
Mayo, 2000 Teng, 2003	X	X	5	Community care	X	X	X	Individual goal setting	Weekly team meeting	Du (LOS, freq), Rea, pQoL (SF-36), cQoL (CGB), CE	Rea, Mor,
Rodericks, 2001	X	X	4	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (BI, Rivermead mobility index), FFP (FAU), Rea, pQoL (SF-36), CE	Rea, Mor,
Croty, 2001, 2003	X	X	7	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (mBI) Du (LOS)	Du (du)
Baurz Holter, 2002	X	X	3	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (TUG, BBS, ABC), pQoL (SF-36), cQoL (SF-36), cQoL (CSI)	Du (du)
Askim, 2004	X	X	4	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (NEADL), Du (LOS), pQoL, (CHQ), Mor	Rea, Mor,
Caniffie, 2004 Miller, 2005	X	X	4	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (mRS, BI), Du (LOS), pQoL (NHP), cQoL (CSI), Mor	Rea, Mor,
Caplan, 2006	X	X	6	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (BI, NEADL), Du (du, freq) Rea, pQoL (EQ5D, CHQ), cQoL (CHQ), Mor	Rea, Mor,
Zidén, 2008, 2010	X	X	4	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (FIM), Du (LOS, du, freq), Mor	Rea, Mor,
Zidén, 2008, 2010	X	X	4	Community care	X	X	X	Individual goal setting	Weekly team meeting	FPa (FAM, TUG-STS), FFP (FAU), pQoL (SF-36)	Rea, Mor,

(Continued)


Table 2. Continued

Author, year (colour indicates Risk of Bias)	Structure				Process				Environment		Outcome											
	Type of acute care setting	OGR starts inpatient	Patients' home	Day care/outpatient clinic	Type of OGR setting	OGR multi-disciplinary team	OGR multi-disciplinary members	Facilities, external health-care	Amount and timing	Coordination of care	Specific intervention	Caregiver	Reported outcome (measurement)									
Crooy, 2008	X	X	X	X	X	8	Facilitated transport	< 4 weeks	4-10 weeks	X	Coordinator	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Du (du), Rea	Fa (AMPS, TUG), Du (LOS, freq), pQoL (SF-36), cQoL (SF-36, CSI)
Eaton, 2008	X	X	X	X	X	X	Cooperation with primary care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Rea, pQoL (SF-36, HADS, CRQ-SI)
Barben, 2011	X	X	X	X	X	7	Specialised from inpatient	< 4 weeks	4-10 weeks	X	Coordinator	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (NEADL), Du (LOS), Rea, pQoL (EQ-5D); cQoL (CHQ-30)	
ESD stroke Bergen study; Hofstad, 2014	X	X	X	X	X	4	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (mRS, BI, AMPS, FAC, TUG, 5mWS), Du (LOS, du)
Gjelsvik, 2014, Tadle, 2015	X	X	X	X	X	4	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (mRS), Du (LOS, du)
Rasmussen, 2016	X	X	X	X	X	4	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (mRS), pQoL (EQ-5D)
Samana, 2017	X	X	X	X	X	4	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (FIM), FPP (FAI), Du (LOS)
Karlson, 2016, 2020, Berggen 2019	X	X	X	X	X	6	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (BI, Katz-ADL, walking ability), Du (du, freq), Rea, Mor
Barben, 2019	X	X	X	X	X	3	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (BI), Du (du, freq)
Vluggen, 2021	X	X	X	X	X	6	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (Katz-15), FPP (FAI), pQoL (SSQoL), cQoL (QoL-scale), Mor
Kam Yuet Wong, 2022	X	X	X	X	X	6	Community care	< 4 weeks	4-10 weeks	X	Case manager	Depending on needs	Weekly team meetings	Individual goal setting	Task, context-specific training	Home visit during inpatient period	Self-training program	Education meeting for patient	Training/education meeting for caregiver	Participation and cooperation	Favours intervention	Fa (mBI)


Fa = functional performance activity, FPP = functional performance participation, Rea = re-admissions, pQoL = patients' quality of life, cQoL = caregivers' quality of life, Mor = mortality, CE = cost-effectiveness, LOS = length of stay, inpatient before start OGR, du = duration of OGR, freq = frequency of therapy session. AMPS = assessment of motor and process skills, (m)BI = (modified) Barthel index, CRQ-SA = chronic respiratory questionnaire-self-administered, EQ5D = EuroQol 5D, FAC = functional ambulatory scale, FAI = Frenchay activity index, FIM = functional impairment scale, GHQ-30 = general health questionnaire, HADS = hospital anxiety and depression scale, IAM = instrumental activity measure, IPA = impact on participation and activity, LOS = length of inpatient stay before intervention, MMSE = mini mental state examination, mRS = modified Rankin scale, NEADL = Nottingham extended ADL, NHP-rest = nine hole peg-test, NRS = numeric rating scale, NHP = Nottingham Health Profile, OARS = older Americans resource scale for instrumental ADL, SF-36 = short form health survey, STS = sit to stand, SSQoL = stroke specific quality of life scale, TUG = timed up and go test, 5mTW = 5 meter timed walk, 10MWS = 10 meter walk speed. <sup>1</sup> Same team inpatient and in OGR, <sup>#</sup>General practitioner provides routine medical care




2A Barthel index. Note: GRADE certainty of evidence rated low due to substantial heterogeneity and high Risk of Bias



2B Length of in-patient stay. Note: GRADE certainty of evidence rated low due to substantial heterogeneity and high Risk of Bias



2C Re-admission. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias




**Figure 2.** Meta-analyses 2A Barthel index. Note: GRADE certainty of evidence rated low due to substantial heterogeneity and high Risk of Bias 2B Length of in-patient stay. Note: GRADE certainty of evidence rated low due to substantial heterogeneity and high Risk of Bias 2C Re-admission. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias 2D SF-36 physical health subscale. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias 2E SF-36 mental health subscale. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias.


**A. Preitschopf et al.**

Effectiveness of Outpatient Ambulatory Geriatric Rehabilitation: A systematic review and Meta-analysis


2D SF-36 physical health subscale. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias



2E SF-36 mental health subscale. Note: GRADE certainty of evidence rated moderate due to high Risk of Bias



**Figure 2.** Continued.



**Figure 3.** Data synthesis of reported outcomes on functional performance participation, caregivers' quality of life, mortality and cost-effectivity in 24 included randomized controlled trials of outpatient rehabilitation programmes designed for geriatric patients.

OGR was assessed in 20 studies of which nine studies [33, 39, 43, 45, 48, 52, 56, 60, 68] were included in the meta-analysis (Figure 2B). The analysis was based on 1,260 participants (696 for the intervention and 564 for control). There is a significant difference between OGR and usual care ( $P=0.03$ , MD =  $-2.41$ , 95% CI:  $[-4.61$  to  $-0.22]$ ). There was significant ( $P=0.0001$ ) and substantial heterogeneity ( $I^2=74\%$ ). When excluding high RoB studies [39, 45, 52, 56, 68], the analysis demonstrates no significant difference ( $P=0.46$ ) with an MD of  $-0.50$  (95% CI:  $[-1.81$ – $0.81]$ ). The heterogeneity reduced to  $I^2=0\%$  ( $P=0.88$ ). The subgroup analyses did not show a significant difference between OGR and usual care (Appendix 4).

### Re-admission

The number of re-admissions to hospital or a nursing home was measured in 11 studies [27, 30, 40, 41, 52, 53, 56, 57, 63, 66, 68] (Figure 2C). The analysis was based on 1,565 participants (804 for the intervention and 761 for control). It shows no significant difference ( $P=0.31$ ) with an RR of 0.90 (95% CI:  $[0.73$ – $1.10]$ ). The heterogeneity was low ( $I^2=15\%$ ,  $P=0.30$ ). The subgroup analyses and the exclusion of studies with a high RoB [40, 41, 52, 53, 56, 57, 68] show similar effect (Appendix 4).

### Patients' QoL

The pQoL was assessed in 18 studies. The effect of OGR on the Short Form health survey (SF-36) [69] was assessed in seven studies of which five [34, 39, 56, 58, 66] were included in the meta-analysis (Figure 2D and E). There is no significant difference between OGR and usual care on the patients' quality of life physical health ( $P=0.84$  MD:  $-0.21$ , 95% CI:  $[-2.30$ – $1.87]$ ) or mental health ( $P=0.86$  MD:  $-0.18$ , 95% CI:  $[-1.87$ – $2.23]$ ) subscales. The subgroup analyses and the exclusion of studies with a high RoB [34, 39, 56] did not alter the effect (Appendix 4).

Appendix 6 shows a narrative synthesis of FPp, LOS, re-admission and pQoL, of all included studies in this review.

### Data synthesis

The harvest plot (Figure 3) shows overall no effect of the OGR intervention. The primary outcome FPp is reported in seven studies of which 43% ( $n=3$ ) demonstrated a favourable effect of the intervention, but 57% ( $n=4$ ) observed a neutral effect. Studies concluded that participants showed better autonomy and participated in more outdoor activities [32, 50]. The cQoL is reported in eleven studies of which 82% ( $n=9$ ) observed a neutral effect. Studies indicated that teaching and consultation possibilities can empower the caregiver to fulfil their supportive role [40, 50]. The outcome of cost-effectiveness is an outlier, and 75% ( $n=6$ ) of the eight studies that reported cost-effectiveness show a favourable effect of the intervention. These studies [28, 31, 34, 40, 41, 54] concluded that a shorter inpatient

period, less need of care and less re-admissions to hospital or nursing home might be causing this effect.

### Organisation of the intervention

The specific structural, procedural and environmental aspects of the OGR programmes of the included 23 studies, plus the reported outcomes and measurements are summarised in Table 2.

### Structure

The acute care setting before admission to OGR was mainly in a hospital setting (83%). The OGR programme started in nine (37.5%) studies during the inpatient rehabilitation period immediately after admission and the same multidisciplinary team followed the participants during the whole rehabilitation trajectory. The mean number of multidisciplinary team members was 4.7 (range: 2–8), which were specialised in GR in 14 (58.3%) studies. Co-operation with primary care had a prominent place in 12 (50%) studies.

### Process

The duration of OGR shows great heterogeneity with a maximum duration ranging from 4 to 44 weeks. A 4-week duration was most common and occurs in eight of the studies (33.3%). Also, in nine (37.5%) of the studies, coordination of the OGR was performed by a team member acting as a case manager. Individual goal setting, mostly together with the patient, was applied in 14 (58.3%) studies.

### Environment

In 15 (62.5%) studies, an active participation role and close cooperation with the caregiver was mentioned. Furthermore, in six (25%) studies, an education or training meeting was provided to the caregiver and focused on disease management, guiding the patient in daily activities and how to perform and supervise the self-training program.

### Discussion

This systematic review aimed to present the effects of OGR on various outcomes and describe the organisation and content of OGR. We can conclude that OGR is as effective as usual care on the primary outcome FPp and secondary outcomes pQoL and re-admission rate. Furthermore, we demonstrated a positive effect on shortening the inpatient LOS; however, this result is not reflected in the subgroup analyses based on diagnosis-group and is based on low certainty evidence.

The data synthesis on the primary outcome FPp shows that the included studies reported comparable effects of OGR versus usual care. Remarkably participation was only used as an outcome in seven of the studies although the main goal of geriatric rehabilitation is to restore functioning and participation levels. It is known that after discharge from

an inpatient GR setting, older people face the consequences of disabilities in functioning and have trouble resuming participation in meaningful activities [15, 70, 71]. Possibly no difference is found because there seems to be too little focus on participation goals during rehabilitation which could enhance outcomes on participation level in outpatient rehabilitation [72, 73]. Therefore, we believe that focussing on participation goals starting inpatient and maintaining these in OGR could enhance outpatient rehabilitation.

In contrast to other outcomes, the data synthesis for cost-effectiveness demonstrated a favourable effect of the OGR intervention. However, the studies describing cost-effectiveness were mostly outdated and different calculation methods were used. Nevertheless, cost-effectiveness is important to demonstrate the added value of OGR [74]. Therefore, we recommend performing an economic evaluation alongside effectiveness studies in the future to gain better insight into the cost-effectiveness of OGR.

Unexpectedly, eHealth was rarely mentioned as a process element in the included studies. eHealth has become increasingly important in GR in recent years, and the added value has been presented in several studies. Pol et al. [75] have demonstrated positive effects using activity monitoring on top of cognitive behavioural OGR therapy on patients' reported daily functioning for people with hip fractures. Kraaijkamp [76] likewise shows positive results of blended eHealth applications and their feasibility within GR.

The reported overview of the organisation elements of OGR commonly used in the included studies can be used to further develop the organisation and content of OGR. The main difference between OGR and usual care is the rehabilitation environment. OGR mainly takes place at the patients' home with the advantage to stimulate and practice ADL activities in the own environment [9]. This could promote mobilisation and participation rehabilitation goals [77]. Still, we lack a precise description of the interventions including therapy exercises, intensity, frequency and place [78]. Moreover, it is not clear how the inpatient process can be adapted to achieve earlier and smooth discharge to OGR. We recommend performing multiple intervention studies to determine which elements are efficient, feasible and cost-effective in OGR. Additionally, future research is needed on structure, process and environmental elements of OGR from the perspective of patients, professionals and policymakers.

### **Strengths and limitations**

The strength of this systematic review is the focus on patients requiring GR with various diagnoses. Many previous studies focused on the specific diagnosis group of stroke [1]; however, various diagnosis groups occur in OGR, and OGR must be applicable for them all [79]. Another strength is the focus on older people aged  $\geq 65$  with multi-morbidities. This group is often excluded from trials because of polypharmacy or multi-morbidities [80]. Nonetheless, these people qualify for OGR [1, 2]. To our knowledge, this review is the first to examine the effects of OGR specially designed for patients requiring GR.

This review also has limitations. First, the meta-analysis was only performed in four of the eight outcomes due to heterogeneity in the measurement scales used and little sufficient data were available. Yet, there is a low certainty of evidence as many studies had a high risk of bias or exhibited some concerns in one or more items, such as deviations from the intended intervention. This phenomenon was also found in the review of Handoll et al. [16] and is inherent to the type of study because blinding of the patients and carers is impossible in a rehabilitation setting. Secondly, the OGR interventions show substantial clinical heterogeneity, which could influence the strength of the meta-analysis [21]. It is questionable whether pooled analyses of and comparison between the different interventions is appropriate. Subgroup analyses on the diagnose-group showed similar results, although the number of studies was small, and therefore, no firm statements can be made [21, 81]. We still believe that it is appropriate to perform the meta-analyses because the wide variety of diseases is a hallmark of GR. In this study, we provide an overview of which elements are interesting to explore further. This study indicates that OGR is as effective as usual care and possibly more cost-effective. Further development of outpatient rehabilitation seems necessary given the challenges for future-proof care of older people.

### **Conclusion**

Our systematic review shows that OGR is as effective as usual care on the primary outcome FPa and the secondary outcomes pQoL, and re-admission rate. Yet, we found low-certainty evidence for OGR being effective in shortening inpatient LOS. The data synthesis showed indications that OGR might be cost-effective. Additionally, it demonstrated various frequently used structural, procedural and environmental elements of OGR: (i) inpatient start and the same team provides rehabilitation at home, (ii) close cooperation with primary care, (iii) an OGR coordinator, (iv) individual goal setting and (v) an educational session to patient and caregiver. Future research is needed to reach consensus on the content and organisation of OGR and to determine which elements are efficient, feasible and cost-effective. In addition, more focus is needed on participation-level outcomes.

---

**Acknowledgements:** Availability of data and materials—the data will be made available, from the corresponding author on reasonable request.

**Declaration of Conflicts of Interest:** None.

**Declaration of Sources of Funding:** None.

---

### **References**

1. van Balen R, Gordon AL, Schols JMGA, Drewes YM, Achterberg WP. What is geriatric rehabilitation and how should it



- be organized? A Delphi study aimed at reaching European consensus. *Eur Geriatr Med* 2019; 10: 977–87.
2. Grund S, Gordon AL, van Balen R *et al.* European consensus on core principles and future priorities for geriatric rehabilitation: consensus statement. *Eur Geriatr Med* 2020; 11: 233–8.
  3. Langhorne P, Baylan S, Early Supported Discharge Trialists. Early supported discharge services for people with acute stroke. *Cochrane Database Syst Rev* 2017; 7: Cd000443. <https://doi.org/10.1002/14651858.CD000443.pub4>.
  4. Achterberg WP, Cameron ID, Bauer JM, Schols JM. Geriatric rehabilitation-state of the art and future priorities. *J Am Med Dir Assoc* 2019; 20: 396–8.
  5. van den Besselaar J, Buurman B, Hartel L *et al.* Navigate on experiences; of care receivers, caregivers and care professionals in geriatric rehabilitation, primary care residence and physician care for specific patient groups [in Dutch]. The Netherlands: ZonMw and the Ministry of Health, Welfare & Sport 2021; 01-01-2021 Report No.
  6. Holstege MS, Bakkers E, van Balen R, Gussekloo J, Achterberg WP, Caljouw MA. Structured scoring of supporting nursing tasks to enhance early discharge in geriatric rehabilitation: the BACK-HOME quasi-experimental study. *Int J Nurs Stud* 2016; 64: 13–8.
  7. Kiel S, Gerhardt S, Meinhold S, Meinhold A, Schmidt CO, Chenot JF. Course of treatment and sustainability of ambulatory geriatric rehabilitation - an analysis of 128 geriatric patients. *Rehabilitation* 2019; 58: 104–11.
  8. Nanninga CS, Postema K, Schönherr MC, van Twillert S, Lettinga AT. Combined clinical and home rehabilitation: case report of an integrated knowledge-to-action study in a Dutch rehabilitation stroke unit. *Phys Ther* 2015; 95: 558–67.
  9. Tijssen LM, Derksen EW, Achterberg WP, Buijck BI. Challenging rehabilitation environment for older patients. *Clin Interv Aging* 2019; 14: 1451–60.
  10. Grund S, van Wijngaarden JP, Gordon AL, Schols J, Bauer JM. EuGMS survey on structures of geriatric rehabilitation across Europe. *Eur Geriatr Med* 2020; 11: 217–32.
  11. Consultzorg Q. Quick Scan Geriatric Rehabilitation Care, Primary Care Residence and Additional Physician Care (in Dutch). The Netherlands: Q-Consultzorg, 2019; 01-05-2019. Report No.
  12. Jesus TS, Hoenig H. Postacute rehabilitation quality of care: toward a shared conceptual framework. *Arch Phys Med Rehabil* 2015; 96: 960–9.
  13. Donabedian A. The quality of care. How can it be assessed. *JAMA* 1988; 260: 1743–8.
  14. WHO. International Classification of Functioning Disability and Health 2020 [cited 2020]. Available from: <https://www.who.int/standards/classifications/international-classification-of-functioning-disability-and-health>.
  15. Pol M, Peek S, van Nes F, van Hartingsveldt M, Buurman B, Kröse B. Everyday life after a hip fracture: what community-living older adults perceive as most beneficial for their recovery. *Age Ageing* 2019; 48: 440–7.
  16. Handoll HH, Cameron ID, Mak JC, Panagoda CE, Finnegan TP. Multidisciplinary rehabilitation for older people with hip fractures. *Cochrane Database Syst Rev* 2021; 11: Cd007125. <https://doi.org/10.1002/14651858.CD007125.pub3>.
  17. Page MJ, Bossuyt PM, Boutron I, Hoffmann TC *et al.* The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews. 2020. Available from: <https://systematicreviewsjournal.biomedcentral.com/articles/10.1186/s13643-021-01626-4>.
  18. Ustün TB, Chatterji S, Kostanjsek N *et al.* Developing the World Health Organization disability assessment schedule 2.0. *Bull World Health Organ* 2010; 88: 815–23.
  19. Boutron IPM, Higgins JPT, Altman DG, Lundh A, Hróbjartsson A. Considering bias and conflicts of interest among the included studies. In: Higgins JPTTJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, 6.2nd edition. The Cochrane collaboration, 2021. Available from <https://training.cochrane.org/handbook/current>.
  20. Higgins JPT, Altman DG, Gøtzsche PC *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343: d5928. <https://doi.org/10.1136/bmj.d5928>.
  21. Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch VA. *Cochrane Handbook for Systematic Reviews of Interventions*. 6.2nd edition. The Cochrane collaboration, 2021. Available from <https://training.cochrane.org/handbook/current>.
  22. Schünemann HJ, Mustafa RA, Brozek J *et al.* GRADE guidelines: 21 part 1. Study design, risk of bias, and indirectness in rating the certainty across a body of evidence for test accuracy. *J Clin Epidemiol* 2020; 122: 129–41.
  23. Schünemann HJ, Mustafa RA, Brozek J *et al.* GRADE guidelines: 21 part 2. Test accuracy: inconsistency, imprecision, publication bias, and other domains for rating the certainty of evidence and presenting it in evidence profiles and summary of findings tables. *J Clin Epidemiol* 2020; 122: 142–52.
  24. Review Manager (Rev Man). vol. 5.4.1, September 2020. Available from <https://training.cochrane.org/online-learning/core-software/revman>.
  25. Campbell M, McKenzie JE, Sowden A *et al.* Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020; 368: l6890. <https://doi.org/10.1136/bmj.l6890>.
  26. Popay J, Roberts H, Sowden A *et al.* Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A Product from the ESRC Methods Programme. UK: Lancaster University, 2006. <https://doi.org/10.13140/2.1.1018.4643>.
  27. Rodgers H, Soutter J, Kaiser W *et al.* Early supported hospital discharge following acute stroke: pilot study results. *Clin Rehabil* 1997; 11: 280–7.
  28. McNamee P, Christensen J, Soutter J *et al.* Cost analysis of early supported hospital discharge for stroke. *Age Ageing* 1998; 27: 345–51.
  29. Widén Holmqvist L, von Koch L, Kostulas V *et al.* A randomized controlled trial of rehabilitation at home after stroke in Southwest Stockholm. *Stroke* 1998; 29: 591–7.
  30. von Koch L, Widén Holmqvist L, Kostulas V, Almazán J, de Pedro-Cuesta J. A randomized controlled trial of rehabilitation at home after stroke in Southwest Stockholm: outcome at six months. *Scand J Rehabil Med* 2000; 32: 80–6.
  31. von Koch L, de Pedro-Cuesta J, Kostulas V, Almazán J, Widén HL. Randomized controlled trial of rehabilitation at home after stroke: one-year follow-up of patient outcome, resource use and cost. *Cerebrovasc Dis* 2001; 12: 131–8.
  32. Thorsén AM, Holmqvist LW, de Pedro-Cuesta J, von Koch L. A randomized controlled trial of early supported discharge and continued rehabilitation at home after stroke: five-year follow-up of patient outcome. *Stroke* 2005; 36: 297–303.

33. Baskett JJ, Broad JB, Reekie G, Hocking C, Green G. Shared responsibility for ongoing rehabilitation: a new approach to home-based therapy after stroke. *Clin Rehabil* 1999; 13: 23–33.
34. Anderson C, Rubenach S, Mhurchu CN, Clark M, Spencer C, Winsor A. Home or hospital for stroke rehabilitation? Results of a randomized controlled trial : I: health outcomes at 6 months. *Stroke* 2000; 31: 1024–31.
35. Indredavik B, Fjaertoft H, Ekeberg G, Løge AD, Mørch B. Benefit of an extended stroke unit service with early supported discharge: a randomized, controlled trial. *Stroke* 2000; 31: 2989–94.
36. Fjaertoft H, Indredavik B, Lydersen S. Stroke unit care combined with early supported discharge: long-term follow-up of a randomized controlled trial. *Stroke* 2003; 34: 2687–91.
37. Fjaertoft H, Indredavik B, Johnsen R, Lydersen S. Acute stroke unit care combined with early supported discharge. Long-term effects on quality of life. A randomized controlled trial. *Clin Rehabil* 2004; 18: 580–6.
38. Fjærtøft H, Rohweder G, Indredavik B. Stroke unit care combined with early supported discharge improves 5-year outcome: a randomized controlled trial. *Stroke* 2011; 42: 1707–11.
39. Mayo NE, Wood-Dauphinee S, Côté R *et al.* There's no place like home : an evaluation of early supported discharge for stroke. *Stroke* 2000; 31: 1016–23.
40. Teng J, Mayo NE, Latimer E *et al.* Costs and caregiver consequences of early supported discharge for stroke patients. *Stroke* 2003; 34: 528–36.
41. Roderick P, Low J, Day R *et al.* Stroke rehabilitation after hospital discharge: a randomized trial comparing domiciliary and day-hospital care. *Age & Ageing* 2001; 30: 303–10.
42. Bautz-Holter E, Svein U, Rygh J, Rodgers H, Wyller TB. Early supported discharge of patients with acute stroke: a randomized controlled trial. *Disabil Rehabil* 2002; 24: 348–55.
43. Askim T, Rohweder G, Lydersen S, Indredavik B. Evaluation of an extended stroke unit service with early supported discharge for patients living in a rural community. A randomized controlled trial. *Clin Rehabil* 2004; 18: 238–48.
44. Hofstad H, Gjelsvik BE, Næss H, Eide GE, Skouen JS. Early supported discharge after stroke in Bergen (ESD stroke Bergen): three and six months results of a randomised controlled trial comparing two early supported discharge schemes with treatment as usual. *BMC Neurol* 2014; 2014: 239.
45. Gjelsvik BE, Hofstad H, Smedal T *et al.* Balance and walking after three different models of stroke rehabilitation: early supported discharge in a day unit or at home, and traditional treatment (control). *BMJ Open* 2014; 4: e004358. <https://doi.org/10.1136/bmjopen-2013-004358>.
46. Taule T, Strand LI, Assmus J, Skouen JS. Ability in daily activities after early supported discharge models of stroke rehabilitation. *Scand J Occup Ther* 2015; 22: 355–65.
47. Rasmussen RS, Østergaard A, Kjær P *et al.* Stroke rehabilitation at home before and after discharge reduced disability and improved quality of life: a randomised controlled trial. *Clin Rehabil* 2016; 30: 225–36.
48. Santana S, Rente J, Neves C *et al.* Early home-supported discharge for patients with stroke in Portugal: a randomised controlled trial. *Clin Rehabil* 2017; 31: 197–206.
49. Rafsten L, Danielsson A, Nordin A *et al.* Gothenburg very early supported discharge study (GOTVED): a randomised controlled trial investigating anxiety and overall disability in the first year after stroke. *BMC Neurol* 2019; 19: 277. <https://doi.org/10.1186/s12883-019-1503-3>.
50. Vluggen T, van Haastregt JCM, Tan FE, Verbunt JA, van Heugten CM, Schols J. Effectiveness of an integrated multidisciplinary geriatric rehabilitation programme for older persons with stroke: a multicentre randomised controlled trial. *BMC Geriatr* 2021; 21: 134. <https://doi.org/10.1186/s12877-021-02082-4>.
51. Kam Yuet Wong F, Wang SL, Ng SSM *et al.* Effects of a transitional home-based care program for stroke survivors in Harbin, China: a randomized controlled trial. *Age Ageing* 2022; 51: afac027. <https://doi.org/10.1093/ageing/afac027>.
52. Donald IP, Baldwin RN, Bannerjee M. Gloucester hospital-at-home: a randomized controlled trial. *Age Ageing* 1995; 24: 434–9.
53. Cunliffe AL, Gladman JR, Husbands SL, Miller P, Dewey ME, Harwood RH. Sooner and healthier: a randomised controlled trial and interview study of an early discharge rehabilitation service for older people. *Age Ageing* 2004; 33: 246–52.
54. Miller P, Gladman JR, Cunliffe AL, Husbands SL, Dewey ME, Harwood RH. Economic analysis of an early discharge rehabilitation service for older people. *Age Ageing* 2005; 34: 274–80.
55. Caplan GA, Williams AJ, Daly B, Abraham K. A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department – the DEED II study. *J Am Geriatr Soc* 2004; 52: 1417–23.
56. Crotty M, Giles LC, Halbert J, Harding J, Miller M. Home versus day rehabilitation: a randomised controlled trial. *Age Ageing* 2008; 37: 628–33.
57. Parker SG, Oliver P, Pennington M *et al.* Rehabilitation of older patients: day hospital compared with rehabilitation at home. *Age Ageing* 2011; 40: 557–62.
58. Crotty M, Whitehead CH, Gray S, Finucane PM. Early discharge and home rehabilitation after hip fracture achieves functional improvements: a randomized controlled trial. *Clin Rehabil* 2002; 16: 406–13.
59. Crotty M, Whitehead C, Miller M, Gray S. Patient and caregiver outcomes 12 months after home-based therapy for hip fracture: a randomized controlled trial. *Arch Phys Med Rehabil* 2003; 84: 1237–9.
60. Zidén L, Frändin K, Kreuter M. Home rehabilitation after hip fracture. A randomized controlled study on balance confidence, physical function and everyday activities. *Clin Rehabil* 2008; 22: 1019–33.
61. Zidén L, Kreuter M, Frändin K. Long-term effects of home rehabilitation after hip fracture - 1-year follow-up of functioning, balance confidence, and health-related quality of life in elderly people. *Disabil Rehabil* 2010; 32: 18–32.
62. Karlsson Å, Berggren M, Gustafson Y, Olofsson B, Lindelöf N, Stenvall M. Effects of geriatric interdisciplinary home rehabilitation on walking ability and length of hospital stay after hip fracture: a randomized controlled trial. *J Am Med Dir Assoc* 2016; 17: 464.e9–e15.
63. Berggren M, Karlsson Å, Lindelöf N *et al.* Effects of geriatric interdisciplinary home rehabilitation on complications and readmissions after hip fracture: a randomized controlled trial. *Clin Rehabil* 2019; 33: 64–73.
64. Karlsson Å, Lindelöf N, Olofsson B *et al.* Effects of geriatric interdisciplinary home rehabilitation on independence



- in activities of daily living in older people with hip fracture: a randomized controlled trial. *Arch Phys Med Rehabil* 2020; 101: 571–8.
65. Karlsson Å, Berggren M, Olofsson B *et al.* Geriatric interdisciplinary home rehabilitation after hip fracture in people with dementia - a subgroup analysis of a randomized controlled trial. *Clin Interv Aging* 2020; 2020: 1575–86.
  66. Eaton T, Young P, Fergusson W *et al.* Does early pulmonary rehabilitation reduce acute health-care utilization in COPD patients admitted with an exacerbation? A randomized controlled study. *Respirology* 2009; 14: 230–8.
  67. Physiopedia. Barthel Index. 2020 [cited 13 September 2019]. Available from: [https://www.physio-pedia.com/index.php?title=Barthel\\_Index&oldid=222910](https://www.physio-pedia.com/index.php?title=Barthel_Index&oldid=222910).
  68. Caplan GA, Coconis J, Board N, Sayers A, Woods J. Does home treatment affect delirium? A randomised controlled trial of rehabilitation of elderly and care at home or usual treatment (the REACH-OUT trial). *Age Ageing* 2006; 35: 53–60.
  69. Physiopedia. 36-item short form survey (SF-36). 2020 [cited 26 May 2020]. Available from: [https://www.physio-pedia.com/index.php?title=36-Item\\_Short\\_Form\\_Survey\\_\(SF-36\)&oldid=239687](https://www.physio-pedia.com/index.php?title=36-Item_Short_Form_Survey_(SF-36)&oldid=239687).
  70. Simning A, Caprio TV, Seplaki CL, Temkin-Greener H, Szanton SL, Conwell Y. Patient-reported outcomes in functioning following nursing home or inpatient rehabilitation. *J Am Med Dir Assoc* 2018; 19: 864–70.
  71. van Seben R, Reichardt LA, Essink DR, van Munster BC, Bosch JA, Buurman BM. "I feel worn out, as if I neglected myself": older Patients' perspectives on post-hospital symptoms after acute hospitalization. *Gerontologist* 2019; 59: 315–26.
  72. Plant SE, Tyson SF, Kirk S, Parsons J. What are the barriers and facilitators to goal-setting during rehabilitation for stroke and other acquired brain injuries? A systematic review and meta-synthesis. *Clin Rehabil* 2016; 30: 921–30.
  73. Vaalburg AM, Wattel E, Boersma P, Hertogh C, Gobbens R. Goal-setting in geriatric rehabilitation: can the nursing profession meet patients' needs? *Nurs Forum* 2021; 56: 648–59.
  74. Ligtenberg G, Staal PC, Goettsch WG, Knies S. Cost-effectivity in healthcare [in Dutch]. The Netherlands: Health insurance college, 2013; 30-09-2013.
  75. Pol MC, Ter Riet G, van Hartingsveldt M, Kröse B, Buurman BM. Effectiveness of sensor monitoring in a rehabilitation programme for older patients after hip fracture: a three-arm stepped wedge randomised trial. *Age Ageing* 2019; 48: 650–7.
  76. Kraaijkamp JJM, Van Dam van Isselt EF, Persoon A, Versluis A, Chavannes NH, Achterberg WP. eHealth in geriatric rehabilitation: systematic review of effectiveness, feasibility, and usability. *J Med Internet Res* 2021; 23: e24015. <https://doi.org/10.2196/24015>.
  77. Ramsey KA, Loveland P, Rojer AGM *et al.* Geriatric rehabilitation inpatients roam at home! A matched cohort study of objectively measured physical activity and sedentary behavior in home-based and hospital-based settings. *J Am Med Dir Assoc* 2021; 22: 2432–2439.e1.
  78. Verweij L, van de Korput E, Daams JG *et al.* Effects of Postacute multidisciplinary rehabilitation including exercise in out-of-hospital settings in the aged: systematic review and meta-analysis. *Arch Phys Med Rehabil* 2019; 100: 530–50.
  79. Becker C, Achterberg W. Quo vadis geriatric rehabilitation. *Age Ageing* 2022; 51: 1–3.
  80. van Deudekom FJ, Postmus I, van der Ham DJ *et al.* External validity of randomized controlled trials in older adults, a systematic review. *PLoS One* 2017; 12: e0174053. <https://doi.org/10.1371/journal.pone.0174053>.
  81. Gagnier JJ, Moher D, Boon H, Beyene J, Bombardier C. Investigating clinical heterogeneity in systematic reviews: a methodologic review of guidance in the literature. *BMC Med Res Methodol* 2012; 12: 111. <https://doi.org/10.1186/1471-2288-12-111>.

Received 6 April 2022; editorial decision 29 October 2022