

RESEARCH PAPER



Microbial production of mevalonate by recombinant *Escherichia coli* using acetic acid as a carbon source

Xin Xu^{a,b}, Meng Xie^a, Qian Zhao^c, Mo Xian^a, and Huizhou Liu^a

^aCAS Key Laboratory of Biobased Materials, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, China; ^bUniversity of Chinese Academy of Sciences, Beijing, China; ^cCollege of Environment and Safety Engineering, Qingdao University of Science and Technology, Qingdao, China

ABSTRACT

We sought to produce mevalonate, an important organic acid, by recombinant *Escherichia coli*, using acetic acid, a less costly alternative feedstock, as a carbon source. In this study, the mevalonate biosynthesis pathway originating with acetate was constructed in recombinant *E. coli*, resulting in the production of 1.06 g L⁻¹ mevalonate with a productivity of 0.03 g L⁻¹ h⁻¹ in a 5-L bioreactor. The mevalonate concentration and productivity were significantly enhanced with increased cell density during 2-stage aerobic fermentation, reaching 7.85 g L⁻¹ and 0.13 g L⁻¹ h⁻¹, respectively. Fed-batch fermentation was further optimized under anaerobic and microaerobic conditions, and mevalonate concentrations reached 3.05 g L⁻¹ and 4.97 g L⁻¹, respectively, indicating that the oxygen supply exerts a large impact on mevalonate production from acetate. This study describes a method with high potential to produce mevalonate with the engineered *E. coli* strain XU143 using the less costly alternative feedstock acetate as a carbon source.

ARTICLE HISTORY

Received 2 March 2017
Revised 12 April 2017
Accepted 24 April 2017

KEYWORDS

Acetate assimilation;
aeration; mevalonate;
two-stage fed-batch;
Escherichia coli

Introduction

Mevalonate (β -hydroxy- β -methyl- δ -valerolate) is a key organic compound that is produced via the mevalonate pathway in most eukaryotes and higher plants. This compound is the precursor for steroids and terpenes, including isoprene, pinene, carotenoid, artemisinin, and paclitaxel, which are widely used as material monomers, advanced fuels and pharmaceuticals. Mevalonate biosynthetic pathways originating from sugar have been constructed in recombinant *E. coli*.^{1,2} However, with global food shortages and increasing demand for mevalonate, non-food substrates have attracted attention as alternative feedstocks. Acetate is a potential low-cost carbon source because it is an ordinary, aqueous material produced by both biologic and chemical processes. (1) Specifically, acetate is a byproduct of anaerobic digestion processes such as acetogenesis, acidogenesis, and methanogenesis.^{3,4} (2) Additionally, biomass hydrolysates contain acetate because hemicelluloses are generally acetylated.⁵ (3) Acetate is also co-produced during syngas fermentation.⁶

Escherichia coli is one of the most commonly used host microorganisms in the biochemical industry. Acetate metabolism pathways in *E. coli* have previously been elucidated. Acetate accumulates due to overflow metabolism when *E. coli* is cultivated on glucose, but acetate is also used as a primary carbon source during glucose starvation.⁷ In *E. coli*, acetate is converted into acetyl-CoA via the AMP-forming acetyl CoA synthetase (*acs*) and phosphotransacetylase/acetate kinase (*pta-ackA*) pathways (Fig. 1).⁸ Subsequently, acetyl-CoA enters the TCA cycle and glyoxylate cycle, providing energy and precursors for cell growth. Recently, free fatty acids were produced from acetate by an engineered *E. coli* strain, and the *acs* pathway was more effective than the *pta-ackA* pathway.⁹ *E. coli* also represents an attractive host to convert acetate into other chemicals, including mevalonate, through various biocatalysis processes. To our knowledge, there have been no previous reports describing mevalonate biosynthesis from acetate.

Central carbon metabolism in *E. coli* strains cultured with acetate differs greatly from glucose-based

CONTACT Mo Xian  xianmo@qibebt.ac.cn; Huizhou Liu  LiuHuizhou@qibebt.ac.cn  Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, 266101, China.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. The metabolic pathways of mevalonate biosynthesis from acetate. Solid line: mevalonate biosynthesis pathway used in this study; dotted line: acetate assimilation pathway in *E. coli*. Genes in upper pathway: *acs*, for acetyl-CoA synthase; Genes in lower pathway: *mvaE*, for acetyl-CoA acetyltransferase; *mvaS*, for HMG-CoA synthase.

metabolism. During acetate assimilation, glycolytic flux and the pentose phosphate pathway (PPP) are almost completely impaired.^{10,11} However, certain important cofactors, such as ATP and NAD(P)H, are generated by both pathways when *E. coli* is cultivated on glucose.^{12,13} This may exert adverse effects on cell growth and product synthesis. Studies investigating fermentation strategies and conditions may address this problem. In this study, the acetate-derived mevalonate biosynthesis pathway was constructed in recombinant *E. coli*, which was cultured in a bioreactor using acetate as the only carbon source. Then, 2-stage fermentation for mevalonate production was performed to improve cell growth and mevalonate concentration. Additionally, mevalonate production and overall conversion from acetate under different aerobic conditions were compared.

Result and discussion

Construction of the mevalonate biosynthesis pathway originating with acetate

The AMP-forming acetyl CoA synthetase from *Escherichia coli*, which catalyzes acetate to acetyl-CoA (upper pathway), and the MVA pathway from

Enterococcus faecalis, which converts acetyl-CoA into mevalonate (lower pathway) (Fig. 1), were assembled in *E. coli* strain XU143 to achieve efficient mevalonate production originating with acetate. This assembly was performed with the following considerations: first, AMP-forming acetyl CoA synthetase was previously verified to be more efficient for acetate assimilation.^{8,9} Second, the MVA pathway from *E. faecalis* was shown to be efficient in our previous study.² After shake-flask cultivation for 48 h in the presence of acetate, 368 g L⁻¹ mevalonate was obtained from *E. coli* strain XU143, while the control strain XU141 failed to form mevalonate. The mevalonate production efficiency in strain XU143 was approximately 1.3-fold greater than that of strain XU142, which only contained the lower pathway (280 g L⁻¹) (Fig. 2). This result indicates acetate-derived mevalonate biosynthesis by XU143 has much developmental potential.

Fed-batch fermentation using acetate as the sole carbon source (FB-1)

As shown in Figure 2, mevalonate production by XU143 using acetate as the sole carbon source was achieved. Next, we grew XU143 in a bioreactor to




Figure 2. Mevalonate production of different strains using acetate as a carbon source when cultured in shaking flasks. When OD_{600} reached 0.6–0.9, cultures were induced for 36 h using IPTG in shake-flasks. All the experiments were performed in triplicates. 368 g L^{-1} mevalonate was obtained from acetate with XU143, while the control strain XU141 failed to form mevalonate from acetate.

obtain larger amounts of mevalonate. pH-coupled fed-batch fermentation was performed with XU143 using acetate as the sole carbon source. Biomass, mevalonate accumulation and acetate consumption were monitored over the course of fermentation (Fig. 3). The mevalonate concentration peaked at 1.06 g L^{-1} after 43.5 h of cultivation. Overall yield and productivity were 0.30 g g^{-1} and $0.03 \text{ g L}^{-1} \text{ h}^{-1}$, respectively. Moreover, the

specific mevalonate production rate exhibited a trend similar to that of cell growth: both reached a maximum during the early stage of induction and then decreased gradually. Thus, mevalonate production from acetate partially correlated with cell growth.

Mevalonate was produced by XU143 from acetate with a high maximum specific production rate ($102 \text{ mg L}^{-1} \text{ DCW h}^{-1}$) compared with that achieved




Figure 3. Fed-batch fermentation using acetate as the sole carbon source. Induction was performed at 9 h at the DCW of 0.4 g L^{-1} . The maximum concentration of mevalonate reached 1.06 g L^{-1} . Overall yield and productivity were 0.30 g g^{-1} and $0.03 \text{ g L}^{-1} \text{ h}^{-1}$, respectively.

with glucose ($50 \text{ mg L}^{-1} \text{ DCW h}^{-1}$).¹ Therefore, XU143 may have great potential for mevalonate biosynthesis. However, low production and biomass may be attributable to 2 factors. First, acetate exerts a negative effect on intracellular protein stability.^{8,9} Second, the overexpression of heterologous genes leads to a burden on cellular metabolism. Thus, a 2-stage fermentation process was designed. The first stage produced a significant amount of biomass derived from glucose, while the second stage facilitated the accumulation of mevalonate following the addition of acetate.

Two-stage fed-batch fermentation (FB-2)

Changes in the glucose, acetate and biomass concentrations observed during the fermentation process are shown in Figure 4. Glucose was completely consumed after culture for 9 h. Then, IPTG was added, and ammonium acetate was used as both the sole carbon and nitrogen source. During the first stage of culture on glucose, the biomass concentration reached 8.6 g L^{-1} with a yield of 0.43 g of biomass g^{-1} of glucose. These values are consistent with those reported in the literature for *E. coli* grown on glucose.^{14,15} When cultivated on acetate, the *E. coli* strain appeared to adapt to this new carbon source, and the biomass decreased slightly. During this period, the mevalonate concentration increased in association with acetate consumption, meaning the strain continued to assimilate the carbon source, but carbon flux was diverted toward mevalonate

synthesis, and cell proliferation halted. After cultivation for 60 h, the mevalonate concentration reached 7.85 g L^{-1} , corresponding to an overall yield of 0.27 g g^{-1} and a productivity of $0.13 \text{ g L}^{-1} \text{ h}^{-1}$ (Fig. 4).

A 2-stage fermentation strategy is conducive to the bioconversion of certain products.^{16,17} However, there have been no reports describing 2-stage fermentation involving a switch in the carbon source. By using glucose during the first stage and acetate during the second stage, the mevalonate concentration and productivity greatly increased by 8.4- and 4.3-fold, respectively, compared with the results achieved in FB-1. Simultaneously, overall yield and the maximum specific mevalonate production rate resembled that of FB-1 (Table 1). This suggests that 2-stage fermentation, which avoids the limitations of acetate assimilation, is an effective strategy for mevalonate bioconversion from acetate.

Two-stage fed-batch fermentation under different aerobic conditions

Aeration conditions play an important role in acetate metabolism in *E. coli*. Acetate accumulation is inversely correlated with dissolved oxygen (DO) levels when *E. coli* is grown on glucose.^{18,19} However, there have been few reports investigating acetate assimilation under different aerobic conditions. Therefore, we next performed 2-stage fed-batch fermentation under anaerobic and microaerobic (with aeration rates of 1.0




Figure 4. Two-stage fermentation using glucose in batch cultivation and acetate in fed-batch cultivation. When glucose was completely consumed, IPTG was added, and ammonium acetate was used as both the carbon and nitrogen source. Mevalonate concentration reached 7.85 g L^{-1} , corresponding to an overall yield of 0.27 g g^{-1} and a productivity of $0.13 \text{ g L}^{-1} \text{ h}^{-1}$.

Table 1. A comparison of mevalonate fermentation parameters among different fermentation modes.

Carbon source	Aeration	DCW _{max} (g L ⁻¹)	Mevalonate (g L ⁻¹)	q _{p max} (mg L ⁻¹ DCW h ⁻¹)	Productivity (g L ⁻¹ h ⁻¹)	Y _p (g g ⁻¹ acetate)
Acetate	Aerobic	0.67	1.06	102	0.03	0.30
Glucose-acetate	Aerobic	8.6	7.85	103	0.13	0.27
Glucose-acetate	Anaerobic	8.5	3.05	16.5	0.05	0.37
Glucose-acetate	Microaerobic	7.6	4.97	60	0.08	0.29

q_{p max} Maximum specific mevalonate production rate;
 Productivity Biosynthesis amount of mevalonate per ferment liquid per hour;
 Yield g mevalonate/ g acetate. The yield was calculated after induced only.

vvm) conditions. During the first stage, fully aerobic conditions were implemented until glucose was depleted. During the second stage, IPTG was added, and anaerobic and microaerobic conditions were implemented using ammonium acetate as the sole carbon source. Figure 5 shows the results from the second stage of fermentation.

Cell mass decreased in all experiments after induction, and the minimum cell mass was below 5.4 g L⁻¹ under anaerobic conditions. Furthermore, the mevalonate concentration was 3.05 g L⁻¹ with a yield of 0.37 g g⁻¹ acetate under anaerobic conditions. Under microaerobic conditions, mevalonate

production improved and reached 4.97 g L⁻¹ with a yield of 0.29 g g⁻¹ acetate at 1.0 vvm (Fig. 5 and Table 1). The maximum mevalonate concentration and minimum yield were observed under the fully aerobic conditions achieved with FB-2. With a lower oxygen supply, mevalonate concentration and productivity decreased, but yield increased. The distribution of catabolic flux changes under aerobic conditions, possibly to adjust the intracellular redox state.²⁰⁻²² Therefore, the yield trend was assumed to be associated with acetate flux via the TCA and mevalonate pathways under different aerobic conditions.




Figure 5. Fed-batch fermentation under different aerobic conditions. A: The cell growths; B: Mevalonate productions; C: Acetate consumptions. During the first stage, fully aerobic conditions were implemented until glucose was depleted. During the second stage, IPTG was added, and anaerobic and microaerobic conditions were implemented using ammonium acetate as the sole carbon source. This figure shows the results from the second stage of fermentation. Mevalonate concentration was 3.05 g L⁻¹ with a yield of 0.37 g g⁻¹ acetate under anaerobic conditions. In microaerobic conditions, mevalonate production was improved and reached 4.97 g L⁻¹ with a yield of 0.29 g g⁻¹ acetate at 1.0 vvm. The maximum mevalonate concentration and minimum yield were observed under fully aerobic conditions achieved with FB-2.

Conclusion

The engineered *E. coli* strain XU143 was capable of using acetate as a sole carbon source to produce 7.85 g L⁻¹ mevalonate during a 2-stage fermentation process. Anaerobic and microaerobic fed-batch fermentation experiments indicated oxygen supply exerts a key impact on mevalonate production from acetate. Further work must be done to improve mevalonate accumulation using different feeding strategies and by optimizing culture conditions. Overall, the method described here demonstrates great potential for mevalonate production by *E. coli* using the less costly feedstock acetate.

Materials and methods

Strain construction

All plasmids and strains used in this study are listed in Table 2. *E. coli* W3110 genomic DNA was amplified by PCR as a template to obtain the *acs* gene using the primers *acs*-F (5'-TAGAGATCTTTACGATGGCATCGC G-3') and *acs*-R (5'-GCCCATATGATGAGCCAAATT CACAAAC-3'). The PCR product was cloned into the pCOLADuet-1 vector between the *Bgl*III and *Nae*I sites to generate the plasmid pCOLADuet-*acs*. The *mvaE* and *mvaS* genes were cloned into pACYCDuet-1, resulting in the plasmid pYJM17 (Yang et al. 2012). pCOLADuet-1 and pACYCDuet-1 were transformed into *E. coli* BL21(DE3) to generate *E. coli* XU141. pYJM17 and pCOLADuet-1 were transformed into *E. coli* BL21(DE3) to generate *E. coli* XU142. pCOLADuet-*acs* and pYJM17 were transformed into *E. coli* BL21(DE3) to generate *E. coli* XU143.

Table 2. Strains and plasmids used in this study.

Strains or plasmids	Characteristic	Reference
Strains		
<i>E. coli</i> W3110	DNA template for amplification	TIANGEN
<i>E. coli</i> DH5 α	Host for general cloning	TIANGEN
<i>E. coli</i> BL21 (DE3)	Host for protein expression	Novagen
XU141	<i>E. coli</i> BL21(DE3)/pCOLA Duet/pACYCDuet	This study
XU142	<i>E. coli</i> BL21(DE3)/pCOLA Duet/pYJM17	This study
XU143	<i>E. coli</i> BL21(DE3)/pCOLA Duet- <i>acs</i> /pYJM17	This study
Plasmids		
pCOLADuet TM -1	ColA origin, Kan ^r	Novagen
pACYCDuet-1	p15A (pACYC184), Cm ^r	Novagen
pCOLADuet- <i>acs</i>	pCOLADuet-1 carrying <i>acs</i> from <i>E. coli</i> W3110	This study
pYJM17	pACYCDuet-1 carrying <i>mvaE</i> and <i>mvaS</i> from <i>Enterococcus faecalis</i>	

Medium

Primary seed cultures of XU101 were grown in LB medium containing 10 g L⁻¹ tryptone, 5 g L⁻¹ yeast extract, and 10 g L⁻¹ NaCl. The secondary seed culture medium was the same as the fermentation medium. The fermentation minimal medium contained (per liter): 2.5 g of KH₂PO₄, 3 g of (NH₄)₂SO₄, 1 g of citric acid·H₂O, 1 g of C₂H₃O₂Na·2H₂O, 1.86 g of KCl, 0.08 g of FeSO₄·7H₂O, 0.24 g of MgSO₄·7H₂O, 1 g of β ine, 2 g of beef powder, and 1 ml of trace elements solution. The initial carbon sources for fed-batch fermentation (acetate as the only carbon source) and 2-stage fermentation (glucose-acetate) were 2 g L⁻¹ sodium acetate and 20 g L⁻¹ glucose, respectively. The trace elements stock solution contained (per liter): 2.9 g of ZnSO₄·7H₂O, 3.7 g of (NH₄)₆Mo₇O₂₄·4H₂O, 24.7 g of H₃BO₃, 2.5 g of CuSO₄·5H₂O, and 15.8 g of MnCl₂·4H₂O.

Cultivation

Fed-batch fermentation was performed to scale up mevalonate production from acetate. Fermentation was performed in a Biostat B plus MO5L fermentor (Sartorius Stedim Biotech GmbH, Germany) containing 2 L of medium as described above. During the fermentation process, pH was automatically maintained at 7.0 with ammonia and sulfuric acid. During both fed-batch fermentation (acetate as the only carbon source) and 2-stage fed-batch fermentation (glucose-acetate), the dissolved oxygen (DO) concentration was maintained at 20% saturation for fully aerobic conditions. Feeding medium contained 6 M ammonium acetate. When initial glucose was depleted, ammonium acetate was fed continuously to maintain an acetate concentration ranging from 25 mmol L⁻¹ to 75 mmol L⁻¹ in fermentation broth. Two-stage fermentation (glucose-acetate) under microaerobic and anaerobic conditions was also performed. During the first stage, fully aerobic conditions were applied as described above until glucose was depleted, and then 6 M ammonium acetate was added followed by the implementation of either microaerobic (with aeration rates of 1.0 vvm) or anaerobic (by flushing with N₂ instead of oxygen) conditions. Samples were withdrawn to determine cell mass and metabolite concentrations.

Analytical methods

Biomass was determined by measuring the optical density of the culture sample at 600 nm (1 OD unit = 0.43 DCW L⁻¹). The concentration of residual glucose

was quantified using a SBA-40D biologic sensing analyzer. Acetate and mevalonate concentrations were determined by performing HPLC with an aminex HPX-87H ion exclusion column (Bio-Rad, USA). To analyze mevalonate, samples were prepared by mixing of 0.5 ml of 0.1- μ m-filtered fermentation broth with 0.5 ml of 3 M H₂SO₄ and again filtering the resulting solution through a 0.1- μ M filter into HPLC vials, no later than 10 min before HPLC detection.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This work was supported by National Natural Science Foundation of China (21572242 and 21376255), and Taishan Scholars Climbing Program of Shandong (No.tspd20150210).

References

- Tabata K, Hashimoto SI. Production of mevalonate by a metabolically-engineered *Escherichia coli*. *Biotechnol Lett* 2004; 26:1487-91; <https://link.springer.com/article/10.1023%2FB%3ABILE.0000044449.08268.7d>; PMID: 15604784; <https://doi.org/10.1023/B:BILE.0000044449.08268.7d>
- Yang J, Xian M, Su S, Zhao G, Nie Q, Jiang X, Zheng Y, Liu W. Enhancing production of bio-isoprene using hybrid MVA pathway and isoprene synthase in *E. coli*. *PLoS ONE* 2012; 7(4):e33509; <http://journals.plos.org/plosone/articleid=10.1371/journal.pone.0033509>; PMID:22558074; <https://doi.org/10.1371/journal.pone.0033509>
- Periana RA, Mironov O, Taube D, Bhalla G, Jones CJ. Catalytic oxidative condensation of CH₄ to CH₃COOH in one step via CH activation, *Science* 2003; 301:814-8; <http://science.sciencemag.org/content/301/5634/814>; PMID:12907796; <https://doi.org/10.1126/science.1086466>
- Ni C, Wu X, Dan J, Du D. Facile recovery of acetic acid from waste acids of electronic industry via a partial neutralization pretreatment (PNP) Distillation strategy. *Sep Purif Technol* 2014; 132:23-6; <http://www.sciencedirect.com/science/article/pii/S138358661400272X>; <https://doi.org/10.1016/j.seppur.2014.04.052>
- Zhu Z, Sathitsuksanoh N, Vinzant T, Schell DJ, McMillan JD, Zhang YHP. Comparative study of corn stover pretreated by dilute acid and cellulose solvent-based lignocellulose fractionation: Enzymatic hydrolysis, supramolecular structure, and substrate accessibility. *Biotechnol Bioeng* 2009; 103:715-24; <http://onlinelibrary.wiley.com/doi/10.1002/bit.22307/abstract;jsessionid=68731E51BDF65BD8357BD8678D5F8B9D.f04t03>; PMID: 19337984; <https://doi.org/10.1002/bit.22307>
- Pradeep CM, Samir K. Biomass-derived syngas fermentation into biofuels: opportunities and challenges. *Bioresour Technol* 2010; 101:5013-22; <http://www.sciencedirect.com/science/article/pii/S0960852409017635>; <https://doi.org/10.1016/j.biortech.2009.12.098>
- Lin H, Natalia M, Castro, George N, Bennett, San KY. Acetyl-CoA synthetase overexpression in *Escherichia coli* demonstrates more efficient acetate assimilation and lower acetate accumulation: a potential tool in metabolic engineering. *Appl Microbiol Biotechnol* 2006; 71:870-4; <https://link.springer.com/article/10.1007%2Fs00253-005-0230-4>; PMID:16496143; <https://doi.org/10.1007/s00253-005-0230-4>
- Wolfe AJ. The acetate switch. *Microbiol Mol Biol Rev* 2005; 69:12-50; <http://mmb.asm.org/content/69/1/12>; PMID:15755952; <https://doi.org/10.1128/MMBR.69.1.12-50.2005>
- Xiao Y, Ruan Z, Liu Z, Wu S, Varman A, Liu Y, Tang Y. Engineering *Escherichia coli* to convert acetic acid to free fatty acids. *Biochem Eng J* 2013; 76:60-9; <http://www.sciencedirect.com/science/article/pii/S1369703X13001174>; <https://doi.org/10.1016/j.bej.2013.04.013>
- Oh MK, Rohlin L, Kao KC, Liao JC. Global expression profiling of acetate-grown *Escherichia coli*. *J Biol Chem* 2002; 277:13175-83; <http://www.jbc.org/content/277/15/13175>; PMID:11815613; <https://doi.org/10.1074/jbc.M110809200>
- Zhao J, and Shimizu K. Metabolic flux analysis of *Escherichia coli* K12 grown on ¹³C-labeled acetate and glucose using GC-MS and powerful flux calculation method. *J Biotechnol* 2003; 101:101-17; <http://www.sciencedirect.com/science/article/pii/S0168165602003164>; PMID:12568740; [https://doi.org/10.1016/S0168-1656\(02\)00316-4](https://doi.org/10.1016/S0168-1656(02)00316-4)
- Zhu G, Golding B, Dean AM. The selective cause of an ancient adaptation. *Science* 2005; 307:1279-82; <http://science.sciencemag.org/content/307/5713/1279>; PMID:15653464; <https://doi.org/10.1126/science.1106974>
- El-Mansi M, Cozzone AJ, Shiloach J, Eikmanns BJ. Control of carbon flux through enzymes of central and intermediary metabolism during growth of *Escherichia coli* on acetate. *Curr Opin Microbiol* 2006; 9:173-9; <http://www.sciencedirect.com/science/article/pii/S1369527406000221>; PMID:16530464; <https://doi.org/10.1016/j.mib.2006.02.002>
- Fu J, Wilson DB, Shuler ML. Continuous, high level production and excretion of a plasmid-encoded protein by *Escherichia coli* in a two-stage chemostat. *Biotechnol Bioeng* 1993; 41:937-46; <http://onlinelibrary.wiley.com/doi/10.1002/bit.260411004/abstract>; PMID:18601275; <https://doi.org/10.1002/bit.260411004>
- Baheri HR, Hill GA, Roesler WJ. Optimum design for production of tryptophan synthetase subunit from recombinant *Escherichia coli* in a two-stage continuous flow stirred tank reactor (growth and production stage). *Chem Eng J* 1996; 62:143-8; <http://www.sciencedirect.com/science/article/pii/0923046795030654>
- Oh JS, Cho D, Park TH. Two-stage continuous operation of recombinant *Escherichia coli* using the bacteriophage lambda Q(-) vector. *Bioprocess Biosyst Eng* 2005; 28:1-7; <https://link.springer.com/article/10.1007%2Fs00449-005-0418-0>; PMID:16096763; <https://doi.org/10.1007/s00449-005-0418-0>

- [17] Choi SJ, Park DH, Jung KH. Development and optimization of two-stage cyclic fed-batch culture for hG-CSF production using L-arabinose promoter of *Escherichia coli*. *Bioprocess Biosyst Eng* 2001; 24:51-8; <https://link.springer.com/article/10.1007%2Fs004490100231>; <https://doi.org/10.1007/s004490100231>
- [18] Alexeeva S, Hellingwerf KJ, Mattos MJT. Requirement of ArcA for redox regulation in *Escherichia coli* under microaerobic but not anaerobic or aerobic conditions. *J Bacteriol* 2003; 185:204-9; <http://jb.asm.org/content/185/1/204>; PMID:12486057; <https://doi.org/10.1128/JB.185.1.204-209.2003>
- [19] Phue JN, Shiloach J. Impact of dissolved oxygen concentration on acetate accumulation and physiology of *E. coli* BL21, evaluating transcription levels of key genes at different dissolved oxygen conditions. *Metab Eng* 2005; 7:353-63; <http://www.sciencedirect.com/science/article/pii/S1096717605000534>; PMID:16099189; <https://doi.org/10.1016/j.ymben.2005.06.003>
- [20] Aleeva S, Kort B, Sawers G, Hellingwerf KJ, Mattos MJT. Effects of limited aeration and of the arcAB system on intermediary pyruvate catabolism in *Escherichia coli*. *J Bacteriol* 2000; 182:4934-40; <http://jb.asm.org/content/182/17/4934>; <https://doi.org/10.1128/JB.182.17.4934-4940.2000>
- [21] Tran QH, Uden G. Changes in the proton potential and the cellular energetics of *Escherichia coli* during growth by aerobic and anaerobic respiration or by fermentation. *Eur J Biochem* 1998; 251:538-43; <http://onlinelibrary.wiley.com/doi/10.1046/j.1432-1327.1998.2510538.x/abstract>; PMID:9492330; <https://doi.org/10.1046/j.1432-1327.1998.2510538.x>
- [22] Paul B, Hirshfield I. The effect of acid treatment on survival and protein expression of a laboratory K-12 strain *Escherichia coli*. *Research in Microbiology* 2003; 154:115-21; <http://www.sciencedirect.com/science/article/pii/S0923250802000116>; PMID:12648726; [https://doi.org/10.1016/S0923-2508\(02\)00011-6](https://doi.org/10.1016/S0923-2508(02)00011-6)