

Tibialis anterior muscle needle biopsy and sensitive biomolecular methods: a useful tool in myotonic dystrophy type 1

S. Iachettini,¹ R. Valaperta,^{2,3}
A. Marchesi,⁴ A. Perfetti,⁵ G. Cuomo,¹
B. Fossati,⁶ L. Vaienti,⁴ E. Costa,^{2,3}
G. Meola,⁶ R. Cardani¹

¹Laboratory of Muscle Histopathology and Molecular Biology, IRCCS Policlinico San Donato, Milan

²Research Laboratories, IRCCS Policlinico San Donato, Milan

³Service of Laboratory Medicine 1 Clinical Pathology, IRCCS Policlinico San Donato, Milan

⁴Department of Plastic and Reconstructive Surgery, IRCCS Policlinico San Donato, University of Milan

⁵Laboratory of Molecular Cardiology, IRCCS Policlinico San Donato, Milan

⁶Department of Biomedical Science for Health, IRCCS Policlinico San Donato, University of Milan, Italy

Abstract

Myotonic dystrophy type 1 (DM1) is a neuromuscular disorder caused by a CTG repeat expansion in 3'UTR of *DMPK* gene. This mutation causes accumulation of toxic RNA in nuclear foci leading to splicing misregulation of specific genes. In view of future clinical trials with antisense oligonucleotides in DM1 patients, it is important to set up sensitive and minimally-invasive tools to monitor the efficacy of treatments on skeletal muscle. A *tibialis anterior* (TA) muscle sample of about 60 mg was obtained from 5 DM1 patients and 5 healthy subjects through a needle biopsy. A fragment of about 40 mg was used for histological examination and a fragment of about 20 mg was used for biomolecular analysis. The TA fragments obtained with the minimally-invasive needle biopsy technique is enough to perform all the histopathological and biomolecular evaluations useful to monitor a clinical trial on DM1 patients.

Introduction

Myotonic dystrophy type 1 (DM1) is the most common muscular dystrophy in adults, affecting 1 in 6-10,000 live birth. Main clinical

features of DM1 are myotonia, weakness and atrophy of skeletal muscles.¹ DM1 is caused by the expansion of a CTG repeat (from 50 to 3000 repeats) in the 3' untranslated region of the *Dystrophia Myotonica Protein Kinase (DMPK)* gene.²⁻⁴ This mutation gives rise to toxic RNAs containing repetitive CUG repeat that accumulate in cell nuclei as ribonuclear inclusions causing changes of alternative splicing for a specific group of transcripts.^{5,6} Expanded CUG repeats interact with RNA-binding proteins such as Muscleblind-like (MBNL) protein family, causing their nuclear sequestration and subsequently resulting in reduced protein activity in several cellular processes.⁷⁻⁹ Another pathogenic effect of *DMPK* mutation is the induction of post-transcriptional upregulation of another RNA binding protein, CUG-binding protein 1 (CUGBP1).¹⁰ MBNLs and CUGBP1 are antagonistic regulators of alternative splicing, and their functional imbalance leads to embryonic patterns of alternative splicing in adult DM1 tissues.¹¹

Currently, there are no disease-modifying therapies for patients with DM1 and treatments are only to manage symptoms. To date, two main experimental therapeutic strategies of targeting expanded repeat RNA in DM1 have been described: i) antisense oligomer-induced degradation of toxic CUG-containing RNA,¹²⁻²⁰ and ii) inhibition of pathogenic interaction of CUG-containing RNA with nuclear proteins without causing significant degradation of targeted transcript, by either antisense oligomers (ASOs) or small compounds that bind to CUG repeat hairpin.²¹⁻²³ The expected effects of these treatments are the prevention of MBNLs sequestration and/or a significant reduction of nuclear foci formation. These results should lead to the correction of alternative splicing abnormalities for several MBNL-sensitive exons. Indeed, alternative splice events have good potential to function as biomarkers of DM severity and therapeutic response since the splicing misregulation is directly connected to RNA toxicity and proteins sequestrations. Moreover, many splicing defects are correlated with muscle weakness and histopathology and some of them directly implicated in symptoms of DM1.²⁴⁻²⁸ However, at skeletal muscle level, still there is no a definitive mechanistic explanation for the histopathological features of this disease.²⁹ Recent studies have indicated that the distal muscle *tibialis anterior* (TA) is the best muscle to be used to test therapeutic interventions in DM1 patients because it is preferentially affected at both histological and functional level. Moreover, in DM1 patients splicing events are more severely affected in TA than in proximal muscles such as *vastus lateralis* or *biceps brachii*, indicating that TA biopsies are suitable for biomarker discovery

Correspondence: Dr. Rosanna Cardani, Laboratory of Muscle Histopathology and Molecular Biology, IRCCS-Policlinico San Donato, Piazza E. Malan 1, 20097 San Donato Milanese (MI), Italy. Tel. +39.02.52774480. E-mail: rosanna.cardani@grupposandonato.it

Key words: Myotonic dystrophy type 1; *tibialis anterior*; needle biopsy; alternative splicing; clinical trial.

Contributions: SI and RV contributed equally to this work; SI, biomolecular analysis, data analysis and interpretation, participating in drafting the manuscript; RV, genetic analysis, data analysis and interpretation, manuscript revision; AM, setting up the *tibialis anterior* needle biopsies technique, data analysis and interpretation, manuscript revision; AP, RNA extraction, manuscript revision; GC, muscle histology and immunohistochemistry, FISH, manuscript revision; BF, clinical evaluation, data analysis, manuscript revision; LV, setting up the *tibialis anterior* needle biopsies technique, manuscript revision; EC, data analysis, manuscript revision; GM, design and conceptualization of the study, histopathological and clinical evaluation, manuscript revision, final approval of the version to be published; RC, design and conceptualization of the study, handling and treatment of *tibialis anterior* needle biopsies, muscle histopathological evaluation, morphometric analysis, data analysis and interpretation, manuscript drafting, final approval of the version to be published.

Funding: this work was supported by unrestricted grant by FMM (*Fondazione Malattie Miotoniche*), San Donato Milanese (MI), Italy.

Conflict of interest: the authors declare no conflict of interest.

Received for publication: 17 August 2015.

Accepted for publication: 3 October 2015.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright S. Iachettini et al., 2015
Licensee PAGEPress, Italy
European Journal of Histochemistry 2015; 59:2562
doi:10.4081/ejh.2015.2562

in DM1.^{27,28} It should be noted that serial muscle samplings will be required to test a therapeutic response in skeletal muscle, thus the analysis must be performed on small tissue samples easily obtained by minimally invasive biopsy procedures such as needle biopsy technique.

Muscle needle biopsy was developed by Bergström in 1962 as an alternative to open biopsy.³⁰ Needle biopsy is a less invasive procedure than open biopsy, presents a low complication rates and the scar is shorter. It is partic-

ularly indicated for the diagnosis of neuromuscular disease in infancy and childhood or in inflammatory myopathy and to evaluate the disease progression and the response to treatment. The sample size obtained by this technique is smaller than that obtained with an open biopsy and some limitations are the impossibility to directly visualize the site of sampling and the difficulty to identify the right orientation of the fibres of the obtained sample. However, a recent study on more than 13,000 patients demonstrated that using needle biopsy it is possible to obtain an appropriate sample for the analyses required in >99% of cases.³¹

In this work we show that, taking advantage of both the minimal invasiveness of needle biopsy and the suitability of TA muscle for the analysis of DM1 biomarkers, the small muscle fragment obtained from a TA needle biopsy is enough to perform an histopathological and biomolecular characterization of skeletal muscle in DM1 patients. This muscle characterization will be used to monitor a DM1 clinical trial by comparing the obtained results before and after the therapeutic treatment.

Materials and Methods

Patients and TA muscle biopsy

TA muscle biopsies were taken under sterile conditions from 5 DM1 patients and from 5 healthy subjects and were used for this study after receiving written informed consent from the patients. The site is prepped with betadine. Local anaesthesia is performed by 2% mepivacaine injection (7-8 cc) in the subcutaneous tissue and intramuscular. A 1 cm transverse skin incision is performed at the junction between the proximal and middle third of the anterior side of the leg, 2 cm laterally to the tibial crest. Minimal blunt dissection is performed to visualize the deep fascia, which is then incised with a blade for 5 mm. An UCH skeletal muscle needle with aspiration has been used (Dixons Surgical Instruments Ltd., Wickford, UK). This needle is composed of two concentric hollow. The outer (5 mm external

diameter) has a blunt point and a side window and the inner, with a cutting edge at the end, slides freely up and down acting as a guillotine. There is a central plunger which is removed during the sampling. A suction-system is connected to the outer cylinder to obtain samples.

The needle is inserted into the *tibialis anterior* muscle with its side window closed and facing laterally. Once in the muscle, suction-system is activated (200 mm Hg) and the inner cylinder is withdrawn slightly, opening the window. With the free hand, pressure is applied on the outside of the thigh to cause the muscle to bulge into the side window. The central cylinder is then pushed home and a sample obtained with the guillotine action of the cutting edge. The biopsy is performed perpendicular to the longitudinal orientation of the myofibers as opposed to open biopsies, and repeated with a 45° angle to the skin, almost parallel to the orientation of the myofibers, similar to open biopsies. If necessary, more muscle may be obtained by subsequent reinsertion of the needle through the same skin incision. After withdrawal of the needle firm pressure is applied to the thigh for 5 min to prevent any hematoma. Since wound tension is critical in lower limbs, the skin edges are closed with 5-0 monofilament absorbable subcutaneous stitches. Compressive dressing is applied for 5 days.

One sample of TA muscle of about 60 mg was taken. One fragment of about 40 mg was used for histological examination and one fragment of about 20 mg was used for biomolecular analysis. The diagnosis of DM1 was based upon the clinical diagnostic criteria set by the International Consortium for Myotonic Dystrophy.³² DM1 genotyping has been performed on genomic DNA extracted from peripheral blood leukocytes as described.³³ Clinical data on patients used in these study are reported in Table 1.

This study was authorized by the Institutional Ethics Committee (ASL MI2, Melegnano, MI, Italy) and was conducted according to the principles expressed in the Declaration of Helsinki, the institutional regulation and Italian laws and guidelines.

Muscle histopathology and immunohistochemistry

To identify the right orientation of the fibers, a muscle sample of about 40 mg was dissected under a stereomicroscope. Muscle tissue was fresh-frozen in isopentane cooled in liquid nitrogen. Histopathological analyses were performed on serial sections (8 µm) processed for routine histological or histochemical stainings. A standard myofibrillar ATPase staining protocol was used after preincubation at pH 4.3, 4.6, and 10.4.³⁵ The most typical alterations, such as nuclear clump fibers (*i.e.*, aggregates of myonuclei with a thin rim of cytoplasm), nuclear centralization and fiber size variability, were analyzed on serial muscle sections. Central nuclei and nuclear clumps were classified as absent or present. The evaluation of the area of the total transverse section was made with Image J (Scion Co., Frederick, MD, USA) on a single digital image of the entire muscle cross-sectional area obtained by tiling slightly overlapping images of the transverse muscle section.

For immunohistochemistry, sections 6 µm thick were air-dried and rehydrated in phosphate buffer pH 7.4 (PBS). Non-specific binding sites were blocked with normal goat serum (NGS; Dako, Denmark) at a dilution 1:20 in PBS containing 2% bovine serum albumin (BSA; Sigma-Aldrich, St. Louis, MO, USA) for 20 min at room temperature (RT). Mouse monoclonal primary antibodies against two different myosin heavy chain (MHC) isotypes were used at the following dilutions: MHCfast, 1:400 in PBS + 2% BSA; MHCslow, 1:400 in PBS + 2% BSA (Sigma-Aldrich). Each antibody was applied for 1 h at RT. After washing in PBS 3 times for 5 min, sections were incubated with goat anti-mouse biotinylated secondary antibody diluted 1:300 in PBS + 2% BSA for 1 h at RT. After PBS washing (3x5 min), sections were incubated with Vectastain Elite ABC kit (Vector Laboratories, Burlingame, CA, USA) for 30 min and then exposed to the 3,3'-diaminobenzidine tetrahydrochloride (DAB; Sigma-Aldrich) chromogen reaction solution for 10 min. Nuclei were counterstained with Mayer's haematoxylin. Quantitative evaluation

Table 1. Clinical data of DM1 patients.

Patients	Age at biopsy	Age at onset	CTG expansion	MRC	MIRS	Myotonia	Arrhythmia	CPK
DM1-1	33	20	360	129	2	Present	Not present	Normal values
DM1-2	21	15	460	105.22	2	Present	Present	203 U/L
DM1-3	27	21	890	105	3	Present	Not present	Normal values
DM1-4	42	18	190	130	2	Not present	Not present	Normal values
DM1-5	26	13	360	111.33	3	Present	Not present	531 U/L

MRC, Medical Research Council, scale for muscle strength; scale (0–5 grade) on 15 muscles at both side in the upper and lower limbs for a total of 150 maximum score; MIRS, muscle impairment rating scale, stage of the disease for myotonic dystrophy type 1 (DM1) patients;³⁹ CPK, creatine phosphokinase blood levels.

of fiber diameter was made as described previously by Vihola *et al.*³⁶ with Image J (Scion Co.) on images taken with a light microscope (160x, original magnification). The size of muscle fibers was assessed by measuring the *smallest fiber diameter*.

All data were elaborated using Microcal Origin (Microcal Software Inc., Northampton, MA, USA). The metahistograms were normalized to normal mean diameter for men and women.

Fluorescence *in situ* hybridization combined with MBNL1 immunofluorescence

The fluorescence *in situ* hybridization (FISH) was performed on DM1 muscle frozen sections using a (CAG)₆-CA probe (IDT, Coralville, IA, USA) as previously reported by Cardani *et al.*³⁷ Briefly, 6 μ m thick transverse cryostatic sections were air dried for 30 min and fixed with 2% paraformaldehyde for 30 min at 4°C. The sections were then washed in PBS and permeabilized for 5 min in 2% acetone in PBS, prechilled at -20°C. After washing in PBS, sections were incubated in 40% formamide and 2x saline solution citrate (SSC) for 10 min at RT, and hybridized for 2 h at 37°C, with 1 ng/L (CAG)₆-CATexas red labeled probe in 30% formamide, 2xSSC, 0.02% BSA, 67 ng/L yeast tRNA, 2 mM vanadyl ribonuclease complex (all these reagents were from Sigma-Aldrich, St. Louis, MO). The sections were washed first in 40% formamide and 2xSSC at 45°C for 30 min, then in 1xSSC at 45°C for 15 min and another 1xSSC wash at RT. The FISH-labeled sections were then processed for the immunofluorescence detection of MBNL1, as follows. The sections were pre-incubated for 20 min at RT with 5% NGS in PBS+2% BSA and then incubated overnight at 4°C with the rabbit polyclonal antibody A2764 for MBNL1 (1:1000 in PBS+2% BSA). The sections were washed in PBS and then incubated for 1 h at RT with a goat anti-rabbit Alexa488-labeled antibody (Molecular Probes, Life Technologies, Milan, Italy), diluted 1:200 in PBS+2% BSA, counterstained with DAPI (Sigma-Aldrich), and mounted with Mowiol (Cabochem, Milan, Italy).

cDNA synthesis

Frozen TA samples (20 mg) were practiced for the extraction of total RNA using TRIzol reagent (Life Technologies) and 20 ng of RNA were reverse transcribed using the WT-Ovation RNA Amplification System (NuGEN, San Carlos, CA, USA) according to the manufacturer's instructions. The resulting cDNAs were used to perform both quantitative real time-PCR (qRT-PCR) and classical PCR.

Quantitative RT-PCR

The expression level of the *DMPK* gene was measured by Quantitative RT-PCR (qRT-PCR) using the Hs01094329_m1 Assay-on-demand gene expression product, labeled with FAM dye. GAPDH was chosen as the housekeeping, internal control and its expression level was measured using the Hs02758991_g1 Assay-on-demand gene expression product, labeled with FAM dye. Each PCR reaction was performed in triplicate using the TaqMan Universal PCR Master Mix - No AmpErase UNG (Applied Biosystems, Life Technologies) and the 7900HT Fast Real Time PCR System (Applied Biosystem, Life Technologies).

Alternative splicing analysis

The RT-PCR splicing analysis for the *INSR* (insulin receptor) was performed using Platinum Taq Polymerase (Invitrogen, Carlsbad, CA, USA) according to manufacturer's protocol. Briefly, the reaction was conducted in a mix containing 150 ng of cDNA, Taq MgCl₂-free buffer, 2 mM MgCl₂, 400 M dNTPs, 800 μ M forward primer and 800 μ M reverse primer. After 2 min at 94°C, 35 cycles of amplification were performed, each consisting of 30 s at 94°C, 30 s at 60°C and 1 min at 72°C. The RT-PCR splicing analysis for *CAMK2G* (calcium/calmodulin-dependent protein kinase II gamma), *CAMK2B* (calcium/calmodulin-dependent protein kinase II beta), *CACNA1S* (dihydropyridine receptor), *CLCN1* (skeletal muscle chloride channel, voltage-sensitive 1), *NFIX* (Nuclear Factor 1X), *PDLIM3* (PDZ and LIM domain 3) and *LDB3* (LIM Domain Binding 3) were performed using My Taq Red Mix (Bioline), according to manufacturer's protocol. The PCR cycling conditions were: 1 min at 95°C followed by 35 cycles of 15 s at 95°C, 15 s at primer specific T_m, 10 s at 72°C. The primers used for all PCR reactions were

listed in Table 2 with their respective temperatures of melting (T_m). Total PCR products were electrophoretically resolved on 2% agarose gel for *INSR*, *CAMK2G*, *CAMK2B*, *CACNA1S*, *LDB3*, *NFIX* and *PDLIM3* genes and 3,5% agarose gel for *CLCN1*.

Qualitative analysis of the amplified products was performed using EtBr-stained gels (Sigma-Aldrich) scanned on a ChemiDoc Universal Hood (Biorad). The quantitative analysis was performed quantifying the intensity of each band with ImageJ software densitometry and calculating the fraction of abnormally spliced (AS) isoform respect to the total amount of isoforms (AS isoform/total). The expression level of GAPDH was used as the housekeeping, internal control.

Results

One aim of this work was to verify if a muscle fragment of about 40 mg obtained from TA needle biopsy of DM1 patients was adequate to evaluate the histopathological alterations and to analyze *in situ* biomolecular markers. The mean area size of transverse muscle sections obtained from 10 TA needle biopsies resulted to be 1.75 \pm 0.74 mm², about 13 fold smaller than the area size of transverse sections obtained from an open biopsy of *biceps brachii* (*data not shown*). Nevertheless, routine stainings performed on TA transverse sections permit to evaluate the presence of the histological alterations commonly observable in DM1 skeletal muscle such as nuclear clump fibers, nuclear centralization and fiber size variability (Figure 1). The results of histopathological analysis of DM1 TA are reported in Table 3. Moreover, the immunohistochemical staining of MHC slow and fast myosin and the subse-




Figure 1. Histological stainings of transverse sections of frozen *tibialis anterior* needle biopsy of DM1 patients. A) Hematoxylin and eosin. B) Gomori trichrome. A high fiber size variability with several atrophic (arrows) and hypertrophic (asterisks) fibers is evident. Scale bars: 50 μ m

quent analysis of type 1 and type 2 fiber diameters allow to obtain metahistograms of fiber diameter distribution (Figure 2) and to verify the dystrophic changes. The results of the analysis of immunostained sections were reported in Table 3.

To date, targeting the mutant RNA appears to hold some of the greatest prospect for therapeutic intervention in DM1. The reduction of CUGexp RNA would be expected to release sequestered MBNL1 protein and to improve its splicing regulatory activity. Fluorescent *in situ* hybridization (FISH) in combination with MBNL1 immunofluorescence performed on muscle sections is a useful method to reveal if a reduction in the average number of foci of CUGexp RNA and of MBNL1 before and after the ASO treatment is evident. This reduction can be evaluated on sections obtained from the small muscle sample of TA needle biopsy since the number of myonuclei that can be screened (388 ± 56 per mm^2) is enough to obtain significant results on changes in the number and size of ribonuclear inclusions or MBNL1 foci. All DM1 patients used in this study showed ribonuclear inclusions colocalizing with MBNL1 foci (Figure 3).

Consistent with MBNL1 activity restoration, alternative splicing of MBNL1-dependent exons should be examined in order to verify

the efficacy of the therapeutic treatment. To do this it is necessary to use an efficient, sensitive and reliable RNA-to-DNA conversion step since the available muscle sample is limited. NuGEN's Ovation qPCR System is a highly efficient system that provides a fast and simple

method for preparing amplified cDNA from a very small input of total RNA. Moreover, 20% more genes are detected using the amplified cDNAs obtained with the Ovation qPCR System than using those obtained with classical RT-PCR techniques. This system amplifies the




Figure 2. A,B Fast myosin immunostaining of *tibialis anterior* transverse sections of a healthy subject (A) and of a DM1 patient (B); type 2 fibers (fast positive fibers) are stained in brown; scale bars: 100 μm . C,D Metahistograms obtained from the analysis of muscle fiber diameters in a healthy subject (C) and in a DM1 patient (D). The results are based on sections immunostained for MHC fast or slow myosin

Table 2. List of PCR primers used for alternative splicing analysis.

Gene	Forward primer	Reverse primer	T _m (°C)
<i>INSR</i>	5'- CCAAAGACAGACTCTCAGAT-3'	5'-AACATCGCCAAGGGACCTGC-3'	60
<i>CAMK2G ex17</i>	5'-CAACGCTACAGATGGGATCA-3'	5'-AAAGTCCCCATTGTTGATGG-3'	60
<i>CAMK2G ex18</i>	5'-CTCCACAGAGAGCTGCAACA-3'	5'-CAGGCTCAAGGAAGTGAGG-3'	60
<i>CAMK2B</i>	5'- CAGGAGACTGTGGAGTGTCTG-3'	5'-AGCGTCTTCATCCTCTATGGTGG-3'	62
<i>CACNA1S</i>	5'-GCTACTTTGGAGACCCTGGAA-3'	5'-AGGAGGGTTTCGACTCCTTCTG-3'	60
<i>CLCN1</i>	5'-GGTTGTCTGAAGGAATACCTCAC-3'	5'-TCCTCTCCAGTAGTCCGAACAG-3'	60
<i>LDB3 ex11</i>	5'-GACTACCAGGAACGCTTCAACC-3'	5'-GACAGAAGGCCGGATGCTG-3'	62
<i>LDB3 ex5</i>	5'-GCGTCAACACAGACACCATGACC-3'	5'-TCATCTGGGCCAGGATGCGGAA-3'	62
<i>NFIX</i>	5'-GAGCCCTGTTGATGACGTGTTCTA-3'	5'-CTGCACAAACTCCTCAGTGAGTC-3'	62
<i>PDLIM3</i>	5'-AGCCCATCCTTTCAAAATCAAC-3'	5'-AGAGCCATCGTCCACCAITC-3'	58
<i>GAPDH</i>	5'-AGCCTCCCCTTCGCTCTCT-3'	5'-GCCAGCATCGCCCCACTTGA-3'	60

T_m, melting temperature.

Table 3. Histopathological features of *tibialis anterior* needle biopsy in DM1 patients.

Patients	Central nuclei	Nuclear clumps	AFast	HFast	ASlow	HSlow
DM1-1	Rare	Rare	+	++	-	++
DM1-2	Absent	Absent	-	+	±	-
DM1-3	Present	Absent	±	+	+	-
DM1-4	Absent	Present	++	-	-	++
DM1-5	Rare	Absent	-	++	±	+

AFast, fast fiber atrophy; HFast, fast fiber hypertrophy; ASlow, slow fiber atrophy; HSlow, slow fiber hypertrophy; -, absent; ±, modest; +, present; ++, prominent (grade based on fiber diameter methahistograms).

entire transcriptome of 5-50 ng total RNA into micrograms of cDNA across a wide range of transcript abundance.

In our study, the total RNA extracted from a TA frozen sample of about 20 mg was around 8 µg. This quantity of RNA is sufficient to perform 400 RT-PCR using the Ovation qPCR System starting from 20 ng of RNA for each reaction. The total cDNA amount obtained was around 2 mg. This quantity is enough to perform serial molecular analysis, such as qRT-PCR and classical PCR, in order to evaluate the expression levels of *DMPK* and the splicing pattern of several genes involved in the multi-systemic phenotype of DM1 patients respectively. The analysis of the expression levels of *DMPK* gene by qRT-PCR in muscle biopsy

showed that while in healthy patients the expression levels of the gene are quite homogeneous, in DM1 patients an evident variability in the amount of *DMPK* mRNA was observable. Indeed in 3/5 DM1 patients the *DMPK* expression is lower than that observed in healthy subjects, while in two patients *DMPK* expression is respectively 4 and 7 fold higher than the media levels observed in healthy controls and 7 and 11 fold higher than media levels observed in the other DM1 patients (Figure 4A). Alternative splicing of several genes involved in the multi-systemic phenotype of DM1 was analyzed by RT-PCR. For each splice event, the fraction of splice products that included or skipped the alternative exon(s) was compared for five healthy controls and five

DM1 subjects. We observed a splicing alteration of *CAMK2G* exon 18, *LDB3* exon 11 and *LDB3* exon 5 in 60% of DM1 patients and a misregulation of *INSR* exon 11, *CAMK2G* exon 17, *CAMK2B* exon 13, *CACNAS1* exon 29, *CLCN1* exon 7a isoforms, *NFIX* exon 7 and *PDLIM3* exon 5 in 80% of DM1 patients (Figure 4 B,C).

Discussion

Clinical trials for a pharmacological treatment need a road map to patient-focused biomarker measures. Therefore, the study of disease mechanisms, pathogenesis and manifestations is required to select biomarker for the conceptualization of treatment benefits. Before treatment, it is also necessary to collect data from registry of patients in order to understand the natural history of muscle impairment through longitudinal study. So it will be easier to develop muscle impairment endpoints at clinical level, which are also very helpful and necessary during clinical trial in DM1. A phase III trial for the ASO ISIS-DMPKrx is on-going in patients affected by myotonic dystrophy type 1 in order to evaluate the safety and tolerability of multiple doses of this compound (www.clinicaltrials.gov). ISIS-DMPKrx was designed to reduce the production of toxic *DMPK* RNA in cells, including muscle cells. Therapeutic application of this strategy to DM1 patients is expected to cause a rapid knockdown of CUGexp RNA in skeletal muscle, correcting the physiological, histopathological and transcriptomic features of the disease. Published *in vitro* and *in vivo* preclinical studies demonstrated that various antisense oligomers targeting CUGexp repeats induce the degradation of mutant *DMPK* transcripts leading to a reduction of MBNL1 sequestration. Consequently, MBNL1 biological function is restored and the splicing alteration of several genes is corrected including *CLCN1* alternative splicing leading to an improvement of myopathic changes.¹²⁻²⁰

In view of a clinical trial on large scale, it appears evident the need to improve specific tools and to define biomolecular markers that can be used for monitoring the success of the therapy. In particular, it is necessary to validate tools characterized by feasibility and the test-retest reliability. The needle biopsy is a minimally invasive procedure that presents a low complication rate, causes a short scar and, therefore, it can be repeated on the same patient and on the same muscle over the time. These features make needle biopsy a good candidate as a tool to follow a clinical trial on DM1 patients. However, one limitation of this procedure is that the tissue samples obtained are




Figure 3. FISH in combination with MBNL1-immunofluorescence reveals the presence of foci (arrows) of toxic RNA (red spots in panel A) and of MBNL1 (green spots in panel B) co-localizing in nuclei (visualized by the DAPI blue fluorescence) of TA muscle fibers. Scale bars: 10 µm.




Figure 4. A) *DMPK* gene expression measured by qRT-PCR; *GAPDH* was chosen as the housekeeping, internal control; each PCR reaction was performed in triplicate. B) RT-PCR splicing analysis of the indicated genes in 5 healthy subjects and in 5 DM1 patients; arrows indicated the different alternative splicing products: the products that include the specific exon (Ex) and those that exclude the specific exon (Δ); the loading control was based on the expression levels of the *GAPDH*. C) Analysis of the percentage of altered isoform expression of the indicated genes. Histograms represent mean ± SD.

very small. The main effort of this work was therefore to set the conditions in order to succeed in performing histopathological examination and biomolecular analysis on small muscle fragments obtained with needle biopsy. All the analysis were performed on TA muscle since it appears to be the best muscle to test therapeutic interventions in DM1 patients.²⁸

The analysis performed on TA muscle section obtained from needle biopsy indicates that, despite the small fragment dimensions, the cross-section area obtained was enough to evaluate structural changes in muscle fibers, central nuclei frequency, muscle fiber diameter and presence of CUGexp RNA and MBNL1 foci. These parameters should be taken in consideration to verify if the therapeutic treatment can protect against structural changes (mainly a prevention of fiber atrophy) and can decrease toxic RNA and MBNL1 accumulation. The use of specific reverse transcription and amplification systems allowed us to obtain a huge amount of cDNA starting from small amount of RNA. This permitted us to analyze *DMPK* expression levels in our TA muscle samples. As previously reported, our results show an inter-individual variability in *DMPK* expression in DM1.³⁸ DM1 is associated with the misregulation of alternative splicing of many genes that can explain the multisystemic phenotype of the disease. For this reason, splicing events in skeletal muscle are useful biomarkers of DM1 therapeutic response. The amount of cDNA obtained also permit us to analyse the splicing alteration of a large number of genes such as *INSR* and *CLCN1* that are involved in insulin resistance and myotonia respectively and *CACNA1S* whose misplacing may contribute to muscle degeneration. The obtained data confirmed that the percentage of splicing misregulation observed in our DM1 patients is similar to the percentage reported by Nakamori *et al.*²⁸ However, it is reasonable to consider that in clinical trial only splice events of genes that correlate with skeletal muscle symptoms will be examined, on the assumption that the correction of these defects may lead to functional improvement.³⁹

The main limitation of this study is that it does not extend our knowledge on DM1 pathomechanism beyond what is already known. Some data presented in this study were already discussed by Nakamory *et al.*²⁸ who studied more muscle samples and more splicing events. However, the aim of our study was to assess if histological and biomolecular biomarkers might be evaluated on a small fragment of TA muscle to obtain the data necessary to evaluate the effects, positive as well as negative, of therapeutic treatment in DM1 patients. Indeed, by using the minimally invasive needle biopsies technique for sampling the small fragment of muscle, the procedure

can be performed before and after the treatment on the same patient.

In conclusion, our analysis demonstrates that using a minimally invasive procedure as needle biopsy it is possible to obtain small TA fragments which are an adequate amount of muscle tissue to perform all the histopathological and biomolecular analysis useful to monitor the success of a clinical trial on DM1 patients.

References

- Harper PS Myotonic dystrophy, 3th edition. WB Sanders, London: 2001.
- Brook JD, McCurrach ME, Harley HG, Buckler AJ, Church D, Aburatani H, et al. Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 39 end of a transcript encoding a protein kinase family member. *Cell* 1992; 69:799-808.
- Fu YH, Pizzuti A, Fenwick RG Jr, King J, Rajnarayan S, Dunne PW, et al. An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science* 1992; 255:1256-8.
- Mahadevan M, Tsilfidis C, Sabourin L, Shutler G, Amemiya C, Jansen G, et al. Myotonic dystrophy mutation: an unstable CTG repeat in the 39 untranslated region of the gene. *Science* 1992; 255:1253-5.
- Davis BM, McCurrach ME, Taneja KL, Singer RH and Housman DE Expansion of a CUG trinucleotide repeat in the 3 untranslated region of myotonic dystrophy protein kinase transcripts results in nuclear retention of transcripts. *Proc Natl Acad Sci USA* 1997;94:7388-93.
- Orengo JP, Cooper TA. Alternative splicing in disease. *Adv Exp Med Biol* 2007;623:212-23.
- Miller JW, Urbinati CR, Teng-Umuay P, Stenberg MG, Byrne BJ, Thornton CA, et al. Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J* 2000; 9:4439-48.
- Wang ET, Cody NA, Jog S, Biancolella M, Wang TT, Treacy DJ, et al. Transcriptome-wide regulation of pre-mRNA splicing and mRNA localization by muscleblind proteins. *Cell* 2012;150:710-724.
- Malatesta M, Giagnacovo M, Costanzo M, Cisterna B, Cardani R, Meola G. Muscleblind-like1 undergoes ectopic relocation in the nuclei of skeletal muscles in myotonic dystrophy and sarcopenia. *Eur J Histochem* 2013;57:e15.
- Kuyumcu-Martinez NM, Wang GS and Cooper TA Increased steady-state levels of CUGBP1 in myotonic dystrophy 1 are due to PKC-mediated hyperphosphorylation. *Mol Cell* 2007;28:68-78.
- Lin X, Miller JW, Mankodi A, Kanadia RN, Yuan Y, Moxley RT, et al. Failure of MBNL1-dependent post-natal splicing transitions in myotonic dystrophy. *Hum Mol Genet* 2006;15:2087-97.
- Krol J, Fiszler A, Mykowska A, Sobczak K, de Mezer M, Krzyzosiak WJ. Ribonuclease dicer cleaves triplet repeat hairpins into shorter repeats that silence specific targets. *Mol Cell* 2007;25:575-86.
- Sobczak K, Wheeler TM, Wang W, Thornton CA RNA interference targeting CUG repeats in a mouse model of myotonic dystrophy. *Mol Ther* 2013;21:380-7.
- Langlois MA, Lee NS, Rossi JJ, Puymirat J. Hammerhead ribozyme-mediated destruction of nuclear foci in myotonic dystrophy myoblasts. *Mol Ther* 2003;7:670-80.
- Francois V, Klein AF, Beley C, Jollet A, Lemercier C, Garcia L, et al. Selective silencing of mutated mRNAs in DM1 by using modified hU7-sRNAs. *Nat Struct Mol Biol* 2011;18:85-7.
- Nakamori M, Gourdon G, Thornton CA. Stabilization of expanded (CTG)ⁿ(CAG) repeats by antisense oligonucleotides. *Mol Ther* 2011;19:2222-7.
- Wheeler TM, Leger AJ, Pandey SK, MacLeod AR, Nakamori M, Cheng SH, et al. Targeting nuclear RNA for in vivo correction of myotonic dystrophy. *Nature* 2012;488:111-5.
- Lee JE, Bennett CF, Cooper TA. RNase H-mediated degradation of toxic RNA in myotonic dystrophy type 1. *Proc Natl Acad Sci USA* 2012;109:4221-6.
- Mulders SA, van den Broek WJ, Wheeler TM, Croes HJ, van Kuik-Romeijn P, de Kimpe SJ, et al. Triplet-repeat oligonucleotide-mediated reversal of RNA toxicity in myotonic dystrophy. *Proc Natl Acad Sci USA* 2009;106:13915-20.
- Gonzalez-Barriga A, Mulders SA, van de Giessen J, Hooijer JD, Bijl S, van Kessel ID, et al. Design and analysis of effects of triplet repeat oligonucleotides in cell models for myotonic dystrophy. *Mol Ther Nucleic Acids* 2013;2:e81.
- Wheeler TM, Sobczak K, Lueck JD, Osborne RJ, Lin X, Dirksen RT et al. Reversal of RNA dominance by displacement of protein sequestered on triplet repeat RNA. *Science* 2009;325:336-9.
- Wojtkowiak-Szlachcic A, Taylor K, Stepniak-Konieczna E, Sznajder LJ, Mykowska A, Sroka J et al. Short antisense-locked nucleic acids (all-LNAs) correct alternative splicing abnormalities in myotonic dystrophy. *Nucleic Acids Res* 2015;43:3318-31.
- Konieczny P, Stepniak-Konieczna E and Sobczak K MBNL proteins and their target

- RNAs, interaction and splicing regulation. *Nucleic Acids Res* 2014;42:10873-87.
24. Fugier C, Klein AF, Hammer C, Vassilopoulos S, Ivarsson Y, Toussaint A, et al. Misregulated alternative splicing of BIN1 is associated with T tubule alterations and muscle weakness in myotonic dystrophy. *Nat Med* 2011;17:720-5.
 25. Savkur RS, Philips AV, Cooper TA. Aberrant regulation of insulin receptor alternative splicing is associated with insulin resistance in myotonic dystrophy. *Nat Genet* 2001;29:40-7.
 26. Wheeler TM, Lueck JD, Swanson M, Dirksen RT, Thornton CA. Correction of CIC-1 splicing eliminates chloride channelopathy and myotonia in mouse models of myotonic dystrophy. *J Clin Invest* 2007;117:3952-7.
 27. Cardani R, Giagnacovo M, Rossi G, Renna LV, Bugiardini E, Pizzamiglio C, et al. Progression of muscle histopathology but not of spliceopathy in myotonic dystrophy type 2. *Neuromuscul Disord* 2014;24:1042-53.
 28. Nakamori M, Sobczak K, Puwanant A, Welle S, Eichinger K, Pandya S, et al. Splicing biomarkers of disease severity in myotonic dystrophy. *Ann Neurol* 2013; 74:862-72.
 29. Renna LV, Cardani R, Botta A, Rossi G, Fossati B, Costa E, et al. Premature senescence in primary muscle cultures of myotonic dystrophy type 2 is not associated with p16 induction. *Eur J Histochem* 2014;58:2444.
 30. Bergström J. Muscle electrolytes in man. *Scand J Clin Lab Med* 1962;14:511-4.
 31. Tarnopolsky MA, Pearce E, Smith K, Lach B. Suction-modified Bergström muscle biopsy technique: experience with 13, 500 procedures. *Muscle Nerve* 2011; 43:717-25.
 32. Moxley RT 3rd, Meola G, Udd B, Ricker K. Report of the 84th ENMC workshop: PROMM (proximal myotonic myopathy) and other myotonic dystrophy-like syndromes: 2nd workshop. 13–15th October 2000, Loosdrecht: The Netherlands. *Neuromuscul Disord* 2002;12:306-17.
 33. Valaperta R, Sansone V, Lombardi F, Verdelli C, Colombo A, Valisi M, et al. Identification and characterization of DM1 patients by a new diagnostic certified assay: neuromuscular and cardiac assessments. *Biomed Res Int* 2013;2013:958510.
 34. Mathieu J, Boivin H, Meunier D, Gaudreault M, Bégin P. Assessment of a disease-specific muscular impairment rating scale in myotonic dystrophy. *Neurology* 2001;56:336-40.
 35. Dubowitz V. Muscle biopsy, p. 19-40. In: V Dubowitz (ed.) *A Practical Approach*. Bailliere Tindall, London: 1985.
 36. Vihola A, Bassez G, Meola G, Zhang S, Haapasalo H, Paetau A, et al. Histopathological differences of myotonic dystrophy type 1 (DM1) and PROMM/DM2. *Neurology* 2003;60:1854-7.
 37. Cardani R, Mancinelli E, Sansone V, Rotondo G, Meola G. Biomolecular identification of (CCTG)_n mutation in myotonic dystrophy type 2 (DM2) by FISH on muscle biopsy. *Eur J Histochem* 2004;48:437-42.
 38. Botta A, Rinaldi F, Catalli C, Vergani L, Bonifazi E, Romeo V, et al. The CTG repeat expansion size correlates with the splicing defects observed in muscles from myotonic dystrophy type 1 patients. *J Med Genet* 2008;45:639-46.
 39. Foff EP and Mahadevan MS. Therapeutics development in myotonic dystrophy type I. *Muscle Nerve* 2011;44:160-9.