


 Cite this: *RSC Adv.*, 2022, 12, 5517

Received 6th January 2022

Accepted 6th February 2022

DOI: 10.1039/d2ra00078d

rsc.li/rsc-advances

Highly efficient synthesis of 3,4-diarylbutadiene sulfones using Heck–Matsuda reaction†

 Olga V. Shurupova, Sergey A. Rzhhevskiy, , Lidiya I. Minaeva, , Maxim A. Topchiy  and Andrey F. Asachenko *

For the first time we describe a general method for the synthesis of previously not synthesized unsymmetrical 3,4-diarylbutadiene sulfones which can be stable convenient precursors for 2,3-diaryl-1,3-butadienes. Our method for arylation of butadiene sulfones *via* Heck–Matsuda reaction allows to obtain unsymmetrical 3,4-diarylbutadiene sulfones with a variety of alkyl, alkoxy, nitro, ethoxycarbonyl, perfluoroalkyl and halogen substituents (30 examples) in very good yields using readily available reagents and catalysts.

Introduction

Conjugated dienes are one of the most important precursors in organic synthesis that allow to obtain much more complex compounds using fewer synthetic steps.^{1–7} Also they are well known to be significant scaffolds in a wide range of natural products (such as trichostatin A, fuzanin C, modiolin and azoxymycin C) (Fig. 1).^{8,9} 2-Aryl- and 2,3-diaryl-1,3-butadienes have attracted the interest of organic chemists due to their ability to combine Diels–Alder^{10,11} and hetero-Diels–Alder cyclization.^{12,13} This type of conjugated unsaturated compound was also investigated in some other reactions, including hydrohydroxymethylation¹⁴ and hydroxymethylation¹⁵ catalyzed by iridium and ruthenium complexes respectively. Regarding all these facts the efficient and approachable preparation method for 2,3-diaryl-1,3-butadienes possessing certain regioselectivity

is of particular interest both from academic and practical points of view.

Among the possible ways to obtain 1,3-dienes, Wittig reactions,¹⁶ reduction of 1,3-diyne or 1,3-enyne,¹⁷ transition metal-catalyzed cross-coupling reactions¹⁸ and 1,2- or 1,4-elimination of allylic alcohols¹⁹ should be named in the first instance. Nevertheless, obtaining unsymmetrical (containing different aryl groups) 2,3-diaryl-1,3-butadienes is much less studied. The preparation of these important compounds can be proceeded using palladium-catalyzed cross-coupling of *p*-tolyl iodide with 3-silyl-3-phenylbuta-1,3-diene (40% yield)²⁰ or styryl triflates with styryl boronates (31–76% yield).²¹ Palladium-catalyzed Stille cross-coupling was applied to obtain an unsymmetrical 2,3-bis-(2-nitroaryl)-1,3-butadiene derivative in 61% yield.²² The two different hydrazones were assayed in cross-coupling reaction to give unsymmetrical dienes as well as symmetrical ones causing modest yields of necessary products.²³ Also the ene–yne cross-metathesis reaction was reported on one example to make 2,3-diphenyl-1,3-butadiene from ethylene as olefin substrate and internal alkyne.²⁴ Then this method was successfully applied to the variety of alkynes, but it became inefficient in case of substituted unsymmetrical phenylacetylene derivatives since *ortho*-substitution in aryl ring might cause the low reactivity of alkynes in the reaction²⁵ due to both steric hindrance and potential catalyst inhibition of by the coordination of the *ortho*-substitutes to the metal center. Obviously, a highly regioselective synthetic methodology for unsymmetrical 2,3-diaryl-1,3-butadienes from simple and readily available starting materials is significantly desirable.

The substituted 3-sulfolenes are well-known key precursors of not only different sulfolanes,^{26,27} polycyclic heterocycles²⁸ but also conjugated unsaturated compounds.^{29,30} The thermal extrusion of SO₂ (temperatures typically >100 °C) from these stable compounds provides corresponding 1,3-dienes.³¹




Fig. 1 Bioactive molecules with 1,3-diene motif.

A.V. Topchiev Institute of Petrochemical Synthesis, Russian Academy of Sciences, Leninsky Prospect 29, Moscow, 119991, Russia. E-mail: asachenko@ips.ac.ru

† Electronic supplementary information (ESI) available: Detailed experimental procedures, ¹H and ¹³C NMR, IR, HRMS data. See DOI: 10.1039/d2ra00078d



In our previous work we have developed simple and efficient method for preparation of 3-arylbutadiene sulfones *via* Heck–Matsuda reaction (Scheme 1).³²

Concerning our interest in Pd-catalyzed chemistry,^{33–38} for the first time we disclose an efficient and facile Pd-catalyzed cross-coupling reaction of 3-arylbutadiene sulfones with aryl-diazonium salts to construct 3,4-diarylbutadiene sulfones with high regioselectivity. Thus, our proposed method discovered the possibility to explore unsymmetrical diaryl sulfones and their transformations, especially substituted 1,3-dienes synthesis.


Results and discussion

3,4-Diarylbutadiene sulfones synthesis *via* Heck–Matsuda reaction


To investigate the possibility of 3-arylbutadiene sulfones arylation we have started using the conditions that had already shown the best results in the synthesis of 3-arylbutadiene sulfones³² (1 eq. of aryldiazonium tetrafluoroborate, 3 mol% Pd(OAc)₂ in methanol). On the first stage compound 3' (Scheme 2) is formed due to the arylpalladium syn addition to the C–C double bond from the less hindered face of the double bond anti to the Ph group followed by syn palladium hydride elimination resulting in the migration of the C–C bond³⁹ in the reaction product. The base induced double-bond isomerization in case of 3,4-diarylbutadiene sulfones require much harsher conditions than in case of 3-arylbutadiene sulfones maybe due to the sterical hindrance provided by an aryl group on C-3.⁴⁰ Refluxing with DBN (1,5-diazabicyclo(4.3.0)non-5-ene) in 1,4-dioxane turned out to be the most suitable system, which leads to formation the only thermodynamically stable 3,4-diarylbutadiene sulfones (Scheme 2).

However the incomplete conversion and moderate product yields caused us to investigate the optimal aryldiazonium salt and Pd(OAc)₂ amounts using the model reaction between 3-phenyl sulfolene and *p*-nitrophenyldiazonium tetrafluoroborate, that turned out to be 1.5 eq. and 5 mol% Pd(OAc)₂ respectively.

After that the impact of aryldiazonium salt structure was studied on the 6 examples of Heck–Matsuda reaction using the 3-phenyl sulfolene as a substrate (Table 1). In all cases the full conversion of 3-arylbutadiene sulfones took place on the cross-coupling stage, including the example with the sterically hindered *o*-methoxyphenyldiazonium tetrafluoroborate. The migration of double-bond always showed quantitative yield, but required more time in case of methoxy substituted products 3e and 3f (2–4 hours for 3a–d vs. 24 hours for 3e and 3f).



Scheme 1 Synthesis of 3-aryl-3-sulfolenes.




Scheme 2 3,4-Diarylbutadiene sulfones synthesis using Heck–Matsuda reaction.

To investigate the scope and the limitations of the considered method the influence of the aryl substitution in the starting 3-arylbutadiene sulfones in the Pd-catalyzed Heck–Matsuda reaction was explored (Table 2).


Series of 3-arylbutadiene sulfones bearing different electron-donating and electron-withdrawing groups were involved and corresponding 3,4-diarylbutadiene sulfones were obtained in good or excellent yields. To show the versatility of the conditions proposed else 18 examples (Table 3) of unsymmetrical 3,4-diarylbutadiene sulfones bearing different donor, acceptor or neutral substitutions on both aryl rings were obtained.

As it can be found from the Tables 2 and 3, introduction of the electron-withdrawing substitutions to the aryldiazonium salt (Table 3, 5f, 5j, 5k and 5r) lead to the slight yield decrease compared with the arylation agents with the electron-donating groups, even in the *ortho*-position (Table 3, 5a, 5d, 5g, 5i, 5l, 5n). On the contrary in case of the starting 3-arylbutadiene sulfones bearing electron-donating groups (Tables 2 and 3, 5k, 5o, 5s) the moderate yields were observed, while introducing EWG substitutions to sulfones lead to the yield increase. Therefore it can be concluded that the aryldiazonium salt should contain electron-donating substitution whereas the 3-arylbutadiene sulfones should bear the electron-withdrawing one to provide the better product yields.

Table 1 Arylation of 3-phenyl-3-sulfolene with different aryldiazonium salts^a



^a Reaction conditions: step 1: 2a (1 mmol), aryldiazonium salt (1.5 mmol), Pd(OAc)₂ (5 mol%), MeOH 5 ml, 2 h; step 2: DBN 0.2 ml, 1,4-dioxane 5 ml, 100 °C, 2–24 h.

Table 2 Scope of 3,4-diarylbutadiene sulfones^a


^a Reaction conditions: step 1: **2** (1 mmol), aryldiazonium salt (1.5 mmol), $\text{Pd}(\text{OAc})_2$ (5 mol%), MeOH 5 ml, 2 h; step 2: DBN 0.2 ml, $1,4\text{-dioxane}$ 5 ml, $100\text{ }^\circ\text{C}$, 2–24 h.

The 3,4-diaryl butadiene sulfones' reactivity investigation

Conversion of 3,4-diarylbutadiene sulfones to 2,3-diaryl-1,3-butadienes. As it has been already mentioned in the introduction, 3,4-diarylbutadiene sulfones can be easily converted to the corresponding 2,3-diaryl-1,3-butadienes *via* the SO_2 thermal extrusion from the sulfone solution at $100\text{--}150\text{ }^\circ\text{C}$.⁴¹ Since no one had ever produced unsymmetrical 3,4-diarylbutadiene sulfones before us, we felt it necessary to try them in the classical sulfone chemistry transformations. Some of the unsymmetrical 3,4-diarylbutadiene sulfones obtained in the first part of this work underwent this transformation in the 1,2-dichlorobenzene solution at $140\text{ }^\circ\text{C}$ resulting with the stable unsymmetrical 2,3-diaryl-1,3-butadienes with the high yields (Scheme 3, **8a–8c**). These 3 examples of 2,3-diaryl-1,3-butadienes have the different electron-donating and electron-withdrawing substituents in the phenyl rings and were synthesized for the first time. It is also worth mentioning that the whole scope of obtained 3,4-diarylbutadiene sulfones (Tables 2 and 3) may be considered as the precursors for the corresponding 2,3-diaryl-1,3-butadienes, and in most cases (except for Table 3, **5o42,43**) after SO_2 thermal extrusion previously unavailable 2,3-diaryl-1,3-butadienes may be originally synthesized.

[4 + 2] cycloaddition reactions. Cyclization reactions are of significant value in the synthesis of complex structures.²³ 3,4-Diarylbutadiene sulfones are shown to be the convenient and stable substrates in place of 1,3-dienes for cycloaddition reactions, most significantly the [4 + 2] Diels–Alder reaction, in various synthetic routes such as the synthesis of natural products,^{44,45} porphyrins^{46,47} and functionalized fullerenes.^{48,49}


Having explored the scope of 3,4-diarylbutadiene sulfones we have carried out a number of them with some well-known dienophiles *via* Diels–Alder reaction. Thus the cycloadduct between maleic anhydride and 3-(*p*-*tert*-butylphenyl)-4-phenyl-

Table 3 Scope of 3,4-diarylbutadiene sulfones^a

^a Reaction conditions: step 1: **2** (1 mmol), aryldiazonium salt (1.5 mmol), $\text{Pd}(\text{OAc})_2$ (5 mol%), MeOH 5 ml, 2 h; step 2: DBN 0.2 ml, $1,4\text{-dioxane}$ 5 ml, $100\text{ }^\circ\text{C}$, 2–24 h.

butadiene sulfone (**3b**) was obtained in good yield (76%) (Scheme 3, 7).

The reaction of 3-(*p*-*OME*-phenyl)-4-(*p*-*tert*-butylphenyl)-butadiene sulfones with dimethyl acetylenedicarboxylate results to the mixture of the cycloadduct and the product of its aromatization. The comprehensive aromatization of the



Scheme 3 Transformation of 3,4-diaryl-3-sulfolenes to different products.

mixture can be catalyzed by treatment with DDQ or palladium on activated charcoal (Scheme 3, 6).

We have also shown that unsymmetrical dienes can react regioselectively with asymmetric dienophiles *via* Diels–Alder reaction. *N*-Tosyl tetrahydropyridine **9** was obtained as the single regioisomer in good yield (Scheme 3).⁵⁰

Thus, the unsymmetrical diaryl sulfones showed good reactivity under Diels–Alder reaction conditions, which significantly expands the potential scope of polyaromatic compounds which can be used in liquid crystals, microelectronics, *etc.*^{51–53}

Conclusions

In summary, the facile, mild and highly regioselective synthesis of previously undescribed unsymmetrical 3,4-diarylbutadiene sulfones bearing donor, acceptor and sterical hindered substituents was developed. It was shown that this approach is applicable for obtaining different previously unavailable unsymmetrical 2,3-diaryl-1,3-butadienes in very good yield. For the first time the reactivity of 3,4-diarylbutadiene sulfones under Diels–Alder reactions conditions has been demonstrated, which makes it possible to explore new, more complicated polyaromatic compounds.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The study was supported by the Russian Science Foundation (project number 19-73-10185). Part of this work was carried out by Andrey F. Asachenko and Sergey A. Rzhavskiy, as part of the A. V. Topchiev Institute of Petrochemical Synthesis (TIPS) Russian Academy of Sciences (RAS) State Program. This work was performed using the equipment of the Shared Research

Center “Analytical center of deep oil processing and petrochemistry of TIPS RAS”.

Notes and references

- 1 F. Erver, J. R. Kuttner and G. Hilt, *J. Org. Chem.*, 2012, **77**, 8375–8385.
- 2 J.-A. Funel and S. Abele, *Angew. Chem., Int. Ed.*, 2013, **52**, 3822–3863.
- 3 M. Hirano, T. Okamoto, N. Komine and S. Komiya, *Organometallics*, 2012, **31**, 4639–4642.
- 4 S. Ogoshi, K.-i. Tonomori, M.-a. Oka and H. Kurosawa, *J. Am. Chem. Soc.*, 2006, **128**, 7077–7086.
- 5 Y. Wen, L. Huang and H. Jiang, *J. Org. Chem.*, 2012, **77**, 5418–5422.
- 6 M. A. Belaya, D. A. Knyazev, D. D. Borisov, R. A. Novikov and Y. V. Tomilov, *J. Org. Chem.*, 2021, **86**, 8089–8100.
- 7 M. A. Belaya, D. A. Knyazev, R. A. Novikov and Y. V. Tomilov, *Tetrahedron Lett.*, 2020, **61**, 151990.
- 8 K. Trisuwan, V. Rukachaisirikul, Y. Sukpondma, S. Preedanon, S. Phongpaichit, N. Rungjindamai and J. Sakayaroj, *J. Nat. Prod.*, 2008, **71**, 1323–1326.
- 9 M. DellaGreca, C. Di Marino, A. Zarrelli and B. D’Abrosca, *J. Nat. Prod.*, 2004, **67**, 1492–1495.
- 10 S. Kotha, M. Meshram and A. Tiwari, *Chem. Soc. Rev.*, 2009, **38**, 2065–2092.
- 11 S. Kotha and V. Seema, *Synlett*, 2011, **2011**, 2329–2334.
- 12 R. B. Lewis and J. Read de Alaniz, *Tetrahedron*, 2017, **73**, 4045–4051.
- 13 H. Srour, K. Abidi, Z. Sahli, S. Basker, N. Hamdi, M. Achard and C. Bruneau, *ChemCatChem*, 2011, **3**, 1876–1879.
- 14 K. D. Nguyen, D. Herkommer and M. J. Krische, *J. Am. Chem. Soc.*, 2016, **138**, 14210–14213.
- 15 T. Smejkal, H. Han, B. Breit and M. J. Krische, *J. Am. Chem. Soc.*, 2009, **131**, 10366–10367.
- 16 B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, 1989, **89**, 863–927.
- 17 K. N. Campbell and L. T. Eby, *J. Am. Chem. Soc.*, 1941, **63**, 216–219.
- 18 K. C. Nicolaou, P. G. Bulger and D. Sarlah, *Angew. Chem., Int. Ed. Engl.*, 2005, **44**, 4442–4489.
- 19 T. Kitahara, T. Matsuoka, H. Kiyota, Y. Warita, H. Kurata, A. Horiguchi and K. Mori, *Synthesis*, 1994, **1994**, 692–694.
- 20 Z. Ikeda, K. Oshima and S. Matsubara, *Org. Lett.*, 2005, **7**, 4859–4861.
- 21 D. Eom, S. Park, Y. Park, T. Ryu and P. H. Lee, *Org. Lett.*, 2012, **14**, 5392–5395.
- 22 N. H. Ansari, C. A. Dacko, N. G. Akhmedov and B. C. G. Söderberg, *J. Org. Chem.*, 2016, **81**, 9337–9349.
- 23 H. Jiang, L. He, X. Li, H. Chen, W. Wu and W. Fu, *Chem. Commun.*, 2013, **49**, 9218–9220.
- 24 G. Hilt and S. Roesner, *Synthesis*, 2011, **2011**, 662–668.
- 25 M. K. Abderrezak, Z. Kabouche, C. Bruneau and C. Fischmeister, *Catalysts*, 2017, **7**, 365.
- 26 K. Meng, J. Xia, Y. Wang, X. Zhang, G. Yang and W. Zhang, *Org. Chem. Front.*, 2017, **4**, 1601–1605.

- 27 Y. Kita, S. Hida, K. Higashihara, H. S. Jena, K. Higashida and K. Mashima, *Angew. Chem., Int. Ed.*, 2016, **55**, 8299–8303.
- 28 K. Ando and H. Takayama, *Heterocycles*, 1994, **37**, 1417–1439.
- 29 R. Bloch, C. Benecou and E. Guibé-Jampel, *Tetrahedron Lett.*, 1985, **26**, 1301–1304.
- 30 R. Bloch, D. Hassan and X. Mandard, *Tetrahedron Lett.*, 1983, **24**, 4691–4694.
- 31 M. G. Brant and J. E. Wulff, *Synthesis*, 2016, **48**, 1–17.
- 32 S. A. Rzhveskiy, V. N. Bogachev, L. I. Minaeva, G. K. Sterligov, M. S. Nechaev, M. A. Topchiy and A. F. Asachenko, *Mendeleev Commun.*, 2021, **31**, 548–549.
- 33 E. V. Bermesheva, A. I. Wozniak, F. A. Andreyanov, G. O. Karpov, M. S. Nechaev, A. F. Asachenko, M. A. Topchiy, E. K. Melnikova, Y. V. Nelyubina, P. S. Gribanov and M. V. Bermeshev, *ACS Catal.*, 2020, **10**, 1663–1678.
- 34 A. I. Wozniak, E. V. Bermesheva, N. N. Gavrilova, I. R. Ilyasov, M. S. Nechaev, A. F. Asachenko, M. A. Topchiy, P. S. Gribanov and M. V. Bermeshev, *Macromol. Chem. Phys.*, 2018, **219**, 1800323.
- 35 A. A. Ageshina, G. K. Sterligov, S. A. Rzhveskiy, M. A. Topchiy, G. A. Chesnokov, P. S. Gribanov, E. K. Melnikova, M. S. Nechaev, A. F. Asachenko and M. V. Bermeshev, *Dalton Trans.*, 2019, **48**, 3447–3452.
- 36 G. A. Chesnokov, A. A. Ageshina, A. V. Maryanova, S. A. Rzhveskiy, P. S. Gribanov, M. A. Topchiy, M. S. Nechaev and A. F. Asachenko, *Russ. Chem. Bull.*, 2020, **69**, 2370–2377.
- 37 S. A. Rzhveskiy, M. A. Topchiy, V. N. Bogachev, L. I. Minaeva, I. R. Cherkashchenko, K. V. Lavrov, G. K. Sterligov, M. S. Nechaev and A. F. Asachenko, *Mendeleev Commun.*, 2021, **31**, 409–411.
- 38 S. A. Rzhveskiy, M. A. Topchiy, V. N. Bogachev, A. A. Ageshina, L. I. Minaeva, G. K. Sterligov, M. S. Nechaev and A. F. Asachenko, *Mendeleev Commun.*, 2021, **31**, 478–480.
- 39 R. C. Larock, E. K. Yum and H. Yang, *Tetrahedron*, 1994, **50**, 305–321.
- 40 T. S. Chou, S. C. Hung and H. H. Tso, *J. Org. Chem.*, 1987, **52**, 3394–3399.
- 41 C.-W. Ko and T.-s. Chou, *J. Org. Chem.*, 1998, **63**, 4645–4653.
- 42 Y. Zhang, J. Li, X. Li and J. He, *Macromolecules*, 2014, **47**, 6260–6269.
- 43 X. Zhao, S. Zhu, F.-L. Qing and L. Chu, *Chem. Commun.*, 2021, **57**, 9414–9417.
- 44 T.-s. Chou, S.-J. Lee and N.-K. Yao, *Tetrahedron*, 1989, **45**, 4113–4124.
- 45 J. Leonard, S. P. Fearnley, M. R. Finlay, J. A. Knight and G. Wong, *J. Chem. Soc., Perkin Trans. 1*, 1994, **17**, 2359–2361.
- 46 O. Finikova, A. Cheprakov, I. Beletskaya and S. Vinogradov, *Chem. Commun.*, 2001, **3**, 261–262.
- 47 M. G. H. Vicente, A. C. Tomé, A. Walter and J. S. Cavaleiro, *Tetrahedron Lett.*, 1997, **38**, 3639–3642.
- 48 H. Ishida, H. Asaji, T. Hida, K. Itoh and M. Ohno, *Tetrahedron Lett.*, 2000, **41**, 2153–2156.
- 49 H. Ishida, K. Itoh, S. Ito, N. Ono and M. Ohno, *Synlett*, 2001, **2001**, 0296–0298.
- 50 R. Tomifuji, K. Maeda, T. Takahashi, T. Kurahashi and S. Matsubara, *Org. Lett.*, 2018, **20**, 7474–7477.
- 51 H. Bock, M. Rajaoarivelo, S. Clavaguera and É. Grelet, *Eur. J. Org. Chem.*, 2006, **2006**, 2889–2893.
- 52 T. Osawa, T. Kajitani, D. Hashizume, H. Ohsumi, S. Sasaki, M. Takata, Y. Koizumi, A. Saeki, S. Seki, T. Fukushima and T. Aida, *Angew. Chem., Int. Ed.*, 2012, **51**, 7990–7993.
- 53 J. Yin, H. Qu, K. Zhang, J. Luo, X. Zhang, C. Chi and J. Wu, *Org. Lett.*, 2009, **11**, 3028–3031.