

# BMJ Open Socioeconomic, religious, spiritual and health factors associated with symptoms of common mental disorders: a cross-sectional secondary analysis of data from Bhutan's Gross National Happiness Study, 2015

Gyambo Sithey,<sup>1</sup> Mu Li,<sup>1</sup> Li Ming Wen,<sup>1,2</sup> Patrick J Kelly,<sup>1</sup> Kelly Clarke<sup>3</sup>

**To cite:** Sithey G, Li M, Wen LM, *et al.* Socioeconomic, religious, spiritual and health factors associated with symptoms of common mental disorders: a cross-sectional secondary analysis of data from Bhutan's Gross National Happiness Study, 2015. *BMJ Open* 2018;**8**:e018202. doi:10.1136/bmjopen-2017-018202

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-018202>).

Received 19 June 2017  
Revised 13 December 2017  
Accepted 16 January 2018



<sup>1</sup>Sydney School of Public Health, The University of Sydney, Sydney, New South Wales, Australia

<sup>2</sup>Health Promotion Unit, Sydney Local Health District, Sydney, New South Wales, Australia

<sup>3</sup>Institute for Global Health, University College London, London, UK

**Correspondence to** Mr. Gyambo Sithey; [gsithey@yahoo.com](mailto:gsithey@yahoo.com)

## ABSTRACT

**Objective** Common mental disorders (CMDs) are a major cause of the global burden of disease. Bhutan was the first country in the world to focus on happiness as a state policy; however, little is known about the prevalence and risk factors of CMDs in this setting. We aim to identify socioeconomic, religious, spiritual and health factors associated with symptoms of CMDs.

**Design and setting** We used data from Bhutan's 2015 Gross National Happiness (GNH) Survey, a multistage, cross-sectional nationwide household survey. Data were analysed using a hierarchical analytical framework and generalised estimating equations.

**Participants** The GNH Survey included 7041 male and female respondents aged 15 years and above.

**Measures** The 12-item General Health Questionnaire was used to measure symptoms of CMDs. We estimated the prevalence of CMDs using a threshold score of  $\geq 12$ .

**Results** The prevalence of CMDs was 29.3% (95% CI 26.8% to 31.8%). Factors associated with symptoms of CMDs were: older age (65+) ( $\beta=1.29$ , 95% CI 0.57 to 2.00), being female ( $\beta=0.70$ , 95% CI 0.45 to 0.95), being divorced or widowed ( $\beta=1.55$ , 95% CI 1.08 to 2.02), illiteracy ( $\beta=0.48$ , 95% CI 0.21 to 0.74), low income ( $\beta=0.37$ , 95% CI 0.15 to 0.59), being moderately spiritual ( $\beta=0.61$ , 95% CI 0.34 to 0.88) or somewhat or not spiritual ( $\beta=0.76$ , 95% CI 0.28 to 1.23), occasionally considering karma in daily life ( $\beta=0.53$ , 95% CI 0.29 to 0.77) or never considering karma ( $\beta=0.80$ , 95% CI 0.26 to 1.34), having poor self-reported health ( $\beta=2.59$ , 95% CI 2.13 to 3.06) and having a disability ( $\beta=1.01$ , 95% CI 0.63 to 1.40).

**Conclusions** CMDs affect a substantial proportion of the Bhutanese population. Our findings confirm the importance of established socioeconomic risk factors for CMDs, and suggest a potential link between spiritualism and mental health in this setting.

## INTRODUCTION

WHO estimates that approximately 450 million people worldwide suffer from a common mental disorder (CMD) and 75% of

## Strengths and limitations of this study

- This is one of the first studies to explore risk factors for common mental disorders (CMDs) in Bhutan using a large nationally representative sample, including all age groups 15 years and above from both rural and urban communities.
- We estimate the prevalence of CMDs and identify potential socioeconomic, religious, spiritual and health risk factors for CMDs in this population.
- Due to the cross-sectional nature of the data, we were unable to establish causal relationships between risk factors and symptoms of CMDs.
- The General Health Questionnaire was used to assess symptoms of CMDs. This tool has not been validated in Bhutan though it has been used internationally and validated in other South Asian settings.
- Data on established risk factors of CMDs, such as alcohol, substance abuse and history of mental illness were not collected.

them live in low-income and middle-income countries (LMICs) where social inequalities are high.<sup>1</sup> CMDs are 'disorders which are commonly encountered in community settings, and whose occurrence signals a breakdown in normal functioning'.<sup>2</sup> WHO refers to CMDs as a range of anxiety and depressive disorders that impact on the mood or feelings of the affected person.<sup>3</sup> CMDs lead to disability and reduced work participation and productivity.<sup>4,5</sup> National-level epidemiological data on CMDs from LMICs are lacking, hindering mental health service and policy development.<sup>6</sup>

In some LMICs, risk factors for CMDs include poverty, lack of education, female sex, marital discord and divorce.<sup>7-9</sup> However, in other settings factors such as education

and income have been found to be less relevant.<sup>10</sup> Similarly, there have been heterogeneous findings regarding the association between rural versus urban residence and CMDs.<sup>11–13</sup> For example, a meta-analysis of studies from high-income countries, including the UK, the Netherlands, Canada and the USA, found a higher risk of CMDs in urban settings.<sup>14</sup> Religion and spiritual factors also contribute to mental health. A meta-analysis of 147 studies that involved nearly 100 000 subjects from mainly high-income settings found that religiousness was associated with fewer depressive symptoms.<sup>15</sup> Conversely, a study from mainland China, a middle-income setting, reported a higher risk of mental disorders among religious individuals.<sup>16</sup>

The aim of this study is to identify factors associated with symptoms of CMDs in Bhutan, a lower middle-income South Asian country. Bhutan was the first country to focus on happiness as a state policy, however, little is known about mental ill health in this setting. Two Gross National Happiness (GNH) Surveys were conducted in 2010 and 2015 to assess happiness.<sup>17</sup> Through these surveys, data were also collected on CMDs. Our study is a secondary analysis of data from the GNH Survey 2015 and aims to examine associations between socioeconomic, religious, spiritual and health factors, and symptoms of CMDs.

## METHODS

### Study setting and participants

Bhutan is a mainly Buddhist country in the Eastern Himalayas. Seventy per cent of its population live in rural areas<sup>18</sup> and 69% of the total population are farmers.<sup>19</sup> The national literacy rate is 63% among people aged 6 years and above.<sup>18</sup> We used data from the second GNH Survey, conducted between January and May 2015 by the Centre for Bhutan Studies and GNH Research, Royal Government of Bhutan. The GNH Survey is a household survey administered verbally via a trained enumerator to minimise potential literacy barriers. A nationally representative household-based sample was created using a stratified, multistage cluster sampling strategy. First, urban and rural areas within each district were identified as the main sampling strata. Second, the blocks were selected systematically using probability proportional to size. Third, households were randomly selected within the designated enumeration areas. Further details of data collection and management procedures are described in the 2015 GNH Survey report.<sup>17</sup>

### Measures

The GNH Survey measured symptoms of CMDs using the embedded 12-item General Health Questionnaire (GHQ-12). The GHQ-12 is a screening tool to detect minor psychological distress in the general population or in a non-clinical setting.<sup>20</sup> It has been extensively used in population studies in diverse cultures and contexts.<sup>20 21</sup> The tool consists of 12 items assessing


the severity of a symptom over the last 4 weeks using a 4-point Likert scale (0-1-2-3). For each participant, the 12 items were summarised into a single score by adding together responses for each item, giving an overall score that could range from 0 to 36. A higher score indicated poorer mental health. The GHQ-12 has not been validated in Bhutan, though it has been validated in other South Asian settings.<sup>22–24</sup>

A participant was classified as having a CMD if he or she had a total GHQ-12 score of 12 or more. This threshold was recommended by Goldberg *et al*<sup>21</sup> and has a sensitivity of 78.9% and specificity of 77.4%. Furthermore, Lundin *et al* reported that this threshold score provides the optimal trade-off between sensitivity and specificity for the GHQ-12 scored using the Likert method.<sup>25</sup> Hence, we have chosen a threshold of 11/12 (a score of 11 or less indicating the absence of CMD and a score of 12 or greater indicating the presence of CMD) to report the prevalence of CMDs in the population.

From the GNH Survey, we also extracted the data on social and economic status (age, gender, residence, income, marital status, literacy, occupation and household size), religion and spirituality (spirituality, karma and meditation) and health (self-reported health status, disability and walking distance to the nearest health centre) to assess the association of these factors with symptoms of CMDs. A person was considered literate if they were able to read and write in English, Dzongkha (National Language) or Nepali. Urban areas were defined as any settlement with a resident population of more than 5000 and more than 50% of the population being dependent on non-primary economic activities such as construction, the service sector and the civil sector. Rural areas were defined as having a smaller and more sparsely distributed population where the main economic activities are agriculture, livestock and forestry.<sup>26 27</sup> Household income measures included income earned by all household members from any sources, including salary, agricultural/livestock/forestry products and non-agricultural activities, adjusted for in-kind payments.<sup>17</sup> We dichotomised income based on mean annual household income of Nu.164,829 (US\$2535).<sup>18</sup>

Respondents were asked which religion they followed: Buddhism, Hinduism, Christianity, other or none. Spirituality was assessed by asking 'How spiritual do you consider yourself to be: very, moderately, somewhat or not at all?' We combined respondents who answered 'not at all' and 'somewhat' due to low numbers in these categories. The GNH Survey collected data on frequency of meditation. Data were also collected on belief in Buddhist concepts of karma by asking 'Do you consider karma in the course of your daily life: regularly, occasionally, rarely or not at all?'

Self-reported health status was determined by a single question 'In general, would you say your health is excellent, very good, good, fair or poor?' Due to a low number of cases and to be consistent with previous studies,<sup>28 29</sup> we combined responses into good health (ie, 'excellent', 'very good' and 'good') and poor health (ie, 'fair' and



**Figure 1** Analytical framework for the analysis. GHQ-12, General Health Questionnaire.

‘poor’). We also used data on disability and walking time to the nearest health centre.

### Analysis

Stata V.13 was used for all analyses. The prevalence of CMDs was estimated as the proportion of respondents classified as having a CMD. Due to the survey design, estimates were obtained using stratification of district and sample weights of the primary sampling units. Internal consistency for the GHQ-12 score was checked using Cronbach’s alpha.

Based on the global literature and on local concepts of mental illness in Bhutan,<sup>30–34</sup> we identified factors from the GNH Survey dataset that were potentially associated with CMDs. We organised factors into an analytical framework (figure 1). This framework draws on socioecological models for CMDs,<sup>35 36</sup> and distinguishes potentially distal socioeconomic factors (level 1) from more proximal religious and spiritual factors (level 2) and health-related factors (level 3).

We analysed GHQ-12 score as a continuous outcome and explored the association of potential risk factors with

GHQ-12 score through univariable and multivariable linear regression models using generalised estimating equations (GEE). GEE models were used to account for the clustering of respondents within the primary sampling units. We assumed an exchangeable correlation structure and applied sandwich estimator to obtain robust SEs. We also included district as an explanatory variable in all models. The regression coefficients ( $\beta$ ) denote the average change in GHQ-12 score.

All factors that showed an association at  $P \leq 0.25$  in the univariable analyses were included in the multivariable modelling stage.<sup>37</sup> Multivariable models included only those respondents with complete data. To guide multivariable analyses, we applied the analytical framework in figure 1, entering groups of factors into the model procedure in a hierarchical order. The socioeconomic variables (level 1) that showed an association at  $P \leq 0.25$  with GHQ-12 score in the univariable analyses were included in model 1. Model 2 included the socioeconomic variables that remained associated with GHQ-12 score at  $P \leq 0.1$  in model 1, plus religious and spiritual variables (level 2) that were associated at  $P \leq 0.25$ . Model 3 included variables that were associated with GHQ-12 score in model 2 ( $P \leq 0.1$ ), along with health-related factors (level 3) from the univariable analysis ( $P \leq 0.25$ ). This approach avoided reliance on statistical associations by considering the hierarchical inter-relationship between risk factors.<sup>38</sup> A  $P \leq 0.25$  was used for inclusion in the multivariable models to ensure that no important variables were missed. However, in the final model (model 3), variables were considered statistically significant if  $P < 0.01$ , due to the large sample size. Collinearity between variables was checked using variance inflation factors and found to be  $< 2$  for all variables.<sup>39</sup> The fit of the final model was checked using residual plots, which indicated model assumptions were adequately satisfied.

We used secondary data that had no identifying information.

## RESULTS

In total there were 7153 respondents, of whom 7041 had complete data. Table 1 presents key characteristics of respondents: 59% were women; 48% were farmers; 58% had no formal education; more than 60% were from the low-income group; three-quarters were married; 46% said they were 'very spiritual' and 72% lived in rural areas.

GHQ-12 scores were normally distributed with a mean of 9.4 (SD  $\pm 4.8$ ). Cronbach's alpha was 0.84 indicating satisfactory internal consistency. The estimated prevalence of CMDs was 29.3%, 95% CI 26.8% to 31.8%. Table 1 also shows the estimated prevalence for each potential factor and results from univariable analyses. All socioeconomic factors were associated ( $P \leq 0.25$ ) with GHQ-12 score. Among the religious and spiritual factors, degree of spirituality and consideration of karma in daily life were associated with higher GHQ-12 scores. Religion was not associated with GHQ-12 score ( $P = 0.67$ ). All

three health-related factors (self-reported health status, disability and walking time to the nearest health centre) were univariably associated with higher GHQ-12 scores ( $P < 0.01$ ).

Table 2 shows the results of the multivariable models. In model 1, older age, being female, being divorced or widowed, illiteracy, low income and occupation were associated with higher GHQ-12 scores, and were therefore retained in model 2. Rural residence and household size were not associated with GHQ-12 score in model 1. In model 2, level of spirituality and consideration of karma were associated with GHQ-12 score along with retained variables from model 1. In model 3, poor self-reported health status and disability were associated with higher GHQ-12 scores along with retained variables from model 2.

In the final model (model 3), factors independently associated ( $P < 0.01$ ) with higher GHQ-12 scores were: age, gender, marital status, literacy, occupation, income, spirituality, karma, self-reported health status and disability. GHQ-12 score was higher for older ( $\beta = 1.29$ , 95% CI 0.57 to 2.00), female ( $\beta = 0.70$ , 95% CI 0.45 to 0.95), divorced/widowed ( $\beta = 1.55$ , 95% CI 1.08 to 2.02), illiterate ( $\beta = 0.48$ , 95% CI 0.21 to 0.74) and low-income respondents ( $\beta = 0.37$ , 95% CI 0.15 to 0.59). Respondents who were moderately spiritual ( $\beta = 0.61$ , 95% CI 0.34 to 0.88), not or somewhat spiritual ( $\beta = 0.76$ , 95% CI 0.28 to 1.23), or who occasionally ( $\beta = 0.53$ , 95% CI 0.29 to 0.77) or never considered karma in daily life ( $\beta = 0.80$ , 95% CI 0.26 to 1.34) had significantly higher GHQ-12 scores. Respondents with poor self-reported health status ( $\beta = 2.59$ , 95% CI 2.13 to 3.06) and with disability ( $\beta = 1.01$ , 95% CI 0.63 to 1.40) also had higher GHQ-12 scores.

## DISCUSSION

In line with previous research, our study suggests that older age, being female, being divorced or widowed, illiteracy, occupation, low income, poor self-reported health status and having a disability are potential risk factors for CMDs in Bhutan.<sup>8 30 40</sup> Conversely, we found that increased spirituality and belief in karma were protective for CMDs.

We report a 29.3% prevalence of CMDs, similar to community-based studies in other South Asian settings.<sup>41 42</sup> Our estimate was higher compared with the estimate in the 2015 GNH Survey report (10.3%) because we used a lower GHQ-12 threshold score ( $\geq 12$  vs  $\geq 16$ ).<sup>17</sup> Our threshold score was selected based on findings from the Goldberg *et al* study<sup>21</sup> and the Lundin *et al* study.<sup>25</sup>

### Socioeconomic factors are important determinants of mental health status

Our findings support existing evidence that social and economic factors are independently associated with CMDs. We found that divorced or widowed respondents were at risk of CMDs. The prevalence of divorce in Bhutan is increasing, leading to a rise in matrimonial cases and single mothers seeking support for their families from

**Table 1** Univariable results of factors associated with GHQ-12 score, GNH Survey 2015

Variables	Prevalence of common mental disorders		Unadjusted mean change in GHQ-12 score		
	Participants* N	n	%† (95% CI)	β‡ (95% CI)	P value
<b>Income</b>					
High income	2711	640	24.0 (21.6 to 26.6)	Reference	<0.001
Low income	4330	1411	32.7 (30.9 to 34.6)	0.98 (0.73 to 1.22)	
<b>Residence</b>					
Urban	1987	472	24.7 (20.9 to 28.8)	Reference	0.006
Rural	5059	1579	31.4 (29.7 to 33.1)	0.71 (0.20 to 1.21)	
<b>Gender</b>					
Male	2923	776	26.9 (24.6 to 29.3)	Reference	<0.001
Female	4120	1273	30.9 (27.8 to 34.2)	0.94 (0.69 to 1.19)	
<b>Age</b>					
15–24	1113	297	26.9 (24.0 to 30.0)	Reference	<0.001
25–34	1800	439	24.3 (21.8 to 26.9)	–0.20 (–0.53 to 0.13)	
35–44	1562	402	26.5 (22.8 to 30.6)	–0.08 (–0.58 to 0.41)	
45–54	1183	376	31.9 (29.0 to 34.9)	0.49 (0.02 to 0.97)	
55–64	776	263	34.4 (30.9 to 38.1)	0.80 (0.25 to 1.34)	
65+	612	274	44.3 (39.8 to 49.0)	2.48 (1.77 to 3.18)	
<b>Marital status</b>					
Married	5328	1498	26.6 (25.6 to 31.0)	Reference	<0.001
Never married	1070	281	28.2 (24.0 to 29.3)	–0.20 (–0.66 to 0.26)	
Divorced/widowed	648	272	42.4 (38.1 to 46.7)	2.35 (1.82 to 2.88)	
<b>Literacy</b>					
Literate	3578	878	25.0 (22.7 to 27.4)	Reference	<0.001
Illiterate	3468	1173	33.9 (31.4 to 36.5)	1.28 (1.01 to 1.56)	
<b>Occupation</b>					
Farmers	3377	1116	33.2 (31.5 to 34.8)	Reference	<0.001
Home makers	1177	327	28.2 (23.4 to 33.6)	–0.17 (–0.68 to 0.33)	
Students	545	145	27.0 (23.9 to 30.4)	–1.01 (–1.46 to –0.55)	
Civil/business§	1541	323	21.7 (18.8 to 24.9)	–1.33 (–1.63 to –1.03)	
Others	405	140	34.2 (28.7 to 40.1)	0.51 (–0.06 to 1.08)	
<b>Household size</b>					
3–4 members	2491	712	29.0 (25.8 to 32.3)	Reference	0.060
1–2 members	910	302	33.0 (29.6 to 36.6)	0.60 (0.16 to 1.04)	
5–6 members	2453	688	28.1 (25.6 to 30.7)	0.03 (–0.23 to 0.28)	
≥7 members	1190	347	29.3 (26.4 to 32.5)	0.05 (–0.32 to 0.43)	
<b>Religion</b>					
Buddhist	5882	1679	28.7 (26.5 to 31.1)	Reference	0.670
Hindu	1000	331	33.0 (28.9 to 37.5)	0.03 (–0.32 to 0.38)	
Others	163	41	25.1 (18.6 to 33.0)	–0.27 (–0.92 to 0.38)	
<b>Spirituality</b>					
Very	3262	889	27.3 (25.1 to 29.7)	Reference	<0.001
Moderately	3174	960	30.3 (27.1 to 33.7)	0.51 (0.27 to 0.74)	
Not at all	610	202	33.4 (29.6 to 37.4)	0.88 (0.44 to 1.32)	
<b>Meditation</b>					
Yes	1291	390	30.7 (28.0 to 3.5)	Reference	0.190
Never	5755	1661	28.9 (26.3 to 31.7)	0.21 (–0.10 to 0.51)	

Continued

Table 1 Continued

Variables	Participants*		Prevalence of common mental disorders		Unadjusted mean change in GHQ-12 score	
	N	n	%† (95% CI)	β‡ (95% CI)	P value	
Consideration of karma in daily life						
Regularly	4048	1107	27.5 (24.9 to 30.3)	Reference		
Occasionally	2219	685	30.9 (27.6 to 34.4)	0.68 (0.43 to 0.94)		
Never	776	258	33.3 (29.7 to 37.0)	0.98 (0.46 to 1.50)		
Self-reported health status						
Good health	6315	1660	26.4 (24.2 to 28.7)	Reference	<0.001	
Poor health	731	391	54.1 (48.9 to 59.3)	3.58 (3.07 to 4.09)		
Disability						
No	5960	1601	27.0 (24.6 to 29.5)	Reference	<0.001	
Yes	1084	448	42.1 (38.6 to 45.7)	2.28 (1.89 to 2.68)		
Walking distance to health centre (min)						
≤30	3529	929	26.6 (24.3 to 29.1)	Reference	<0.001	
31–60	1379	408	29.5 (25.4 to 34.1)	0.21 (–0.15 to 0.57)		
≥61	2137	714	33.5 (30.9 to 36.2)	0.78 (0.43 to 1.13)		

\*Missing values for all factors were ≤112.

†Percentage may not match n/N due to adjustment for the survey design using Stata's SVY command.

‡The regression coefficients (β) denote the average change in GHQ-12 score.

§Includes civil servants, local government, corporate, business and armed forces.

GHQ-12, General Health Questionnaire; GNH, Gross National Happiness.

local non-governmental organisations.<sup>43 44</sup> Social stigma associated with divorce and court proceedings, coupled with the financial stress of raising a child as a single parent could partly explain the link between divorce and CMDs in Bhutan.

We found that older respondents as well as those who are illiterate or have a low income are more at risk of CMDs, in line with studies from other LMICs.<sup>7 8 30 40 45 46</sup> In Bhutan, 37% of adults are illiterate, however, this is lower among younger generations due to improved utilization of educational services. Illiteracy may be a marker of relative poverty and marginalisation, as well as low utilization of health and education services. WHO reports that CMDs are 1.5–2 times more prevalent among low-income groups.<sup>47</sup> This may be because people living in poverty lack the financial means to education and employment opportunities, perpetuating a negative cycle between poverty and CMDs.<sup>47</sup> Older people may be more at risk of CMDs compared with other age groups due to increased social isolation and susceptibility to non-communicable disease.<sup>48</sup>

We did not find any association between residence (rural/urban) and CMDs. This could be due to the massive rural to urban migration in Bhutan over recent years,<sup>49</sup> masking any potential association. It could also be due to the inclusion of other more distal markers of socioeconomic status in the model, such as income and occupation. Findings from other settings have been heterogeneous. Some studies in USA and Australia found no difference in risk of CMDs between rural and urban areas.<sup>50 51</sup> Studies in the UK<sup>14 52</sup> and Canada<sup>46 53</sup> reported

higher rates of CMDs in urban areas compared with rural areas, whereas studies in China<sup>54</sup> and India<sup>55</sup> reported lower or similar rates of CMDs in rural and urban areas. Such mixed findings could be due to different definitions of rural and urban areas which may incorporate population density,<sup>46</sup> remoteness and accessibility to health services.<sup>13 53</sup>

### Are spirituality and religious involvement beneficial for mental health?

We found that spirituality was associated with higher GHQ-12 scores. This is consistent with findings from other studies suggesting that spirituality is associated with mental health.<sup>56–58</sup> We also found that respondents who occasionally and never considered karma in their daily lives reported higher GHQ-12 scores compared with respondents who regularly considered karma. Regular consideration of karma may relate to an individual's level of religious involvement. Raphael *et al* found a significant positive association between religious involvement and mental health in a review of 43 studies.<sup>32</sup> Similarly, a recent reviews reported that religious participation was inversely associated with depressive symptoms, anxiety, stress and suicide.<sup>59 60</sup>

Bhutan is a predominantly Buddhist country and more than 90% of its population report that they are spiritual.<sup>17</sup> In this setting, spirituality and religious involvement may promote mental health through supportive faith-based community networks, and by providing individuals with a sense of inclusion and community acceptance, as well as meaning and goals for their lives.<sup>57 59 61</sup> Religious

**Table 2** Multivariable analyses of factors associated with GHQ-12 score, GNH Survey 2015

Socioeconomic factors	Model 1 (Socioeconomic factors)		Model 2 (Socioeconomic, spiritual and religious factors)		Model 3 (Socioeconomic, spiritual, religious and health factors)	
	Mean change $\beta^*$ (95% CI)	P value	Mean change $\beta^*$ (95% CI)	P value	Mean change $\beta^*$ (95% CI)	P value
Age (years)		<0.001		<0.001		<0.001
15–24	Reference		Reference		Reference	
25–34	–0.28 (–0.73 to 0.18)		–0.18 (–0.64 to 0.28)		–0.28 (–0.73 to 0.16)	
35–44	–0.21 (–0.73 to 0.32)		–0.05 (–0.56 to 0.46)		–0.28 (–0.77 to 0.21)	
45–54	0.24 (–0.34 to 0.82)		0.47 (–0.12 to 1.06)		0.08 (–0.48 to 0.65)	
55–64	0.52 (–0.11 to 1.14)		0.80 (0.18 to 1.42)		0.32 (–0.26 to 0.90)	
65+	1.90 (1.14 to 2.65)		2.24 (1.46 to 3.02)		1.29 (0.57 to 2.00)	
Gender		<0.001		<0.001		<0.001
Male	Reference		Reference		Reference	
Female	0.84 (0.59 to 1.10)		0.82 (0.57 to 1.08)		0.70 (0.45 to 0.95)	
Marital status		<0.001		<0.001		<0.001
Married	Reference		Reference		Reference	
Never married	0.26 (–0.23 to 0.74)		0.18 (–0.30 to 0.66)		0.13 (–0.34 to 0.60)	
Divorced/widowed	1.64 (1.12 to 2.17)		1.65 (1.13 to 2.17)		1.55 (1.08 to 2.02)	
Literacy		<0.001		<0.001		<0.001
Literate	Reference		Reference		Reference	
Illiterate	0.61 (0.33 to 0.88)		0.57 (0.29 to 0.84)		0.48 (0.21 to 0.74)	
Occupation		0.002		<0.001		0.003
Farmers	Reference		Reference		Reference	
Home makers	0.02 (–0.43 to 0.47)		0.04 (–0.39 to 0.48)		–0.01 (–0.46 to 0.43)	
Students	0.02 (–0.83 to 0.16)		–0.38 (–0.88 to 0.12)		–0.42 (–0.92 to 0.09)	
Civil/business†	0.02 (–0.67 to –0.01)		–0.35 (–0.66 to –0.04)		–0.31 (–0.62 to –0.00)	
Others	0.02 (0.13 to 1.31)		0.73 (0.16 to 1.30)		0.64 (0.10 to 1.17)	
Income		<0.001		<0.001		<0.001
High income	Reference		Reference		Reference	
Low income	0.45 (0.23 to 0.68)		0.44 (0.22 to 0.67)		0.37 (0.15 to 0.59)	
Residence		0.507				
Urban	Reference					
Rural	0.12 (–0.24 to 0.48)					
Household size	0.03 (–0.03 to 0.08)	0.374				
Religious and spiritual factors						
Spirituality				<0.001		<0.001
Very spiritual			Reference		Reference	
Moderately spiritual			0.58 (0.30 to 0.87)		0.61 (0.34 to 0.88)	
No and somewhat			0.79 (0.29 to 1.29)		0.76 (0.28 to 1.23)	
Consider karma in daily life				<0.001		<0.001
Regularly			Reference		Reference	
Occasionally			0.52 (0.28 to 0.77)		0.53 (0.29 to 0.77)	
Never			0.81 (0.26 to 1.35)		0.80 (0.26 to 1.34)	
Meditation				0.146		
Yes			Reference			
No			–0.22 (–0.52 to 0.08)			
Health-related factors						
Self-reported health status						<0.001

Continued

Table 2 Continued

Socioeconomic factors	Model 1 (Socioeconomic factors)		Model 2 (Socioeconomic, spiritual and religious factors)		Model 3 (Socioeconomic, spiritual, religious and health factors)	
	Mean change $\beta^*$ (95% CI)	P value	Mean change $\beta^*$ (95% CI)	P value	Mean change $\beta^*$ (95% CI)	P value
Good health					Reference	
Poor health					2.59 (2.13 to 3.06)	
Disability						<0.001
No disability					Reference	
Disability					1.01 (0.63 to 1.40)	
Walking distance to health centre (min)						0.155
≤30					Reference	
31–60					−0.01 (−0.30 to 0.32)	
≥61					0.30 (0.01 to 0.60)	

\*The regression coefficients ( $\beta$ ) denote the average change in GHQ-12 score.

†Includes civil servants, local government, corporate, business and arm force.  
GHQ-12, General Health Questionnaire; GNH, Gross National Happiness.

involvement is found to be associated with a better ability to cope with stress, and depression, suicide, anxiety and substance abuse.<sup>61</sup> In Bhutan, Buddhist values encompass acceptance, forgiveness and law of karma, which could help to build tolerance and reduce tensions in communities, with benefits for individual mental health. On average, people spend 51 min per day on religious activities.<sup>17</sup> Since the majority of the population is involved in religious activities, the national mental health strategy and action plan (2015–2023) includes working with the monastic and religious institutions in the country.<sup>62</sup> Spiritual beliefs and practices may enable people to face difficulties and provide guidelines for individuals to live and work together. In other words, development policies need to consider religious amenities like temples, monastic schools and access to spiritual leaders as basic consideration in any settlement plans.

Our study did not find an association between frequency of meditation and GHQ-12 score. This may be due to the small proportion (7.5%) of respondents who reported meditating. Meditation is a higher spiritual Buddhist practise uncommon among lay people.

### Interdependency of physical and mental health

In line with several previous studies, we found that individuals with poor self-reported health status or a disability had higher GHQ-12 scores.<sup>46 63–65</sup> Due to the cross-sectional nature of the data, it is not possible to determine the direction of effect between physical health and mental health. However, a bi-directional relationship in which mental ill health exacerbates physical illness which further impacts on mental health is highly plausible.<sup>66</sup> At present, there is no legislation or national policy on disability in Bhutan.<sup>67</sup> Findings from our study suggest that any future disability initiative in this setting should consider incorporating a mental health component.

Our study identifies possible targets for mental health promotion strategies, which could be delivered by non-specialised mental health workers. Potential strategies include provision of mental health support to older people, and mental health literacy programmes for non-specialist health workers to aid recognition, management and prevention of CMDs among individuals with poor general health.<sup>68 69</sup> In rural Bhutan, task shifting to non-specialised health workers like village health volunteers will be essential in providing access to mental health services. At present, there are only four psychiatrists in the country and only 1% of the total expenditure of health is directed towards mental health.<sup>70</sup> This study highlights the need for more research to help advocate for increased resources and political commitment for a national mental health programme.<sup>71 72</sup>

### Strengths and limitations

The main strengths of this study are the large nationally representative sample including all age groups of 15 years and above from both rural and urban communities of Bhutan. In addition, a range of social, economic, religious and health factors were included in the multivariable analysis. However, the study has several limitations. First, it used cross-sectional data making it impossible to establish causal relationships. Second, the GHQ-12 has not been validated in Bhutan, however we used a continuous outcome to conduct the analyses in order to avoid over-reliance on a non-validated threshold score. The GHQ-12 is a self-reported screening tool. Although it is not diagnostic of CMDs, it is a feasible approach to assess mental health in a large national survey. The prevalence of CMDs reported in this study is an estimate due to the lack of a locally validated threshold score. Third, the GHQ-12 was not translated into the local language in the questionnaire. Enumerators translated the items from



English into the appropriate language for each respondent and this could potentially have led to some inconsistencies in translation. Finally, data on other established risk factors of CMDs, such as alcohol, substance abuse and history of mental illness were not collected.

## CONCLUSIONS

Findings from this study highlight the importance of established socioeconomic factors of CMDs in Bhutan and suggest that religious involvement and spirituality may be protective factors for mental health in this setting. Further studies are needed to understand causal pathways to CMDs and to provide evidence to support mental health policy decisions and investment.

**Acknowledgements** The authors would like to thank Dasho Karma Ura, Dorji Penjore and Tshoki Zangmo of the Centre for Bhutan Studies and GNH Research, Royal Government of Bhutan, for providing the data and rendering all possible support.

**Contributors** GS, ML and LMW conceived and designed the concept of the paper. GS analysed the data and wrote the manuscript. PJK and LMW contributed to the data analysis and to the interpretation of the results. ML and KC contributed to the content and editing of the manuscript. ML supervised the whole process. All authors reviewed and approved the final manuscript.

**Funding** The 2015 GNH study was funded by the Royal Government of Bhutan and Japan International Cooperation Agency.

**Competing interests** None declared.

**Patient consent** Obtained.

**Ethics approval** The National Statistics Bureau of the Royal Government of Bhutan reviewed and approved the study protocol.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

- World Health Organization. *World health report*. Geneva: World Health Organization, 2001.
- Goldberg DP, Huxley P. *Common mental disorders: a bio-social model*. New York; London: Tavistock/Routledge, 1992.
- World Health Organization. *Depression and other common mental disorders: Global health estimates*. Geneva: World Health Organization, 2017.
- Patel V, Chisholm D, Parikh R, et al. DCP MNS Author Group. Addressing the burden of mental, neurological, and substance use disorders: key messages from Disease Control Priorities, 3rd edition. *Lancet* 2016;387:1672–85.
- Chisholm D, Sweeny K, Sheehan P, et al. Scaling-up treatment of depression and anxiety: a global return on investment analysis. *Lancet Psychiatry* 2016;3:415–24.
- Saxena S, Paraje G, Sharan P, et al. The 10/90 divide in mental health research: trends over a 10-year period. *Br J Psychiatry* 2006;188:81–2.
- Patel V, Kleinman A. Poverty and common mental disorders in developing countries. *Bull World Health Organ* 2003;81:609–15.
- Lund C, Breen A, Flisher AJ, et al. Poverty and common mental disorders in low and middle income countries: A systematic review. *Soc Sci Med* 2010;71:517–28.
- Affifi TO, Cox BJ, Enns MW. Mental health profiles among married, never-married, and separated/divorced mothers in a nationally representative sample. *Soc Psychiatry Psychiatr Epidemiol* 2006;41:122–9.
- Das J, DoQT, Friedman J, et al. Mental health and poverty in developing countries: revisiting the relationship. *Soc Sci Med* 2007;65:467–80.
- Peen J, Schoevers RA, Beekman AT, et al. The current status of urban-rural differences in psychiatric disorders. *Acta Psychiatr Scand* 2010;121:84–93.
- Caldwell TM, Jorm AF, Dear KB. Suicide and mental health in rural, remote and metropolitan areas in Australia. *Med J Aust* 2004;181:S10–14.
- Murray G, Judd F, Jackson H, et al. Rurality and mental health: the role of accessibility. *Aust N Z J Psychiatry* 2004;38:629–34.
- Paykel ES, Abbott R, Jenkins R, et al. Urban-rural mental health differences in great Britain: findings from the national morbidity survey. *Psychol Med* 2000;30:269–80.
- Smith TB, McCullough ME, Poll J. Religiousness and depression: evidence for a main effect and the moderating influence of stressful life events. *Psychol Bull* 2003;129:614–36.
- Wang Z, Koenig HG, Zhang Y, et al. Religious involvement and mental disorders in mainland china. *PLoS One* 2015;10:e0128800.
- Centre for Bhutan Studies & GNH Research. *2015 GNH Survey Report. A compass towards a just and harmonious society*. Thimphu, Bhutan: Centre for Bhutan Studies & GNH Research, Bhutan, 2016.
- National Statistics Bureau Bhutan and Asian Development Bank. *Bhutan Living Standards Survey*. Philippines and Bhutan: National Statistics Bureau, Bhutan and Asian Development Bank, 2012.
- Katwal T, Dorji S, Dorji R, et al. Community Perspectives on the On-Farm Diversity of Six Major Cereals and Climate Change in Bhutan. *Agriculture* 2015;5:2–16.
- Goldberg D, Williams P. *A user's guide to the General Health Questionnaire NFER-NELSON*: Windsor, 1988.
- Goldberg DP, Gater R, Sartorius N, et al. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol Med* 1997;27:191–7.
- Koirala N, Regmi S, Sharma V, et al. Sensitivity and validity of the General Health Questionnaire (GHQ-12) in a rural community setting in Nepal. *Nepalese J Psychiatry* 1999;1:34–40.
- Bandyopadhyay G, Sinha S, Sen B, et al. Validity of General Health Questionnaire (GHQ-36/GHQ-12) in the psychiatric O.P.D. of a general hospital—a pilot study. *Int J Soc Psychiatry* 1988;34:130–4.
- Montazeri A, Harirchi AM, Shariati M, et al. The 12-item General Health Questionnaire (GHQ-12): translation and validation study of the Iranian version. *Health Qual Life Outcomes* 2003;1:66.
- Lundin A, Hallgren M, Theobald H, et al. Validity of the 12-item version of the General Health Questionnaire in detecting depression in the general population. *Public Health* 2016;136:66–74.
- Schuelka MJ, Maxwell T. *Education in bhutan: culture, schooling, and gross national happiness*. Germany: Springer, 2016.
- Royal Government of Bhutan. *The thromde act of Bhutan*. Bhutan: National Assembly of Bhutan, 2007.
- Wu S, Wang R, Zhao Y, et al. The relationship between self-rated health and objective health status: a population-based study. *BMC Public Health* 2013;13:320.
- Sithey G, Wen LM, Kelly P, et al. Association between Sleep Duration and Self-Reported Health Status: Findings from the Bhutan's Gross National Happiness Study. *J Clin Sleep Med* 2017;13:33–8.
- World Health Organization and Calouste Gulbenkian Foundation. *Social determinants of mental health*. Geneva: World Health Organization, 2014.
- Weber SR, Pargament KI. The role of religion and spirituality in mental health. *Curr Opin Psychiatry* 2014;27:358–63.
- Bonelli RM, Koenig HG. Mental disorders, religion and spirituality 1990 to 2010: a systematic evidence-based review. *J Relig Health* 2013;52:657–73.
- Calabrese JD, Dorji C. Traditional and modern understandings of mental illness in Bhutan: Preserving the benefits of each to support Gross National Happiness. *Journal of Bhutan Studies* 2014;30.
- Tshomo D. *Focus is on the rising menace of depression*. Thimphu, Bhutan: Kuensel, 2017.
- Clarke K, Saville N, Shrestha B, et al. Predictors of psychological distress among postnatal mothers in rural Nepal: a cross-sectional community-based study. *J Affect Disord* 2014;156:76–86.
- Shidhaye R, Patel V. Association of socio-economic, gender and health factors with common mental disorders in women: a

- population-based study of 5703 married rural women in India. *Int J Epidemiol* 2010;39:1510–21.
37. Sithey G, Wen LM, Kelly P, et al. Association between Sleep Duration and Self-Reported Health Status: Findings from the Bhutan's Gross National Happiness Study. *J Clin Sleep Med* 2017;13:33–38.
  38. Victora CG, Huttly SR, Fuchs SC, et al. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* 1997;26:224–7.
  39. O'brien RM. A caution regarding rules of thumb for variance inflation factors. *Qual Quant* 2007;41:673–90.
  40. Seedat S, Scott KM, Angermeyer MC, et al. Cross-national associations between gender and mental disorders in the World Health Organization World Mental Health Surveys. *Arch Gen Psychiatry* 2009;66:785–95.
  41. Khattri JB, Poudel BM, Thapa P, et al. An epidemiological study of psychiatric cases in a rural community of nepal. *Nepal Journal of Medical Sciences* 2013;2:52–6.
  42. Islam MM, Ali M, Ferroni P, et al. Prevalence of psychiatric disorders in an urban community in Bangladesh. *Gen Hosp Psychiatry* 2003;25:353–7.
  43. Bhutan Broadcasting Service. *Bhutan sees increasing divorce*. Thimphu, Bhutan: Bhutan Broadcasting Service, 2015.
  44. Namgyal G. *Dzongkhag court sees rise in matrimonial cases, people blame development Bhutan Oberver*, 2011.
  45. Patel V. Mental health in low- and middle-income countries. *Br Med Bull* 2007;81-82:81–96.
  46. Romans S, Cohen M, Forte T. Rates of depression and anxiety in urban and rural Canada. *Soc Psychiatry Psychiatr Epidemiol* 2011;46:567–75.
  47. World Health Organization. *Breaking the vicious cycle between mental ill-health and poverty*. Geneva: Mental health core to development information sheet, 2007. sheet 1.
  48. World Health Organization. *Risk to mental health: an overview of vulnerabilities and risk factors. Background paper by who secretariat for the development of a comprehensive mental health action plan*. Geneva: World Health Organization, 2012.
  49. National Statistic Bureau and World Bank. *Bhutan poverty assessment 2014*. Bhutan: National Statistic Bureau, Royal Government of Bhutan, 2014.
  50. Breslau J, Marshall GN, Pincus HA, et al. Are mental disorders more common in urban than rural areas of the United States? *J Psychiatry Res* 2014;56:50–5.
  51. Andrews G, Henderson S, Hall W. Prevalence, comorbidity, disability and service utilisation. Overview of the Australian National Mental Health Survey. *Br J Psychiatry* 2001;178:145–53.
  52. Weich S, Twigg L, Lewis G. Rural/non-rural differences in rates of common mental disorders in Britain: Prospective multilevel cohort study. *Br J Psychiatry* 2006;188:51–7.
  53. Wang JL. Rural-urban differences in the prevalence of major depression and associated impairment. *Soc Psychiatry Psychiatr Epidemiol* 2004;39:19–25.
  54. Phillips MR, Zhang J, Shi Q, et al. Prevalence, treatment, and associated disability of mental disorders in four provinces in China during 2001-05: an epidemiological survey. *Lancet* 2009;373:2041–53.
  55. Nayak MB, Patel V, Bond JC, et al. Partner alcohol use, violence and women's mental health: population-based survey in India. *Br J Psychiatry* 2010;196:192–9.
  56. Lee JS, Auyeung TW, Leung J, et al. Long sleep duration is associated with higher mortality in older people independent of frailty: a 5-year cohort study. *J Am Med Dir Assoc* 2014;15:649–54.
  57. Moreira-Almeida A, Neto FL, Koenig HG. Religiousness and mental health: a review. *Rev Bras Psiquiatr* 2006;28:242–50.
  58. Brown DR, Carney JS, Parrish MS, et al. Assessing Spirituality: The Relationship Between Spirituality and Mental Health. *J Spiritual Ment Health* 2013;15:107–22.
  59. AbdAleati NS, Zaharim NM, Mydin YO. *Religiousness and mental health: systematic review study*. Germany: Springer, 2016.
  60. Gupta U, Gupta BS. Religious Involvement, Well-being and Mental Health in Men and Women. *Journal of Psychosocial Research* 2014;9:179.
  61. Koenig HG. Research on religion, spirituality, and mental health: a review. *Can J Psychiatry* 2009;54:283–91.
  62. Ministry of Health. *Mental Health for All: Bhutan mental health strategy and action plan (2015-2023)*. Bhutan: Mental Health Program, Department of Public Health, Ministry of Health, Royal Government of Bhutan, 2015.
  63. Ahmad F, Hhajj AK, Stewart DE, et al. Single item measures of self-rated mental health: a scoping review. *BMC Health Serv Res* 2014;14:398.
  64. Prince M, Patel V, Saxena S, et al. No health without mental health. *The Lancet* 2007;370:859–77.
  65. Mulsant BH, Ganguli M, Seaberg EC. The relationship between self-rated health and depressive symptoms in an epidemiological sample of community-dwelling older adults. *J Am Geriatr Soc* 1997;45:954–8.
  66. Honey A, Emerson E, Llewellyn G, et al. *Mental health and disability. International Encyclopedia of Rehabilitation*. New York: State University of New York Press, 2013.
  67. National Statistic Bureau. *Health MoEaMoTwo-staged child disability study among children 2-9 years*. In: . eds. Thimphu: National Statistic Bureau, 2012.
  68. Kumar ND, Durham Judith C, Kk L. Balancing traditional beliefs and medical science: mental health care in Bhutan. *Bhutan Health Journal* 2015;1:66–9.
  69. Petersen I, Evans-Lacko S, Semrau M, et al. Promotion, prevention and protection: interventions at the population- and community-levels for mental, neurological and substance use disorders in low- and middle-income countries. *Int J Ment Health Syst* 2016;10:30.
  70. Pelzang R. *Mental health care in Bhutan: policy and issues*. Geneva: WHO, 2012.
  71. Lhengye Zhungtshog. *Suicide prevention in Bhutan-A three year action plan (July 2015- June 2018)*. Bhutan: Lhengye Zhungtshog, The Cabinet of Ministers, Royal Government of Bhutan, 2015. 74th Session.
  72. Ministry of Health. *Mental health for all: Bhutan mental health strategy and action plan (2015-2023)*. Bhutan: Mental Health Program, Department of Public Health, Ministry of Health, Royal Government of Bhutan, 2015.