

Corrigendum

Single-nucleotide-resolution mapping of DNA gyrase cleavage sites across the *Escherichia coli* genome

Dmitry Sutormin^{1,2}, Natalia Rubanova¹, Maria Logacheva¹, Dmitry Ghilarov^{1,3,*} and Konstantin Severinov^{1,4,*}

¹Centre for Life Sciences, Skolkovo Institute of Science and Technology, 143026 Moscow, Russia, ²Department of Bioengineering and Bioinformatics, Lomonosov Moscow State University, 119991 Moscow, Russia, ³Malopolska Centre of Biotechnology, Jagiellonian University, 30387 Cracow, Poland and ⁴Waksman Institute for Microbiology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854, USA

Nucleic Acids Research, 2018, <https://doi.org/10.1093/nar/gky1222>

The Authors apologize for an error in reference (98) which should be:

REFERENCE

98. Keane, O.M. and Dorman, C.J. (2003) The *gyr* genes of *Salmonella enterica* serovar Typhimurium are repressed by the factor for inversion stimulation, *Fis. Mol. Gen. Genomics*, **270**, 56–65.

*To whom correspondence should be addressed. Tel: +1 732 445 6095; Fax: +1 732 445 5735; Email: severik@waksman.rutgers.edu
Correspondence may also be addressed to Dmitry Ghilarov. Email: dmitry.gilyarov@uj.edu.pl