

Personality, Socio-Economic Status and Inflammation: Cross-Sectional, Population-Based Study

Keith Millar¹, Suzanne M. Lloyd², Jennifer S. McLean^{3*}, G. David Batty^{4,5}, Harry Burns⁶, Jonathan Cavanagh¹, Kevin A. Deans^{7,8}, Ian Ford², Alex McConnachie², Agnes McGinty⁹, Réne Möttus¹⁰, Chris J. Packard⁹, Naveed Sattar¹¹, Paul G. Shiels¹², Yoga N. Velupillai¹³, Carol Tannahill³

1 College of Medical, Veterinary and Life Sciences, Institute of Mental Health and Wellbeing, University of Glasgow, Glasgow, Scotland, **2** Robertson Centre for Biostatistics, University of Glasgow, Glasgow, Scotland, **3** Glasgow Centre for Population Health, Glasgow, Scotland, **4** MRC Social and Public Health Sciences Unit, Glasgow, Scotland, **5** Department of Epidemiology and Public Health, University College London, London, England, **6** Scottish Government, Edinburgh, Scotland, **7** Department of Vascular Biochemistry, NHS Greater Glasgow and Clyde, Glasgow Royal Infirmary, Glasgow, Scotland, **8** Department of Clinical Biochemistry, NHS Grampian, Aberdeen Royal Infirmary, Aberdeen, Scotland, **9** Glasgow Clinical Research Facility, Tennent Building, Western Infirmary, Glasgow, Scotland, **10** Centre for Cognitive Ageing and Cognitive Epidemiology, Department of Psychology, University of Edinburgh, Edinburgh, Scotland, **11** Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, Scotland, **12** Institute of Cancer Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, Scotland, **13** Graduate Entry Medical School, University of Limerick, Limerick, Ireland

Abstract

Background: Associations between socio-economic status (SES), personality and inflammation were examined to determine whether low SES subjects scoring high on neuroticism or hostility might suffer relatively higher levels of inflammation than affluent subjects.

Methods: In a cross-sectional design, 666 subjects were recruited from areas of high (most deprived – “MD”) and low (least deprived – “LD”) deprivation. IL-6, ICAM-1, CRP and fibrinogen were measured along with demographic and health-behaviour variables, and personality traits of neuroticism, extraversion and psychoticism (hostility). Regression models assessed the prediction of inflammation as a function of personality, deprivation and their interaction.

Results: Levels of CRP and IL-6 were an increasing function of neuroticism and extraversion only in LD subjects opposite trends were seen in MD subjects. The result was ascribed parsimoniously to an inflammatory ceiling effect or, more speculatively, to SES-related health-behaviour differences. Psychoticism was strongly associated with ICAM-1 in both MD and LD subjects.

Conclusions: The association between neuroticism, CRP and IL-6 may be reduced in MD subjects confirming speculation that the association differs across population sub-groups. The association between psychoticism and ICAM-1 supports evidence that hostility has adverse effects upon the endothelium, with consequences for cardiovascular health. Health interventions may be more effective by accounting for personality-related effects upon biological processes.

Citation: Millar K, Lloyd SM, McLean JS, Batty GD, Burns H, et al. (2013) Personality, Socio-Economic Status and Inflammation: Cross-Sectional, Population-Based Study. PLoS ONE 8(3): e58256. doi:10.1371/journal.pone.0058256

Editor: Cordula M. Stover, University of Leicester, United Kingdom

Received: November 7, 2012; **Accepted:** February 5, 2013; **Published:** March 13, 2013

Copyright: © 2013 Millar et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Glasgow Centre for Population Health which is a partnership between National Health Service (NHS) Greater Glasgow and Clyde, Glasgow City Council and the University of Glasgow, supported by the Scottish Government. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Jennifer.mclean@drs.glasgow.gov.uk

Introduction

Individuals who display certain personality characteristics are more likely to indulge in harmful health behaviours [1–8] and to have increased risk of morbidity and mortality [7,9–11]. We have previously extended such findings to examine the association between personality, mental well-being and health behaviours as a function of socio-economic status (SES) [12]. Here, we consider the further association between SES, personality and inflammation.

There is growing evidence to link personality characteristics to inflammatory processes. For example, high levels of neuroticism (N) and low conscientiousness (C) have been associated with elevated levels of C-reactive protein (CRP) and Interleukin-6 (IL-6)

[13], and “pessimistic worry” (a feature of neuroticism) has been linked to high levels of CRP [14]. Similar associations have been shown between high levels of hostility and CRP and IL-6 [15]. Higher levels of inflammation are often consequences of harmful health behaviours such as smoking, poor diet and lack of exercise, and it is significant that the latter behaviours are also characteristic of high N, low C and psychoticism (P) [1–6,8]. However, the fact that the relationship between high N and mortality has been shown in one study to be independent of smoking and exercise [16] might imply the role of factors other than health behaviours. Most recently, one of our groups [17] has confirmed the negative relationship between C and CRP, and that the association was mediated by body mass index (BMI) but not by common health

behaviours of smoking, physical activity and alcohol consumption. Moreover, whilst N has been associated with adverse effects on health [11,16], a protective role for the factor has also been reported [18,19] when neuroticism-related health anxiety leads to positive health behaviours that may have beneficial consequences for health [20].

An association between personality and inflammation might have particular implications for socially-disadvantaged individuals. Low SES is associated with high levels of N [21,22], low levels of C [22], higher hostility [23] and depression [24]. In separate research, low SES was also associated with high levels of inflammatory markers that are in part consequences of harmful health behaviours and a stressful social environment associated with deprivation [25–28]. Given the latter associations, there may then be concern that deprived individuals having neurotic or hostile traits may be at risk of disproportionately greater levels of inflammatory activity. To the present authors' knowledge, no study has considered the interaction between personality factors, SES and inflammation.

The present study examines the association between SES, personality and inflammation in a cohort recruited from the most affluent and most deprived areas of a large British city. The cohort comprises the Psychological, Social and Biological Determinants of Ill Health ("pSoBid") [29,30] study which provides a broad range of biological, social and psychological variables which permit evaluation of the association between personality, inflammation and SES. The database also allows adjustment for covariates such as depression and BMI that are associated with inflammation but which have been omitted from some previous investigations of personality and inflammation [31], and for the factor of intelligence which may influence mortality and morbidity [32–34] in part via the health behaviours that are also associated with high inflammation.

Methods

Ethical Approval

The study was approved by the Glasgow Royal Infirmary Research Ethics Committee. Participants gave their written informed consent.

Study Population and Protocol

The design of the Psychological, Social and Biological Determinants of Ill Health (pSoBid) study has been described in detail elsewhere [29]. Briefly, selection of subjects was based on the Scottish Index of Multiple Deprivation (SIMD) 2004 [35] which rank small areas on the basis of multiple deprivation indicators. Subjects were recruited from five general medical practices in the city of Glasgow, Scotland, that served the bottom 5% of SIMD (i.e. relatively deprived) and five practices in areas classified as the top 20% of the SIMD (i.e. relatively affluent). Between December 2005 and May 2007 we recruited approximately equal numbers from both areas, equal numbers of men and women and equal numbers from each age group (35–44, 45–54 and 55–64 years old).

The original focus of the study was upon factors explaining the marked SES gradient in health, and in particular the role of inflammatory processes and their associated consequences for cardiovascular function. Sample size was therefore determined by the numbers required to detect differences between deprived and affluent groups in mean CRP levels (84% power to detect a 30% difference) and carotid intima media thickness (c-IMT: 82% power to detect a 0.04 mm difference), and did not take specific account

of the small effect sizes associated with personality variables [13]: this issue is considered further below.

Inflammatory Markers

High sensitivity C-reactive protein (CRP) was measured by an immunoturbidimetric assay (Roche Diagnostics Ltd., Burgess Hill, United Kingdom). Interleukin-6 (IL-6) and Intercellular Adhesion Molecule-1 (ICAM) were measured by sandwich ELISA (R&D Systems Europe Ltd., Abingdon, United Kingdom). Fibrinogen was measured on an automated coagulometer (MDA-180, Organon Teknika, Cambridge, United Kingdom).

Psychological, Lifestyle and Clinical Assessment

The short scale of the Eysenck Personality Questionnaire [36] assessed neuroticism (N), psychoticism (P) and extraversion (E) and included a "lie" scale to detect those seeking to present themselves in a socially ideal light. The National Adult Reading Test [37] (NART) defined the individual's peak achieved level of intellectual function [38], or pre-morbid intelligence [39]: the error score correlates negatively with intellectual function or "IQ". The depression sub-scale of the General Health Questionnaire [40] measured low mood. A lifestyle questionnaire assessed health-related variables including regular exercise, alcohol intake, dietary score [12] and smoking. In addition, subjects attended for a detailed clinical assessment which included blood pressure, body mass index (BMI) and c-IMT measurement [29].

Statistical Analysis

Descriptive statistics for continuous variables are presented as mean and standard deviation (SD) or median and interquartile range (IQR), as appropriate, and for categorical variables as frequencies and percentages.

Baseline differences between the groups were tested using linear and (binary or ordinal) logistic regression models for continuous and categorical variables respectively and adjusted for age and sex.

Inflammatory markers were tested for associations with the variables of age, gender, deprivation group, current smoker, alcohol consumption, physical activity, diet score, BMI, GHQ depression score, years of education and NART error score in separate linear regression models. Logarithmic transformations were applied to CRP, IL-6 and ICAM-1. Results are reported as the estimated regression coefficient, its 95% confidence interval (CI) and p-value.

Associations between each inflammatory marker and N, E and P were also tested with linear regression models which included terms for main effects and their interactions with deprivation. Results are reported as the effect estimate for the personality factor (with 95% CI and p-value) within each group, and the p-value for the interaction term testing whether the separate associations are different.

Then, for each inflammatory marker, all the predictor variables were entered into a backward stepwise regression; for the personality factors, the starting model included terms for interactions with deprivation. Terms were removed if they did not improve the fit of the model at a 5% significance level (subject to not removing the main effects of deprivation or a psychological variable whilst their interaction remained in the model). Those factors remaining in the final model are reported as above. This process was repeated, starting with factors found to be significantly associated with the inflammatory marker on univariate analysis, which gave the same results in each case. For those personality measures that were associated with inflammatory markers, the results are also presented graphically as the predicted level of each

marker in relation to the personality variable, with a 95% CI, in the most and least deprived groups.

In addition, we examined whether the LD and MD groups differed in their associations between E and N and health behaviours (smoking, alcohol consumption, diet and physical exercise), and other salient factors of BMI, GHQ depression scores, years of education and NART error scores.

Results

Table 1 confirms that the least deprived (LD) and most deprived (MD) groups were well matched on age and gender, but differed on a range of socioeconomic and lifestyle characteristics. Table 1 also shows, as expected, higher levels of CRP, IL-6, ICAM-1 and fibrinogen in the MD group, as well as higher scores on N and P compared to the LD group (adjusted for age and sex). The groups did not differ significantly in mean E or “Lie” scores.

Personality, SES and Inflammation

Tables 2a to 2d show the outcome of analyses to examine the associations between the inflammatory markers and the above-listed independent variables, allowing for possible interactions with deprivation group.

IL-6 showed significant univariate associations with deprivation, greater age, current smoking, poorer diet, less physical exercise, greater BMI, higher depression GHQ score, fewer years of education and NART error score (denoting relatively lower intellectual function: Table 2a). With respect to personality, only N showed a univariate association with IL-6, the significant positive relationship implying that levels of the marker were an increasing function of N score across the sample. However, further analysis as a function of deprivation grouping showed that the positive relationship was true only in the LD group ($p = 0.02$) whilst a non-significant trend to a negative relationship was seen in MD subjects ($p = 0.058$) so that the interaction between N and deprivation grouping was significant ($p = 0.003$). These relationships, and the interaction, were confirmed in the multivariate analysis (Table 2a) and are illustrated in Figure 1a. The figure shows that whilst levels of IL-6 were overall higher in MD subjects they declined significantly as a function of N in contrast to the positive function seen in LD subjects so that there was therefore no disproportionate increase in levels of the marker amongst high-N-scoring MD subjects. The multivariate analysis also showed that deprivation grouping, current smoking and BMI remained significant predictors of IL-6.

CRP showed a similar interaction between N and deprivation grouping as in the case of IL-6 but without evidence of a significant negative association between the marker and N in the MD subjects (Table 2b). Multivariate analysis showed that whilst levels of CRP were overall higher in MD subjects, the between-group difference was not significant and that levels of the marker were an increasing function of N only in LD subjects ($p = 0.002$) so that the interaction between N and deprivation grouping was significant ($p = 0.004$). As shown in Figure 1b, there was again no evidence of a disproportionate increase in levels of the marker in high-N-scoring MD subjects. CRP was also associated with greater age, fewer years of education, current smoking and BMI.

The multivariate analysis also showed that neither the positive association between E and CRP in LD subjects, nor the opposite relationship in MD subjects was significant ($p = 0.090$ and $p = 0.240$, respectively: Table 2b and Figure 1c), although the interaction between E and deprivation grouping did reach significance ($p = 0.044$).

ICAM-1 (Table 2c) showed univariate associations with all variables except gender and alcohol consumption. In the multivariate analysis, only deprivation group, BMI, fewer years of education and current smoking remained significant. P was the only personality variable to be significantly associated with ICAM-1 in both the univariate and multivariate analyses, and with an equivalent positive association in both groups indicating that hostile and aggressive characteristics were associated with higher levels of the marker (Figure 1d).

In univariate analyses, fibrinogen (Table 2d) was associated with all variables except alcohol, diet and physical activity. Only age, gender, smoking and BMI were independently predictive in the multivariate model. In univariate analyses, whilst N was positively associated with fibrinogen in the population as a whole, no association was found in the multivariate model.

Whilst higher-order interactions have been shown between N and E in studies of the three-factor model [41] where, for example, high E scores might moderate the effects of high N, the present study found no such interactions.

Table 3 shows associations between the personality factors and health behaviours, and BMI, GHQ depression score, years of education and NART error score for MD and LD groups, and the interaction. E was positively associated with the NART error score in LD subjects but negatively associated in MD subjects so that the interaction was significant ($p = 0.032$). The factor of E was also strongly associated with fewer years of education only in the LD group so that the interaction was significant ($p = 0.010$). It is then evident that higher E scores in LD subjects were associated with lower intellectual status and fewer years of education relative to the associations seen in their MD counterparts. In the case of N, a similar but much weaker pattern of associations was observed so that only in the case of years of education was the interaction marginally significant ($p = 0.058$). Further weak associations were seen in the case of health behaviours. Table 3 shows that E was associated with a relatively greater propensity to take physical exercise in MD subjects (interaction: $p = 0.090$), with lower BMI (interaction: $p = 0.055$) and with a higher diet score. Strongest, and contrasting, associations were seen in the case of depression scores: in MD subjects, E was strongly associated with lower depression scores whilst N was strongly associated with higher scores (both interactions: $p < 0.001$).

Summary of Principal Findings

Levels of inflammatory markers were overall higher in MD subjects, as were scores on N and P. However, levels of IL-6 and CRP were an increasing function of N only in LD subjects so that there was no disproportionate increase in the latter inflammatory markers in the MD group. A similar but weaker effect was seen between E and CRP. In the case of ICAM-1, the significant positive association between the marker and P was equivalent in both groups. In MD subjects, higher scores on E were associated with higher intellectual status, whilst an opposite relationship was seen in LD subjects.

Discussion

The results confirm the established separate associations between low SES and high inflammation, and between low SES and high scores on N and P [15,21–28]. Furthermore, to our knowledge, this study is the first to show that the relationship between personality and IL-6, and CRP, differs as a function of SES. Levels of the inflammatory markers were overall higher in the MD group, but as N scores increased, they were associated with increased levels of the marker only in the LD group. There

Table 1. Description of basic demographics, socioeconomic status, personality and markers of inflammation by area deprivation category.

	Least Deprived (n = 342)	Most Deprived (n = 324)	p ^b
Age (years)	51.8 (8.0) ^a	51.5 (8.5)	0.63
Gender			
Male	171 (50.0%)	156 (48.1%)	0.64
Female	171 (50.0%)	168 (51.9%)	
Household income	£41,699 (11,921)	£16,461 (10,056)	<0.001
Education (total years)	16.1 (3.6)	11.8 (2.5)	<0.001
Residential status			
Owner	334 (97.7%)	97 (29.9%)	<0.001
Tenant	8 (2.3%)	227 (70.1%)	
Occupation category ^c			
I & II	251 (73.4%)	62 (19.1%)	<0.001
III	77 (22.5%)	139 (42.9%)	
IV & V	12 (3.5%)	105 (32.4%)	
Unemployed	1 (0.3%)	2 (0.6%)	
Current Smoker			
Yes	28 (8.2%)	161 (49.7%)	<0.001
No	314 (91.8%)	163 (50.3%)	
Alcohol (weekly units)	11.1 (12.7)	11.1 (21.5)	0.906
Diet score ^d	95.7 (51.4)	59.9 (50.4)	<0.001
Activity ^e			
Moderately Active – Active	176 (51.5%)	127 (39.2%)	<0.001
Moderately Inactive – Inactive	166 (48.5%)	197 (60.8%)	
Body mass index (BMI)	26.9 (4.5)	28.7 (6.3)	<0.001
Inflammatory Biomarkers			
C-reactive protein (CRP) (mg/l) ^f	1.09 [0.51–2.27]	2.12 [1.07–4.32]	<0.001
Interleukin-6 (IL-6) (pg/ml) ^f	1.25 [0.87–1.95]	2.28 [1.34–3.13]	<0.001
Intercellular adhesion molecule (ICAM-1) (ng/ml) ^f	229.4 [207.5–263.6]	297.8 [241.9–391.7]	<0.001
Fibrinogen (g/l)	3.23 (0.64)	3.50 (0.76)	<0.001
Eysenck Personality scores (EPQ-R) ^g			
Neuroticism (N)	4.06 (3.19)	5.96 (3.79)	<0.001
Extraversion (E)	7.49 (3.41)	7.34 (3.61)	0.558
Psychoticism (P)	1.26 (1.30)	2.58 (2.02)	<0.001
Lie scale	5.35 (2.68)	5.34 (2.78)	0.962
GHQ depression score ^h	0.17 (0.79)	0.76 (1.71)	<0.001
NART error score ⁱ	7.16 (5.27)	15.61 (9.01)	<0.001

^aValues are presented as Mean (SD) for all participants; or as percentages for categorical variables, adjusted for age and sex;

^bP relates the comparison between the two groups. Categorical variables were compared using Fisher’s exact test, continuous variables were compared using t-tests or Wilcoxon tests as appropriate.

^cThe occupational category could not be determined for n = 1 (0.3%) and n = 16 (5%) of the LD and MD groups respectively. Occupation classified using Registrar General Social Class Classification on basis of current job or, if not currently working, on the basis of participants’ last paid job. Only those who had never been in paid employment were classed as “unemployed.” I - professional occupations; II - managerial and technical occupations; III - manual and non-manual skilled occupations; IV - partly skilled occupations; V - unskilled occupations.

^dDiet score is the participants self-reported consumption of fruit and vegetables (fresh, cooked and raw) per month.

^ePhysical activity level is a combination of activity at work and recreational exercise.

^fData log transformed prior to regression analysis.

^gPersonality trait scores were self-reported, each on a scale of 1 to 12.

^hDepression = depression sub-scale of GHQ-28.

ⁱNART (National Adult Reading Test).

doi:10.1371/journal.pone.0058256.t001




Figure 1. Levels of inflammatory markers as a function of the interaction between personality and deprivation group with 95% confidence bands.
doi:10.1371/journal.pone.0058256.g001

was therefore no evidence that MD subjects having high N scores might suffer disproportionately higher levels of inflammation. A similar, although much weaker, association was observed between E and CRP.

A parsimonious explanation for these effects would be that, at high levels of inflammation, there is simply little scope for the known small and subtle influences of personality upon inflammation [13] to exert a detectable further increase in marker levels: in other words, a ceiling effect. However, whilst plausible, this proposal does not address the evidence that levels of IL-6 and, to a lesser extent, CRP declined as a function of high N in the MD group and with a similar trend being seen in the case of E and CRP.

The latter apparently beneficial effect of high N upon inflammation might be explained if neuroticism-related health anxiety exerted a protective effect through the adoption of positive health behaviours [17–19]. However, as no measure of health anxiety was included in the present methodology, this proposal cannot be assessed; nor is there obvious reason why these MD subjects might have been more prone to health anxieties. Moreover, there was no evidence in Table 3 of any significant associations between N and health behaviours in either of the two groups.

An alternative explanation may be found in differences between the groups in associations between certain health-related factors and N and E that might influence levels of inflammation. Fundamental individual characteristics that might exert such an

influence would be those of intellectual status and years of education. Intellect is known to promote health by influencing self-care and health awareness [43], for example by taking greater dietary care and more exercise to the benefit of BMI, and smoking cessation. The latter factors are known to influence levels of inflammatory activity. Longer years of education may benefit health for similar reasons although the effects are less clear-cut [44]. The analysis showed that high E was associated with relatively higher intellectual function in MD subjects, whilst the opposite association was seen in the LD group. The high-E-scoring LD subjects also had fewer years of education and a similar although weaker association in LD subjects was seen in the case of N. A relatively greater intellect and longer years of education would plausibly tend to foster positive health behaviours. However, whilst high E in MD subjects was associated in Table 3 with a greater propensity to engage in physical activity, a lower BMI and a higher diet score than in high-E-scoring LD subjects, the associations were not formally significant. Moreover, as no adjustments were made in Table 3 for multiple statistical comparisons there is scope for Type-1 error: all p values must be considered as descriptive measures of strength of evidence for the observed associations. E scores were also associated with lower depression in MD subjects, but N scores showed an opposite relationship. The evidence that N was associated with higher depression scores in MD subjects would seem inconsistent with the motivation and drive to adopt positive health behaviours.

Table 2. Univariate and multivariate associations between inflammatory markers and health-related variables, personality traits and deprivation grouping, and the interaction of personality and deprivation.

Independent variables	Univariate analysis			Multivariate analysis		
	Estimate (95% CI)	P	P _{interaction}	Estimate (95% CI)	P	P _{interaction}
2a Interleukin-6 (IL-6)						
Deprivation status (MD vs LD)	0.428 (0.326, 0.529)	<0.001		0.205 (0.098, 0.311)	<0.001	
Age	0.221 (0.159, 0.284)	<0.001		0.192 (0.137, 0.247)	<0.001	
Gender (Male vs. Female)	0.053 (−0.053, 0.160)	0.327				
Education (total years)	−0.038 (−0.052, −0.024)	<0.001				
Current smoker (Yes vs. No)	0.392 (0.277, 0.508)	<0.001		0.385 (0.268, 0.502)	<0.001	
Alcohol (units per week)	0.002 (−0.001, 0.005)	0.252				
Diet score	−0.002 (−0.003, −0.001)	<0.001				
Activity (Active vs. Inactive)	−0.224 (−0.329, −0.118)	<0.001				
BMI	0.048 (0.039, 0.057)	<0.001		0.046 (0.037, 0.055)	<0.001	
GHQ Depression score	0.067 (0.026, 0.108)	0.001				
NART error score	0.018 (0.012, 0.024)	<0.001				
Neuroticism (N)	MD: −0.019 (−0.039, 0.001)	0.058	0.003	MD: −0.022 (−0.040, −0.005)	0.012	0.001
	LD: 0.026 (0.004, 0.048)	0.020		LD: 0.021 (0.001, 0.040)	0.035	
Extraversion (E)	MD: −0.008 (−0.029, 0.013)	0.436	0.215			
	LD: 0.010 (−0.011, 0.031)	0.329				
Psychoticism (P)	MD: 0.034 (−0.002, 0.071)	0.067	0.568			
	LD: 0.015 (−0.039, 0.069)	0.581				
2b C-reactive protein (CRP)						
Deprivation status (MD vs LD)	0.581 (0.411, 0.750)	<0.001		0.102 (−0.107, 0.311)	0.339	
Age	0.225 (0.120, 0.331)	<0.001		0.173 (0.077, 0.269)	<0.001	
Gender (Male vs. Female)	−0.138 (−0.313, 0.038)	0.123				
Education (total years)	−0.079 (−0.102, −0.056)	<0.001		−0.027 (−0.054, −0.001)	0.040	
Current smoker (Yes vs. No)	0.364 (0.170, 0.559)	<0.001		0.369 (0.166, 0.572)	<0.001	
Alcohol (units per week)	−0.003 (−0.009, 0.002)	0.176				
Diet score	−0.002 (−0.004, −0.0003)	0.021				
Activity (Active vs. Inactive)	−0.228 (−0.403, −0.053)	0.011				
BMI	0.089 (0.075, 0.104)	<0.001		0.087 (0.072, 0.102)	<0.001	
GHQ Depression score	0.129 (0.063, 0.195)	<0.001				
NART error score	0.033 (0.023, 0.043)	<0.001				
Neuroticism (N)	MD: 0.005 (−0.028, 0.038)	0.779	0.012	MD: −0.016 (−0.050, 0.018)	0.351	0.004
	LD: 0.069 (0.032, 0.106)	<0.001		LD: 0.054 (0.020, 0.089)	0.002	
Extraversion (E)	MD: −0.027 (−0.062, 0.008)	0.132	0.014	MD: −0.021 (−0.057, 0.014)	0.240	0.044
	LD: 0.035 (0.000, 0.070)	0.048		LD: 0.028 (−0.004, 0.060)	0.090	
Psychoticism (P)	MD: 0.043 (−0.019, 0.105)	0.173	0.809			
	LD: 0.029 (−0.062, 0.121)	0.529				
2c. Intercellular Adhesion Molecule-1 (ICAM-1)						
Deprivation status (MD vs LD)	0.248 (0.210, 0.287)	<0.001		0.079 (0.031, 0.128)	0.001	
Age	0.033 (0.007, 0.059)	0.013				
Gender (Male vs. Female)	0.006 (−0.037, 0.050)	0.768				
Education (total years)	−0.025 (−0.030, −0.020)	<0.001		−0.007 (−0.013, −0.002)	0.011	
Current smoker (Yes vs. No)	0.309 (0.267, 0.351)	<0.001		0.243 (0.196, 0.290)	<0.001	
Alcohol (units per week)	0.000 (−0.001, 0.001)	0.944				
Diet score	−0.001 (−0.002, −0.001)	<0.001				
Activity (Active vs. Inactive)	−0.082 (−0.125, −0.039)	<0.001				
BMI	0.006 (0.002, 0.010)	0.002		0.006 (0.003, 0.010)	<0.001	
GHQ Depression score	0.037 (0.021, 0.054)	<0.001				

Table 2. Cont.

Independent variables	Univariate analysis		Multivariate analysis			
	Estimate (95% CI)	p	P _{interaction}	Estimate (95% CI)	p	P _{interaction}
NART error score	0.009 (0.007, 0.012)	<0.001				
Neuroticism (N)	MD: 0.003 (−0.005, 0.010)	0.443	0.982			
	LD: 0.003 (−0.006, 0.011)	0.508				
Extraversion (E)	MD: −0.006 (−0.014, 0.002)	0.148	0.100			
	LD: 0.004 (−0.004, 0.012)	0.376				
Psychoticism (P)	MD: 0.022 (0.009, 0.036)	0.002	0.947	All: 0.016 (0.006, 0.027)	0.002	
	LD: 0.023 (0.003, 0.044)	0.026				
2d. Fibrinogen						
Deprivation status (MD vs LD)	0.265 (0.155, 0.375)	<0.001				
Age	0.167 (0.100, 0.233)	<0.001		0.142 (0.079, 0.205)	<0.001	
Gender (Male vs. Female)	−0.136 (−0.247, −0.025)	0.017		−0.141 (−0.244, −0.038)	0.007	
Education (total years)	−0.029 (−0.043, −0.014)	<0.001				
Current smoker (Yes vs. No)	0.0252 (0.128, 0.376)	<0.001		0.340 (0.222, 0.457)	<0.001	
Alcohol (units per week)	−0.002 (−0.005, 0.001)	0.256				
Diet score	0.000 (−0.001, 0.001)	0.890				
Activity (Active vs. Inactive)	−0.103 (−0.215, 0.008)	0.069				
BMI	0.039 (0.029, 0.049)	<0.001		0.041 (0.031, 0.050)	<0.001	
GHQ Depression score	0.054 (0.011, 0.097)	0.014				
NART error score	0.013 (0.006, 0.019)	<0.001				
Neuroticism (N)	MD: 0.001 (−0.020, 0.022)	0.943	0.089			
	LD: 0.028 (0.005, 0.052)	0.018				
Extraversion (E)	MD: −0.020 (−0.043, 0.002)	0.074	0.246			
	LD: −0.002 (−0.024, 0.021)	0.882				
Psychoticism (P)	MD: 0.027 (−0.013, 0.067)	0.182	0.523			
	LD: 0.004 (−0.054, 0.063)	0.890				

The outcome of initial univariate analysis shows associations between inflammatory marker and independent variables including the interactions of neuroticism (N), extraversion (E) and psychoticism (P) with deprivation status. Multivariate model determined by backward stepwise regression.

^aBMI (body mass index).

^bDepression = depression sub-scale of GHQ-28.

^cNART (National Adult Reading Test).

doi:10.1371/journal.pone.0058256.t002

The analysis therefore provided some evidence that a number of health-related factors relevant to inflammation had different associations with E and N in MD and LD subjects. It must be speculative given the caveats above, but such differences may contribute in part to the contrasting associations seen between inflammation and personality in the two groups. Other relevant factors might be those of individual variation in coping strategies and resilience to socio-economic adversity, but we omitted to assess such characteristics.

More generally, the results parallel those of Chapman et al. [31] who reported no effect of N, and an inverse relationship between E and IL-6, in their study involving predominantly low SES subjects (their design precluded examining effects in higher SES subjects). The reduction in IL-6 was associated with the activity facet of extraversion (assessed by the NEO-FFI [42]) but we could not confirm this specific association due to limitations of the EPQ. Our findings have the advantage that the supposed low-SES-related association can be confirmed by comparison with an affluent subgroup. Overall, such results are consistent with proposals that the effects of personality may be absent, or of a different nature, in low SES groups and those of ethnic minorities [21,31], leading to

speculation that some traits of N and E may be associated with inflammation only in population sub-groups [45].

The significant association between P and ICAM-1 in both the deprived and affluent groups confirms existing evidence of an adverse influence of hostility upon endothelial function [46], and that cynical hostility is related to greater cytokine production [47] which, significantly in view of the present results, has been shown independent of adjustment for covariates including SES [48]. Our result would confirm that individuals high in hostility and aggression are at increased risk of inflammatory disease [15] and that this vulnerability is evident across the SES continuum.

Our failure to find a significant multivariate association between fibrinogen and the personality factors assessed here is similar to a recent negative result of Möttus et al. [17] who note that fibrinogen is implicated in coagulation and is therefore a more peripheral and less sensitive inflammatory marker.

A differential association between SES groups in terms of personality and health-related factors may have implications for health promotion and intervention [12]. A strategy of adapting interventions to the behaviours and beliefs that characterise particular personality types may improve the implementation of

Table 3. Associations between personality traits and health behaviours in affluent and deprived groups, and their interactions.

	Least deprived	Most deprived	P Interaction
Years of education			
Neuroticism	-0.182 (-0.286, -0.078)	-0.050 (-0.142, 0.042)	0.058
Extraversion	-0.178 (-0.276, -0.081)	0.004 (-0.093, 0.100)	0.010
Psychoticism	-0.104 (-0.361, 0.154)	-0.072 (-0.246, 0.101)	0.840
Current smoker (Yes vs. No)			
Neuroticism	0.957 (0.842, 1.086)	1.048 (0.985, 1.116)	0.200
Extraversion	1.026 (0.911, 1.154)	0.981 (0.921, 1.045)	0.518
Psychoticism	1.317 (1.007, 1.724)	1.061 (0.945, 1.191)	0.144
Alcohol (weekly units)			
Neuroticism	-0.259 (-0.841, 0.323)	-0.221 (-0.736, 0.294)	0.923
Extraversion	0.565 (0.031, 1.100)	0.672 (0.149, 1.195)	0.779
Psychoticism	0.477 (-0.926, 1.879)	0.951 (0.006, 1.896)	0.577
Diet score (fruit and vegetable consumption per month)			
Neuroticism	-0.206 (-1.954, 1.542)	-1.460 (-3.005, 0.085)	0.283
Extraversion	1.078 (-0.526, 2.683)	2.431 (0.861, 4.000)	0.237
Psychoticism	-1.323 (-5.522, 2.877)	-0.987 (-3.817, 1.843)	0.895
Activity (Active vs. Inactive)			
Neuroticism	0.992 (0.925, 1.063)	0.950 (0.892, 1.012)	0.363
Extraversion	1.007 (0.944, 1.074)	1.091 (1.021, 1.167)	0.090
Psychoticism	0.993 (0.839, 1.177)	0.852 (0.749, 0.969)	0.151
BMI^a			
Neuroticism	0.181 (-0.007, 0.368)	0.077 (-0.090, 0.243)	0.406
Extraversion	0.142 (-0.032, 0.317)	-0.097 (-0.268, 0.074)	0.055
Psychoticism	0.379 (-0.079, 0.838)	-0.068 (-0.378, 0.241)	0.109
GHQ depression score^b			
Neuroticism	0.053 (0.012, 0.094)	0.184 (0.148, 0.220)	<0.001
Extraversion	-0.019 (-0.060, 0.022)	-0.128 (-0.168, -0.088)	<0.001
Psychoticism	0.037 (-0.068, 0.143)	0.054 (-0.018, 0.126)	0.792
NART error score^c			
Neuroticism	-0.035 (-0.272, 0.202)	0.083 (-0.127, 0.294)	0.455
Extraversion	0.168 (-0.048, 0.383)	-0.162 (-0.373, 0.049)	0.032
Psychoticism	-0.241 (-0.805, 0.322)	0.434 (0.053, 0.814)	0.049

^aBMI (body mass index).

^bDepression = depression sub-scale of GHQ-28.

^cNART (National Adult Reading Test).

doi:10.1371/journal.pone.0058256.t003

intervention programmes. There have been cogent proposals that interventions targeted at low SES groups may be refined to account for dispositional differences [31], and that the identification of individuals whose personality styles render them vulnerable to particular health risks would allow them to benefit from closer monitoring which might result in earlier detection and treatment [45]. Our findings confirm that account should be taken of SES when exploring relationships between personality and biological factors relating to health: indeed, the overall main effect of N gives the misleading impression of a positive relationship between neuroticism and CRP and IL-6 across the sample. In fact, the relationship may be reduced or reversed in deprived individuals, confirming proposals that the association between personality and inflammation may differ amongst population sub-groups [21,31] and implying that targeted interventions may require particular subtlety.

The present study has a number of limitations. The most significant is the cross-sectional design which precludes any attribution of causality. The three-factor assessment of personality precluded analysis of some facets of N and E that may determine the expression, or inhibition, of particular health-related behaviours [6]. The sample size, while large by comparison with some other studies, was determined by the numbers needed to detect between-group differences in two biological variables but, as the effect sizes due to personality variables are typically small [13], the sample size may have been insufficient to detect subtle effects of personality at high levels of inflammation. Moreover, whilst the sampling procedure was successful in recruiting two samples which differed widely in terms of SES, their composition was, self-evidently, not fully representative of the Scottish population. These considerations lead to the more general question as to whether the present results are representative of “deprived” individuals within

the population. We acknowledge that our MD sample may not represent an extreme of socio-economic deprivation for the simple reason that such individuals do not readily volunteer as research subjects [49]. We therefore concede that the present MD subjects may represent a sub-group whose relative intellect and years of education have encouraged an interest in health and well-being, and provided the confidence to volunteer and engage in the study. The outgoing and sociable characteristics of those higher in E would further encourage this tendency. Finally, the exclusively white Caucasian sample also precluded examination of ethnic differences that have been shown relevant in other investigations of personality and inflammation [21]. We also note the high number of statistical tests performed, which might have resulted in an increased type I error rate.

Despite these limitations, the demonstration of an interaction between personality, deprivation and inflammation is a novel finding. Evidence that the expression of personality influences upon inflammation may differ within sub-groups of the population provides a caveat that misleading conclusions may be drawn as to

the relationship between personality and biological variables if the factor of SES is neglected.

Acknowledgments

Thanks are due to administrative staff in the Glasgow Centre for Population Health; Robertson Centre for Biostatistics, University of Glasgow for data management, statistical support and analysis; Health Information & Technology section of NHS Greater Glasgow & Clyde for sample selection and analysis; Alasdair Buchanan for GPASS data and analysis; all members of GP Practices who participated in the study and the participants themselves.

Author Contributions

Conceived and designed the experiments: KM SML JSM GDB HB JC KAD IF A. McConnachie A. McGinty RM CJP NS PGS YNV CT. Performed the experiments: YNV KAD A. McGinty. Analyzed the data: KM SML A. McConnachie. Contributed reagents/materials/analysis tools: KM SML KAD A. McConnachie CJP. Wrote the paper: KM SML JSM GDB HB JC KAD IF A. McConnachie A. McGinty RM CJP NS PGS YNV CT.

References

1. Arai Y, Hosokawa T, Fukao A, Izumi Y, Hisamichi S (1997) Smoking behaviour and personality: a population-based study in Japan. *Addiction* 92: 1023–1033.
2. Berlin I, Singleton EG, Pedarriose AM, Lancrenon S, Rames A, et al. (2003) The modified reasons for smoking scale: factorial structure, gender effects and relationship with nicotine dependence and smoking cessation in French smokers. *Addiction* 98: 1575–1583.
3. Chapman BP, Fiscella K, Kawachi I, Duberstein PR (2009a) Personality, socioeconomic status, and all-cause mortality in the United States. *Am J Epidemiol* 171: 83–92.
4. De Moor MHM, Beem AL, Stubbe JH, Boomsma DI, De Geus EJC (2006) Regular exercise, anxiety, depression and personality: a population-based study. *Prev Med* 42: 273–279.
5. Raynor DA, Levine H (2009) Associations between the five-factor model of personality and health behaviors among college students. *J Am Coll Health* 58: 73–81.
6. Terracciano A, Costa PT Jr (2004) Smoking and the five-factor model of personality. *Addiction* 99: 472–481.
7. Terracciano A, Sutin AR, McCrae RR, Deiana B, Ferrucci L, et al. (2009) Facets of personality linked to underweight and overweight. *Psychosom Med* 71: 682–689.
8. Yeung RR, Hemsley DR (1997) Personality, exercise and psychological well-being: static relationships in the community. *Person Ind Diff* 22: 47–53.
9. Mroczak DK, Spiro A 3rd (2007) Personality change influences mortality in older men. *Psychol Sci* 18: 371–376.
10. Nabi H, Kivimäki M, Zins M, Elovainio M, Consoli SM, et al. (2008) Does personality predict mortality? Results from the GAZEL French prospective cohort study. *Int J Epidemiol* 37: 386–396.
11. Roberts BW, Kuncel NR, Shiner B, Caspi A, Goldberg LR (2007) The power of personality. The comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspect Psychol Sci* 2: 313–345.
12. Packard CJ, Cavanagh J, McLean JS, McConnachie A, Messow CA, et al. (2012) Interaction of personality traits with social deprivation in determining mental wellbeing and health behaviours. *J Pub Health* 34: 615–624.
13. Sutin AR, Terracciano A, Deiana B, Naitza S, Ferrucci L, et al. (2010) High neuroticism and low conscientiousness are associated with interleukin-6. *Psychol Med* 40: 1485–1493.
14. Henningson S, Baghaei F, Rosmond R, Holm G, Anckarsäter H, et al. (2008) Association between serum levels of C-reactive protein and personality traits in women. *Behav Brain Funct* 4: 16.
15. Marsland AL, Prather AA, Petersen KL, Cohen S, Manuck SB (2008) Antagonistic characteristics are positively associated with inflammatory markers independently of trait negative emotionality. *Brain Behav Immun* 22: 753–761.
16. Shipley BA, Weiss A, Der G, Taylor MD, Deary IJ (2007) Neuroticism, extraversion and mortality in the UK Health and Lifestyle Survey: a 21 year prospective cohort study. *Psychosom Med* 69: 923–931.
17. Möttus R, Luciano M, Starr JM, Pollard MC, Deary IJ (2013). Personality traits and inflammation in men and women in their early 70 s: the Lothian Birth Cohort 1936 Study of Healthy Aging. *Psychosom Med* 75: 11–19.
18. Korten AE, Jorm AF, Jiao Z, Letenneur L, Jacomb PA, et al. (1999) Health, cognitive and psychosocial factors as predictors of mortality in an elderly community sample. *J Epidemiol Comm Health* 53: 83–88.
19. Weiss A, Costa PT (2005) Domain and facet personality predictors of all-cause mortality among Medicare patients aged 65 to 100. *Psychosom Med* 67: 724–733.

20. Roberts BW, Smith J, Jackson JJ, Edmonds G (2009) Compensatory conscientiousness and health in older couples. *Psychol Sci* 20: 553–559.
21. Jonassaint CR, Siegler IC, Barefoot JC, Edwards CL, Williams RB (2009) Low life course socioeconomic status (SES) is associated with negative NEO PI-R personality patterns. *Int J Behav Med* 18: 13–21.
22. Bosma H, van de Mheen HD, Mackenbach JP (1999) Social class in childhood and general health in adulthood: questionnaire study of contribution of psychological attributes. *BMJ* 318: 18–22.
23. Kubzansky L, Kawachi I, Sparrow D (1999) Socioeconomic status, hostility, and risk factor clustering in the normative aging study: any help from the concept of allostatic load? *Ann Behav Med* 21: 330–338.
24. Harper S, Lynch J, Hsu WL, Everson SA, Hillemeier MM, et al. (2002) Life course socioeconomic conditions and adult psychosocial functioning. *Int J Epidemiol* 31: 395–403.
25. Jousilahti P, Salomaa V, Rasi V, Vahtera E, Palosuo T (2003) Association of markers of systematic inflammation, C-reactive protein, serum amyloid A, and fibrinogen, with socioeconomic status. *J Epidemiol Comm Health* 57: 730–733.
26. Lubbock LA, Goh A, Ali S, Ritchie J, Whoolley MA (2005) Relation of low socioeconomic status to C-reactive protein in patients with coronary heart disease (from the Heart and Soul Study). *Am J Cardiol* 96: 1506–1511.
27. Nazmi A, Victora CG (2007) Socioeconomic and racial/ethnic differential of C-reactive protein levels: a systematic review of population based studies. *BMC Public Health* 7: 212.
28. Pollitt RA, Kaufman JS, Rose KM, Diez-Roux AV, Zeng D, et al. (2008) Cumulative life course and adult socioeconomic status and markers of inflammation in adulthood. *J Epidemiol Comm Health* 62: 484–491.
29. Velupillai YN, Packard CJ, Batty GD, Bezyak V, Burns H, et al. (2008) Psychological, social and biological determinants of ill health (pSoBid): study protocol of a population based study. *BMC Public Health* 8: 126–137.
30. Deans KA, Bezyak V, Ford I, Batty GD, Burns H, et al. (2009) Differences in atherosclerosis according to area level socioeconomic deprivation: cross sectional, population based study. *BMJ* 339: b4170.
31. Chapman BP, Khan A, Harper M (2009b) Gender, race/ethnicity, personality and interleukin-6 in urban primary care patients. *Brain Behav Immun* 23: 636–642.
32. Batty GD, Wennerstad KM, Davey-Smith G, Gunnell GD, Smith GD, et al. (2009) IQ in early adulthood and mortality by middle-age: cohort study of 1 million Swedish men. *Epidemiol* 20: 100–109.
33. Batty GD, Deary IJ, Benzeval M, Der G (2010) Does IQ predict cardiovascular disease mortality as strongly as established risk factors? Comparison of effect estimates using the West of Scotland Twenty-07 cohort study. *Eur J Cardiovasc Prev Rehabil* 17: 24–27.
34. Hart CL, Taylor MD, Smith GD, Whalley LJ, Starr JM, et al. (2004). Childhood IQ and cardiovascular disease in adulthood: prospective observational study linking the Scottish Mental Survey 1932 and the Midspan studies. *Soc Sci Med* 59: 2131–2138.
35. Scottish Index for Multiple Deprivation (2004) Scottish Government. Available: <http://www.scotland.gov.uk/Topic/Statistics/SIMD/Overview>.
36. Eysenck HJ, Eysenck SBG (1991) Manual of the Eysenck Personality Scales. London: Hodder and Stoughton.
37. Nelson HE, Willison J (1991) National Adult Reading Test (NART): test manual. Windsor, UK: NFER-Nelson.
38. Crawford JR, Deary IJ, Starr J, Whalley LJ (2001) The NART as an index of prior intellectual functioning: a retrospective validity study covering a 66-year interval. *Psychol Med* 31: 451–458.

39. Bright P, Jaldow E, Kopelman MD (2002) The National Adult Reading Test as a measure of premorbid intelligence: a comparison with estimates derived from demographic variables. *J Int Neuropsych Soc* 8: 847–854.
40. Goldberg D (1978) *Manual of the General Health Questionnaire*. Windsor, UK: NFER-Nelson.
41. Parkes KR (1984) Smoking and the Eysenck personality dimensions: an interactive model. *Psychol Med* 14: 825–834.
42. Costa PT, McCrae RR (1992) *Revised NEO Personality Inventory (NEO PI-R) and NEO-Five-Factor Inventory*. Odessa: Psychological Assessment Resources Inc.
43. Gottfredson LS, Deary IJ (2004) Intelligence predicts health and longevity, but why? *Curr Direct Psychol Sci* 13: 1–4.
44. Singh-Manoux A, Ferrie JE, Lynch JW, Marmot M (2005) The role of cognitive ability (intelligence) in explaining the association between socioeconomic position and health: evidence from the Whitehall II Prospective Cohort Study. *Am J Epidemiol* 161: 831–839.
45. Chapman BP, van Wijngaarden E, Seplaki CL, Talbot N, Duberstein P, et al. (2011) Openness and conscientiousness predict 34-week patterns of Interleukin-6 in older persons. *Brain Behav Immun* 25: 667–73.
46. Kop WJ (2003) The integration of cardiovascular behavioural medicine and psychoneuroimmunology: new developments based on converging research fields. *Brain Behav Immun* 17: 233–237.
47. Miller GE, Freedlan KE, Carney RM, Stetler CA, Banks WA (2003) Cynical hostility, depressive symptoms, and the expression of inflammatory risk markers for coronary heart disease. *J Behav Med* 26: 501–515.
48. Janicki-Deverts D, Cohen S, Doyle WJ (2010) Cynical hostility and stimulated Th1 and TH2 cytokine production. *Brain Behav Immun* 24: 58–63.
49. Goodman A, Gatward R (2008) Whom are we missing? Area deprivation and survey participation. *Eur J Epidemiol* 23: 379–387.