

Genome Sequence of *Lactococcus lactis* subsp. *lactis* NCDO 2118, a GABA-Producing Strain

Letícia C. Oliveira,^a Tessália D. L. Saraiva,^a Siomar C. Soares,^{a*} Rommel T. J. Ramos,^b Pablo H. C. G. Sá,^b Adriana R. Carneiro,^a Fábio Miranda,^b Matheus Freire,^b Wendel Renan,^b Alberto F. O. Júnior,^a Anderson R. Santos,^{a*} Anne C. Pinto,^a Bianca M. Souza,^a Camila P. Castro,^a Carlos A. A. Diniz,^a Clarissa S. Rocha,^a Diego C. B. Mariano,^a Edgar L. de Aguiar,^a Edson L. Folador,^a Eudes G. V. Barbosa,^a Flavia F. Aburjaile,^a Lucas A. Gonçalves,^a Luís C. Guimarães,^a Marcela Azevedo,^a Pamela C. M. Agresti,^a Renata F. Silva,^a Sandeep Tiwari,^a Sintia S. Almeida,^a Syed S. Hassan,^a Vanessa B. Pereira,^a Vinicius A. C. Abreu,^a Ulisses P. Pereira,^{a*} Fernanda A. Dorella,^c Alex F. Carvalho,^c Felipe L. Pereira,^c Carlos A. G. Leal,^c Henrique C. P. Figueiredo,^c Artur Silva,^b Anderson Miyoshi,^a Vasco Azevedo^a

Laboratory of Cellular and Molecular Genetics, Institute of Biologic Sciences, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil[†]; Institute of Biologic Sciences, Federal University of Pará, Belém, PA, Brazil[‡]; AQUACEN, National Reference Laboratory for Aquatic Animal Diseases, Ministry of Fisheries and Aquaculture, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil[§]

* Present address: Siomar C. Soares, AQUACEN, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil; Anderson R. Santos, Federal University of Uberlândia, Minas Gerais, MG, Brazil; Ulisses P. Pereira, Federal University of Uberlândia, Minas Gerais, MG, Brazil.

***Lactococcus lactis* subsp. *lactis* NCDO 2118 is a nondairy lactic acid bacterium, a xylose fermenter, and a gamma-aminobutyric acid (GABA) producer isolated from frozen peas. Here, we report the complete genome sequence of *L. lactis* NCDO 2118, a strain with probiotic potential activity.**

Received 21 August 2014 Accepted 26 August 2014 Published 2 October 2014

Citation Oliveira LC, Saraiva TDL, Soares SC, Ramos RTJ, Sá PHCG, Carneiro AR, Miranda F, Freire M, Renan W, Júnior AFO, Santos AR, Pinto AC, Souza BM, Castro CP, Diniz CAA, Rocha CS, Mariano DCB, de Aguiar EL, Folador EL, Barbosa EGV, Aburjaile FF, Gonçalves LA, Guimarães LC, Azevedo M, Agresti PCM, Silva RF, Tiwari S, Almeida SS, Hassan SS, Pereira VB, Abreu VAC, Pereira UP, Dorella FA, Carvalho AF, Pereira FL, Leal CAG, Figueiredo HCP, Silva A, Miyoshi A, Azevedo V. 2014. Genome sequence of *Lactococcus lactis* subsp. *lactis* NCDO 2118, a GABA-producing strain. *Genome Announc.* 2(5):e00980-14. doi:10.1128/genomeA.00980-14.

Copyright © 2014 Oliveira et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Vasco Azevedo, vasco@icb.ufmg.br.

Lactic acid bacteria (LAB), in general, acquire energy from the conversion of sugars into lactic acid (1) and are used for production of many fermented products, such as cheese, yogurt, butter, and wine. Food conservation is due to the medium acidification and production of molecules that inhibit the growth of undesirable microbiota, contributing to the development of desirable organoleptic properties in the final product (2). Moreover, some specific LAB strains produce bioactive molecules such as gamma-aminobutyric acid (GABA) (3), a product of glutamate decarboxylation by the glutamic acid decarboxylase (GAD) enzyme. Usually, GABA acts by modulating the central nervous system, contributing to smooth muscle relaxation and presenting hypotensor activity (4). Also, GABA can immunomodulate the immune system (5). Therefore, GABA-producing bacteria generally present probiotic properties (6). *Lactococcus lactis* NCDO 2118 is a nondairy strain, a xylose fermenter (a common trait of plant-associated strains), and a GABA producer isolated from frozen peas (6, 7).

L. lactis NCDO 2118 was sequenced three times, due to assembling complexity. First, the genome was decoded with the SOLiD 5500 platform with mate-paired libraries, generating a total of 5,133,057,360 bp, (coverage of 2,053 times). The reads were subjected to a Phred 20 quality filter using Quality Assessment software (8) and assembled with the CLC Genomics Workbench, generating a total of 1,641 overlapping sequences. These sequences were removed with the Simplifier (9), ordered and oriented based on the reference *L. lactis* KF147 genome sequence (a plant-

associated strain, accession number CP001834). Then manual curation was performed using Artemis (10), and SSPACE (11) and Gapfiller (12) were used to generate the scaffold and resolve gaps, respectively. At the end of curation and sequence assembly, a total of 409 scaffolds (2,874,854 bp) were obtained.

L. lactis NCDO 2118 was then decoded with the Ion PGM platform with fragment libraries generating a total of 187,303,001 bp (coverage of ~71 times). Genome assembly was performed using Mira 3.9 (13), and the assembled genome sequence was reference aligned with CONTIGuator (14). The redundant overlapping sequences were removed with “in-house scripts,” closing the remnant gaps. Annotation and frameshifts curation were then performed using Artemis and CLC, reducing the initial 1821 pseudogenes to 480.

Finally, the DNA was sequenced using the Ion Torrent PGM with fragment libraries, yielding a total of ~1,249,154,478 bp (coverage of 474 times). Assembly was performed with Mira 4.0.1 and Newbler 2.9 (15). We used CONTIGuator and FGAP 1.7 (16) to perform the alignment and gap closure steps, respectively. We followed the same previously explained steps for annotation and frameshift curation, reducing the pseudogenes to 52.

The complete genome of *L. lactis* NCDO 2118 consists of a single circular chromosome of 2,554,693 bp, containing 2,386 coding sequences (CDS), which had 52 pseudogenes, 66 tRNA genes, and 6 rRNA operons, with a G+C content of 34.9%. There is one plasmid, pNCDO2118 (37,571 bp), with 48 CDS, from which 4 are pseudogenes with a G+C content of 32.33%.

Nucleotide sequence accession numbers. The *Lactococcus lactis* NCDO 2118 chromosome and the plasmid were deposited at DDBJ/EMBL/GenBank under the accession numbers [CP009054](#) and [CP009055](#), respectively.

ACKNOWLEDGMENTS

This work was supported by Rede Paraense de Genômica e Proteômica, Ministério da Pesca e Aquicultura (MPA), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG). We also acknowledge the support from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

REFERENCES

- Carr FJ, Chill D, Maida N. 2002. The lactic acid bacteria: a literature survey. *Crit. Rev. Microbiol.* 28:281–370. <http://dx.doi.org/10.1080/1040-840291046759>.
- van de Guchte M, Ehrlich SD, Maguin E. 2001. Production of growth-inhibiting factors by *Lactobacillus delbrueckii*. *J. Appl. Microbiol.* 91: 147–153. <http://dx.doi.org/10.1046/j.1365-2672.2001.01369.x>.
- Zareian M, Ebrahimpour A, Bakar FA, Mohamed AK, Forghani B, Ab-Kadir MS, Saari N. 2012. A glutamic acid-producing lactic acid bacteria isolated from Malaysian fermented foods. *Int. J. Mol. Sci.* 13: 5482–5497. <http://dx.doi.org/10.3390/ijms13055482>.
- Inoue K, Shirai T, Ochiai H, Kasao M, Hayakawa K, Kimura M, Sansawa H. 2003. Blood-pressure-lowering effect of a novel fermented milk containing gamma-aminobutyric acid (GABA) in mild hypertensives. *Eur. J. Clin. Nutr.* 57:490–495. <http://dx.doi.org/10.1038/sj.ejcn.1601555>.
- Jin Z, Mendu SK, Birnir B. 2013. GABA is an effective immunomodulatory molecule. *Amino Acids* 45:87–94. <http://dx.doi.org/10.1007/s00726-011-1193-7>.
- Mazzoli R, Pessione E, Dufour M, Laroute V, Giuffrida MG, Giunta C, Coccagn-Bousquet M, Loubière P. 2010. Glutamate-induced metabolic changes in *Lactococcus lactis* NCDO 2118 during GABA production: combined transcriptomic and proteomic analysis. *Amino Acids.* 39:727–737. <http://dx.doi.org/10.1007/s00726-010-0507-5>.
- Siezen RJ, Starrenburg MJ, Boekhorst J, Renckens B, Molenaar D, van Hylckama Vlieg JE. 2008. Genome-scale genotype-phenotype matching of two *Lactococcus lactis* isolates from plants identifies mechanisms of adaptation to the plant niche. *Appl. Environ. Microbiol.* 74:424–436. <http://dx.doi.org/10.1128/AEM.01850-07>.
- Ramos RT, Carneiro AR, Baumbach J, Azevedo V, Schneider MP, Silva A. 2011. Analysis of quality raw data of second generation sequencers with Quality Assessment Software. *BMC Res. Notes* 4:130. <http://dx.doi.org/10.1186/1756-0500-4-130>.
- Ramos RT, Carneiro AR, Azevedo V, Schneider MP, Barh D, Silva A. 2012. Simplifier: a web tool to eliminate redundant NGS contigs. *Bioinformatics* 8:996–999. <http://dx.doi.org/10.6026/97320630008996>.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945. <http://dx.doi.org/10.1093/bioinformatics/16.10.944>.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding preassembled contigs using SSPACE. *Bioinformatics* 27: 578–579. <http://dx.doi.org/10.1093/bioinformatics/btq683>.
- Nadalin F, Vezzi F, Policriti A. 2012. GapFiller: a *de novo* assembly approach to fill the gap within paired reads. *BMC Bioinformatics* 13:S8. <http://dx.doi.org/10.1186/1471-2105-13-S14-S8>.
- Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *Comput. Sci. Biol.: Proc. German Conference on Bioinformatics GCB'99 GCB*. Hannover, Germany.
- Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol. Med.* 6:11. <http://dx.doi.org/10.1186/1751-0473-6-11>.
- Ohnishi N, Maruyama F, Ogawa H, Kachi H, Yamada S, Fujikura D, Nakagawa I, Hang'ombe MB, Thomas Y, Mweene AS, Higashi H. 2014. Genome sequence of a *Bacillus anthracis* outbreak strain from Zambia, 2011. *Genome Announc.* 2(2):e00116-14. <http://dx.doi.org/10.1128/genomeA.00116-14>.
- Piro VC, Faoro H, Weiss VA, Steffens MB, Pedrosa FO, Souza EM, Raittz RT. 2014. FGAP: an automated gap closing tool. *BMC Res. Notes* 7:371. <http://dx.doi.org/10.1186/1756-0500-7-371>.