

Late-Stage Molecular Editing Enabled by Ketone Chain-Walking Isomerization

Yannick Brägger, Ori Green, Benjamin N. Bhawal,* and Bill Morandi*

Cite This: *J. Am. Chem. Soc.* 2023, 145, 19496–19502

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Herein, a method for the isomerization of ketones in a manner akin to the chain-walking reaction of alkenes is described. Widely available and inexpensive pyrrolidine and elemental sulfur are deployed as catalysts to achieve this reversible transformation. Key to the utility of this approach was the elucidation of a stereochemical model to determine the thermodynamically favored product of the reaction and the kinetic selectivity observed. With the distinct selectivity profile of our ketone chain-walking process, the isomerization of various steroids was demonstrated to rapidly access novel steroids with “unnatural” oxidation patterns.

The relative arrangement of functional groups within a molecule imparts its physical and biological properties.¹ Thus, controlling the position and orientation of functional groups is a fundamental goal in organic synthesis. Typically, this is realized through the deployment of transformations that selectively introduce functional groups with the desired arrangement.² Alternatively, isomerization reactions offer the opportunity to “correct” either the position or the orientation of functional groups within a molecule. This approach is particularly attractive to facilitate the editing of complex molecules that feature a high density of functionality (Scheme 1A).^{3,4} Aside from the atom economy of this approach,⁵ it is also a step-economical strategy,⁶ enabling access to complex chemical entities through a single synthetic procedure rather than a lengthy *de novo* synthesis. Applying this to inexpensive and readily available biomass feedstocks, such as sugars and steroids, also represents a sustainable approach to access analogues of these privileged scaffolds. In 2020, the Wendlandt group provided a compelling demonstration of the utility of this approach (Scheme 1B).⁷ Using a photocatalytic reaction manifold, they accessed rare monosaccharides of biological importance from more readily available monosaccharides through selective epimerization.

While previous examples clearly show the feasibility and synthetic utility of late-stage epimerization reactions, amending the location of functional groups is arguably even more challenging, as it requires reversible cleavage and transposition of strong bonds. Alkene chain-walking is a rare example and proceeds via a series of β -hydride eliminations and hydride insertions.⁸ This process has greatly impacted organic synthesis by enabling the synthesis of unusual building blocks and unlocking cascade reactions. Achieving a directly analogous process with ketones, another ubiquitous and versatile functional group, would be highly desirable. However, transition-metal-mediated elementary steps to manipulate the C=O bond of ketones for a chain-walking process are lacking.⁹ To circumvent this challenge, the Dong group designed a Catellani-type process to realize a carbonyl 1,2-

transposition (Scheme 1C).¹⁰ This elegant process is kinetically controlled but is irreversible and limited to 1,2-transpositions. Thus, a reversible carbonyl chain-walking process akin to alkene chain-walking would open new synthetic opportunities.¹¹


Given our group's interest in reversible catalytic reactions^{12,13} and molecular editing processes,¹⁴ we sought to develop such a process. Cognizant of the challenges of achieving this through transition-metal catalysis, we looked for a mechanistically distinct approach to perform the desired transformation. We took inspiration from the Willgerdt–Kindler reaction (Scheme 1D),¹⁵ wherein aliphatic ketones are transformed to either an amide or a thioamide, typically by refluxing the ketone with elemental sulfur¹⁶ in a solution of the amine, with the carbonyl group migrating down the aliphatic chain to the terminal position. Notably, the carbonyl group is able to “walk” along a variety of chain lengths, suggesting that a chain-walking process is operative. We hypothesized that, for cyclic ketones, wherein the carbonyl group is unable to migrate to a terminal position and undergo subsequent oxidation, a carbonyl chain-walking process could be realized using substoichiometric amounts of an amine and elemental sulfur. Previously, the isomerization of cyclic ketones has been demonstrated using stoichiometric quantities of both amine and sulfur.¹⁷ Herein, we report a simple procedure for the isomerization of cyclic ketones and demonstrate its application in the late-stage editing of complex steroids to rapidly access unnatural isomers of this biologically prevalent class of natural products (Scheme 1E).

Received: May 31, 2023

Published: August 28, 2023




Scheme 1. Context of This Work



At the outset of this work, we were conscious that, as the reaction would be reversible, a suitable driving force to favor product formation would be necessary. Thus, we targeted synthetically relevant ring systems in which certain isomers would be favored thermodynamically. We were intrigued by geminally disubstituted cyclohexanones, as one of the substituents must occupy an axial position. Consequently, 3,3-geminally disubstituted cyclohexanones are more thermodynamically stable than their 2,2- and 4,4-disubstituted counterparts due to a reduction in the number of 1,3-diaxial interactions (Scheme 2A). Furthermore, this structural pattern is frequently embedded in naturally occurring steroids, offering the prospect to apply our method in the late-stage isomerization of bioactive compounds. To evaluate this hypothesis, DFT calculations were performed to determine the relative ground-state energies of 4,4-, 3,3-, and 2,2-dimethylcyclohex-

Scheme 2. Model for Predicting the Thermodynamic Selectivity of the Reversible Isomerization of Geminally Substituted Dimethylcyclohexanones and Theoretical and Experimental Verification



^aGround-state energy calculations were performed at the B3LYP/def2-QZVPP level of theory. ^bAll yields were determined by GC-FID with an internal standard.


anone (Scheme 2B). This showed that, as expected, the 3,3-dimethylcyclohexanone isomer should be favored.

Subsequently, we assessed reaction conditions to facilitate the isomerization of dimethylcyclohexanones. Critically, the isomerization can be performed with substoichiometric amounts of pyrrolidine and sulfur, in contrast to the Willgerdt–Kindler reaction and previous work.¹⁷ We also observed pyrrolidine to be markedly superior in relation to other amines (see the Supporting Information for further information). Under the optimized conditions, reactions with 4,4- and 3,3-dimethylcyclohexanone (1a and 1b, respectively) yielded a nearly identical mixture of isomers, indicative that an equilibrium is achieved, with 1b being the major product, thereby verifying its greater thermodynamic stability (Scheme 2C). Notably, formation of 2,2-dimethylcyclohexanone 1c was not observed, and no isomerization of 2,2-dimethylcyclohex-

anone occurred when it was submitted to the optimized reaction conditions (Scheme 2D). Together, these results imply that the ketone chain-walking process is reversible, thermodynamically selective for the 3,3-disubstituted isomer, and not applicable to the formation or reaction of sterically encumbered ketones such as 2,2-dimethylcyclohexanone.

The selectivity of the reaction was further probed through the isomerization of a range of mono-substituted cyclohexanones (Scheme 3). With 2-substituted cyclohexanones

Scheme 3. Isomerization of Mono-substituted Cyclohexanones^a




^aIsomer distributions were determined by GC-FID with an internal standard.

3c and 4c, limited reactivity was observed, and with 2-*tert*-butylcyclohexanone 2c, no isomerization was observed at all. Conversely, isomerization of 4-substituted cyclohexanones 2a, 3a, and 4a or 3-substituted cyclohexanones 2b, 3b, and 4b gave limited or no formation of the 2-substituted isomers. With 4-*tert*-butylcyclohexanone 2a and 3-*tert*-butylcyclohexanone 2b, no formation of 2-*tert*-butylcyclohexanone 2c was observed, and the reaction exhibited selectivity for the 3-substituted isomer. The origin of this selectivity remains unclear. It should also be noted that the isomerization of 4-methylcyclohexanone 3a yielded a mixture of all three isomers, showcasing the potential for longer range walking of the carbonyl group.

Collectively, these results provide further insights into the kinetic selectivity of our isomerization process (Scheme 4). Ketones bearing bulky α -substituents (e.g., *tert*-butyl) or α,α -disubstituted ketones neither form nor react under these conditions. However, with less bulky substituents (e.g., methyl), the α -substituted isomer will participate in the reaction. However, this reversible process is considerably


Scheme 4. Kinetic Selectivity of Ketone Isomerization




depressed in comparison to the interconversion of ketone isomers bearing no substituents at the α -position. This selectivity profile is reminiscent of that observed in enamine catalysis, which is typically ineffective with bulkier cycloketones.¹⁸

With a greater understanding of the isomerization process, we sought to apply this methodology to more complex targets, as we anticipated our mild reaction conditions to be compatible with a range of substrates. Various derivatives of the Wieland–Miescher ketone, which has been extensively used as a key intermediate in natural product synthesis,^{19,20} were investigated (Scheme 5A). Isomerization with keto-alcohol 5a proceeded smoothly, affording the 2-oxo-isomer 5b as the major product, with no other isomers detected. Notably, reaction with diketone 6a led to the formation of only one new isomer, 6b. This result underlines the critical role of the selectivity of this transformation; otherwise, a mixture of 16 different regioisomers could be formed. Due to the selectivity profile, the desired product could be obtained in 44% yield. Similarly, reaction with the cis-fused diketone 7a also afforded a single isomeric product, 7b, albeit with reduced selectivity. Further derivatives were subjected to the reaction conditions to evaluate the functional group tolerance of the procedure, and substrates incorporating esters (8a), silyl ethers (9a), alkenes (10a), carbamates (11a), and imides (12a) were all tolerated. While the separation of isomers can potentially present a challenge, we found this could be readily accomplished using preparative HPLC. Combined yields of the reactant and product isomers in these reactions were typically around 80%. However, attempts to identify any byproducts formed were thwarted by the complex mixtures obtained. Derivatives of the Hajos–Parrish ketones, 13a and 14a, were also subjected to the reaction conditions (Scheme 5B). With the 6,5-fused ring system, two new isomeric products were formed, with the 4-oxo-isomers 13c and 14c being formed as minor products, in addition to the 2-oxo-isomers 13b and 14b. Previous syntheses of these ketone isomers have relied on multistep *de novo* synthesis.²¹ For instance, the groups of Wijnberg and de Groot undertook a 7-step synthesis to realize formal isomerization of the Wieland–Miescher ketone derivative 5a.²² This synthetic sequence uses several stoichiometric reagents and utilizes protecting group logic to circumvent selectivity issues. With our method, the same isomerization takes just one single step and avoids the use of protecting groups and stoichiometric reagents, thus constituting a powerful atom- and step-economic upgrade.

We next performed the isomerization of naturally occurring steroids (Scheme 6A).²³ As expected, 3-oxo-steroids smoothly underwent isomerization to afford 2-keto-steroids as the major

Scheme 5. Isomerization of Bicyclic Ketones by Carbonyl Chain-Walking^{a,b,c,d}

^aReactions were performed on 1.0 mmol scale. ^bUnless otherwise stated, products were initially isolated as a mixture of isomers by flash column chromatography. Combined yields of these isomeric mixtures are reported. ^cIsomer ratios were determined by ¹H NMR spectroscopy. ^dUnless otherwise stated, isolated yields following purification by preparative HPLC are reported. ^eIsomer **7b** was directly isolated by flash column chromatography. ^fOnly a small sample was purified by HPLC.

Scheme 6. Isomerization of 3-Oxo-Steroids by Carbonyl Chain-Walking^{a,b,c}


^aReactions were performed on 1.0 mmol scale. ^bIsolated yields following purification by preparative HPLC are reported. ^cIsomer ratios were determined by quantitative ¹³C NMR spectroscopy.

product. Due to the kinetic selectivity of the reaction, neither the 1- nor 4-keto-steroid isomers were detected in any of the crude reaction mixtures. Thus, the isomerization of androstanolone **15a** and mestanolone **16a** afforded exclusively their respective 2-oxo-isomers, **15b** and **16b**. In the isomerization of the diketosteroids, androstanedione **17a** and allopregnenedione **18a**, only isomerization of the A-ring ketone was observed, with the other ketone remaining untouched. This selectivity is particularly imperative in the case of **18a**, because if the other ketone were able to undergo reaction, this would likely lead to the undesired occurrence of a Willgerdt–Kindler rearrangement, forming an amide species at the terminal position and ultimately inhibiting the desired isomerization process. Furthermore, the synthesis of **18b** has previously been reported in an 8-step sequence starting from pregnenolone (Scheme 6B).^{24,25} The isomerization is achieved through several synthetic steps, and further steps are

necessitated by the protecting group strategy and redox manipulations employed to navigate the challenge of achieving selective isomerization. Thus, this truly highlights the synthetic virtues of our new ketone isomerization, especially given its distinct selectivity profile. It should also be noted that the isomerization of diketones **17a** and **18a** would be challenging with the protocol from Dong's group, as this would require selective triflate formation.¹⁰

Having established a protocol to achieve the chain-walking isomerization of ketones, preliminary mechanistic experiments were performed. Based on studies on the Willgerodt–Kindler reaction^{17b,26} and related transformations,^{16,27} we surmised that the reaction likely proceeds through the formation of an enamine intermediate. To test this hypothesis, 4,4-dimethylcyclohexanone **1a** was submitted to the standard reaction conditions but using D₂O as additive and CD₃OD as solvent (Scheme 7). This led to deuterium incorporation detected at

Scheme 7. Deuterium Labeling



^aPositions of deuterium incorporation were determined by ¹H and ²H NMR spectroscopy.

all methylene carbons in both **d-1a** and **d-1b**. Since this includes both of the methylene carbons adjacent to the quaternary carbon, this further affirms the reversibility of the system. Subsequently, this reaction was repeated but without the addition of S₈. No isomerization was observed, and deuterium incorporation was only detected alpha to the ketone, but not at the beta methylene carbons. This result reflects the fact that S₈ is necessary to isomerize the putative enamine intermediate.

In conclusion, we report a novel and simple process for the isomerization of cyclic ketones. Based on well-established concepts in the stereochemistry of cyclohexanes, a model to control this reversible process was devised. Moreover, we found that the process was not amenable to the formation of sterically hindered ketones. Thus, the selective isomerization of 3-oxo-steroids to their 2-oxo-steroid analogues was executed. This process is a rare example of a polar group chain-walking reaction that parallels the venerable alkene chain-walking reaction, which has been a key platform in homogeneous catalysis. We thus expect that our new carbonyl chain-walking will open a wealth of new opportunities for organic synthesis. In a wider context, this work provides compelling motivation for the discovery of new isomerization processes and underlines their value for the late-stage editing of natural products and other complex molecular architectures.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.3c05680>.

Experimental procedures and characterization data for all compounds (PDF)

Accession Codes

CCDC 2266361 and 2285025–2285027 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

Benjamin N. Bhawal – ETH Zürich, 8093 Zürich, Switzerland; School of Chemistry, University of Edinburgh, Edinburgh EH9 3FJ, U.K.; orcid.org/0000-0002-8552-1547; Email: ben.bhawal@ed.ac.uk

Bill Morandi – ETH Zürich, 8093 Zürich, Switzerland; orcid.org/0000-0003-3968-1424; Email: morandib@ethz.ch

Authors

Yannick Brägger – ETH Zürich, 8093 Zürich, Switzerland
Ori Green – ETH Zürich, 8093 Zürich, Switzerland

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.3c05680>

Funding

B.N.B. gratefully acknowledges the University of Edinburgh for funding. B.M. acknowledges ETH Zurich, the European Research Council under the European Union's Horizon 2020 research and innovation program (Shuttle Cat, Project ID: 757608), and the Swiss National Science Foundation (SNSF 213812) for funding. O.G. thanks the International Human Frontier Science Program Organization (LT000861/2020-L).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the NMR service, X-ray service (SMoCC), and the Molecular and Biomolecular Analysis Service (MoBiAS) of ETH Zürich for technical assistance. We thank the Morandi group for helpful discussions and critical proofreading of the manuscript.

■ REFERENCES

- (1) Szpilman, A. M.; Carreira, E. M. Probing the Biology of Natural Products: Molecular Editing by Diverted Total Synthesis. *Angew. Chem., Int. Ed.* **2010**, *49*, 9592–9628.
- (2) (a) Mahatthanachai, J.; Dumas, A. M.; Bode, J. W. Catalytic Selective Synthesis. *Angew. Chem., Int. Ed.* **2012**, *51*, 10954–10990. (b) Hong, B.; Luo, T.; Lei, X. Late-Stage Diversification of Natural Products. *ACS Cent. Sci.* **2020**, *6*, 622–635. (c) Guillemard, L.; Kaplaneris, N.; Ackermann, L.; Johansson, M. J. Late-Stage C–H Functionalization Offers New Opportunities in Drug Discovery. *Nat. Rev. Chem.* **2021**, *5*, 522–545. (d) Zhang, L.; Ritter, T. A Perspective on Late-Stage Aromatic C–H Bond Functionalization. *J. Am. Chem. Soc.* **2022**, *144*, 2399–2414.
- (3) For selected recent examples see: (a) Metternich, J. B.; Gilmour, R. A Bio-Inspired, Catalytic E→Z Isomerization of Activated Olefins.

- J. Am. Chem. Soc.* **2015**, *137*, 11254–11257. (b) Hill, C. K.; Hartwig, J. F. Site-selective oxidation, amination and epimerization reactions of complex polyols enabled by transfer hydrogenation. *Nat. Chem.* **2017**, *9*, 1213–1221. (c) Hölzl-Hobmeier, A.; Bauer, A.; Silva, A. V.; Huber, S. M.; Bannwarth, C.; Bach, T. Catalytic Deracemization of Chiral Allenes by Sensitized Excitation with Visible Light. *Nature* **2018**, *564*, 240–243. (d) Oswood, C. J.; MacMillan, D. W. Selective Isomerization via Transient Thermodynamic Control: Dynamic Epimerization of trans to cis Diols. *J. Am. Chem. Soc.* **2022**, *144*, 93–98. (e) Zhao, K.; Knowles, R. R. Contra-Thermodynamic Positional Isomerization of Olefins. *J. Am. Chem. Soc.* **2022**, *144*, 137–144. (f) Occhialini, G.; Palani, V.; Wendlandt, A. E. Catalytic, Contra-Thermodynamic Positional Alkene Isomerization. *J. Am. Chem. Soc.* **2022**, *144*, 145–152. (g) Zhang, Y.-A.; Palani, V.; Seim, A. E.; Wang, Y.; Wang, K. J.; Wendlandt, A. E. Stereochemical Editing Logic Powered by the Epimerization of Unactivated Tertiary Stereocenters. *Science* **2022**, *378*, 383–390. (h) Carder, H. M.; Wang, Y.; Wendlandt, A. E. Selective Axial-to-Equatorial Epimerization of Carbohydrates. *J. Am. Chem. Soc.* **2022**, *144*, 11870–11877.
- (4) For selected reviews see: (a) Dugave, C.; Demange, L. Cis–Trans Isomerization of Organic Molecules and Biomolecules: Implications and Applications. *Chem. Rev.* **2003**, *103*, 2475–2532. (b) Molloy, J. J.; Morack, T.; Gilmour, R. Positional and Geometrical Isomerisation of Alkenes: The Pinnacle of Atom Economy. *Angew. Chem., Int. Ed.* **2019**, *58*, 13654–13664. (c) Nevesely, T.; Wienhold, M.; Molloy, J. J.; Gilmour, R. Advances in the E → Z Isomerization of Alkenes Using Small Molecule Photocatalysts. *Chem. Rev.* **2022**, *122*, 2650–2694. (d) Suh, C. E.; Carder, H. M.; Wendlandt, A. E. Selective Transformations of Carbohydrates Inspired by Radical-Based Enzymatic Mechanisms. *ACS Chem. Biol.* **2021**, *16*, 1814–1828.
- (5) (a) Trost, B. The Atom Economy—A Search for Synthetic Efficiency. *Science* **1991**, *254*, 1471–1477. (b) Trost, B. M. Atom Economy—A Challenge for Organic Synthesis: Homogeneous Catalysis Leads the Way. *Angew. Chem., Int. Ed.* **1995**, *34*, 259–281.
- (6) Newhouse, T.; Baran, P. S.; Hoffmann, R. W. The Economies of Synthesis. *Chem. Soc. Rev.* **2009**, *38*, 3010–3021.
- (7) Wang, Y.; Carder, H. M.; Wendlandt, A. E. Synthesis of rare sugar isomers through site-selective epimerization. *Nature* **2020**, *578*, 403–408.
- (8) (a) Vasseur, A.; Bruffaerts, J.; Marek, I. Remote Functionalization through Alkene Isomerization. *Nat. Chem.* **2016**, *8*, 209–219. (b) Sommer, H.; Juliá-Hernández, F.; Martin, R.; Marek, I. Walking Metals for Remote Functionalization. *ACS Cent. Sci.* **2018**, *4*, 153–165. (c) Larionov, E.; Li, H.; Mazet, C. Well-Defined Transition Metal Hydrides in Catalytic Isomerizations. *Chem. Commun.* **2014**, *50*, 9816–9826.
- (9) (a) Fu, G. C.; Grubbs, R. H. Synthesis of Cycloalkenes via Alkylidene-Mediated Olefin Metathesis and Carbonyl Olefination. *J. Am. Chem. Soc.* **1993**, *115*, 3800–3801. (b) Park, Y. J.; Park, J.-W.; Jun, C.-H. Metal–Organic Cooperative Catalysis in C–H and C–C Bond Activation and Its Concurrent Recovery. *Acc. Chem. Res.* **2008**, *41*, 222–234.
- (10) Wu, Z.; Xu, X.; Wang, J.; Dong, G. Carbonyl 1,2-Transposition through Triflate-Mediated α -Amination. *Science* **2021**, *374*, 734–740.
- (11) During the preparation of this manuscript, an elegant strategy for the isomerization of amides to ketones via an Ir-catalyzed 1,2-carbonyl transposition was published as a preprint: Shennan, B. D. A.; Sánchez Alonso, S.; Dixon, D. J. 1,2-Redox Transpositions of Tertiary Amides. *ChemRxiv Preprint*, DOI: 10.26434/chemrxiv-2023-rchb2.
- (12) For reviews see: (a) Bhawal, B. N.; Morandi, B. Catalytic Isofunctional Reactions—Expanding the Repertoire of Shuttle and Metathesis Reactions. *Angew. Chem., Int. Ed.* **2019**, *58*, 10074–10103. (b) Bhawal, B. N.; Morandi, B. Isodesmic Reactions in Catalysis – Only the Beginning? *Isr. J. Chem.* **2018**, *58*, 94–103. (c) Bhawal, B. N.; Morandi, B. Catalytic Transfer Functionalization through Shuttle Catalysis. *ACS Catal.* **2016**, *6*, 7528–7535.
- (13) For selected examples see: (a) Fang, X.; Yu, P.; Morandi, B. Catalytic Reversible Alkene–Nitrile Inter-conversion through Controllable Transfer Hydrocyanation. *Science* **2016**, *351*, 832–836. (b) Fang, X.; Cacherat, B.; Morandi, B. CO- and HCl-Free Synthesis of Acid Chlorides from Unsaturated Hydrocarbons via Shuttle Catalysis. *Nat. Chem.* **2017**, *9*, 1105–1109. (c) Lian, Z.; Bhawal, B. N.; Yu, P.; Morandi, B. Palladium-Catalyzed Carbon–Sulfur or Carbon–Phosphorus Bond Metathesis by Reversible Arylation. *Science* **2017**, *356*, 1059–1063. (d) Lee, Y. H.; Morandi, B. Metathesis-Active Ligands Enable a Catalytic Functional Group Metathesis between Aryl Chlorides and Aryl Iodides. *Nat. Chem.* **2018**, *10*, 1016–1022. (e) Delcaillau, T.; Bismuto, A.; Lian, Z.; Morandi, B. Nickel-Catalyzed Inter- and Intramolecular Aryl Thioether Metathesis by Reversible Arylation. *Angew. Chem., Int. Ed.* **2020**, *59*, 2110–2114. (f) Bhawal, B. N.; Reisenbauer, J. C.; Ehinger, C.; Morandi, B. Overcoming Selectivity Issues in Reversible Catalysis: A Transfer Hydrocyanation Exhibiting High Kinetic Control. *J. Am. Chem. Soc.* **2020**, *142*, 10914–10920. (g) Dong, X.; Roeckl, J. L.; Waldvogel, S. R.; Morandi, B. Merging Shuttle Reactions and Pzid Electrocatalysis for Reversible Vicinal Dihalogenations. *Science* **2021**, *371*, 507–514. (h) Delcaillau, T.; Boehm, P.; Morandi, B. Nickel-Catalyzed Reversible Functional Group Metathesis between Aryl Nitriles and Aryl Thioethers. *J. Am. Chem. Soc.* **2021**, *143*, 3723–3728. (i) Lutz, M. D. R.; Gasser, V. C. M.; Morandi, B. Shuttle Arylation by Rh(I) Catalyzed Reversible Carbon–Carbon Bond Activation of Unstrained Alcohols. *Chem* **2021**, *7*, 1108–1119. (j) Reisenbauer, J. C.; Bhawal, B. N.; Jelmini, N.; Morandi, B. Development of an Operationally Simple, Scalable, and HCN-Free Transfer Hydrocyanation Protocol Using an Air-Stable Nickel Precatalyst. *Org. Process Res. Dev.* **2022**, *26*, 1165–1173. (k) Boehm, P.; Müller, P.; Finkelstein, P.; Rivero-Crespo, M. A.; Ebert, M.-O.; Trapp, N.; Morandi, B. Mechanistic Investigation of the Nickel-Catalyzed Metathesis between Aryl Thioethers and Aryl Nitriles. *J. Am. Chem. Soc.* **2022**, *144*, 13096–13108.
- (14) (a) Reisenbauer, J. C.; Green, O.; Franchino, A.; Finkelstein, P.; Morandi, B. Late-stage diversification of indole skeletons through nitrogen atom insertion. *Science* **2022**, *377*, 1104–1109. (b) Finkelstein, P.; Reisenbauer, J. C.; Botlik, B. B.; Green, O.; Florin, A.; Morandi, B. Nitrogen Atom Insertion into Indenes to Access Isoquinolines. *Chem. Sci.* **2023**, *14*, 2954–2959.
- (15) (a) Willgerodt, C. Ueber Die Einwirkung von Gelbem Schwefelammonium Auf Ketone Und Chinone. *Ber. Dtsch. Chem. Ges.* **1887**, *20*, 2467–2470. (b) Kindler, K. Studien über den Mechanismus chemischer Reaktionen. Erste Abhandlung. Reduktion von Amidinen und Oxydation von Aminen. *Liebigs Ann. Chem.* **1923**, *431*, 187–230. (c) Priebbenow, D. L.; Bolm, C. Recent Advances in the Willgerodt–Kindler Reaction. *Chem. Soc. Rev.* **2013**, *42*, 7870–7880.
- (16) Nguyen, T. B. Recent Advances in Organic Reactions Involving Elemental Sulfur. *Adv. Synth. Catal.* **2017**, *359*, 1066–1130.
- (17) (a) Sejbal, J.; Klinot, J.; Protiva, J.; Vystrčil, A. Reactions of Triterpenoid Ketones with Sulfur and Morpholine under the Conditions of Willgerodt–Kindler Reaction. *Collect. Czech. Chem. Commun.* **1986**, *51*, 118–127. (b) Carmack, M.; Behforouz, M.; Berchtold, G. A.; Berkowitz, S. M.; Wiesler, D.; Barone, R. The Willgerodt–Kindler Reactions. 6. Isomerization of the Carbonyl Group in Alkanones and Cycloalkanones. *J. Heterocycl. Chem.* **1989**, *26*, 1305–1318.
- (18) (a) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric Enamine Catalysis. *Chem. Rev.* **2007**, *107*, 5471–5569. (b) Pihko, P. M.; Majander, I.; Erkkilä, A. Enamine Catalysis. *Top. Curr. Chem.* **2010**, *291*, 145–200. (c) Sulzer-Mossé, S.; Alexakis, A. Chiral Amines as Organocatalysts for Asymmetric Conjugate Addition to Nitroolefins and Vinyl Sulfones via Enamine Activation. *Chem. Commun.* **2007**, 3123–3135.
- (19) Bradshaw, B.; Bonjoch, J. The Wieland–Miescher Ketone: A Journey from Organocatalysis to Natural Product Synthesis. *Synlett* **2012**, *23*, 337–356.
- (20) For selected examples in total synthesis, see: (a) Danishefsky, S. J.; Masters, J. J.; Young, W. B.; Link, J. T.; Snyder, L. B.; Magee, T. V.; Jung, D. K.; Isaacs, R. C. A.; Bornmann, W. G.; Alaimo, C. A.; Coburn, C. A.; Di Grandi, M. J. Total Synthesis of Baccatin III and Taxol. *J. Am. Chem. Soc.* **1996**, *118*, 2843–2859. (b) Stork, G.; West,

F.; Lee, H. Y.; Isaacs, R. C. A.; Manabe, S. The Total Synthesis of a Natural Cardenolide: (+)-Digitoxigenin. *J. Am. Chem. Soc.* **1996**, *118*, 10660–10661.

(21) Qian, M.; Engler-Chiurazzi, E. B.; Lewis, S. E.; Rath, N. P.; Simpkins, J. W.; Covey, D. F. Structure–activity studies of non-steroid analogues structurally-related to neuroprotective estrogens. *Org. Biomol. Chem.* **2016**, *14*, 9790–9805.

(22) Orru, R. V. A.; Wijnberg, J. B. P. A.; Bouwman, C. T.; De Groot, A. Rearrangement vs Homofragmentation: Chemical Consequences of Different σ -Relays on the Heterolysis of Sulfonate Esters Induced by Through-Bond Interactions. *J. Org. Chem.* **1994**, *59*, 374–382.

(23) Varela, C.; Tavares da Silva, E. J.; Amaral, C.; Correia da Silva, G.; Baptista, T.; Alcaro, S.; Costa, G.; Carvalho, R. A.; Teixeira, N. A. A.; Roleira, F. M. F. New Structure–Activity Relationships of A- and D-Ring Modified Steroidal Aromatase Inhibitors: Design, Synthesis, and Biochemical Evaluation. *J. Med. Chem.* **2012**, *55*, 3992–4002.

(24) Clegg, A. S.; Denny, W. A.; Jones, E. R. H.; Kumar, V.; Meakins, G. D.; Thomas, V. E. M. Studies in the Steroid Group. Part LXXXII. The Preparation of Nine Mono- and Eight Di-Oxoandrostanes, 5α -Estran-17-One, and 5α -Pregnane-2,20-Dione. *J. Chem. Soc., Perkin Trans. 1* **1972**, 492–498.

(25) Wong, F. F.; Chen, C.-Y.; Chen, T.-H.; Huang, J.-J.; Fang, H.-P.; Yeh, M.-Y. Synthesis of 3α -Hydroxy-21-(1'-Imidazolyl)- 3β -Methoxyl-Methyl- 5α -Pregnan-20-One via Lithium Imidazole with 17α -Acetylbromopregnanone. *Steroids* **2006**, *71*, 77–82.

(26) (a) Wang, Z. *Comprehensive Organic Name Reactions and Reagents*; John Wiley & Sons, Inc., 2010; pp 3018–3025. DOI: 10.1002/9780470638859.conrr672. (b) Bible, R. H., Jr. *J. Am. Chem. Soc.* **1957**, *79*, 3924–3925. (c) Carmack, M.; Behforouz, M.; Bechtold, G. A.; Berkowitz, S. M.; Wiesler, D.; Barone, R. The Willgerodt Kindler Reactions. 7. The Mechanisms. *J. Heterocycl. Chem.* **1989**, *26*, 1319–1323. (d) Dauben, W. G.; Rogan, J. B. The Willgerodt Reaction with Acylmesitylenes. The Mechanism of the Reaction. *J. Am. Chem. Soc.* **1956**, *78*, 4135–4139.

(27) (a) Mayer, R.; Gewald, K. The Action of Carbon Disulfide and Sulfur on Enamines, Ketimines, and CH Acids. *Angew. Chem., Int. Ed.* **1967**, *6*, 294–306. (b) Solovyev, A. Y.; Androsov, D. A.; Neckers, D. C. One-Pot Synthesis of Substituted 2-Aminobenzo[b]Thiophenes. *J. Org. Chem.* **2007**, *72*, 3122–3124. (c) Androsov, D. A.; Solovyev, A. Y.; Petrov, M. L.; Butcher, R. J.; Jasinski, J. P. A Convenient Approach towards 2- and 3-Aminobenzo[b]Thiophenes. *Tetrahedron* **2010**, *66*, 2474–2485.